The Ankyrin Repeat-containing Protein Akrlp Is Required for the Endocytosis of Yeast Pheromone Receptors

Scott A. Givan and George F. Sprague, Jr.*

Institute of Molecular Biology and Department of Biology, University of Oregon, Eugene, Oregon 97403

Submitted April 22, 1997; Accepted April 25, 1997
Monitoring Editor: David Botstein

The *Saccharomyces cerevisiae* α -factor receptor (Ste3p) requires its C-terminal cytoplasmic tail for endocytosis. Wild-type receptor is delivered to the cell surface via the secretory pathway but remains there only briefly before being internalized and delivered to the vacuole for degradation. Receptors lacking all or part of the cytoplasmic tail are not subject to this constitutive endocytosis. We used the cytoplasmic tail of Ste3p as bait in the two-hybrid system in an effort to identify other proteins involved in endocytosis. One protein identified was Akrlp, an ankyrin repeat-containing protein. We applied three criteria to demonstrate that Akrlp is involved in the constitutive endocytosis of Ste3p. First, when receptor synthesis is shut off, *akr1* Δ cells retain the ability to mate longer than do *AKR1* cells. Second, Ste3p half-life is increased by greater than 5-fold in *akr1* Δ cells compared with *AKR1* cells. Third, after a pulse of synthesis, newly synthesized receptor remains at the cell surface in *akr1* Δ mutants, whereas it is rapidly internalized in *AKR1* cells. Specifically, in *akr1* Δ mutants, newly synthesized receptor is accessible to exogenous protease, and by indirect immunofluorescence, the receptor is located at the cell surface. *akr1* Δ cells are also defective for endocytosis of the α -factor receptor (Ste2p). Despite the block to constitutive endocytosis exhibited by *akr1* Δ cells, they are competent to carry out ligand-mediated endocytosis of Ste3p. In contrast, *akr1* Δ cells cannot carry out ligand-mediated endocytosis of Ste2p. We discuss the implications for Akrlp function in endocytosis and suggest a link to the regulation of ADP-ribosylation proteins (Arf proteins).

INTRODUCTION

Endocytosis provides one means for a cell to govern the steady-state abundance of plasma membrane proteins. For example, a protein destined for the plasma membrane may be synthesized at a significant rate but, nonetheless, be present at the cell surface in relatively low amounts as a result of rapid constitutive endocytosis. Endocytosis also provides a means to adjust the abundance of a plasma membrane protein in response to physiological stimuli. The cell biology of the yeast pheromone receptors illustrates both effects of endocytosis. Newly synthesized receptors are delivered to the cell surface by the secretory pathway

but reside there only a short time before being internalized and delivered to the vacuole for degradation (Davis *et al.*, 1993; Schandel and Jenness, 1994). As a result of this constitutive endocytosis, the receptors exhibit a half-life of 15–30 min (Davis *et al.*, 1993; Schandel and Jenness, 1994). The receptors are also subject to ligand-mediated endocytosis. Binding of pheromone accelerates the rate of degradation of wild-type receptors and overcomes the internalization block of some mutant receptors defective for constitutive endocytosis (Davis *et al.*, 1993; Rohrer *et al.*, 1993; Schandel and Jenness, 1994). Thus, there may be mechanistic differences between constitutive and ligand-mediated endocytosis. In this report, we sought to identify proteins that interact with the pheromone

* Corresponding author.

receptors with the expectation that some of these proteins could play a role in endocytosis.

The pheromone receptors, Ste2p (the α -factor receptor) and Ste3p (the a-factor receptor), are members of the serpentine G protein-coupled receptor family. Ste2p and Ste3p share no significant amino acid identity, but each has an N-terminal hydrophobic domain that presumably spans the membrane seven times and a C-terminal hydrophilic domain that is displayed on the cytoplasmic face of the plasma membrane. The cytoplasmic domain (or tail) mediates endocytosis. Removal of part of the cytoplasmic tail blocks constitutive endocytosis. For example, in the case of Ste3p, deletion of the C-terminal 105 amino acids causes a defect in constitutive endocytosis and, hence, the receptor accumulates at the cell surface (Davis *et al.*, 1993). The truncated receptor remains competent for ligand-mediated endocytosis, however. Binding of a-factor causes internalization of the truncated receptor and it is delivered to the vacuole where it accumulates if the strain is deficient for vacuole protease activity (i.e., in a *pep4* mutant; Davis *et al.*, 1993). Similarly, deletion of part of the cytoplasmic tail of Ste2p blocks constitutive internalization of that receptor but does not block ligand-mediated internalization (Konopka *et al.*, 1988; Reneke *et al.*, 1988; Rohrer *et al.*, 1993; Schandel and Jenness, 1994). In the context of a truncated Ste2p, amino acid substitutions in the residual Ste2p cytoplasmic tail have been identified that block ligand-mediated endocytosis (Rohrer *et al.*, 1993). Similar experiments have recently been published for Ste3p (Tan *et al.*, 1996). Together, these studies imply that the pheromone receptor cytoplasmic tails contain two, and perhaps more, signals that mediate endocytosis. Recent studies indicate that at least one of these signals involves the ubiquitination of specific amino acids within the cytoplasmic tails (Hicke and Riezman, 1996; Roth and Davis, 1996). However, aside from the sequence element defined by amino acid substitutions in truncated Ste2p and Ste3p, attempts to define specific cis-acting sequences required for either constitutive or ligand-mediated endocytosis have not yet been successful. It is possible that there are redundant elements within the cytoplasmic tails.

Genetic analysis in yeast has identified a number of gene products required for the efficient operation of the endocytic pathway and subsequent characterization has identified crudely the step in the pathway governed by some of these gene products. In some cases, this genetic analysis has identified gene products or subcellular structures that have mammalian counterparts also thought to act in the endocytic pathway. For example, abrogation of clathrin function causes a defect, albeit a modest one, in the initial

internalization of the yeast pheromone receptors (Payne *et al.*, 1988; Tan *et al.*, 1993). Similarly, the actin cytoskeleton is required for internalization of the pheromone receptors (Kubler and Riezman, 1993). In keeping with the involvement of the actin cytoskeleton in endocytosis, several genes required for the proper organization of the cytoskeleton have been identified by mutations that perturb endocytosis. These include SAC6 (which encodes a fimbrin homologue; Kubler and Riezman, 1993), SLA2/END4 (Raths *et al.*, 1993), CMD1 (which encodes calmodulin; Kubler *et al.*, 1994), END3 (Bénédetti *et al.*, 1994), END5/VRP1 (which encodes verprolin; Munn *et al.*, 1995), END6/RVS161 (Munn *et al.*, 1995), RVS167 (Munn *et al.*, 1995), ARP2 (Moreau *et al.*, 1996), and myosin (Geli and Riezman, 1996; Goodson *et al.*, 1996). However, it is unclear whether these gene products have a direct effect on endocytosis or simply influence endocytosis due to their role in regulating actin organization. Several members of the Rab GTPase family have been implicated in endocytic vesicular transport. These include Ypt51p, Ypt52p, and Ypt53 (each of which is homologous to mammalian Rab5 and thought to act at an early post-plasma-membrane step; Singer-Kruger *et al.*, 1994, 1995) and Ypt7, which is homologous to mammalian Rab7 and thought to act at a later endocytic step (Wichmann *et al.*, 1992; Schimmoller and Riezman, 1993), possibly controlling traffic from the early to late endosome. Finally, REN1/VPS2 and VPS27 are thought to control traffic from the late endosome to the vacuole (Davis *et al.*, 1993; Piper *et al.*, 1995). In summary, genetic analysis has provided a sketch of the yeast endocytic pathway, but it is apparent that much remains to be learned about how particular proteins are selected for internalization and the mechanism by which trafficking in the pathway is orchestrated.

We wished to gain a better understanding of the initial events in endocytosis and sought to identify proteins that interact with the a-factor receptor (Ste3p) cytoplasmic tail. We identified Akr1p in a two-hybrid screen of yeast genomic libraries by using the Ste3p cytoplasmic tail as bait. Akr1p is an ankyrin repeat-motif-containing protein that has previously been suggested to regulate signaling in the pheromone response pathway, perhaps by interacting with subunits of the heterotrimeric G protein (Kao *et al.*, 1996; Pryciak and Hartwell, 1996). We show that cells lacking Akr1p are defective for constitutive endocytosis of Ste3p and appear to be blocked at an early step in the endocytic pathway. Mutants lacking Akr1p can carry out ligand-mediated endocytosis of Ste3p. This phenotype distinguishes Akr1p from other gene products thought to act early in the endocytic pathway. Additionally, *akr1* Δ cells are defective for both constitutive and α -factor induced internalization of Ste2p.

MATERIALS AND METHODS

Plasmids and Strains

To construct pSL2194, which encodes the LexA DNA-binding domain fused to the 183 C-terminal residues of Ste3p (Ste3Tailp), the *STE3* gene in pSL1078 (Davis and Sprague, unpublished data) was first subjected to site-directed mutagenesis to introduce a *NcoI* site at nucleotide 836 of the open reading frame (ORF; causing Gly₂₈₀Leu₂₈₁ to Ala₂₈₀Met₂₈₁ substitutions), creating plasmid pSL1372 (Davis and Sprague, unpublished data). A 567-bp *NcoI*(Klenow-treated)-*PstI* fragment of pSL1372 was then inserted into *SmaI*-*PstI*-cut pBTM116 (Bartel *et al.*, 1993) to create pSL2179. Finally, to minimize the number of amino acids added to the Ste3p C terminus as a result of these manipulations, T4 DNA polymerase was used to remove the 5' overhang of the 3' *PstI* site and the resulting linear plasmid was religated. This places the first stop codon after the *STE3* insert in-frame with the fusion protein ORF and results in the addition of Pro-Asp to the C terminus of Ste3Tailp. The gene fusion is constitutively expressed from the *Saccharomyces cerevisiae* *ADHI* promoter.

Yeast strains SY1426 and SY1610 have been described previously (Davis *et al.*, 1993) and carry *GAL1-STE3* and *GAL1-STE3Δ365* alleles, respectively, in place of *STE3*. SY2841 is a derivative of SY1426 containing an *akr1::URA3* allele created by one-step gene replacement (Rothstein, 1991) using plasmid pPB590 (Kao *et al.*, 1996); correct replacement was confirmed by Southern blot analysis. Yeast strain SY3106 is isogenic to SY1426 except that it carries the native *STE3* allele and an integrated *mfa2-Δ1::FUS1-lacZ* allele. Yeast strain SY3251 is SY3106 into which the *akr1::URA3* allele is incorporated. Yeast strain SY1960 has been described previously (Davis *et al.*, 1993). Yeast strain SY3273 contains an *akr1::LEU2* disruption created by integrating a *ura3::LEU2* fragment produced by polymerase chain reaction into SY1960 carrying an *akr1::URA3* disruption created by integrating pPB590. Yeast strain CTY10-5D has been described previously and carries a genomic LexA_{Operator}-*β*-galactosidase reporter gene (Bartel *et al.*, 1993). Yeast strain YY2126 is CTY10-5D transformed with the two-hybrid bait plasmid pSL2194.

To immunolocalize Ste3p, we transformed yeast strain SY1426 or SY3243, which is an *akr1::ura3* derivative of SY2841 created by passaging on 5-fluoroorotic acid plates and screening for stable *ura*⁻ colonies, with plasmid pSL2624 encoding *GAL1-STE3(3Xmyc)* (generously provided by P. Kinsey, University of Oregon, Eugene, OR).

Two-Hybrid Screen

We screened yeast two-hybrid genomic DNA libraries essentially as described previously (Chien *et al.*, 1991; Bartel *et al.*, 1993). Yeast strain YY2126 was transformed with the genomic DNA libraries carried in the pGAD1F, pGAD2F, and pGAD3F vectors and plated on medium selecting for both the bait and library plasmids. The pGADF vectors are identical except for the reading frame of the library inserts (Chien *et al.*, 1991; Bartel *et al.*, 1993). We screened for blue colonies among approximately 400,000 transformants distributed equally among the three libraries. Putative Ste3Tailp interacting candidates were subjected to further examination. Candidate plasmids that activated the reporter gene without pSL2194 were eliminated from the collection. Candidate plasmids that required pSL2194 for reporter gene activity were subsequently screened against several other bait plasmids, including LexA-Fus1 (B. Ferguson, University of Oregon), LexA-p53 and LexA-lamin (LexA-p53 and LexA-lamin were generously provided by S. Fields, University of Washington, Seattle, WA). Clones interacting only with Ste3Tailp bait were subjected to DNA sequencing analysis using a primer corresponding to nucleotides 2570–2586 of the *GAL4* ORF. The library plasmid identified in our screen encoding the Gal4_{AD}-Akr1 fusion protein was stored in *Escherichia coli* strain CY3090.

Nitrocellulose Filter *β*-Galactosidase Assay

Detection of *β*-galactosidase activity was visualized essentially as described previously (Casadaban *et al.*, 1983) except that nitrocellulose filters were frozen at -70°C instead of in liquid N₂.

Mating Assays

Yeast patch mating assays using a *GAL1*-regulated *STE3* (*GAL1-STE3*) were performed as described previously (Davis *et al.*, 1993). For quantitative mating analysis of strains SY1426, SY1610 and SY2841, the following protocol was used (Sprague, 1991). Approximately 1×10^7 logarithmic phase cells were mixed with 2×10^6 SY1426, SY1610, or SY2841 cells grown to early logarithmic phase in either YEPGalactose (2% galactose, final concentration) medium or YEPGalactose medium to which dextrose was added (3%, final concentration) 1 h before the mating reaction. After mixing, cells were collected onto sterile 0.45- μm (pore size) nitrocellulose filter discs, and the discs were transferred to either YEPGalactose or YEPDextrose plates for 3 h. Cells were resuspended in water, lightly sonicated, and plated onto medium selecting for *a/a* diploids or the *a* haploids. Mating efficiency is expressed as the ratio of the number of *a/a* diploids produced to the number of *a* haploid cells present in the mating reaction.

Analysis of Ste3p Turnover and Cell Surface Abundance

To determine the half-life of Ste3p, we used two different pulse-chase protocols. The first relied on conventional radioactive labeling to examine a pulse of Ste3p expressed from its native promoter. The second relied on carbon source manipulation to provide a pulse of Ste3p synthesis from a *GAL1-STE3* construct. This latter protocol was combined with extracellular protease treatment to assess the dynamics of receptor delivery to the cell surface and its subsequent internalization.

In the radioactive pulse-chase protocol, exponentially growing cells were labeled for 15 min with [³⁵S]methionine (1 mCi/ml) as described previously (Davis *et al.*, 1993; Roth and Davis, 1996). Unlabeled methionine was added to 100 $\mu\text{g}/\text{ml}$ to initiate the chase. At 0 (before unlabeled methionine was added), 15, 45, and 90 min of chase, approximately 4×10^7 cells were collected by centrifugation and resuspended in 20 μl of BO buffer (8 M urea, 50 mM Tris-HCl, pH 8.0, 0.1 mM EDTA, 1% 2-mercaptoethanol). Glass beads were added and extracts prepared by vortexing for 15 s. Twenty microliters of FB buffer (5% SDS, 8 M urea, 50 mM Tris-HCl, pH 8.0, 0.1 mM EDTA, 1% 2-mercaptoethanol) were added and vortexing continued for an additional 60 s. Finally, the extracts were incubated at 100°C for 5 min. Immunoprecipitation of Ste3p was carried out as described previously (Davis *et al.*, 1993), except that we used polyclonal antiserum 348–4 preadsorbed to IgG-sorb. Precipitated proteins were subjected to SDS-PAGE.

In the alternative pulse-chase protocol, we used a modification of a previously published protease shaving procedure (Davis *et al.*, 1993). We grew *GAL1-STE3* cells in YEP Raffinose (2%) until early logarithmic phase ($0.5\text{--}1.0 \times 10^7$ cells/ml), at which time galactose was added to a final concentration of 2% to induce synthesis of Ste3p. After 30 min, dextrose was added to a final concentration of 3%, which rapidly inhibits *GAL1-STE3* expression (as judged by Northern blot analysis). The amount of Ste3p produced from such an induction is similar to the steady-state level of Ste3p expressed from its native promoter (examined by Western blot analysis). Samples were taken from the raffinose culture prior to addition of galactose and at various times after addition of galactose and dextrose as follows. Approximately 1×10^8 cells were collected by gentle centrifugation over crushed ice, washed once in 1.5 ml of chilled Pronase buffer II (PBII; 1.4 M sorbitol, 25 mM Tris-HCl, pH 7.5, 2 mM MgCl₂, 20 mM Na₂S₂O₃, 0.5% 2-mercaptoethanol), and resuspended in 350 μl of chilled PBII. Cells were stored on ice until all samples were collected. After a brief incubation at 30°C , samples

Figure 1. Two-hybrid interaction between Ste3Tailp and Akr1p. Yeast cells were grown on buffered sucrose minimal medium selecting for LexA-Ste3Tail and library plasmids (A–C). The cells were transferred to nitrocellulose and exposed to 5-bromo-4-chloro-3-indolyl β -D-galactoside (D–F) as described in MATERIALS AND METHODS. Cell patches in A and D carry LexA-Ste3Tail bait plasmid and negative control plasmid GAD424. Cell patches in sections B and E carry negative control plasmid LexA-Lamin and GAL4_{AD}-Akr1 library plasmid. Cell patches in C and F carry LexA-Ste3Tail and GAL4_{AD}-Akr1 library plasmids; significant β -galactosidase activity is only observed when LexA-Ste3Tailp and GAL4_{AD}-Akr1 plasmids are present in yeast strain CTY10–5D.

were split into two 170- μ l aliquots. One aliquot of each pair was treated with a broad-spectrum protease by adding 42.5 μ l of Pronase solution (Calbiochem, San Diego, CA; 2500 U/ml in PBII); the other aliquot was treated with a mock Pronase mixture, which was simply 42.5 μ l of PBII lacking Pronase. Cells were incubated for 60 min at 37°C. After Pronase treatment, cells were collected by gentle centrifugation, resuspended in PBII containing 17% trichloroacetic acid, and stored at –70°C until further analysis. Approximately 280 μ l of acid-washed glass beads were added to frozen samples, which were then vortexed at high speed for 10 min. The glass beads were pelleted by a quick centrifugation and washed twice with 280 μ l of 5% trichloroacetic acid. The washes were pooled and centrifuged at 4°C for 10 min to pellet cellular debris. This pellet was resuspended in 40 μ l of SDS-polyacrylamide gel sample buffer (Davis *et al.*, 1993) and subjected to SDS-PAGE and Western blot analysis.

To examine a-factor induced endocytosis of Ste3p, the carbon source shift pulse-chase protocol was used except that one volume of a-factor-containing culture supernatant, purified as described previously (Davis *et al.*, 1993), was added in conjunction with the addition of dextrose to 3%. Protein extracts were prepared as described above and subjected to SDS-PAGE and Western blot analysis.

Analysis of Ste2p Turnover

To examine the cellular degradation of Ste2p, we used cycloheximide to block new Ste2p synthesis (Schandel and Jenness, 1994) and followed the disappearance of Ste2p by Western blot analysis. Cultures of yeast strains SY1960 or SY3273 carrying plasmid pAB524 (*CEN URA3 STE2*; Anne Burkholder, University of Washington) were grown to early logarithmic phase ($0.5\text{--}1.0 \times 10^7$ cells/ml) in minimal medium selecting for the *STE2* plasmid. Cycloheximide was added to a final concentration of 10 μ g/ml and growth continued for 7 min. After removing 5 ml ($t = 0$), the culture was split in half. To one half, α -factor was added to a final concentration of 1×10^6 M. Five-milliliter samples were removed from both cultures at 10, 30, 60, and 90 min. Cell extracts were prepared as above for Ste3p analysis. The protein concentration of each cell extract was determined by bicinchoninic acid assay (Pierce, Rockford, IL). Each sample was diluted to a final protein concentration of 1 μ g/ μ l, incubated for 5 min at 37°C, and subjected to SDS-PAGE. Quantification of Ste2p-specific bands was performed with a Relysis RELI4816

flatbed scanner and NIH Image 1.60 software on films exposed to blots containing dilutions of each time point to verify linearity. Both strains used in these experiments displayed similar growth-inhibitory sensitivities to cycloheximide.

Indirect Immunofluorescence Staining

To immunolocalize Ste3p, we transformed SY1426 or SY3241 with pSL2624 and grew cells to early logarithmic phase in minimal medium lacking uracil and containing 2% raffinose as the carbon source (SRaff – ura). Galactose was added to 2%, followed 3 h later by the addition of dextrose to 3%. Cells were allowed to grow for 1 h after dextrose addition before being collected and prepared for antibody staining as described previously (Davis *et al.*, 1993).

Western Blot Analysis

After SDS-PAGE, cell extracts were transferred to a nitrocellulose membrane. For detection of Ste3p, the membrane was probed with a cell culture supernatant containing monoclonal antibody directed against the C-terminal tail of Ste3p at dilutions between 1:50 and 1:100. Antigen–antibody complexes were visualized using horseradish peroxidase (HRP)-conjugated donkey anti-mouse antibody (Jackson ImmunoResearch, West Grove, PA) in combination with either ECL (Amersham, Arlington Heights, IL) or SuperSignal CL-HRP substrate system (Pierce) fluorescent imaging as per the manufacturers' instructions. For detection of the cytoplasmic protein phosphoglycerate kinase (Pgk), polyclonal anti-Pgk antibody (generously provided by T. Stevens, University of Oregon) was used at a 1:10,000 dilution and detected with alkaline phosphatase-conjugated anti-rabbit (Promega, Madison, WI) by colorimetric means. For detection of Ste2p we used polyclonal anti-Ste2 antibody generously provided by Nick Davis (Wayne State University) at a 1:2000 dilution and visualized with HRP-conjugated anti-rabbit (Promega) in combination with SuperSignal CL-HRP substrate system (Pierce) as per the manufacturers' instructions.

RESULTS

Akr1p Displays a Two-Hybrid Interaction with Ste3p Intracellular Tail

As outlined in the INTRODUCTION, the intracellular tail of Ste3p is required for receptor endocytosis. To identify additional factors in this process, we used plasmid pSL2194 to express Ste3Tailp as bait for a two-hybrid screen of yeast genomic libraries. Sixty-nine plasmids that conferred β -galactosidase production were identified from a screen of approximately 400,000 colonies (see MATERIALS AND METHODS). The DNA sequence of each library insert was determined and nine distinct genes were chosen for further analysis. One of these genes, *AKR1*, is the subject of this report; analysis of the other genes will be reported elsewhere. As illustrated in Figure 1, only cells carrying both pSL2194 and the Gal4_{AD}-Akr1 fusion protein library plasmid activated the β -galactosidase reporter gene (see Figure 1, C and F). The fragment of *AKR1* isolated begins at nucleotide 252 of the ORF, which corresponds to amino acid 85 of Akr1p, midway through the first ankyrin repeat motif (Kao *et al.*, 1996).

akr1Δ Cells Exhibit Prolonged Mating Ability after Ste3p Receptor Synthesis Inhibition

As mentioned in the INTRODUCTION, Akr1p influences signaling in the pheromone response pathway in an as yet undefined way (Kao *et al.*, 1996; Pryciak and Hartwell, 1996). We asked whether Akr1p might also play a role in the endocytosis of Ste3p. As a first test of this possibility, we investigated the mating properties of wild-type and *akr1Δ* mutant cells that carry *GAL1-STE3* as the sole means to express Ste3p. *AKR1 GAL1-STE3* cells quickly lost the ability to mate 1 h after transfer from galactose to dextrose medium (Table 1). This condition blocks new receptor synthesis, and preexisting surface receptor is rapidly internalized and degraded in the vacuole as a result of constitutive endocytosis (Davis *et al.*, 1993). The cells are effectively receptorless and are, therefore, deficient for mating. In contrast, *akr1Δ* cells retained some mating capacity after inhibition of *GAL1-STE3* expression, suggesting that *akr1Δ* cells are at least partially defective for constitutive endocytosis (Table 1). However, the mating proficiency of *akr1Δ GAL1-STE3* cells is not as great as wild-type *AKR1* cells carrying a *STE3* partial truncation allele, *GAL1-STE3Δ365* (Table 1), which encodes a receptor known to be defective for constitutive endocytosis (Davis *et al.*, 1993).

akr1Δ Cells Display a Ste3p Degradation Defect

The experiments described above are consistent with the possibility that *akr1Δ* cells are defective for endocytosis of Ste3p, but other possibilities exist. For example, a mutation that resulted in an elevated basal signal in the pheromone pathway or that increased sensitivity to pheromone could allow mating by effectively amplifying the signal generated by the few Ste3p molecules that remain at the cell surface 1 h after repression of Ste3p synthesis. Indeed, published reports indicate that *akr1Δ* cells exhibit an elevated basal signal and are supersensitive to added pheromone (Kao *et al.*, 1996; Pryciak and Hartwell, 1996). However, in our strain background deletion of *AKR1* did not increase basal pheromone response as measured by expression of *FUS1-lacZ* in strains SY3106 (*AKR1*; 5.8 U of β -galactosidase activity) and SY3251 (*akr1Δ*; 1.3 U), nor did it increase α -factor-induced *FUS1-lacZ* levels (*AKR1*, 38.6 U; *akr1Δ*, 30.5 U), which is a sensitive indicator of pathway activity. In addition, *akr1Δ* cells were only slightly more sensitive to α -factor than *AKR1* cells, as measured by halo assay, and this apparent increase in sensitivity may simply reflect the slower growth rate of *akr1Δ* cells. In our strain background, *akr1Δ* did cause other phenotypes congruent with those previously reported; slow growth at 30°C, no growth at 37°C, and abnormal cell morphology at 37°C (Kao *et al.*, 1996; Pryciak and Hartwell, 1996).

Table 1. Quantification of mating efficiency

Strain	Galactose		Dextrose	
	Mating Eff ^a	Relative to <i>AKR1</i>	Mating Eff	Relative to <i>AKR1</i>
<i>AKR1</i>	0.2965	1.0	0.00096	1.0
<i>STE3Δ365</i>	0.4026	1.4	0.33690	350
<i>akr1Δ</i>	0.2290	0.8	0.0361	38

^a Eff, efficiency.

Given that deletion of *AKR1* does not substantially affect pheromone signaling in our strains, we pursued the possibility that the mating phenotype of *akr1Δ* cells reveals a defect in endocytosis. Because endocytosis from the plasma membrane ultimately leads to vacuolar degradation of Ste3p (Davis *et al.*, 1993), cells compromised for Ste3p endocytosis should degrade the receptor more slowly. We initially followed the degradation of Ste3p by Western blot analysis of samples isolated after a short burst of *GAL1-STE3* expression (see MATERIALS AND METHODS). Ste3p was degraded rapidly in *AKR1* cells (see Figures 3A and 4A, minus Pronase lanes), but in contrast, Ste3p was quite stable in *akr1Δ* cells after inhibition of *GAL1-STE3* (see Figures 3B and 4B, minus Pronase lanes).

As a second measure of Ste3p turnover kinetics, we performed a [³⁵S]methionine pulse-chase experiment using cells that express Ste3p from its native promoter. After a 15-min pulse and an extended chase in excess unlabeled methionine, Ste3p was observed to have a half-life of approximately 18 min in *AKR1* cells (Figure 2). In contrast, Ste3p was degraded much more slowly in *akr1Δ* cells, displaying a half-life of approximately 95 min. These results are consistent with previous estimates of Ste3p half-life in wild-type cells (Davis *et al.*, 1993; Gammie *et al.*, 1995) and support the idea that *akr1Δ* cells have a defect in endocytosis.

akr1Δ Cells Display an Early Block in Ste3p Endocytosis

In principle, the reduced degradation of Ste3p seen in *akr1Δ* cells could reflect a block anywhere along the endocytic pathway from the cell surface to the vacuole, or it could even reflect a partial block in the secretory pathway. To localize more precisely the defect in cellular trafficking of Ste3p, we measured the kinetics of its delivery to the cell surface and subsequent internalization by assessing susceptibility to exogenous protease (the pulse-chase Pronase, or PCP assay). To provide a pulse of receptor synthesis, we used the *GAL1-STE3* allele. Galactose was added to a raffinose culture of *GAL1-STE3 AKR1* or *akr1Δ* cells and incubation continued for 30 min.

Figure 2. $[^{35}\text{S}]$ Methionine pulse-chase analysis of Ste3p half-life in *AKR1* and *akr1Δ* cells. Strains SY3106 (*AKR1*) and SY3251 (*akr1Δ*) were labeled with $[^{35}\text{S}]$ methionine for 15 min and then excess unlabeled methionine was added to initiate the chase. Ste3p was immunoprecipitated from samples collected at the indicated times and subjected to SDS-PAGE as described in MATERIALS AND METHODS. (A) Autoradiograph exposure of SDS-PAGE gel illustrating degradation kinetics of Ste3p in SY3106 and SY3251. (B) Quantitation of the Ste3p bands from a similar exposure of the gel in A. Quantitation of the radioactivity in each band was obtained with the Molecular Dynamics Storm phosphorimaging system and ImageQuant software.

Dextrose was then added to block transcription of *GAL1-STE3*. Samples were taken at intervals over this pulse-chase time course and added to chilled buffer containing NaN_3 to halt membrane trafficking. The intact cells were then treated with a broad-spectrum protease (Pronase), or mock treated, and extracts were prepared and subjected to SDS-PAGE and Western blot analysis.

Figure 3 represents a PCP comparison of *AKR1* and *akr1Δ* cells. In *AKR1* cells, Ste3p is synthesized upon addition of galactose. By the end of the 30-min pulse of receptor synthesis in wild-type cells, a fraction of the receptor has reached the cell surface, as revealed by susceptibility to Pronase (Figure 3A, lanes 3–6); Pronase digestion reduces the amount of 48-kDa full-length Ste3p and causes the concomitant appearance of a 32-kDa Ste3p intracellular tail digestion product (Figure 3A). We interpret the protease-resistant Ste3p

Figure 3. PCP analysis of Ste3p subcellular location. *GAL1-STE3* was expressed in strains SY1426 (*AKR1*; A) and SY2841 (*akr1Δ*; B) as described in MATERIALS AND METHODS. Lanes raff, samples collected when cells were growing in YEP-raffinose; lanes 30G, samples collected 30 min after galactose was added; lanes 5D and subsequent lanes, samples collected at the indicated times (minutes) after dextrose was added. Therefore, cells from the D time points see three carbon sources: raffinose (*GAL1-STE3* noninducing), galactose (*GAL1-STE3* inducing) and dextrose (*GAL1-STE3* repressing). Pron (-/+) indicates absence or addition of Pronase. Samples were removed after the indicated growth conditions and treated to block further endocytosis of Ste3p. Protein extracts were then prepared and subjected to SDS-PAGE and Western blot analysis as described in MATERIALS AND METHODS. Blots were initially probed with monoclonal antibody recognizing Ste3p and subsequently probed with polyclonal antiserum recognizing the cytoplasmic protein Pgk. Molecular weights are indicated to the left of the gels. The cytoplasmic protein Pgk serves as loading control and to verify the integrity of the cells.

present at the 30G (30 min after galactose was added) and 5D (5 min after dextrose was added) time points (Figure 3A) to be molecules in transit to the cell surface because by 15 min of chase (Figure 3A, lanes 15D) essentially all receptor was protease sensitive and inferred to be at the cell surface. As the chase progressed, the abundance of Ste3p decreased, reflecting internalization and delivery to the vacuole, where it was completely degraded. The transit time from the plasma membrane to the vacuole must be rapid, be-

cause we fail to see an accumulation of Pronase-insensitive Ste3p at later dextrose time points.

As illustrated in Figure 3B, Ste3p was synthesized and reached the cell surface in *akr1Δ* cells with kinetics similar to that seen in *AKR1* cells (compare Figure 3B, lanes 3–8, to Figure 3A, lanes 3–8), indicating that *Akr1p* is not involved in transport of Ste3p to the plasma membrane. However, unlike the observation for *AKR1* cells, Ste3p abundance did not diminish during the chase and essentially all the Ste3p remained susceptible to exogenous protease. These data therefore indicate that Ste3p remains at the cell surface in *akr1Δ* mutants and imply that *AKR1* function is required early in the endocytic pathway. It should be noted that general membrane flow from the plasma membrane to the vacuole appears to be normal since we observed normal uptake of the fluorescent dye FM 4–64 (Vida and Emr, 1995) at 30°C (Givan and Sprague, unpublished observations). In addition, sorting of vacuolar proteins appears to be normal because we failed to detect significant secretion of the vacuolar peptidase carboxypeptidase Y (Givan and Sprague, unpublished observations).

We considered the possibility that the presence of cell surface receptor in *akr1Δ* mutants at late chase times could result from a late block in the endocytic pathway rather than from an early block. For example, perhaps the accumulation of an intermediate endocytic compartment indirectly affects early steps in endocytosis by mass action or some other means. Such a defect could simply slow Ste3p internalization. As a test of this possibility, we used the PCP assay to determine whether Ste3p became Pronase resistant at more extended chase times in *akr1Δ* cells. However, even 180 min after receptor synthesis was shut-off, a substantial amount of Pronase-sensitive Ste3p remained and no significant accumulation of Pronase-protected Ste3p was observed in *akr1Δ* cells (Figure 4B, lanes 9–14). The PCP assay can detect Pronase-protected Ste3p in mutants known to be defective at a late endocytic step, for example, *ren1/vps2* (Davis *et al.*, 1993; Givan and Sprague, unpublished observations). Therefore, these results indicate that the major endocytic block in *akr1Δ* cells occurs at an early step.

Pheromone-induced Endocytosis of Ste3p Is Not Blocked in *akr1Δ* Cells

As mentioned in the INTRODUCTION, Ste3p is subject to at least two modes of endocytosis: constitutive and pheromone induced (Davis *et al.*, 1993). To explore whether *Akr1p* is involved in the pheromone-induced endocytosis of Ste3p, we used a modification of the PCP assay. Specifically, both dextrose and a culture supernatant containing *a*-factor (see MATERIALS AND METHODS) were added to a culture of *GAL1-STE3 AKR1* or *GAL1-STE3 akr1Δ* cells after the

Figure 4. PCP analysis of Ste3p subcellular location at extended time points. Cells were grown and treated as in Figure 3. (A) Strain SY1426 (*AKR1*). (B) Strain SY2841 (*akr1Δ*).

30-min galactose induction pulse. Samples were removed at intervals and analyzed as described above for the PCP assay. Figure 5A illustrates Ste3p internalization and degradation in *AKR1* cells. The majority of Ste3p is degraded by 120 min after the addition of dextrose and the *a*-factor supernatant (Figure 5A, lanes 9 and 10). The slightly delayed internalization and degradation of Ste3p in Figure 5A, as compared to Figure 4A, appear to be caused by the addition of exhausted culture supernatant, not the presence of *a*-factor per se, because the addition of a similar supernatant lacking *a*-factor caused a comparable delay in Ste3p turnover (Givan and Sprague, unpublished observation). Figure 5B illustrates *a*-factor-induced internalization and degradation of Ste3p in *akr1Δ* cells. The presence of *a*-factor caused a gradual reduction in receptor abundance over the course of the dextrose chase (compare to Figure 4B). The kinetics of this pheromone-induced internalization of Ste3p in *akr1Δ* mutants is similar to the kinetics of pheromone-induced internalization in wild-type cells of a truncated Ste3p (Ste3Δ365p) that is refractory to constitutive endocytosis (as judged by PCP analysis).

Figure 5. PCP analysis of Ste3p subcellular location after a-factor treatment. Cells were grown and treated as in Figure 3 except that a culture supernatant containing a-factor was added concomitant with dextrose. (A) Strain SY1426 (*AKR1*). (B) Strain SY2841 (*akr1Δ*).

akr1Δ Cells Display More Surface-localized Ste3p after Receptor Synthesis Inhibition

As a further test of the cellular location of Ste3p after inhibition of *GAL1-STE3*, we performed indirect immunofluorescence with a strain containing a 3X-myc-tagged version of Ste3p under galactose control [*GAL1-STE3(3X-myc)*]. *AKR1* and *akr1Δ* strains containing *GAL1-STE3(3X-myc)* were subjected to a galactose to dextrose growth regimen and stained with a monoclonal antibody recognizing the myc epitope. After growth on dextrose for 1 h, *AKR1* cells displayed primarily punctate intracellular staining of Ste3(3X-myc)p (Figure 6C). This punctate staining presumably reflects receptors present in endocytic compartments in transit to the vacuole. Vacuolar staining of Ste3(3X-myc)p is not observed because receptors are degraded in these *PEP4* cells. In contrast, *akr1Δ* cells displayed primarily cell-surface staining of Ste3(3X-myc)p (Figure 6D). Moreover, no punctate staining was observed in the *akr1Δ* cells (Figure 6D), presumably reflecting the lack of inter-

Figure 6. Indirect immunofluorescent localization of Ste3(3X-myc)p. *GAL1-STE3(3X-myc)* containing cells were subjected to a galactose-pulse/dextrose-chase regimen and prepared for antibody staining following 1 h of growth on dextrose. (A and B) Representative *AKR1* cells. (C and D) Representative *akr1Δ* cells. Approximately 85% of the *AKR1* cells displaying significant staining with the monoclonal anti-myc antibody appear as in B ($n = 100$), with the remainder having a diffuse cytoplasmic staining. Approximately 65% of the *akr1Δ* cells displaying significant Ste3(3X-myc)p staining appear as in D ($n = 100$), with the remainder displaying diffuse cytoplasmic staining.

nalization and incorporation of Ste3(3X-myc)p into endosomal organelles.

akr1Δ Cells Are Defective for Ste2p Endocytosis

Although Ste3p and Ste2p, the two pheromone receptors, do not share significant amino acid similarities, the same mechanisms are thought to govern their endocytosis. C-terminal truncation mutants of each receptor are defective for constitutive endocytosis (Konopka *et al.*, 1988; Reneke *et al.*, 1988; Davis *et al.*, 1993). Moreover, mutations in several genes, including *REN1/VPS2* (Davis *et al.*, 1993), *END4/SLA2* (Roth and Davis, 1996; P. Kinsey, personal communication), *DNM1* (Gammie *et al.*, 1995), and *CHC1* (Tan *et al.*, 1993), block the endocytosis of each receptor. To examine Akr1p's role in Ste2p endocytosis, we followed the degradation kinetics of Ste2p in *AKR1* and *akr1Δ* cells. As Figure 7A, lanes 2–6, illustrates, Ste2p disappears quite rapidly after cycloheximide (10 μ g/ml) is added to a logarithmic-phase culture of *AKR1* cells. In contrast, Ste2p remains stable in *akr1Δ* cells after cycloheximide addition (see Figure 7A, lanes 8–12). Additionally, *akr1Δ* cells accumulate a greater amount of steady-state Ste2p (Figure 7A, compare lanes 2 and 8), consistent with a defect in the constitutive internalization and degradation of Ste2p. The half-life of Ste2p calculated from gels similar to those shown in Figure 7A is 27 min in *AKR1* cells and 109 min in *akr1Δ* cells, representing a fourfold increase for *akr1Δ* cells.

We also investigated the degradation of Ste2p in *AKR1* and *akr1Δ* cells in response to α -factor binding. As is observed in Figure 7B (lanes 2–6), *AKR1* cells degrade Ste2p more rapidly in the presence of α -fac-

Figure 7. Analysis of Ste2p degradation kinetics. (A) Cycloheximide was added to a logarithmic-phase culture of strain SY1960 (AKR1) or SY3273 (*akr1Δ*) carrying a *STE2* plasmid and incubated for 7 min ($t = 0$). Samples in B were exposed to α -factor immediately after $t = 0$. Samples were removed at the indicated times (minutes) after $t = 0$ and protein extracts were prepared as described in MATERIALS AND METHODS. Ten micrograms of total cellular protein were loaded in each lane and subjected to SDS-PAGE and Western blot analysis. Nitrocellulose membranes were probed with polyclonal anti-Ste2p as described in MATERIALS AND METHODS. Lanes – represent protein extracts from SY1960 or SY3273 cells lacking the *STE2* plasmid.

tor. In contrast, Ste2p remains stable in *akr1Δ* cells even in the presence of α -factor (see Figure 7B, lanes 8–12). Therefore, Akrlp appears to be required for both constitutive and pheromone-induced endocytosis of Ste2p.

DISCUSSION

The C-terminal intracellular tail of the yeast α -factor receptor is required for constitutive endocytosis. To understand better the mechanisms underlying this mode of endocytosis, we wished to identify other proteins involved. Using the two-hybrid system to screen yeast genomic libraries, we identified Akrlp as a protein that interacts with the intracellular tail of Ste3p. Whether Akrlp and Ste3p interact directly or only as part of a larger complex cannot be discerned from these two-hybrid results. Nonetheless, analysis of the phenotype of *akr1Δ* mutants reveals that Akrlp plays a role in executing the constitutive endocytosis function of the Ste3p intracellular tail. Specifically, we find that newly synthesized Ste3p is retained at the cell surface in *akr1Δ* mutants for a protracted time compared with wild-type cells. Likewise, *akr1Δ* mutants are defective for the constitutive endocytosis of the α -factor receptor.

Akrlp may play a subtle role in ligand-mediated endocytosis of Ste3p in addition to its more profound role in constitutive endocytosis. This possibility follows from the observation that α -factor stimulated internalization and delivery of Ste3p to the vacuole in

akr1Δ mutants appears to be slow compared with wild-type cells. Wild-type cells expressing a version of Ste3p (Ste3 Δ 365p) that is not subject to constitutive endocytosis but is still competent to undergo ligand-mediated endocytosis show similar kinetics of internalization (Givan and Sprague, unpublished observations). Perhaps Akrlp, acting through the Ste3p tail, facilitates the merging of the ligand-stimulated endocytic pathway with the constitutive pathway. Alternatively, perhaps Akrlp functions at more than one step in the endocytic pathway, but its involvement in a later step is apparent only when its role at an early stage in constitutive endocytosis is bypassed by activation of the ligand-induced pathway. In any event, the finding that loss of Akrlp leads to a more pronounced block in constitutive endocytosis of Ste3p than in ligand-mediated endocytosis of Ste3p supports the idea that early events in the two pathways are mechanistically distinct. As alluded to above, this possibility was suggested based on the properties of C-terminal truncated versions of Ste3p. The phenotype of *akr1Δ* mutants now extends this mechanistic distinction to include a trans-acting protein.

Our results suggest that the ligand-induced endocytic paths traversed by the two pheromone receptors may differ. Specifically, we find that ligand-mediated internalization of Ste3p can occur in an Akrlp-independent manner whereas ligand-mediated endocytosis of Ste2p requires Akrlp. Recently, an amino acid sequence unique to Ste3p that can mediate the pheromone induced endocytosis of a truncated Ste3p (Ste3 Δ 365p) has been identified (Tan *et al.*, 1996). Ste2p lacks a similar sequence, so perhaps the two receptors use different signals for ligand-mediated endocytosis. In light of this possibility, it is interesting that the sequencing of the complete *S. cerevisiae* genome has uncovered a homologue of Akrlp. Akrlp, encoded by ORF YOR034c, shares 41% overall amino acid identity with Akrlp. Due to this high degree of similarity between the two proteins, we have begun to investigate Akrlp's role in endocytosis. Preliminary results indicate that Akrlp is involved in Ste3p constitutive endocytosis, although to a lesser extent than Akrlp. Perhaps Akrlp plays a specific role in Ste3p α -factor-induced endocytosis.

Our experiments imply that Akrlp acts at an early step in endocytosis, potentially at or near the plasma membrane, to mediate constitutive internalization of both Ste3p and Ste2p from the cell surface. What might be Akrlp's function in endocytosis? Preliminary results place limits on the type of functions to be considered and raise an intriguing possibility for future investigation. Many mutants that exhibit endocytosis defects display abnormal actin localization (Kubler and Riezman, 1993; Raths *et al.*, 1993; Bénédicti *et al.*, 1994; Kubler *et al.*, 1994; Munn *et al.*, 1995; Moreau *et al.*, 1996). However, as judged by

rhodamine-phalloidin staining, the actin cytoskeleton is normal in *akr1Δ* mutants (Givan and Sprague, unpublished observations). On the basis of hydrophobicity and α -helix amphiphilicity analyses, Akr1p has several potential membrane spanning regions (Kao *et al.*, 1996), suggesting it may be an integral membrane protein. Protease shaving experiments similar to those reported herein for Ste3p failed to detect Akr1p at the cell surface (Givan and Sprague, unpublished observations). However, indirect immunolocalization experiments suggest that Akr1p resides near the plasma membrane (A. Bender, personal communication). Hence, Akr1p could be a membrane spanning protein associated with early endocytic vesicular compartments, where it may regulate import of plasma membrane proteins into these vesicles.

Other studies have identified interactions between Akr1p and proteins involved in regulating budding and pheromone signaling (Bem1p and the G protein subunits Ste4p, Ste18p, and Gpa1p), whereas our studies identified an interaction with the pheromone receptor that appears to mediate endocytosis. Is there a common theme that unites these seemingly disparate Akr1p activities? Ste4p and Gpa1p are demonstrably not required for receptor endocytosis (Jenness and Spatrick, 1986; Zanolari *et al.*, 1992; Davis *et al.*, 1993), nor has Bem1p been implicated in Ste3p or Ste2p endocytosis. Given the array of potential Akr1p roles, it is intriguing that Akr1p displays a two-hybrid interaction with Gcs1p (Kao *et al.*, 1996), a protein that shows sequence similarity to GTPase-activating proteins for ADP ribosylation factor proteins (Arf proteins). Recent biochemical evidence indicates that Gcs1p can activate the intrinsic GTPase activity of mammalian and yeast Arf proteins (Poon *et al.*, 1996). Arf proteins are implicated in several vesicular transport processes, including endocytosis (Moss and Vaughan, 1995; Whitney *et al.*, 1995), and Arf proteins are also thought to participate with some trimeric G proteins in regulating vesicular traffic (Colombo *et al.*, 1995). In light of the possibility that Arfs may participate in endocytosis, it is noteworthy that *gcs1Δ* cells exhibit an endocytosis defect (Wang *et al.*, 1996; Petsko and Gray, personal communication); *akr1Δ* strains show similar defects (Givan and Sprague, unpublished observation). Perhaps Akr1p plays a role in the regulation of Arf GTPase proteins in yeast through interactions with Arf-GTPase-activating protein proteins such as Gcs1p. In such a scenario, the lack of Akr1p could lead to defects in endocytosis due to misregulation of one or more Arf proteins. Identification of a yeast Arf protein involved in endocytosis would lend support to this model. On the basis of amino acid homology, two candidates for such an Arfp exist in yeast that are encoded by the *ARL2/ARF3/YOR094w* and *ARL1/YBR164c* ORFs. *ARL2/ARF3/YOR094w* encodes an Arfp that has been shown

to function outside of Golgi transport (Lee *et al.*, 1994), where Arf1p and Arf2p are thought to act (Stearns *et al.*, 1990a,b). Interestingly, the *Arl2/Arf3/YOR094w* protein is most similar in amino acid sequence to human Arf6p (Lee *et al.*, 1994), which has been implicated in the regulation of early endocytic transport (D'Souza-Schorey *et al.*, 1995; D'Souza-Schorey and Stahl, 1995; Peters *et al.*, 1995). Further work will determine whether Akr1p and Gcs1p are acting together to regulate the endocytosis of plasma membrane proteins, possibly through the control of Arf proteins.

ACKNOWLEDGMENTS

We thank Phil Kinsey for innumerable discussions and suggestions, Joe Horecka for invaluable technical assistance, and all members of the Sprague laboratory for interest and encouragement during this work. We thank Nick Davis for assistance with Ste3p immunoprecipitations and Joe Horecka and Sylvia Gibson for purification of the Ste3p monoclonal antibody. We also thank Alan Bender and Peter Pryciak for reagents, helpful discussions, and communication of results prior to publication, and Pak Phi Poon, Gerald Johnston and Greg Petsko for communication of results prior to publication. This work was supported by research (GM-30027) and training (HD-07348) grants from the U.S. Public Health Service and a grant from the American Cancer Society (CB-76).

REFERENCES

- Bartel, P.L., Chien, C., Sternglanz, R., and Fields, S. (1993). Using the two-hybrid system to detect protein-protein interactions. *Cellular Interactions in Development: A Practical Approach*, Oxford, United Kingdom: IRL Press.
- Bénédetti, H., Rath, S., Crausaz, F., and Riezman, H. (1994). The *END3* gene encodes a protein that is required for the internalization step of endocytosis and for actin cytoskeleton organization in yeast. *Mol. Biol. Cell* 5, 1023–1037.
- Casadaban, M.J., Martinez-Arias, A., Shapira, S.K., and Chou, J. (1983). β -Galactosidase gene fusions for analyzing gene expression in *Escherichia coli* and yeast, *Methods in Enzymology: Recombinant DNA*, Part B, New York: Academic Press, 293–308.
- Chien, C.-T., Bartel, P.L., Sternglanz, R., and Fields, S. (1991). The two-hybrid system: a method to identify and clone genes for proteins that interact with a protein of interest. *Proc. Natl. Acad. Sci. USA* 88, 9578–9582.
- Colombo, M.I., Inglese, J., D'Souza, S.C., Beron, W., and Stahl, P.D. (1995). Heterotrimeric G proteins interact with the small GTPase ARF. Possibilities for the regulation of vesicular traffic. *J. Biol. Chem.* 270, 24564–24571.
- D'Souza-Schorey, C., Li, G., Colombo, M.I., and Stahl, P.D. (1995). A regulatory role for ARF6 in receptor-mediated endocytosis. *Science* 267, 1175–1178.
- D'Souza-Schorey, C., and Stahl, P.D. (1995). Myristoylation is required for the intracellular localization and endocytic function of ARF6. *Exp. Cell. Res.* 221, 153–159.
- Davis, N.G., Horecka, J.L., and Sprague, G.F., Jr. (1993). Cis- and trans-acting functions required for endocytosis of the yeast pheromone receptors. *J. Cell Biol.* 122, 53–65.
- Gammie, A.E., Kurihara, L.J., Vallee, R.B., and Rose, M.D. (1995). *DNM1*, a dynamin-related gene, participates in endosomal trafficking in yeast. *J. Cell Biol.* 130, 553–566.

- Geli, I.M., and Riezman, H. (1996). Role of type I myosins in receptor-mediated endocytosis in yeast. *Science* 272, 533–535.
- Goodson, H.V., Anderson, B.L., Warrick, H.M., Pon, L.A., and Spudis, J.A. (1996). Synthetic lethality screen identifies a novel yeast myosin I gene (*MYO5*): myosin I proteins are required for polarization of the actin cytoskeleton. *J. Cell Biol.* 133, 1277–1291.
- Hicke, L., and Riezman, H. (1996). Ubiquitination of a yeast plasma membrane receptor signals its ligand-stimulated endocytosis. *Cell* 84, 277–287.
- Jenness, D.D., and Spatrick, P. (1986). Down regulation of the alpha-factor pheromone receptor in *S. cerevisiae*. *Cell* 46, 345–353.
- Kao, L.-R., Peterson, J., Ruiru, J., Bender, L., and Bender, A. (1996). Interactions between the ankyrin repeat-containing protein Akr1p and the pheromone response pathway in *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 16, 168–178.
- Konopka, J.B., Jenness, D.D., and Hartwell, L.H. (1988). The C-terminus of the *S. cerevisiae* alpha-pheromone receptor mediates an adaptive response to pheromone. *Cell* 54, 609–620.
- Kubler, E., and Riezman, H. (1993). Actin and fimbrin are required for the internalization step of endocytosis in yeast. *EMBO J.* 12, 2855–2862.
- Kubler, E., Schimmoller, F., and Riezman, H. (1994). Calcium-independent calmodulin requirement for endocytosis in yeast. *EMBO J.* 13, 5539–5546.
- Lee, F.J., Stevens, L.A., Kao, Y.L., Moss, J., and Vaughan, M. (1994). Characterization of a glucose-repressible ADP-ribosylation factor 3 (ARF3) from *Saccharomyces cerevisiae*. *J. Biol. Chem.* 269, 20931–20937.
- Moreau, V., Madania, A., Martin, R.P., and Winson, B. (1996). The *Saccharomyces cerevisiae* actin-related protein Arp2 is involved in the actin cytoskeleton. *J. Cell Biol.* 134, 117–132.
- Moss, J., and Vaughan, M. (1995). Structure and function of ARF proteins: activators of cholera toxin and critical components of intracellular vesicular transport processes. *J. Biol. Chem.* 270, 12327–12330.
- Munn, A.L., Stevenson, B.J., Geli, M.I., and Riezman, H. (1995). *end5*, *end6*, and *end7*: mutations that cause actin delocalization and block the internalization step of endocytosis in *Saccharomyces cerevisiae*. *Mol. Cell Biol.* 15, 1721–1742.
- Payne, G.S., Baker, D., van Tuinen, E., and Schekman, R. (1988). Protein transport to the vacuole and receptor-mediated endocytosis by clathrin heavy chain-deficient yeast. *J. Cell Biol.* 106, 1453–1461.
- Peters, P.J., Hsu, V.W., Ooi, C.E., Finazzi, D., Teal, S.B., Oorschot, V., Donaldson, J.G., and Klausner, R.D. (1995). Overexpression of wild-type and mutant *ARF1* and *ARF6*: distinct perturbations of nonoverlapping membrane compartments. *J. Cell Biol.* 128, 1003–1017.
- Piper, R.C., Cooper, A.A., Yang, H., and Stevens, T.H. (1995). *VPS27* controls vacuolar and endocytic traffic through a prevacuolar compartment in *Saccharomyces cerevisiae*. *J. Cell Biol.* 131, 603–617.
- Poon, P.P., Wang, X., Rotman, M., Huber, I., Cukierman, E., Cassel, D., Singer, R.A., and Johnston, G.C. (1996). *Saccharomyces cerevisiae* Gcs1 is an ADP-ribosylation factor GTPase-activating protein. *Proc. Natl. Acad. Sci. USA* 93, 10074–10077.
- Pryciak, P.M., and Hartwell, L.H. (1996). *AKR1* encodes a candidate effector of the G beta gamma complex in the *Saccharomyces cerevisiae* pheromone response pathway and contributes to control of both cell shape and signal transduction. *Mol. Cell Biol.* 16, 2614–2626.
- Raths, S., Rohrer, J., Crausaz, F., and Riezman, H. (1993). *end3* and *end4*: two mutants defective in receptor-mediated and fluid-phase endocytosis in *Saccharomyces cerevisiae*. *J. Cell Biol.* 120, 55–65.
- Reneke, J.E., Blumer, K.J., Courchesne, W.E., and Thorner, J. (1988). The carboxy-terminal segment of the yeast alpha-factor receptor is a regulatory domain. *Cell* 55, 221–234.
- Rohrer, J.B., Bénédetti, H., Zanolari, B., and Riezman, H. (1993). Identification of a novel sequence mediating regulated endocytosis of the G protein-coupled alpha-pheromone receptor in yeast. *Mol. Cell Biol.* 13, 511–521.
- Roth, A.F., and Davis, N.G. (1996). Ubiquitination of the yeast alpha-factor receptor. *J. Cell Biol.* 134, 661–674.
- Rothstein, R. (1991). Targeting, disruption, replacement, and allele rescue: integrative DNA transformation in yeast, *Methods in Enzymology: Guide to Yeast Genetics and Molecular Biology*, San Diego, CA: Academic Press, 281–301.
- Schandel, K.A., and Jenness, D.D. (1994). Direct evidence for ligand-induced internalization of the yeast alpha-factor pheromone receptor. *Mol. Cell Biol.* 14, 7245–7255.
- Schimmoller, F., and Riezman, H. (1993). Involvement of Ypt7p, a small GTPase, in traffic from late endosome to the vacuole in yeast. *J. Cell Sci.* 106, 823–830.
- Singer-Kruger, B., Stenmark, H., Dusterhoft, A., Philippsen, P., Yoo, J.S., Gallwitz, D., and Zerial, M. (1994). Role of three rab5-like GTPases, Ypt51p, Ypt52p, and Ypt53p, in the endocytic and vacuolar protein sorting pathways of yeast. *J. Cell Biol.* 125, 283–298.
- Singer-Kruger, B., Stenmark, H., and Zerial, M. (1995). Yeast Ypt51p and mammalian Rab5: counterparts with similar function in the early endocytic pathway. *J. Cell Sci.* 108, 3509–3521.
- Sprague, G.F., Jr. (1991). Assay of yeast mating reaction, *Methods in Enzymology: Guide to Yeast Genetics and Molecular Biology*, San Diego, CA: Academic Press, 77–93.
- Stearns, T., Kahn, R.A., Botstein, D., and Hoyt, M.A. (1990a). ADP ribosylation factor is an essential protein in *Saccharomyces cerevisiae* and is encoded by two genes. *Mol. Cell Biol.* 10, 6690–6699.
- Stearns, T., Willingham, M.C., Botstein, D., and Kahn, R.A. (1990b). ADP-ribosylation factor is functionally and physically associated with the Golgi complex. *Proc. Natl. Acad. Sci. USA* 87, 1238–1242.
- Tan, P.K., Davis, N.G., Sprague, G.F., and Payne, G.S. (1993). Clathrin facilitates the internalization of seven transmembrane segment receptors for mating pheromones in yeast. *J. Cell Biol.* 123, 1707–1716.
- Tan, P.K., Howard, J.P., and Payne, G.S. (1996). The sequence NPF_{XD} defines a new class of endocytosis signal in *Saccharomyces cerevisiae*. *J. Cell Biol.* 135, 1789–1800.
- Vida, T.A., and Emr, S.D. (1995). A new vital stain for visualizing vacuolar membrane dynamics and endocytosis in yeast. *J. Cell Biol.* 128, 779–792.
- Wang, X., Hoekstra, M.F., DeMaggio, A.J., Dhillon, N., Vancura, A., Kuret, J., Johnston, G.C., and Singer, R.A. (1996). Prenylated isoforms of yeast casein kinase I, including the novel Yck3p, suppress the *gcs1* blockage of cell proliferation from stationary phase. *Mol. Cell Biol.* 16, 5375–5385.
- Whitney, J.A., Gomez, M., Sheff, D., Kreis, T.E., and Mellman, I. (1995). Cytoplasmic coat proteins involved in endosome function. *Cell* 83, 703–713.
- Wichmann, H., Hengst, L., and Gallwitz, D. (1992). Endocytosis in yeast: evidence for the involvement of a small GTP-binding protein (Ypt7p). *Cell* 71, 1131–1142.
- Zanolari, B., Raths, S., Singer-Kruger, B., and Riezman, H. (1992). Yeast pheromone receptor endocytosis and hyperphosphorylation are independent of G protein-mediated signal transduction. *Cell* 71, 755–763.