Tetracycline Inhibition of Cell-Free Protein Synthesis

II. Effect of the Binding of Tetracycline to the Components of the System

L. E. DAY

Pfizer Medical Research Laboratories, Chas. Pfizer & Co., Inc., Groton, Connecticut

Received for publication 10 May 1966

ABSTRACT

DAY, L. E. (Chas. Pfizer & Co., Inc., Groton, Conn.). Tetracycline inhibition of cell-free protein synthesis. II. Effect of the binding of tetracycline to the components of the system. *J. Bacteriol.* **92**:197-203. 1966.—When tetracycline, an inhibitor of cell-free protein synthesis, was preincubated with each component of the *Escherichia coli* cell-free system, i.e., ribosomes, soluble ribonucleic acid (sRNA), polyuridylic acid (poly U), and S-100 (supernatant enzymes), only the ribosomal-bound antibiotic was inhibitory to the cell-free assay. Experiments designed to further localize the site of inhibition to either the 50S (Svedberg) or the 30S ribosomal subunit were not conclusive. Tritiated tetracycline (7-³H-tetracycline) was bound to isolated 50S ribosomes, and these were recombined with 30S subunits to form 70S ribosomes. When these ribosomes were dissociated and the subunits reisolated, the antibiotic was found with both the 50S and the 30S particles. The same results were observed when the tetracycline was initially bound to the 30S subunit.

The previous report in this series (3) described studies which demonstrated the binding of tetracycline to several of the various nucleic acid components of the *Escherichia coli* cell-free protein-synthesizing system. Although the antibiotic was observed to bind to polyuridylic acid (poly U) and polyadenylic acid (poly A), to soluble ribonucleic acid (sRNA), and to both subunits of 70S (Svedberg) ribosomes, no evidence was given that indicated which, if any, of these binding sites represented the primary locus of inhibition of the cell-free system. Connamacher and Mandel (2) suggested that the antibiotic inhibits protein synthesis by binding directly to the messenger ribonucleic acid (mRNA) and preventing the attachment of sRNA to the mRNA-ribosome complex, whereas Suarez and Nathans (18) postulated that tetracycline binds to the ribosomes, thus obstructing one of the two aminoacyl-sRNA binding sites.

This report contains the results of experiments designed to resolve the question of the locality of the site of inhibition in the cell-free system. This was approached by preincubating each of the components of cell-free assay with tetracycline, followed by extensive dialysis or centrifugation through 10% sucrose to remove unbound antibiotic, and utilization of the component in the poly U-directed C¹⁴-phenylalanine incorpora-

tion assay. The preincubation was carried out with poly U, sRNA, supernatant enzyme fraction (S-100), and ribosomes. Only the ribosomal-bound antibiotic was inhibitory to the cell-free system. The antibiotic was then preincubated with isolated ribosomal subunits, and these were recombined with untreated subunits to form 70S ribosomes with the tetracycline bound to either the 50S or the 30S particle. This was done in an effort to localize the site of inhibition to one or the other subunit. It was observed that the tetracycline was equally as effective as an inhibitor of the cell-free system when bound to the 50S or to the 30S portion of the 70S ribosome. Further experimentation, however, indicated that when 7-³H-tetracycline was initially bound to one species of isolated ribosomal subunits, subsequent manipulations with the other species to form 70S ribosomes resulted in the antibiotic being associated with the both subunits. It is not possible by this technique to resolve the location of the site of inhibition on the ribosome.

MATERIALS AND METHODS

Culture and cultural methods. *E. coli* W was maintained and cultured as before (3).

Preparation of cell-free extracts, ribosomes, and hybrid ribosomes. The preparation of cell-free extracts, ribosomes, and ribosomal subunits has been described

previously (3). Ribosomes [70S (TC)] were prepared by incubating 360 μ g of tetracycline with 18 mg of 70S ribosomes (ribosomal protein) in 0.01 M tris(hydroxymethyl)aminomethane-HCl (Tris), pH 7.3, and 0.01 M $MgCl_2$ at 37 C for 15 min. This mixture was then layered over 10% sucrose in the Tris-HCl and Mg^{++} buffer and was centrifuged at 5×10^4 rev/min for 1.5 hr in the 50 rotor of the Spinco model L2-50 at 3 C. The pellet was homogenized in 0.01 M Tris-HCl (pH 7.3) and 0.01 M $MgCl_2$, and the centrifugation through sucrose was repeated. Control ribosomes were treated in the same manner, except that tetracycline was omitted from the initial reaction mixture.

Isolated 50S ribosomal subunits (19.5 mg of protein) and 30S subunits (16 mg of protein) were incubated with 390 and 320 μ g of tetracycline, respectively, under the conditions described above. Both reaction mixtures were then dialyzed at 5 C against 0.01 M Tris-HCl (pH 7.9), 0.01 M $MgCl_2$, and 0.02 M NH_4Cl for 4 hr, and against fresh buffer for 16 hr. The ribosomal suspensions were then removed from the dialysis tubing, assayed for protein, and recombined with each other and with control subunits to form the 70S ribosomes. Control subunits were treated in the same manner as the tetracycline-treated subunits, except that tetracycline was omitted from the initial reaction mixture. The subunits were recombined to form 70S ribosomes by mixing 2 parts of 50S subunits (based on protein) to 1 part of 30S subunit in the presence of 0.01 M $MgCl_2$. Sucrose density-gradient centrifugation analysis indicated that the treated subunits recombined to form typical 70S ribosomes. The reconstituted ribosomes were designated as follows: 50S(TC)-30S, 70S ribosomes with the tetracycline bound to the 50S subunit; 50S-30S(TC), 70S ribosomes with the tetracycline bound to the 30S subunit; 50S(TC)-30S(TC), 70S ribosomes with tetracycline bound to both subunits; and 50S-30S, reconstituted 70S control ribosomes with no bound tetracycline.

Preparation of poly U, sRNA, and S-100 preincubated with tetracycline. The following reaction mixtures were incubated at 37 C for 10 min: (i) poly U, 2.0 mg; tetracycline HCl, 1.0 mg, in 5.0 ml of 0.01 M Tris-HCl (pH 7.3) and 0.01 M $MgCl_2$; (ii) sRNA (21), 10.0 mg; tetracycline HCl, 133 μ g, in 1.0 ml of 0.01 M Tris-HCl (pH 7.3) and 0.01 M $MgCl_2$; (iii) S-100 protein (19), 23.3 mg; and tetracycline HCl, 233 μ g in a final volume of 1.0 ml. A control for each reaction mixture was treated in an identical manner except that the tetracycline was omitted. Mixtures (i) and (ii) and controls were dialyzed at 3 C against three changes of 0.01 M Tris-HCl and 0.01 M $MgCl_2$ for 8, 16, and 8 hr, respectively. Mixture (iii) and its control were dialyzed twice against 1 liter of 0.02 M Tris-HCl (pH 7.9), 0.01 M $MgCl_2$, and 0.005 M mercaptoethylamine for 16 and 4 hr, respectively. The dialyzed mixtures were removed and frozen in liquid nitrogen until used. The tetracycline-treated components were designated S-100(TC), sRNA(TC), and poly U(TC).

Poly U-directed C^{14} -phenylalanine incorporation studies. The method of Szer and Ochoa (19) was used for experiments involving cell-free polymerization of phenylalanine with use of poly U as the synthetic

mRNA. The reaction mixtures contained 0.5 mg of ribosomal protein, 1.0 mg of S-100 protein, and 0.75 mg of *E. coli* sRNA in a final volume of 250 μ liters. The radioactivity of the C^{14} -phenylalanine incorporated was assayed by the method of Nirenberg (13).

Binding of C^{14} -phenylalanyl-sRNA by ribosomes. C^{14} -phenylalanyl-sRNA was prepared as described by Nirenberg, Matthaei, and Jones (15) and had a final specific activity of 4,650 count/min per OD_{260} . The binding of C^{14} -phenylalanyl-sRNA to ribosomes in the presence of poly U was achieved by use of the method of Nirenberg and Leder (14). These reaction mixtures contained, in a final volume of 100 μ liters, the following: 50 $m\mu$ moles of poly U, 2.0 OD_{260} C^{14} -phenylalanyl-sRNA, 4.0 OD_{260} ribosomes, 10 μ moles of Tris-HCl (pH 7.2), 2 μ moles of Mg acetate, and 5 μ moles of KCl. The radioactivity was assayed in scintillation vials (14).

The sedimentation profile of ribosomal-bound, C^{14} -phenylalanyl-sRNA was established by use of density gradient centrifugation. The reaction mixtures contained: 1.0 mg of ribosomes, 4.0 OD_{260} C^{14} -phenylalanyl-sRNA, 5.0 μ g of poly U in 200 μ liters of 0.01 M Tris-HCl (pH 7.2) containing 0.01 M $MgCl_2$ and 0.16 M NH_4Cl . The control contained 70S ribosomes, and the test reaction contained 70S(TC) ribosomes. Both mixtures were incubated at 25 C for 10 min, and then analyzed by sucrose density gradient centrifugation as described (3).

Distribution of tetracycline in ribosomal subunits after initial binding to either 50S or to 30S particles. The initial reaction mixture contained: 22 mg of 50S ribosomes and 16 μ g of 7- H^3 -tetracycline (specific activity of 80 mc/mmole) in 500 μ liters of 0.01 M Tris-HCl (pH 7.3), and 0.01 M $MgCl_2$. The mixture was incubated at 37 C for 30 min. The reaction mixture was then dialyzed at 3 C for two intervals of 24 hr each against 4 liters of 0.01 M Tris-HCl (pH 7.3) containing 0.01 M $MgCl_2$. A 100- μ liter volume of the dialyzed suspension was removed and layered over a 4.6-ml sucrose density gradient (5 to 20%) containing Tris and Mg^{++} buffer at the above concentrations. The gradient was centrifuged at 3.5×10^4 rev/min for 2 hr in the SW-39 rotor. The gradient was fractionated and assayed as described (3). The remainder of the dialyzed reaction mixture was then incubated at 37 C for 30 min with 20 mg of 30S ribosomal subunits in a final volume of 1.0 ml of 0.01 M Tris-HCl (pH 7.3) buffer containing 0.01 M $MgCl_2$. A 100- μ liter sample of this reaction mixture was removed and analyzed by density gradient centrifugation as described above. The remainder of the reaction mixture was then dialyzed for two periods of 6 hr each at 3 C against 2 liters of 0.01 M Tris-HCl buffer (pH 7.3) containing 5×10^{-4} M $MgCl_2$ to dissociate the ribosomal subunits. The entire reaction mixture was layered over a 54-ml sucrose gradient (5 to 20%) in 0.01 M Tris buffer containing 5×10^{-4} M $MgCl_2$ and centrifuged at 2.5×10^4 rev/min for 10 hr in the SW-25.2 rotor of the L2-50 ultracentrifuge. This gradient was fractionated into 1.0-ml fractions, and 50 μ liters was removed from each fraction for OD_{260} assay and 50 μ liters was removed for radioactivity determinations as described (3). Those fractions con-

taining the 50S subunits were pooled (Fig. 2C) as were those containing 30S subunits. These pools were centrifuged in the SW-39 rotor overnight (19 hr) at 3.7×10^4 rev/min. The pellets were taken up in 100 μ liters of Tris buffer containing 5×10^{-4} M Mg^{++} , layered over 4.6-ml sucrose gradients (5 to 20%) in the same buffer, and centrifuged at 3.5×10^4 rev/min. The gradient containing the 50S particles was centrifuged for 2 hr; the gradient containing the 30S particles, for 4 hr. The gradients were assayed for OD₂₆₀ and radioactivity (3).

The same experiment as described in the previous paragraph was repeated, except that the initial binding reaction was carried out with 30S ribosomal subunits and 7-H³-tetracycline. The only variation in procedure was that the initial series of dialyses of the 30S-tetracycline reaction mixture was carried out for a total of 46 hr, the dialyzing buffer being changed after 6, 16, and 8 hr.

Protein assay. The method of Lowry et al. (12) was used for estimating the protein content of ribosomal and S-100 suspensions.

Materials. Poly U and tetracycline were the same as used previously (3). Uniformly labeled C¹⁴-phenylalanine with a specific activity of 300 mc/mmole and 7-H³-tetracycline with a specific activity of 80 mc/mmole were purchased from New England Nuclear Corp., Boston, Mass.

RESULTS

The effects of the tetracycline antibiotics on the cell-free incorporation of amino acids into peptides has been documented by others (4, 11, 16), and it has been observed in this laboratory that the amount of tetracycline required to inhibit the reaction by 50% is about 20 μ g/mg of ribosomal protein (*unpublished data*). This concentration varies somewhat depending on the activity of the extract being used, the more active preparations being more sensitive to the antibiotic. To standardize the amount of tetracycline in the reaction mixtures being assayed, the antibiotic was preincubated with each component at the ratio that would normally result in 50% inhibition in the cell-free assay. The results tabulated in Table 1 indicate that only ribosomes which have been pretreated with the antibiotic are inhibitory to the cell-free incorporation of amino acids into peptides, even though it has been observed that the tetracycline binds to the sRNA and to the poly U (3). The preincubation of sRNA both with and without the antibiotic results in a large decrease in the amount of the radioactive label incorporated. We also observed this when the sRNA and its appropriate control were preincubated with tetracycline, and the polynucleotide and unbound antibiotic were separated by gel filtration through Sephadex G-25 (*unpublished data*). The reason for this

TABLE 1. *Effect of preincubation of the individual components of the Escherichia coli cell-free system with tetracycline on the poly U-directed incorporation of C¹⁴-phenylalanine into polypeptide*

Preincubated component	Phenylalanine incorporated ^a	Inhibition
	<i>μmoles</i>	%
S-100 control ^b	17.2	
S-100(TC).....	17.3	0
sRNA control.....	2.8	
sRNA(TC).....	3.4	0
Poly U control.....	11.9	
Poly U(TC).....	11.3	5
70S ribosomes.....	12.2	
70S(TC) ribosomes.....	6.1	50

^a The values tabulated are the average of duplicate assays.

^b Control components were preincubated and recovered in the same manner as the tetracycline-treated component except tetracycline was not present in the control reaction mixture.

loss of stimulatory activity of the sRNA is not apparent at this time.

Since the data thus far obtained indicated that the ribosome was the site of inhibition by tetracycline, subsequent studies were directed toward localization of this sensitive site on either one or the other of the ribosomal subunits. Prior data (3) demonstrated that the antibiotic was bound by both the 50S and the 30S subunit; therefore hybrid 70S ribosomes were prepared as described under Materials and Methods, and utilized in the poly U-C¹⁴-phenylalanine assay. These results (Table 2) do not indicate that either subunit possesses the sensitive site of tetracycline inhibition, but rather that tetracycline bound to either unit is equally effective as an inhibitor of this reaction. When the antibiotic is bound to both subunits, the level of inhibition is higher but is not additive. It is also of interest to note that the control ribosomes prepared in this manner are more sensitive to the effects of the antibiotic, since the level that normally results in 50% inhibition gives about 80% with these ribosomes.

The rapid method of Nirenberg and Leder (14) for assaying the binding of aminoacyl-sRNA to ribosomes in the presence of mRNA was used as a technique to determine if ribosomal-bound tetracycline is as effective as an inhibitor of this reaction as tetracycline added directly to the reaction mixture. It was also used as an assay

method, because this reaction does not require an energy-generating system, as is necessary in the cell-free peptide synthetic reaction. Table 3 indicates that the levels of inhibition with the hybrid ribosomes is about the same as with the control ribosomes to which the inhibitor was added at the time the reaction was initiated. The results with this assay also suggest that the binding of tetracycline to either one or the other subunit is an event which is deleterious to ribosomal function, and the sensitivity of these structures to tetracycline is a characteristic of both. It should be pointed out that 2 μ g of tetracycline added to the 50S-30S control ribosomes (equiva-

lent to 20 μ g of tetracycline per mg of ribosomal protein) inhibits this reaction by 50%, the level of inhibition observed when this concentration of antibiotic is added to the poly U- C^{14} -phenylalanine system. The separation of ribosomal subunits, preincubation, and recombination to form 70S ribosomes resulted in only a slight loss in activity (about 12%) as evidenced by the comparison of the amount of radioactivity bound by the 70S ribosomes (untreated control) and by the 50S-30S ribosomes.

TABLE 2. *Effect of tetracycline bound to the 50S subunit or to the 30S subunit of Escherichia coli 70S ribosomes on the poly U-directed incorporation of C^{14} -phenylalanine into polypeptide*

Ribosomes used	Phenylalanine incorporated ^a	Inhibition
	μ moles	%
50S-30S control.....	2.9	
50S-30S control + 10 μ g of tetracycline.....	.6	79
50S(TC)-30S(TC).....	.9	69
50S(TC)-30S.....	1.2	59
50S-30S(TC).....	1.4	52

^a The values tabulated are the average of duplicate assays.

TABLE 3. *Effect of tetracycline bound to the 50S subunit or to the 30S subunit of Escherichia coli ribosomes on the poly U-directed binding of C^{14} -phenylalanyl-sRNA to these ribosomes*

Ribosomes used	C^{14} -phenylalanyl-sRNA bound ^a	Inhibition
	count/min	%
70S untreated control.....	1,050	
50S-30S control.....	896	
50S-30S control + 2 μ g of tetracycline.....	461	49 ^b
50S(TC)-30S(TC).....	413	54
50S(TC)-30S.....	384	57
50S-30S(TC).....	429	52

^a The values tabulated are the average of duplicate assays.

^b The per cent inhibition was calculated with the 50S-30S control as 100% incorporation.

FIG. 1. *Distribution of radioactivity in reconstituted ribosomes and ribosomal subunits after initial binding of 7- H^3 -tetracycline to 50S subunits. Sedimentation profile of (A) 50S subunits after incubation with 7- H^3 -tetracycline, (B) reconstituted 70S ribosomes after addition of unlabeled 30S subunits to 50S-7- H^3 -tetracycline reaction mixture, (C) reconstituted 70S ribosomes after dissociation to 50S and 30S subunits, (D) 50S subunits isolated from dissociated 70S ribosomes [fractions 18 to 23, (C)], and (E) 30S subunits isolated from dissociated 70S ribosomes [fractions 27 to 35, (C)]. Solid line, OD_{260} ; broken line, radioactivity.*

FIG. 2. Distribution of radioactivity in reconstituted ribosomes and ribosomal subunits after initial binding of 7-³H-tetracycline to 30S subunits. Sedimentation profile of (A) 30S subunits after incubation with 7-³H-tetracycline, (B) reconstituted 70S ribosomes after addition of unlabeled 50S subunits to 30S-7-³H-tetracycline reaction mixture, (C) reconstituted 70S ribosomes after dissociation to 50S and 30S subunits, (D) 50S subunits isolated from dissociated 70S ribosomes [fractions 20 to 24, (C)], and (E) 30S subunits isolated from dissociated 70S ribosomes [fractions 28 to 36, (C)]. Solid line, OD₂₆₀; broken line, radioactivity.

FIG. 3. Binding of C¹⁴-phenylalanyl-sRNA to *Escherichia coli* ribosomes. (A) 70S control ribosomes, (B) 70S (TC) ribosomes. Solid line, OD₂₆₀; broken line, radioactivity.

Since the preparation of ribosomes with the antibiotics bound to either one or the other of the ribosomal subunits did not demonstrate a unique site of inhibition on either one or the other subunit, the suggestion was made that the tetracycline might not remain with the subunit to which it was initially bound during subsequent manipulations and reconstitution of the 70S ribosome. Investigation of this possibility was carried out by binding labeled tetracycline to 50S subunits, adding excess unlabeled 30S subunits in 10^{-2} M Mg^{++} to reconstitute the 70S ribosomes, and then reisolating the individual subunits to determine the distribution of the

radioactive antibiotic. The results of this study (Fig. 1) indicate that the antibiotic is bound by both subunits during these manipulations. The initial binding reaction was carried out in 10^{-2} M Mg^{++} because this is the concentration required during the cell-free assay with poly U and phenylalanine, although previous studies (3) indicated more antibiotic is bound by individual subunits at 5×10^{-4} M Mg^{++} . When the initial binding was carried out with the labeled antibiotic and the 30S particles (Fig. 2), essentially the same final distribution of the antibiotic was observed in both subunits. Whether the antibiotic is being dissociated from the subunit to which it was initially bound, making it available for binding to the other subunit at the time the 70S ribosomes are reformed, or whether excess, unbound antibiotic not removed by extensive dialysis (Fig. 1A and 2A) is being bound by the other subunit, is not clear. The latter seems to be the more likely possibility.

The inhibition of binding of aminoacyl-sRNA to ribosomes by tetracycline antibiotics has been previously reported; Suarez and Nathans (18), using the technique of Nirenberg and Leder (14) and Hierowski (7), established the sedimentation profile in a sucrose-density gradient. In both instances, the antibiotic was added directly to the reaction mixture. Figure 3 is the result of a similar experiment with density-gradient centrifugation; however, the tetracycline was preincubated with the ribosomes, the ribosomes washed twice to remove unbound

antibiotic and then used to bind the C^{14} -phenylalanyl-sRNA. The 70S(TC) ribosomes bind less phenylalanyl-sRNA (Fig. 3B) than do the control ribosomes (Fig. 3A).

DISCUSSION

The recent literature (1, 8, 9, 17) indicates that the tetracycline antibiotics function by several different modes of action in the growing cell, depending on the concentration of the antibiotic, i.e., bacteriostatic or bactericidal, conditions of aeration of the culture, pH of the medium, etc. A series of careful studies by Morrison (1, 8, 9) demonstrated that three modes of action may result in the inhibition of growth of *Aerobacter aerogenes*. One of these mechanisms was the derangement of protein synthesis; however, it was suggested that all three modes of action were the result of inhibition of hydrogen-transfer reactions. The observation that ribosomal-bound tetracycline is inhibitory to the binding of aminoacyl-sRNA to ribosomes does not lend support to the view that the interference with protein synthesis is the secondary result of inhibition of hydrogen transfer or other energy-generating reactions (1, 17). The binding of the aminoacyl-sRNA by ribosomes is not energy-dependent, and the interference with this binding by tetracycline suggests that the antibiotic occupies a site on the ribosome which obstructs the attachment of the aminoacyl-sRNA, as previously pointed out by Suarez and Nathans (18).

Since extensive dialysis of the reaction mixtures described here did not completely remove excess unbound tetracycline (Fig. 1A and 2A), the possibility exists that the inhibition observed with the reconstituted ribosomes (Tables 2 and 3) might actually result from this unbound antibiotic acting at some other point. This does not appear to be likely, since the same dialysis technique was used after preincubation of tetracycline with the S-100 fraction, poly U, and the sRNA, and no inhibition was observed in these instances, even though unbound tetracycline was undoubtedly present after dialysis. Another pertinent point is that the 70S(TC) ribosomes were inhibitory to both the cell-free incorporation of phenylalanine into peptides and to the binding of C^{14} -phenylalanyl-sRNA to the ribosomes. These ribosomes were preincubated with the antibiotic and then washed twice through 10% sucrose. It does not seem likely that the excess unbound antibiotic would sediment in this solution and be available as free tetracycline when the ribosomal pellet was resuspended in buffer. On this basis it is felt that ribosomal binding of the antibiotic is the inhibitory event in the cell-free protein synthetic system.

Calculations based on the previous results (3) indicated that about 1 molecule of tetracycline was bound to each ribosomal subunit when the separated subunits were incubated with the antibiotic in 5×10^{-4} M Mg^{++} . Similar calculations based on the data shown in Fig. 1D and 1E and Fig. 2D and 2E demonstrate the binding of the antibiotic to the individual subunits to be significantly less than this after extensive dialysis, centrifugations, and other manipulations. This may be attributed to the fact that the initial binding was carried out in 10^{-2} M Mg^{++} , and that these manipulations may have removed some of the bound antibiotic. About three times as much tetracycline was bound by the 30S subunit as by the 50S, even when the initial binding reaction was carried out with the 50S particles. It is beyond the scope of this paper to consider possible mechanisms of inhibition other than those concerned with protein synthesis, but it is reasonable to believe that the tetracyclines have many activities in growing cells in addition to the interference with protein synthetic reactions. The chelating properties of tetracycline undoubtedly result in its association with many macromolecules of the bacterial cell through metal complexes (10), and it is therefore difficult to assign to this group of antibiotics a primary locus of inhibition in the intact cell. The observation that tetracycline binds to both subunits of *E. coli* ribosomes (3) and that this ribosomal-bound tetracycline is inhibitory to cell-free protein synthetic reactions and to the binding of aminoacyl-sRNA to these ribosomes is evidence that the primary site of inhibition by tetracycline in the cell-free system is the ribosome, although the location of the inhibitory site has not been further restricted to either the 50S or to the 30S subunit by the technique used. Yokota and Akiba (20) reported the isolation of a tetracycline-resistant mutant which was also resistant to the antibiotic in cell-free protein synthetic assays. By use of the supernatant enzymes from a sensitive strain of *E. coli* and the ribosomes from the resistant culture, it was determined that the site of resistance was on the ribosome. This type of mutant can be further utilized to localize the resistance on the ribosomal subunits, thereby resolving the site of inhibition to one subunit or the other.

ACKNOWLEDGMENT

I thank Ernest Grant for his excellent technical assistance.

LITERATURE CITED

1. BENBOUGH, J., AND G. A. MORRISON. 1965. Bacteriostatic actions of some tetracyclines. *J. Pharm. Pharmacol.* 17:409-422.

2. CONNAMACHER, R. H., AND H. G. MANDEL. 1965. Binding of tetracycline to the 30S ribosomes and to polyuridylic acid. *Biochem. Biophys. Res. Commun.* **20**:98-103.
3. DAY, L. E. 1966. Tetracycline inhibition of cell-free protein synthesis. I. Binding of tetracycline to components of the system. *J. Bacteriol.* **91**: 1917-1923.
4. FRANKLIN, T. J. 1963. The inhibition of incorporation of leucine into protein of cell-free systems from rat liver and *Escherichia coli* by chlortetracycline. *Biochem. J.* **87**:449-453.
5. GALE, E. F., AND J. P. FOLKES. 1963. The assimilation of amino acids by bacteria. *Biochem. J.* **53**:493-498.
6. HASH, J. H., M. WISHNICK, AND P. A. MILLER. 1964. On the mode of action of the tetracycline antibiotics in *Staphylococcus aureus*. *J. Biol. Chem.* **239**:2070-2078.
7. HIEROWSKI, M. 1965. Inhibition of protein synthesis by chlortetracycline in the *E. coli* in vitro system. *Proc. Natl. Acad. Sci. U.S.* **53**:594-599.
8. JONES, J. G., AND G. A. MORRISON. 1962. The bacteriostatic actions of tetracycline and oxytetracycline. *J. Pharm. Pharmacol.* **14**:808-824.
9. JONES, J. G., AND G. A. MORRISON. 1963. Inhibition by tetracycline and oxytetracycline of the consumption of pyruvate by *Aerobacter aerogenes*. *J. Pharm. Pharmacol.* **15**:34-44.
10. KOHN, K. W. 1961. Mediation of divalent metal ions in the binding of tetracycline to macromolecules. *Nature* **191**:1156-1158.
11. LASKIN, A. I., AND W. M. CHAN. 1964. Inhibition by tetracyclines of polyuridylic acid directed phenylalanine incorporation in *Escherichia coli* cell-free systems. *Biochem. Biophys. Res. Commun.* **14**:137-142.
12. LOWRY, O. H., N. J. ROSEBROUGH, A. L. FARR, AND R. J. RANDALL. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**:265-275.
13. NIRENBERG, M. W. 1964. Cell-free protein synthesis directed by messenger RNA, p. 17-23. In S. P. Colowick and N. O. Kaplan [ed.], *Methods in enzymology*, vol. 6. Academic Press, Inc., New York.
14. NIRENBERG, M. W., AND P. LEDER. 1964. RNA codewords and protein synthesis. *Science* **145**: 1399-1407.
15. NIRENBERG, M. W., J. H. MATTHAEI, AND O. W. JONES. 1962. An intermediate in the biosynthesis of polyphenylalanine directed by synthetic template RNA. *Proc. Natl. Acad. Sci. U.S.* **48**:104-109.
16. RENDI, R., AND S. OCHOA. 1962. Effect of chloramphenicol on protein synthesis in cell-free preparations of *Escherichia coli*. *J. Biol. Chem.* **237**:3711-3713.
17. SHAPOSHNIKOV, V. N., AND V. K. BAKULEV. 1964. Interrelations between biosynthesis of proteins, of ribonucleic acid, and of certain coenzymes in the action of tetracyclines on *Staphylococcus aureus* 209P. *Dokl. Akad. Nauk Uz.SSR* **158**: 987-989.
18. SUAREZ, G., AND D. NATHANS. 1965. Inhibition of aminoacyl-sRNA binding to ribosomes by tetracycline. *Biochem. Biophys. Res. Commun.* **18**:743-750.
19. SZER, W., AND S. OCHOA. 1964. Complexing ability and coding properties of synthetic polynucleotides. *J. Mol. Biol.* **8**:823-834.
20. YOKOTA, T., AND T. AKIBA. 1962. Studies on the mechanism of transfer of drug-resistance in bacteria. 23. Mechanisms of the antibacterial action of tetracycline and the tetracycline resistance in the artificial, TC-resistant strain of *E. coli*. *Med. Biol. (Tokyo)* **64**:39-44.
21. ZUBAY, G. 1962. The isolation and fractionation of soluble ribonucleic acid. *J. Mol. Biol.* **4**: 347-356.