Published in final edited form as:

Pharmacol Biochem Behav. 2008 October ; 90(4): 614–624. doi:10.1016/j.pbb.2008.05.004.

Fetal Alcohol Spectrum Disorder-associated depression: evidence for reductions in the levels of brain-derived neurotrophic factor in a mouse model

Kevin K. Caldwell^a, S. Sheema^a, Rodrigo D Paz^{a,b}, Sabrina L. Samudio-Ruiz^a, Mary H. Laughlin^c, Nathan E. Spence^c, Michael J Roehl^c, Sara N. Alcon^a, and Andrea M. Allan^a

^a Department of Neurosciences, School of Medicine, University of New Mexico, Albuquerque, New Mexico, 87131, USA

^b Departamento de Psiquiatría y Neurociencias, Universidad Diego Portales, Santiago de Chile

^c School of Medicine, University of New Mexico, Albuquerque, New Mexico, 87131, USA

Abstract

Prenatal ethanol exposure is associated with an increased incidence of depressive disorders in patient populations. However, the mechanisms that link prenatal ethanol exposure and depression are unknown. Several recent studies have implicated reduced brain-derived neurotrophic factor (BDNF) levels in the hippocampal formation and frontal cortex as important contributors to the etiology of depression. In the present studies, we sought to determine whether prenatal ethanol exposure is associated with behaviors that model depression, as well as with reduced BDNF levels in the hippocampal formation and/or medial frontal cortex, in a mouse model of fetal alcohol spectrum disorder (FASD). Compared to control adult mice, prenatal ethanol-exposed adult mice displayed increased learned helplessness behavior and increased immobility in the Porsolt forced swim test. Prenatal ethanol exposure was associated with decreased BDNF protein levels in the medial frontal cortex, but not the hippocampal formation, while total BDNF mRNA and BDNF transcripts containing exon III, IV or VI were reduced in both the medial frontal cortex and the hippocampal formation of prenatal ethanol-exposed mice. These results identify reduced BDNF levels in the medial frontal cortex and hippocampal formation as potential mediators of depressive disorders associated with FASD.

Keywords

BDNF; hippocampus; frontal cortex; prenatal; ethanol; depression

INTRODUCTION

Prenatal ethanol exposure has been shown to be associated with a spectrum of neurochemical, physiologic, and behavioral dysfunctions that affect various aspects of an individual's life. The range of clinical abnormalities that are associated with prenatal ethanol exposure is termed

Corresponding author: Kevin K. Caldwell, Ph.D., Department of Neurosciences, MSC 08 4740, 1 University of New Mexico, Albuquerque, NM 87131, KCaldwell@salud.unm.edu, telephone: 505-272-0033; fax: 505-272-8082.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

fetal alcohol spectrum disorder (FASD). The incidence of FASD is estimated to be as high as 1 in 100 births (Sampson et al., 1997). Although the most common manifestation of FASD is a reduction in cognitive abilities (Sampson et al., 1997; Streissguth and O'Malley, 2000), FASD is also comorbid with various psychiatric illnesses, including depression (Famy et al., 1998; O'Connor and Kasari, 2000; O'Connor et al., 2002; O'Connor and Paley, 2006; Roebuck et al., 1999). Several genetic and biochemical abnormalities have been shown to be associated with depression (Berton and Nestler, 2006; Duman and Monteggia, 2006; Hasler et al., 2007; Jans et al., 2007; Levinson, 2006); however, none of these has been implicated as a mechanism linking prenatal ethanol exposure and depressive disorders. Therefore, we sought to determine whether depressive-like behaviors are exhibited by a mouse model of FASD and, if so, to investigate the relationship between these behaviors and a mechanism that has been proposed to contribute to depressive behavior, reductions in hippocampal formation and medial frontal cortical brain- derived neurotrophic factor (BDNF) levels.

Several lines of evidence indicate that reduced BDNF levels in the prefrontal cortex, hippocampus, or both play an important role in depression. BDNF expression is diminished in postmortem hippocampus and prefrontal cortex collected from suicide cases with untreated depression compared to patients who were being treated with antidepressants at the time of suicide (Chen et al., 2001; Karege et al., 2005). Experimental paradigms, such as learned helplessness and the forced swim task, which induce behaviors modeling depression in laboratory animals, have been shown to cause reductions in BDNF protein and mRNA levels in the frontal cortex and/or hippocampus (Itoh et al., 2004; Russo-Neustadt et al., 2001; Song et al., 2006; Takeda et al., 2006). Additionally, elevated BDNF protein levels have been shown to block depressive-like behaviors in these same paradigms (Shirayama et al., 2002; Siuciak et al., 1997). Finally, long-term antidepressant therapy has been shown to increase BDNF levels in the hippocampus (Nibuya et al., 1995; Okamoto et al., 2003; Dwivedi et al., 2006) and frontal cortex (Altar et al., 2003; Okamoto et al., 2003). These results, combined with the observation that therapeutic responses to antidepressants generally require chronic administration of the drugs, have led to the "neurotrophic hypothesis" of depression (reviewed in Duman and Monteggia, 2006).

Prenatal exposure to ethanol has been reported to either decrease or have no effect on BDNF protein levels measured in the postnatal rat brain. Heaton et al. (2000) reported that prenatal ethanol exposure did not affect BDNF protein levels measured in hippocampal formation or the cortex/striatum on postnatal day 1. Feng et al. (2005) found that prenatal ethanol exposure reduced BDNF protein levels in postnatal day 7 and 8 cortex and hippocampal formation. Similarly, Climent et al. (2002) reported that prenatal ethanol exposure reduced BDNF protein levels in the cortex, when measured on postnatal days 5–21, while pups were being nursed by ethanol-consuming moms; however, levels were not different from controls on postnatal day 35 after weaning. We are unaware of any published studies on the effects of prenatal ethanol exposure on BDNF levels in the brains of adult rodents.

The studies described herein demonstrate that adult mice that were exposed to ethanol throughout gestation display depressive-like behaviors, manifested as augmented learned helplessness and immobility in the forced swim test. In addition, we found that prenatal exposure to ethanol was associated with reduced levels of BDNF protein, as well as mRNA, in the frontal cortex and reduced BDNF mRNA levels in the hippocampal formation. These studies demonstrate that a mouse model of FASD displays depressive-like behaviors that share similarities with clinical depression exhibited by diagnosed FASD patients and, thus, should serve as a useful tool for the identification of neurochemical alterations that underlie this disorder.

MATERIALS AND METHODS

Animals

All animal procedures employed in these studies were approved by the University of New Mexico Institutional Animal Care and Use Committee. To reduce possible litter effects, only one mouse was used from each litter for each of the assays conducted, thus the unit of analysis was litter. The offspring from 19 ethanol-consuming and 16 saccharin-consuming dams were used in these studies.

Prenatal Ethanol Exposure Paradigm

The procedures employed for prenatal ethanol exposure were essentially as described in Allan et al. (2003). Briefly, female C57BL/6J mice (The Jackson Laboratory; Bar Harbor, ME) were offered 22 hr free access to either 0.066% saccharin or water for 1 week. Following this period, ethanol was introduced to the saccharin tube for the ethanol group, while the control group continued to have access to saccharin alone. The ethanol concentration was increased from 0% to 5% in a stepwise fashion over five days. After two weeks of stable drinking, females were placed on breeder chow and a male was introduced into the female's cage. Once the female was determined to be pregnant, the male was removed, nesting material was added to the cage, and the female continued to consume ethanol or saccharin. Ethanol consumption was measured every other day and the g ethanol consumed/kg body weight/day were calculated for each dam. Within one day of birth, the ethanol and the saccharin concentrations were reduced by one-half every two days until the mice were consuming water postpartum. We have found that this procedure yields maternal care behaviors that are indistinguishable between saccharin- and ethanol-consuming moms (Allan et al., 2003) and, thus, avoids the introduction of the confound of poor maternal care that can occur with postpartum maternal consumption of ethanol or sudden ethanol withdrawal. Pups were weaned at 23–25 days and maintained in same-sex and litter-mate cages, 3–4 mice per cage, with free access to water and chow until adulthood.

Measurement of Blood Ethanol Concentrations

Maternal blood ethanol concentrations were determined as previously described (Allan et al., 2003). Blood samples were collected from the saphenous vein of mice that were approximately 14 days pregnant during the middle of the drinking period. These measures were performed on a separate group of dams that were drinking at a rate similar to that of the dams whose offspring were used in the present studies.

Behavioral Testing Procedures

Mice were maintained on a 12 hr light: 12 hr dark cycle (lights on 7:00 AM). Behavioral testing was conducted between 9 AM and 3 PM. All studies were performed using female mice between 60 and 90 days of age at the time of testing. Animals were used in only one behavioral test.

Learned helplessness

The apparatus for the learned helplessness task consisted of a CoulbournTM Habitest[©] shuttlebox with a stainless steel grid floor for administration of the footshock. The apparatus was located within a sound-attenuated chamber. 70% isopropanol was used to clean the walls and floor after the removal of each mouse from the shuttlebox.

The method for assessing learned helplessness was modified from Caldarone et al. (2004). On Day 1, animals received 20 uncontrollable and unpredictable footshocks (0.5 mA, 2 second duration); the probability for delivery of the footshock was assigned at 0.5 every 15 seconds. The animal was removed 30 seconds after the delivery of the last shock and returned to its

home cage. We note that, although many published procedures for the production of learned helplessness employ more than 100 inescapable footshocks to an animal (Caldarone et al., 2000; Caldarone et al., 2004; MacQueen et al., 2001; Newton et al., 2002; Ridder et al., 2005; Schulte-Herbrüggen et al., 2006), we were able to identify increased levels of learned helplessness in prenatal ethanol-exposed mice with the use of this milder procedure. On the testing day (Day 2), animals experienced 28 test trials with an intertrial interval of 30 seconds. For the first five trials, a footshock (0.5 mA) and auditory cue (80 dB, 6 Hz clicker) both began at the start of the trial; after a one second delay, the guillotine door separating the two chambers of the shuttlebox opened. The last 23 trials initiated with the tone and the door rising, and the onset of the footshock was delayed for three seconds. Sensors in each compartment of the shuttlebox turned off the footshock and tone and closed the guillotine door, if the animal escaped through the raised door. If an escape was not made within 24 seconds from the initiation of the trial, the shock and tone terminated and the door closed. The subject's response in each trial was scored as an "escape", if it fled through the door to the non-shock side of the chamber within the 24 sec test session, or as a "failure to escape", if the animal remained in the side of the apparatus in which the shock was being delivered. In addition, latency for the mouse to escape through the door was measured as the time from the door rising to the time it closed.

Forced swim test

Mice were placed in a 30 cm diameter, 46 cm tall cylinder of water (22–25°C, depth 26 cm) for three minutes, as described previously (Porsolt et al., 1977; Sunal et al., 1994). Six behaviors were assessed every five seconds during the last two minutes of the session, as described by Schramm et al. (2001): floating, twitching, kicking, swimming, climbing, and thrashing. The combination of floating, twitching and kicking behaviors was used as an index of immobility, with the remaining behaviors representing escape-directed behavior.

Basal Open Field Activity

Mice were placed in a dim red light illuminated open field apparatus (102.5 × 102.5 × 47.5 cm; Opto-Varimex, Columbus Instruments, Columbus, OH) for 15 minutes on two consecutive days. The first day was used for acclimation to the task. The open field apparatus was equipped with a series of photo-beams (15 × 15). Open field locomotor (horizontal) activity was measured as the number of photo beam interruptions recorded on the second day.

General information related to animals used in the biochemical and molecular studies

Mice were maintained on the same light:dark cycle and were of the same age range as used for the behavioral studies. All studies were performed using female mice that were euthanized between 9 AM and 3 PM. Additionally, none of the animals used in the behavioral studies were used for the biochemical and molecular analyses, which were performed on naïve mice.

Determination of BDNF protein levels

BDNF protein levels were determined in 96-well ELISA plates (Chemicon, Temecula, CA), as described by the manufacturer.

Determination of proBDNF levels

The hippocampal formation and medial frontal cortex were isolated, then homogenized in 0.5 mL ice-cold homogenization buffer (Buckley et al., 2004) and centrifuged (1,000 $\times$ g_{max}, 6 min, 4 °C). The supernatant was decanted and the pellet resuspended, homogenized and centrifuged as before. The supernatant from the second centrifugation step was combined with the initial supernatant and frozen in aliquots at –80 °C until use. Samples (30 µg medial frontal cortex; 70 µg hippocampal formation) were thawed, diluted in 4X SDS-PAGE Sample Buffer containing NuPAGE Reducing Agent (Invitrogen Corporation, Carlsbad, CA) and separated

on 12% NuPAGE Bis-Tris Polyacrylamide Gels (Invitrogen Corporation) using MOPS SDS Running Buffer containing NuPAGE Antioxidant (Invitrogen Corporation). Proteins were transferred to PVDF (Bio-Rad Laboratories, Hercules, CA) membranes in NuPAGE Transfer Buffer (Invitrogen Corporation) containing NuPAGE Antioxidant and methanol. The membranes were blocked with 3% (w/v) BSA in TBST [150 mM NaCl, 20 mM Tris (pH 7.5), 0.05% (v/v) Tween 20] for 60 minutes at room temperature, then incubated overnight with affinity purified anti-proBDNF rabbit antibody (Chemicon #AB5613P; 0.25 µg/mL) in BSA/TBST at 4 °C. The membrane was washed 4 × 5 min with TBST and incubated with HRP-conjugated secondary antibody (HRP-Goat-Anti-Mouse-Anti-IgG (H + L); Jackson ImmunoResearch) in BSA/TBST for 90 minutes at room temp. The membrane was washed 4 × 5 min with TBST and immunoreactivity detected using a Supersignal West Pico Chemiluminescent Kit (Pierce Biotechnology Inc). Luminescence was detected with film, digitized and quantified. Arbitrary units of immunoreactivity were assigned based on the measured object density divided by the amount of protein loaded onto the gel.

mRNA isolation

mRNA was isolated using the Oligotex Direct mRNA Mini Kit (Qiagen, Valencia, CA) according to the manufacturer's protocol. The mRNA concentration was determined (OD 260 nm) using a NanoDrop® ND-1000 spectrophotometer (NanoDrop Technologies, Wilmington, DE). Purified mRNA was stored at -80 °C

cDNA synthesis

Reverse transcription reactions were performed in 20 µL following the manufacturer's protocol (Invitrogen Corporation). Briefly, a 10 µL reaction mixture containing 10 ng mRNA, 500 ng oligo dT, 1 mM each dNTP was heated at 65 °C for 5 minutes and chilled on ice for 5 minutes, followed by addition of 1 µL (200 units) M-MLV reverse transcriptase, 1 µL RNaseOUT (40 units/µL), 2µL DTT (0.1M) and sterile RNAase free water to 20µL. The whole reaction mixture was incubated at 37 °C for 50 minutes, then 75 °C for 10 minutes. The synthesized cDNA was stored at -20 °C until used. The suitability of the cDNA for PCR was determined using 2 µL of cDNA added to a 40 µL PCR. A control reaction containing 2µl of purified mRNA was run without reverse transcription to test for presence of DNA contamination in the purified mRNA samples. The PCR was analyzed on a 2% (w/v) agarose gel.

Primers

Oligonucleotide sequences (Table 1) specific to mouse BDNF Exons I-IV, VI and IX, and to β actin, cyclophilin A, hypoxanthine phosphoribosyltransferase 1 (HPRT1), and 18S rRNA were designed using Primer Express software (Applied Biosystems, Foster City, CA). The primers used for BDNF exon I, II, IV, and VI were based on sequences reported by Tsankova et al (2006). The specificity of each primer pair was confirmed by the identification of a single PCR product of predicted size on agarose gels followed by sequencing of the excised band (sequencing performed by the UNM Proteomic and Molecular Biology Core facility).

Semi-quantitative determination of transcript levels by Real Time-PCR

Real time PCR was conducted in a Gene Amp 7700 sequence detection system (Applied Biosystems) using a 96-well plate (MicroAmp™ Fast Optical 96-well Reaction plates and MicroAmp™ Optical Adhesive Film, both from Applied Biosystems). Reactions were performed in triplicate. Each 20 µL reaction contained forward and reverse primers (100 nM each, final concentration), 10 µL Power Sybr® Green PCR Master Mix reaction buffer (Applied Biosystems) and 2 µL cDNA. Following one initial step of 95 °C for 10 minutes, the cycling parameters were 95 °C for 15 sec, 60 °C for 1 min, and 40 cycles. The PCR products were monitored by measuring the increase in fluorescence caused by binding of Sybr® Green

Dye to the double-stranded DNA; Ct (cycle threshold) values were calculated by SDS software v1.9 (Applied Biosystems). The absence of nonspecific amplification was confirmed by analyzing PCR products by 2% (w/v) agarose gel.

In studies designed to identify an internal control to be used in analyses of BDNF gene expression, we assessed the expression of several housekeeping genes (β -actin, cyclophilin A, HPRT1, and 18S rRNA) using Ct values derived from Real-Time PCR. Reactions were performed in triplicate. The relative quantification of the BDNF gene in different tissue samples was determined using the $2^{-\Delta\Delta C_t}$ method described by Livak and Schmittgen (2001). Initially, the ΔC_t value, the difference between the average of the triplicate Ct values for the BDNF and the internal control β actin, for each sample was calculated ($\Delta C_{t_{\text{sample}}} = \Delta C_{t_{\text{BDNF}}} - \Delta C_{t_{\beta \text{ actin}}}$). Next, the average of the ΔC_t values for the saccharin control animals ($\Delta C_{t_{\text{SAC av}}}$) was determined and subtracted from each sample ($\Delta C_{t_{\text{sample}}} - \Delta C_{t_{\text{SAC av}}}$), giving a $\Delta\Delta C_t$ value for each. This, in turn, was used to calculate the $2^{-\Delta\Delta C_t}$ value for each. Because 2^0 is one, the mean $2^{-\Delta\Delta C_t}$ value for the saccharin control samples was approximately one, depending on the variability in the samples, and the mean $2^{-\Delta\Delta C_t}$ value for the ethanol-exposed mouse samples was expressed relative to one.

Statistical Analysis

All data were analyzed using a Student two-tailed t-test except the analyses of the regression lines and curves for the learned helplessness data, which were analyzed by ANOVA, using GraphPad Prism (version 4.0).

RESULTS

Fetal Alcohol Exposure Paradigm

The present studies were conducted using C57BL/6J mice and a voluntary two-bottle choice paradigm for prenatal ethanol exposure (Allan et al. 2003). The average consumption of ethanol by the mothers of the mice used in the present studies was $9.55 (\pm 2.57)$ g ethanol/kg body weight/day. In a concurrent study, we found that the average maternal blood ethanol concentration was 0.78 mg/mL in a group of C57BL/6 dams ($n=7$) whose average daily consumption of ethanol was 8 g/kg body weight. Across six breeding rounds, including the four from which the animals used in the present studies were derived, we observed that there were no differences between the weight gained by saccharin control or ethanol-consuming dams, nor were there differences in litter size (Table 2).

Prenatal ethanol-exposed mice display increased learned helplessness

Learned helplessness is a long-standing test for assessing depressive-like behavior (Maier, 1984; Caldarone et al., 2000; El Yacoubi and Vaugeois, 2007; McArthur and Borsini, 2006; Nestler et al., 2002b). The learned helplessness procedure that was employed in the present studies exposed saccharin control and ethanol-exposed mice to a series of inescapable footshocks, then in a subsequent test session assessed whether the animals escaped the footshock, when given the opportunity to do so; if escape occurred, the latency to escape was also recorded. The data for the % of test subjects that escaped (Figure 1, left) were best fit by the equation for a hyperbola, whereas the data for the escape latency (Figure 1, right) were best fit by a straight line.

Both saccharin and prenatal ethanol-exposed animals displayed increased escape behavior across the 28 testing trials, as demonstrated by an increased % of animals that escaped (Figure 1, left) and decreased mean escape latencies (Figure 1, right) as testing progressed. The percentage of animals that escaped was greater for the saccharin control mice than for the prenatal ethanol-exposed mice throughout the trials. Whereas nearly all of the control animals

escaped from trial 15 onward, approximately 40 % of the prenatal ethanol-exposed mice failed to escape in these later trials, demonstrating learned helplessness, or depressive-like behavior. The best-fit curves for the saccharin control and prenatal ethanol-exposed data sets were significantly different from each other [$F(2,52)=44.0$, $p<0.0001$] (Fig 1 Left). The similar slopes of the regression lines fitting the escape latency data indicate that the rate of improvement in escape behavior was similar in both groups [$F(1,445)=33.0$, $p<0.01$] (Fig. 1 Right). However, the line fitting the prenatal ethanol-exposed mouse data is elevated relative to that for the saccharin data, indicating that prenatal ethanol-exposed mice displayed increased learned helplessness behavior throughout the testing session.

Prenatal ethanol-exposed mice display increased immobility in the Porsolt forced swim test

As a second test for depressive-like behavior, we assessed the behavior of prenatal ethanol-exposed and saccharin control mice in the forced swim test, originally described by Porsolt et al. (1977). In this test, an animal is placed into a container of water, with the level of water sufficiently high to prevent the animal from touching the bottom of the container without submerging its body. Initially, the test subject exhibits gross motor movements (swimming, climbing, and thrashing) that are indicative of escape behavior, then eventually becomes relatively immobile, either floating or making only small motor movements (twitching and kicking), which are indicative of a state of despair (Porsolt et al., 1977). Figure 2 shows that, compared to saccharin control mice, mice exposed to ethanol prenatally displayed increased immobility (i.e., increased depressive behavior) in the forced swim test ($t(14)=8.41$, $p<0.0001$).

Prenatal ethanol-exposed and saccharin control mice do not differ in locomotor activity

Behavioral responses in both the learned helplessness and forced swim tests are dependent, in part, on an animal's motor skills; therefore, the observed depressive-like behaviors of the prenatal ethanol-exposed mice in these tests could, at least in part, be due to reduced locomotor activity in these animals, compared to saccharin control mice. To address this issue, we assessed locomotor activity in the two groups of mice under non-anxiogenic conditions (i.e., dim light illumination). We found no difference between the locomotor activity of prenatal ethanol-exposed and saccharin control mice (Figure 3).

Basal BDNF protein levels are reduced in the medial frontal cortex, but not the hippocampal formation, of ethanol-exposed mice

Elevated learned helplessness susceptibility has been correlated with reduced basal BDNF levels in the frontal cortex and hippocampal formation (Ridder et al., 2005); further, infusions of BDNF into the hippocampus (Shirayama et al., 2002), midbrain (Siuciak et al., 1997) or cerebral ventricles (Hoshaw et al., 2005) produce antidepressant effects when animals are subsequently tested in the learned helplessness (Shirayama et al., 2002; Siuciak et al., 1997) or forced swim (Hoshaw et al., 2005; Shirayama et al., 2002; Siuciak et al., 1997) tests. Thus, we sought to determine whether the observed increases in learned helplessness behavior and immobility in the forced swim test of prenatal ethanol-exposed mice were accompanied by decreased BDNF protein in the medial frontal cortex and/or hippocampal formation. Figure 4 shows that, compared to saccharin control mice, BDNF levels were reduced in the prenatal ethanol-exposed mouse frontal cortex ($t(13)=2.7$, $p<0.02$), but not in the hippocampal formation.

Prenatal alcohol exposure reduces proBDNF levels in the medial frontal cortex

BDNF, like other neurotrophins, is synthesized in a precursor form, termed proBDNF (Lu 2003), a 32-kDa protein that is cleaved to the 14-kDa mature BDNF (Mowla et al., 2001). BDNF is formed extracellularly, by the action of plasmin (Pang et al., 2004), and intracellularly,

both in the trans-Golgi network by furin and in secretory granules by convertases, including PACE4 and PACE5/6-B (Mowla et al., 2001; Seidah et al., 1996).

In order to begin addressing the mechanism(s) that underlies the observed decrease in BDNF protein in the medial frontal cortex, we analyzed proBDNF protein levels in this brain region, as well as in the hippocampal formation (Figure 5). Immunoblotting data showed that anti-proBDNF immunoreactivity was decreased in the medial frontal cortex ($t(17)=2.44$, $p<0.02$), but not the hippocampal formation, of mice exposed prenatally to ethanol, thus paralleling the data for BDNF protein levels.

Prenatal alcohol exposure is associated with reduced BDNF mRNA transcript levels in the medial frontal cortex and hippocampal formation

We sought to determine whether reduced BDNF mRNA levels may contribute to the observed decrease in medial frontal cortical proBDNF and BDNF protein. For these studies, it was necessary to identify a suitable internal standard for semi-quantitative (relative) Real-Time PCR analyses. Commonly used internal controls include β -actin, glyceraldehyde-3-phosphate dehydrogenase (GAPDH), cyclophilin A/B, HPRT1, and 18S rRNA. Several reports have found effects of experimental manipulations on the expression of these genes (e.g., Maier et al., 1999; Jang et al., 2005; Mogal and Abdulkadir, 2006; Verma and Shapiro, 2006; Tanic et al., 2007; Nishimura et al., 2008). Prenatal ethanol exposure has been reported to alter β -actin expression in embryos (Green et al., 2007) and neonatal cerebellum (Zoeller et al., 1994), but not neonatal or adolescent intestine (Bhalla et al., 2004). We compared cyclophilin A, HPRT1, 18S rRNA, and β -actin expression in saccharin control and prenatal ethanol-exposed adult hippocampal formation and medial frontal cortex (Figure 6). Prenatal exposure to ethanol did not alter the levels of β -actin or HPRT1 in either brain region, whereas it was associated with significantly increased 18S rRNA levels in the hippocampal formation ($t(14) = 3.64$, $p<0.01$) and reduced cyclophilin A levels in the medial frontal cortex ($t(14) = 5.16$, $p<0.01$). Based on these data, we chose to use β -actin as an internal control during analyses of BDNF transcript expression.

The mouse BDNF gene consists of multiple noncoding exons upstream of a common coding exon (Figure 7A). Until recently, most studies had identified six exons in the mouse gene, whereas Liu et al. (2006) identified eight exons and Aid et al. (2007) identified nine exons in the murine BDNF gene. We chose to follow the nomenclature of Aid et al. (2007). Exons are designated using Roman numerals, according to their relative position, with I being the 5'-most exon. Exons I–IXa are noncoding exons, while exon IX contains the open reading frame that encodes proBDNF. In order to measure individual BDNF mRNA transcripts, we used PCR primer pairs consisting of a forward primer specific for the upstream, non-coding exon I–VI, excluding exon V, and a reverse primer specific for the coding exon IX. Figure 7B shows that, in both the medial frontal cortex and hippocampal formation, prenatal ethanol exposure was associated with significantly decreased levels of BDNF total mRNA (exon IX-containing transcripts; medial frontal cortex, $t(14)=3.7$, $p<0.01$; hippocampal formation, $t(14)=3.9$, $p<0.01$) and BDNF mRNA transcripts containing exon III (medial frontal cortex, $t(14)=2.8$, $p<0.02$; hippocampal formation, $t(14)=4.2$, $p<0.01$), IV (medial frontal cortex, $t(14)=8.1$, $p<0.01$; hippocampal formation, $t(14)=6.2$, $p<0.01$), or VI (medial frontal cortex, $t(14)=8.6$, $p<0.01$; hippocampal formation, $t(14)=3.6$; $p<0.01$), but not those containing exon I or II.

CONCLUSIONS

FASD mice display depressive-like behaviors

We have identified increased learned helplessness, as well as increased immobility in a forced swim test, in adult mice prenatally exposed to moderate levels of ethanol. Thus, using two well

accepted tests for behaviors that model depression, we found that prenatal ethanol exposure is associated with depressive-like behavior in a mouse FASD model. These findings in prenatal ethanol-exposed mice confirm and extend those of Slone and Redei (2002), Wilcoxon et al. (2005), and Carneiro et al. (2005), who reported that adult rats that were exposed prenatally to ethanol displayed increased immobility in the forced swim test.

The escape response in the learned helplessness test includes a learning component; thus, it can be argued that the observed increase in escape latency and escape failures in prenatal ethanol-exposed mice is a consequence of their known learning impairments (Allan et al., 2003). However, this is refuted by our finding that the slopes of the regression lines fitting the escape latency data for the ethanol-exposed and saccharin mice were similar, indicating that the two groups learned the escape response at comparable rates. The observed increased immobility in the forced swim test, which does not involve learning, also supports our conclusion that prenatal ethanol-exposed mice display behaviors that model depression. It can also be argued that the observed increase in escape latency in the learned helplessness task and increased immobility in the forced swim task are due to reduced locomotor activity in the prenatal ethanol-exposed mice. However, these mice do not differ from saccharin control mice in their locomotor activity in an open field, further supporting the conclusion that the behaviors in the learned helplessness and forced swim tests represent depressive-like behaviors.

BDNF protein and mRNA levels are reduced in the brain of prenatal ethanol-exposed adult mice

Several hypotheses have been put forth regarding the genetic and biochemical bases of depression (Hayley et al., 2005; Jans et al., 2007; Levinson, 2006; Nestler et al., 2002a). Many of these have focused on mechanisms underlying the compromised functioning of the medial frontal cortex and hippocampal formation which is observed in depressed individuals (Berton and Nestler, 2006; Drevets, 2000; Johnstone et al., 2007; Phillips et al., 2003a,b). Consistent with the neurotrophic hypothesis of depression, we found that proBDNF and BDNF protein levels were reduced in the medial frontal cortex of prenatal ethanol-exposed mice compared to saccharin control mice; however, proBDNF and BDNF protein levels were not different in the hippocampal formation of these same animals. We also found that prenatal ethanol exposure was associated with reduced BDNF total mRNA levels in both the medial frontal cortex and hippocampal formation. In both brain regions, the levels of exon III-, IV-, and VI-containing transcripts were decreased, while the levels of exon I- and II-containing transcripts were not different between prenatal ethanol-exposed and saccharin control mice. Thus, the observed decreases in BDNF mRNA transcript levels in prenatal ethanol-exposed mice correlated with reduced proBDNF and BDNF protein levels in the medial frontal cortex, but not in the hippocampal formation. This indicates that proBDNF and BDNF protein levels in prenatal ethanol-exposed mice are likely controlled primarily by transcription in the medial frontal cortex, but post-transcriptionally in the hippocampal formation. A similar lack of correlation between hippocampal BDNF mRNA and protein levels has been reported by Nawa et al. (1995) and Greisen et al. (2005). Studies conducted by Nawa et al. (1995) using two-month old male Sprague-Dawley rats show that hippocampal BDNF protein levels are elevated four days following limbic seizures induced by electrolytic lesion of the dentate gyrus hilus. In contrast, Isackson et al. (1991), using the same procedure to induce limbic seizures in adult male Sprague-Dawley rats, reported that BDNF mRNA levels in the hippocampus were returned to baseline levels by 4 days following seizure induction. Greisen et al. (2005) reported that, compared to control animals, hippocampal formation BDNF protein levels were increased 63%, while BDNF mRNA levels were unchanged, in three-month old male Sprague Dawley rats that had experienced daily maternal separation on postnatal days 2 through 14. In addition to alterations in BDNF mRNA and protein metabolism, these investigators propose that the disparity between BDNF protein and mRNA levels may be accounted for by anterograde

transport of BDNF protein to the hippocampus by afferent neuronal projections from other brain regions, such as the medial septal nucleus. Finally, a lack of agreement between BDNF mRNA and protein levels has been observed in the striatum and prefrontal cortex following chronic cocaine administration in rats (Fumagalli et al., 2007).

What mechanisms may underlie the observed decreases in specific BDNF transcripts?

Chronic defeat stress has been shown to decrease the expression of exon III- and IV-containing transcripts in the hippocampal formation (Tsankova et al., 2006). Therefore, increased chronic stress levels may contribute to the observed reductions in exon III- and IV-containing transcripts in prenatal ethanol-exposed mice, at least in the hippocampal formation. Interestingly, we have reported that prenatal ethanol-exposed female mice have higher plasma corticosterone levels than do saccharin controls (Allan et al., 2003), which could also account for the observed decrease in exon VI-containing transcripts in the medial frontal cortex and hippocampal formation, where the expression of exon VI-containing transcripts is repressed by corticosterone (Dwivedi et al., 2006; Hansson et al., 2006). The expression of BDNF exon IV-containing transcripts is controlled by the methylation of CpG dinucleotides in the promoter linked to this exon (Chen et al., 2003; Dennis and Levitt, 2005; Martinowich et al., 2003). Maier et al. (1999) reported that prenatal ethanol exposure was associated with reduced BDNF mRNA levels in the postnatal day 10 Sprague-Dawley rat olfactory bulb, and this reduction was correlated with hypermethylation of the BDNF gene. Although these investigators did not demonstrate specific reductions in exon IV-containing species or methylation of the exon IV promoter, their results suggest that the observed decrease in BDNF mRNA containing exon IV in prenatal ethanol-exposed mice may be due to increased methylation of CpG dinucleotides preceding the transcriptional initiation site for exon IV. In addition to being controlled by methylation, the promoter for exon IV, as well as the promoter associated with exon I, has a cAMP-responsive element (CRE) that binds cAMP-responsive element binding protein (CREB; Fang et al., 2003; Ou and Gean, 2007; Shieh et al., 1998; Tabuchi et al., 2002; Tao et al., 1998). Thus, reduced levels of CREB activation in mice that were exposed to ethanol prenatally could contribute to the observed reduction in the expression of exon IV-containing transcripts in these animals. However, the absence of an observed effect of prenatal ethanol exposure on exon I-containing transcripts does not support a role for global reductions in the level of activated CREB. The expression of BDNF transcripts containing exon VI is regulated by extracellular signal-regulated protein kinase 2 (ERK2; Park et al., 2006; Takeuchi et al., 2002). We have found that receptor-dependent activation of ERK2 in hippocampal formation slices is impaired (Samudio-Ruiz, Valenzuela, Allan, and Caldwell; unpublished observation). Therefore, reduced ERK2 activation could play a role in the observed decrease in exon VI-containing transcripts in the hippocampal formation of ethanol-exposed mice. Finally, acetylated histone H4 has been shown to regulate the expression of promoter VI-containing transcripts in the medial frontal cortex (Bredy et al., 2007). We have found that global levels of acetyl (Lys5, Lys8, Lys12, Lys16) histone H4 are reduced in the medial frontal cortex, but not hippocampal formation, of prenatal ethanol-exposed mice (Caldwell and Allan, unpublished observation), implicating reduced binding of acetylated histone H4 at the promoter for BDNF exon VI as a candidate mechanism underlying the observed reduced level of exon VI-containing transcripts in the medial frontal cortex of the prenatal ethanol-exposed mice.

Evidence exists that does not support a role for BDNF in depressive-like behaviors

Although several lines of research indicate that BDNF plays an important role in depression and depressive-like behaviors, it should be noted that evidence exists that does not support this conclusion. Angelucci et al. (2000) reported that, compared to the Flinders Resistant Line (FRL), the Flinders Sensitive Line (FSL) rats, which display a variety of depressive-like behaviors, have increased basal BDNF protein levels in the frontal cortex, but not the hippocampal formation. Greisen et al. (2005) found that adult male Sprague-Dawley rats

having reduced hippocampal formation, and unchanged frontal cortical, BDNF protein levels display normal behavior in the forced swim test. Hellweg et al. (2007) have reported that olfactory bulbectomy, which has been proposed as a rodent model of depression, of adult male C57BL/6N mice leads to increased BDNF protein levels in the hippocampal formation and frontal cortex. Although this study may appear to be contradictory to our findings, it is important to note that the C57BL/6J mouse strain, which we employed, and the C57BL/6N mouse strain differ in behavioral responses, such as extinction of fear-conditioned responding (Stiedl et al., 1999), and physiologic responses, such as effects of anesthetics on cardiac function (Roth et al., 2002). Thus, as Hellweg et al. (2007) have proposed, BDNF may contribute only to certain types of depressive states or only under certain condition (e.g., certain mouse strains or substrains) in laboratory models of depression. Finally, Greenwood et al. (2007) have found that learned helplessness in adult, male Fischer F344 rats does not depend on hippocampal formation BDNF. This last observation is consistent with our finding of unaltered BDNF protein in the hippocampal formation of prenatal ethanol-exposed mice compared to saccharin controls, yet FASD mice display elevated learned helplessness behavior.

Relevance to pharmacologic management of clinical depression in FASD patients

The present studies may have relevance to the choice of antidepressant for the treatment of FASD-associated depression. Chronic tranylcypromine treatment has been shown to increase BDNF protein levels in the rat frontal cortex, but not the hippocampal formation, whereas fluoxetine and desipramine did not affect medial frontal cortical BDNF protein levels (Altar et al., 2003). Dwivedi et al. (2006) reported that chronic phenelzine and desipramine increased BDNF mRNA in both the frontal cortex and hippocampal formation, while fluoxetine increased BDNF only in the hippocampal formation. Dias et al. (2003) found that chronic desipramine increased the levels of BDNF exon III-containing transcripts in the piriform cortex and inner and outer layers of the neocortex; in contrast, fluoxetine was without effect in these regions. Okamoto et al. (2003) found that chronic amitriptyline increased BDNF protein levels in the hippocampal formation and frontal cortex. Thus, drugs shown to increase BDNF mRNA or protein levels in the frontal cortex, such as tranylcypromine, amitriptyline, phenelzine, and possibly desipramine, may be effective in treating depressive symptoms in FASD patients, whereas fluoxetine may be less effective.

Summary

These studies demonstrate that prenatal ethanol-exposed female mice display depressive-like behaviors, when tested as adults. It is important to note that all of the studies described in this report were conducted in female offspring. The choice of conducting these studies in females was based, in part, on the observation that the incidence of clinical depression is higher in females than in males (Kessler et al. 1994, Weissman and Olfson 1995, Piccinelli and Wilkinson 2000, Kessler 2003, Kessler et al. 2006; Grigoriadis and Robinson 2007). Additionally, Calderone et al. (2000) have reported that female C57BL/6J mice are more prone to develop learned helplessness than are male C57BL/6J mice. It will be interesting to determine if a similar correlation between prenatal ethanol exposure, depression, and brain BDNF levels is found in male mice. Monteggia et al (2007) have reported that, while conditional knockout of forebrain BDNF is associated with depressive-like behavior in female mice, male conditional knockout mice do not display increased depressive behavior.

In addition to displaying depressive-like behaviors, adult female offspring that had been exposed to ethanol prenatally had reduced BDNF mRNA and protein levels in the medial frontal cortex and BDNF mRNA levels in the hippocampal formation. These data demonstrate that depressive-like behaviors in a mouse FASD model and alterations in brain BDNF are correlated. Although they do not provide direct evidence that these reductions in BDNF are an underlying cause of the observed depressive-like behaviors, the data are consistent with the

neurotrophic hypothesis of depression and indicate that the hypothesis may be extended to include FASD-associated depression. In order to determine whether reductions in BDNF play a causative role in prenatal ethanol-associated depressive-like behaviors, future studies will need to identify the developmental stage at which prenatal ethanol exposure exerts its effects on BDNF expression and test the behavioral consequence of reversing these effects (e.g., through targeted overexpression of BDNF). The results of the present study do, however, support the use of antidepressants that have been shown to increase frontal cortical BDNF in the treatment of FASD-associated depression.

It is possible that the effects of prenatal ethanol exposure on BDNF mRNA are, at least in part, mediated by, or secondary to, changes in serotonergic neurotransmission. Several investigators have reported that developing serotonergic neurons are highly vulnerable to the damaging effects of ethanol (Rathbun and Druse, 1985; Sari and Zhou, 2004; Tajuddin, and Druse, 1993, 1999; Zhou et al., 2002, 2005). These ethanol-induced decreases in 5-HT levels and neurons are likely to have a significant cascading effect on the developing nervous system as a whole, thereby impacting BDNF. Additionally, as noted previously, several antidepressants, including serotonin-selective reuptake inhibitors (SSRIs), have been shown to modify BDNF levels in the prefrontal cortex and hippocampus. Thus, future studies will aim to determine the contribution of reduced serotonergic neurotransmission to the observed alterations in BDNF levels.

Acknowledgments

We thank Julie Chynoweth for technical assistance and Fernando Valenzuela for thoughtful evaluation of the manuscript. Sources of support: Chilean Government Scholarship for postdoctoral training (MIDEPLAN) to RDP; U.S. Department of Army Award #DAMD17-01-1-0680 to AMA; SLS-R was supported by NIH-NIAAA award #1T32AA014127.

References

- Aid T, Kazantseva A, Piirsoo M, Palm K, Timmusk T. Mouse and rat BDNF gene structure and expression revisited. *J Neurosci Res* 2007;85:525–535. [PubMed: 17149751]
- Allan AM, Chynoweth J, Tyler LA, Caldwell KK. A mouse model of prenatal ethanol exposure using a voluntary drinking paradigm. *Alcohol Clin Exp Res* 2003;27:2009–2016. [PubMed: 14691390]
- Altar CA, Whitehead RE, Chen R, Wortwein G, Madsen TM. Effects of electroconvulsive seizures and antidepressant drugs on brain-derived neurotrophic factor protein in rat brain. *Biol Psychiatry* 2003;54:703–709. [PubMed: 14512210]
- Angelucci F, Aloe L, Vasquez PJ, Mathe AA. Mapping the differences in the brain concentration of brain-derived neurotrophic factor (BDNF) and nerve growth factor (NGF) in an animal model of depression. *NeuroReport* 2000;11:1360–1374.
- Berton O, Nestler EJ. New approaches to antidepressant drug discovery: beyond monoamines. *Nat Rev Neurosci* 2006;7:137–151. [PubMed: 16429123]
- Bhalla S, Mahmood S, Mahmood A. Effect of prenatal exposure to ethanol on postnatal development of intestinal transport functions in rats. *Eur J Nutr* 2004;43:109–115. [PubMed: 15083318]
- Bredy TW, Wu H, Crego C, Zellhoefer J, Sun YE, Barad M. Histone modifications around individual BDNF gene promoters in prefrontal cortex are associated with extinction of conditioned fear. *Learn Mem* 2007;14:268–276. [PubMed: 17522015]
- Buckley CT, Sekiya F, Kim YJ, Rhee SG, Caldwell KK. Identification of phospholipase C- γ 1 as a mitogen-activated protein kinase substrate. *J Biol Chem* 2004;279:41807–41814. [PubMed: 15258148]
- Caldarone BJ, George TP, Zachariou V, Picciotto MR. Gender differences in learned helplessness behavior are influenced by genetic background. *Pharmacol Biochem Behav* 2000;66:811–817. [PubMed: 10973520]

- Caldarone BJ, Harrist A, Cleary MA, Beech RD, King SL, Picciotto MR. High-affinity nicotinic acetylcholine receptors are required for antidepressant effects of amitriptyline on behavior and hippocampal cell proliferation. *Biol Psychiatry* 2004;56:657–664. [PubMed: 15522249]
- Carneiro LMV, Diogenes JPL, Vasconcelos SMM, Aragao GF, Noronha EC, Gomes PB, Viana GSB. Behavioral and neurochemical effects on rat offspring after prenatal exposure to ethanol. *Neurotoxicol Teratol* 2005;27:585–592. [PubMed: 16039829]
- Chen B, Dowlatsahi D, MacQueen GM, Wang J-F, Young LT. Increased hippocampal BDNF immunoreactivity in subjects treated with antidepressant medication. *Biol Psychiatry* 2001;50:260–265. [PubMed: 11522260]
- Chen WG, Chang Q, Lin Y, Meissner A, West AE, Griffith EC, Jaenisch R, Greenberg ME. Derepression of BDNF transcription involves calcium-dependent phosphorylation of MeCP2. *Science* 2003;302:885–889. [PubMed: 14593183]
- Climent E, Pascual M, Renau-Piqueras J, Guerri C. Ethanol exposure enhances cell death in the developing cerebral cortex: role of brain-derived neurotrophic factor and its signaling pathways. *J Neurosci Res* 2002;68:213–225. [PubMed: 11948666]
- Dennis KE, Levitt P. Regional expression of brain derived neurotrophic factor (BDNF) is correlated with dynamic patterns of promoter methylation in the developing mouse forebrain. *Mol Brain Res* 2005;140:1–9. [PubMed: 16054725]
- Dias BG, Banerjee SB, Duman RS, Vaidya VA. Differential regulation of brain derived neurotrophic factor transcripts by antidepressant treatments in the adult rat brain. *Neuropharmacology* 2003;45:553–563. [PubMed: 12907316]
- Drevets WC. Functional anatomical abnormalities in limbic and prefrontal cortical structures in major depression. *Prog Brain Res* 2000;126:413–431. [PubMed: 11105660]
- Duman RS, Monteggia LM. A neurotrophic model for stress-related mood disorders. *Biol Psychiatry* 2006;59:1116–1127. [PubMed: 16631126]
- Dwivedi Y, Rizavi HS, Pandey GN. Antidepressants reverse corticosterone-mediated decrease in brain-derived neurotrophic factor expression: differential regulation of specific exons by antidepressants and corticosterone. *Neuroscience* 2006;139:1017–1029. [PubMed: 16500030]
- El Yacoubi M, Vaugeois JM. Genetic rodent models of depression. *Curr Opin Pharmacol* 2007;7:3–7. [PubMed: 17169613]
- Fang H, Chartier J, Sodja C, Desbois A, Ribocco-Lutkiewicz M, Walker PR, Sikorska M. Transcriptional activation of the human brain-derived neurotrophic factor gene promoter III by dopamine signaling in NT2/N neurons. *J Biol Chem* 2003;278:26401–26409. [PubMed: 12738784]
- Famy C, Streissguth AP, Unis AS. Mental illness in adults with fetal alcohol syndrome or fetal alcohol effects. *Am J Psychiatry* 1998;155:552–554. [PubMed: 9546004]
- Feng MJ, Yan SE, Yan QS. Effects of prenatal alcohol exposure on brain-derived neurotrophic factor and its receptor tyrosine kinase B in offspring. *Brain Res* 2005;1042:125–132. [PubMed: 15854584]
- Fumagalli F, Di Pasquale L, Caffino L, Racagni G, Riva MA. Repeated exposure to cocaine differently modulates BDNF mRNA and protein levels in rat striatum and prefrontal cortex. *Eur J Neurosci* 2007;26:2756–2763. [PubMed: 18001273]
- Green ML, Singh AV, Zhang Y, Nemeth KA, Sulik KK, Knudsen TB. Reprogramming of genetic networks during initiation of the Fetal Alcohol Syndrome. *Dev Dyn* 2007;236:613–631. [PubMed: 17200951]
- Greenwood BN, Strong PV, Foley TE, Thompson RS, Fleshner M. Learned helplessness is independent of levels of brain-derived neurotrophic factor in the hippocampus. *Neuroscience* 2007;144:1193–1208. [PubMed: 17161541]
- Greisen MH, Altar CA, Bolwig TG, Whitehead R, Wortwein G. Increased adult hippocampal brain-derived neurotrophic factor and normal levels of neurogenesis in maternal separation rats. *J Neurosci Res* 2005;79:772–778. [PubMed: 15690366]
- Grigoriadis S, Robinson GE. Gender issues in depression. *Ann Clin Psychiatry* 2007;19:247–255. [PubMed: 18058282]
- Hansson AC, Sommer WH, Metsis M, Stromberg I, Agnati LF, Fuxe K. Corticosterone actions on the hippocampal brain-derived neurotrophic factor expression are mediated by exon IV promoter. *J Neuroendocrinol* 2006;18:104–114. [PubMed: 16420279]

- Hasler G, van der Veen JW, Tumonis T, Meyers N, Shen J, Drevets WC. Reduced prefrontal glutamate/ glutamine and γ -aminobutyric acid levels in major depression determined using proton magnetic resonance spectroscopy. *Arch Gen Psychiatry* 2007;64:193–200. [PubMed: 17283286]
- Hayley S, Poulter MO, Merali Z, Anisman H. The pathogenesis of clinical depression: stressor- and cytokine-induced alterations of neuroplasticity. *Neuroscience* 2005;135:659–678. [PubMed: 16154288]
- Heaton MB, Mitchell JJ, Paiva M, Walker DW. Ethanol-induced alterations in the expression of neurotrophic factors in the developing rat central nervous system. *Dev Brain Res* 2000;121:97–107. [PubMed: 10837897]
- Hellweg R, Zueger M, Fink K, Hörtnagl H, Gass P. Olfactory bulbectomy in mice leads to increased BDNF levels and decreased serotonin turnover in depression-related brain areas. *Neurobiol Dis* 2007;25:1–7. [PubMed: 16990008]
- Hoshaw BA, Malberg JE, Lucki I. Central administration of IGF-I and BDNF leads to long-lasting antidepressant-like effects. *Brain Res* 2005;1037:204–208. [PubMed: 15777771]
- Isackson PJ, Huntsman MM, Murray KD, Gall CM. BDNF mRNA expression is increased in adult rat forebrain after limbic seizures: temporal patterns of induction distinct from NGF. *Neuron* 1991;6:937–948. [PubMed: 2054188]
- Itoh T, Tokumura M, Abe K. Effects of rolipram, a phosphodiesterase 4 inhibitor, in combination with imipramine on depressive behavior, CRE-binding activity and BDNF level in learned helplessness rats. *Eur J Pharmacol* 2004;498:135–142. [PubMed: 15363987]
- Jans LA, Riedel WJ, Markus CR, Blokland A. Serotonergic vulnerability and depression: assumptions, experimental evidence and implications. *Mol Psychiatry* 2007;12(6):522–543. [PubMed: 17160067]
- Johnstone T, van Reekum CM, Urry HL, Kalin NH, Davidson RJ. Failure to regulate: counterproductive recruitment of top-down prefrontal-subcortical circuitry in major depression. *J Neurosci* 2007;27:8877–8884. [PubMed: 17699669]
- Karege F, Vaudan G, Schwald M, Perroud N, La Harpe R. Neurotrophin levels in postmortem brains of suicide victims and the effects of antemortem diagnosis and psychotropic drugs. *Mol Brain Res* 2005;136:29–37. [PubMed: 15893584]
- Kessler RC. Epidemiology of women and depression. *J Affect Disord* 2003;74:5–13. [PubMed: 12646294]
- Kessler RC, Akiskal HS, Ames M, Birnbaum H, Greenberg P, A RM, Jin R, Merikangas KR, Simon GE, Wang PS. Prevalence and effects of mood disorders on work performance in a nationally representative sample of U.S. workers. *Am J Psychiatry* 2006;163:1561–1568. [PubMed: 16946181]
- Kessler RC, McGonagle KA, Nelson CB, Hughes M, Swartz M, Blazer DG. Sex and depression in the National Comorbidity Survey. II: Cohort effects. *J Affect Disord* 1994;30:15–26. [PubMed: 8151045]
- Levinson DF. The genetics of depression: a review. *Biol Psychiatry* 2006;60:84–92. [PubMed: 16300747]
- Liu Q-R, Lu L, Zhu X-G, Gong J-P, Shaham Y, Uhl GR. Rodent BDNF genes, novel promoters, novel splice variants, and regulation by cocaine. *Brain Res* 2006;1067:1–12. [PubMed: 16376315]
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the $2^{-\Delta\Delta C_T}$ method. *Methods* 2001;25:402–408. [PubMed: 11846609]
- Lu B. Pro-region of neurotrophins: role in synaptic modulation. *Neuron* 2003;39:735–738. [PubMed: 12948441]
- MacQueen GM, Ramakrishnan K, Croll SD, Siuciak JA, Yu G, Young LT, Fahnstock M. Performance of heterozygous brain-derived neurotrophic factor knockout mice on behavioral analogues of anxiety, nociception, and depression. *Behav Neurosci* 2001;115:1145–1153. [PubMed: 11584927]
- Maier SE, Cramer JA, West JR, Sohrabji F. Alcohol exposure during the first two trimesters equivalent alters granule cell number and neurotrophin expression in the developing rat olfactory bulb. *J Neurobiol* 1999;41:414–423. [PubMed: 10526319]
- Maier SF. Learned helplessness and animal models of depression. *Prog Neuropsychopharmacol Biol Psychiatry* 1984;8:435–446. [PubMed: 6385140]
- Martinowich K, Hattori D, Wu H, Fouse S, He F, Hu Y, Fan G, Sun YE. DNA methylation-related chromatin remodeling in activity-dependent *Bdnf* gene regulation. *Science* 2003;302:890–893. [PubMed: 14593184]

- McArthur R, Borsini F. Animal models of depression in drug discovery: a historical perspective. *Pharmacol Biochem Behav* 2006;84:436–452. [PubMed: 16844210]
- Mogal A, Abdulkadir SA. Effects of Histone Deacetylase Inhibitor (HDACi); Trichostatin-A (TSA) on the expression of housekeeping genes. *Mol Cell Probes* 2006;20:81–86. [PubMed: 16326072]
- Monteggia LM, Luikart B, Barrot M, Theobald D, Malkovska I, Nef S, Parada LF, Nestler EJ. Brain-derived neurotrophic factor conditional knockouts show gender differences in depression-related behaviors. *Biol Psychiatry* 2007;61:187–197. [PubMed: 16697351]
- Mowla SJ, Farhadi HF, Pareek S, Atwal JK, Morris SJ, Seidah NG, Murphy RA. Biosynthesis and post-translational processing of the precursor to brain-derived neurotrophic factor. *J Biol Chem* 2001;276:12660–12666. [PubMed: 11152678]
- Nawa H, Carnahan J, Gall C. BDNF protein measured by a novel enzyme immunoassay in normal brain and after seizure: partial disagreement with mRNA levels. *Eur J Neurosci* 1995;7:1527–1535. [PubMed: 7551179]
- Nestler EJ, Barrot M, DiLeone RJ, Eisch AJ, Gold SJ, Monteggia LM. Neurobiology of depression. *Neuron* 2002a;34:13–25. [PubMed: 11931738]
- Nestler EJ, Gould E, Manji H, Bucan M, Duman RS, Gershenfeld HK, Hen R, Koester S, Lederhendler I, Meaney M, Robbins T, Winsky L, Zalcman S. Preclinical models: status of basic research in depression. *Biol Psychiatry* 2002b;52:503–528. [PubMed: 12361666]
- Newton SS, Thome J, Wallace TL, Shirayama Y, Schlesinger L, Sakai N, Chen J, Neve R, Nestler EJ, Duman RS. Inhibition of cAMP response element-binding protein or dynorphin in the nucleus accumbens produces an antidepressant-like effect. *J Neurosci* 2002;22:10883–10890. [PubMed: 12486182]
- Nibuya M, Morinobu S, Duman RS. Regulation of BDNF and trkB mRNA in rat brain by chronic electroconvulsive seizure and antidepressant drug treatments. *J Neurosci* 1995;15:7539–7547. [PubMed: 7472505]
- Nishimura M, Nikawa T, Kawano Y, Nakayama M, Ikeda M. Effects of dimethyl sulfoxide and dexamethasone on mRNA expression of housekeeping genes in cultures of C2C12 myotubes. *Biochem Biophys Res Commun*. 2008In press
- O'Connor MJ, Kasari C. Prenatal alcohol exposure and depressive features in children. *Alcohol Clin Exp Res* 2000;24:1084–1092. [PubMed: 10924014]
- O'Connor MJ, Paley B. The relationship of prenatal alcohol exposure and the postnatal environment to child depressive symptoms. *J Pediatr Psychol* 2006;31:50–64. [PubMed: 15802607]
- O'Connor MJ, Shah B, Whaley S, Cronin P, Gunderson B, Graham J. Psychiatric illness in a clinical sample of children with prenatal alcohol exposure. *Am J Drug Alcohol Abuse* 2002;28:743–754. [PubMed: 12492268]
- Okamoto H, Shino Y, Hashimoto K, Kumakiri C, Shimizu E, Shirasawa H, Iyo M. Dynamic changes in AP-1 transcription factor DNA binding activity in rat brain following administration of antidepressant amitriptyline and brain-derived neurotrophic factor. *Neuropharmacology* 2003;45:251–259. [PubMed: 12842131]
- Ou LC, Gean PW. Transcriptional regulation of brain-derived neurotrophic factor in the amygdala during consolidation of fear memory. *Mol Pharmacol* 2007;72:350–358. [PubMed: 17456785]
- Pang PT, Teng HK, Zaitsev E, Woo NT, Sakata K, Zhen S, Teng KK, Yung W-H, Hempstead BL, Lu B. Cleavage of proBDNF by tPA/plasmin is essential for long-term hippocampal plasticity. *Science* 2004;306:487–491. [PubMed: 15486301]
- Park SY, Lee JY, Choi JY, Park MJ, Kim DS. Nerve growth factor activates brain-derived neurotrophic factor promoter IV via extracellular signal-regulated protein kinase 1/2 in PC12 cells. *Mol Cells* 2006;21:237–243. [PubMed: 16682819]
- Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception I: The neural basis of normal emotion perception. *Biol Psychiatry* 2003a;54:504–514. [PubMed: 12946879]
- Phillips ML, Drevets WC, Rauch SL, Lane R. Neurobiology of emotion perception II: Implications for major psychiatric disorders. *Biol Psychiatry* 2003b;54:515–528. [PubMed: 12946880]
- Piccinelli M, Wilkinson G. Gender differences in depression. Critical review. *Br J Psychiatry* 2000;177:486–492. [PubMed: 11102321]

- Porsolt RD, Le Pichon M, Jalfre M. Depression: a new animal model sensitive to antidepressant treatments. *Nature* 1977;266:730–732. [PubMed: 559941]
- Rathbun W, Druse MJ. Dopamine, serotonin and acid metabolites in brain regions from the developing offspring of ethanol treated rats. *J Neurochem* 1985;44:57–62. [PubMed: 2578063]
- Ridder S, Chourbaji S, Hellweg R, Urani A, Zacher C, Schmid W, Zink M, Hortnagl H, Flor H, Henn FA, Schutz G, Gass P. Mice with genetically altered glucocorticoid receptor expression show altered sensitivity for stress-induced depressive reactions. *J Neurosci* 2005;25:6243–6250. [PubMed: 15987954]
- Roebuck TM, Mattson SN, Riley EP. Behavioral and psychosocial profiles of alcohol-exposed children. *Alcohol Clin Exp Res* 1999;23:1070–1076. [PubMed: 10397293]
- Roth DM, Swaney JS, Dalton ND, Gilpin EA, Ross J Jr. Impact of anesthesia on cardiac function during echocardiography in mice. *Am J Physiol Heart Circ Physiol* 2002;282:H2134–H2140. [PubMed: 12003821]
- Russo-Neustadt A, Ha T, Ramirez R, Kesslak JP. Physical activity-antidepressant treatment combination: impact on brain-derived neurotrophic factor and behavior in an animal model. *Behav Brain Res* 2001;120:87–95. [PubMed: 11173088]
- Sampson PD, Streissguth AP, Bookstein FL, Little RE, Clarren SK, Dehaene P, Hanson JW, Graham JM. Incidence of fetal alcohol syndrome and prevalence of alcohol-related neurodevelopmental disorder. *Teratology* 1997;56:317–326. [PubMed: 9451756]
- Sari Y, Zhou FC. Prenatal alcohol exposure causes long-term serotonin neuron deficit in mice. *Alcohol Clin Exp Res* 2004;28:941–948. [PubMed: 15201637]
- Schramm NL, McDonald MP, Limbird LE. The α_2A -adrenergic receptor plays a protective role in mouse behavioral models of depression and anxiety. *J Neurosci* 2001;21:4875–4882. [PubMed: 11425914]
- Schulte-Herbrüggen O, Chourbaji S, Muller H, Danker-Hopfe H, Brandwein C, Gass P, Hellweg R. Differential regulation of nerve growth factor and brain-derived neurotrophic factor in a mouse model of learned helplessness. *Exp Neurol* 2006;202:404–409. [PubMed: 16914143]
- Seidah NG, Benjannet S, Pareek S, Chretien M, Murphy RA. Cellular processing of the neurotrophin precursors of NT3 and BDNF by the mammalian proprotein convertases. *FEBS Lett* 1996;379:247–250. [PubMed: 8603699]
- Shieh PB, Hu S-C, Bobb K, Timmusk T, Ghosh A. Identification of a signaling pathway involved in calcium regulation of *BDNF* expression. *Neuron* 1998;20:727–740. [PubMed: 9581764]
- Shirayama Y, Chen AC-H, Nakagawa S, Russell DS, Duman RS. Brain-derived neurotrophic factor produces antidepressant effects in behavioral models of depression. *J Neurosci* 2002;22:3251–3261. [PubMed: 11943826]
- Siuciak JA, Lewis DR, Wiegand SJ, Lindsay RM. Antidepressant-like effect of brain-derived neurotrophic factor (BDNF). *Pharmacol Biochem Behav* 1997;56:131–137. [PubMed: 8981620]
- Slone JL, Redei EE. Maternal alcohol and adrenalectomy: asynchrony of stress response and forced swim behavior. *Neurotoxicol Teratol* 2002;24:173–178. [PubMed: 11943505]
- Song L, Che W, Min-Wei W, Murakami Y, Matsumoto K. Impairment of the spatial learning and memory induced by learned helplessness and chronic mild stress. *Pharmacol Biochem Behav* 2006;83:186–193. [PubMed: 16519925]
- Stiedl O, Radulovic J, Lohmann R, Birkenfeld K, Palve M, Kammermeier J, Sananbenesi F, Spiess J. Strain and substrain differences in context- and tone-dependent fear conditioning of inbred mice. *Behav Brain Res* 1999;104:1–12. [PubMed: 11125727]
- Streissguth AP, O'Malley K. Neuropsychiatric implications and long-term consequences of fetal alcohol spectrum disorders. *Semin Clin Neuropsychiatry* 2000;5:177–190. [PubMed: 11291013]
- Sunal R, Gumusel B, Kayaalp SO. Effect of changes in swimming area on results of “behavioral despair test”. *Pharmacol Biochem Behav* 1994;49:891–896. [PubMed: 7886103]
- Tabuchi A, Sakaya H, Kisukeda T, Fushiki H, Tsuda M. Involvement of an upstream stimulatory factor as well as cAMP-responsive element-binding protein in the activation of brain-derived neurotrophic factor gene promoter I. *J Biol Chem* 2002;277:35920–35931. [PubMed: 12114522]
- Tajuddin NF, Druse MJ. Treatment of pregnant alcohol-consuming rats with buspirone: effects on serotonin and 5-hydroxyindoleacetic acid content in offspring. *Alcohol Clin Exp Res* 1993;17:110–114. [PubMed: 7680841]

- Tajuddin NF, Druse MJ. In utero ethanol exposure decreased the density of serotonin neurons. Maternal ipsapirone treatment exerted a protective effect. *Dev Brain Res* 1999;117:91–97. [PubMed: 10536236]
- Takeda H, Tsuji M, Yamada T, Masuya J, Matsushita K, Tahara M, Iimori M, Matsumiya T. Caffeic acid attenuates the decrease in cortical BDNF mRNA expression induced by exposure to forced swimming stress in mice. *Eur J Pharmacol* 2006;534:115–121. [PubMed: 16497295]
- Takeuchi Y, Miyamoto E, Fukunaga K. Analysis on the promoter region of exon IV brain-derived neurotrophic factor in NG108-15 cells. *J Neurochem* 2002;83:67–79. [PubMed: 12358730]
- Tanic N, Perovic M, Mladenovic A, Ruzdijic S, Kanazir S. Effects of aging, dietary restriction and glucocorticoid treatment on housekeeping gene expression in rat cortex and hippocampus-evaluation by real time RT-PCR. *J Mol Neurosci* 2007;32:38–46. [PubMed: 17873286]
- Tao X, Finkbeiner S, Arnold DB, Shaywitz AJ, Greenberg ME. Ca^{2+} influx regulates *BDNF* transcription by a CREB family transcription factor-dependent mechanism. *Neuron* 1998;20:709–726. [PubMed: 9581763]
- Tsankova NM, Berton O, Renthal W, Kumar A, Neve RL, Nestler EJ. Sustained hippocampal chromatin regulation in a mouse model of depression and antidepressant action. *Nat Neurosci* 2006;9:519–525. [PubMed: 16501568]
- Verma AS, Shapiro BH. Sex-dependent expression of seven housekeeping genes in rat liver. *J Gastroenterol Hepato* 2006;21:1004–1008.
- Weissman MM, Olfson M. Depression in women: implications for health care research. *Science* 1995;269:799–801. [PubMed: 7638596]
- Wilcoxon JS, Kuo AG, Disterhoft JF, Redei EE. Behavioral deficits associated with fetal alcohol exposure are reversed by prenatal thyroid hormone treatment: a role for maternal thyroid hormone deficiency in FAE. *Mol Psychiatry* 2005;10:961–971. [PubMed: 15940294]
- Zhou FC, Sari Y, Powrozek TA. Fetal alcohol exposure reduces serotonin innervation and compromises development of the forebrain along the serotonergic pathway. *Alcohol Clin Exp Res* 2005;29:141–149. [PubMed: 15654302]
- Zhou FC, Sari Y, Li TK, Goodlett C, Azmitia EC. Deviations in brain early serotonergic development as a result of fetal alcohol exposure. *Neurotox Res* 2002;4:337–342. [PubMed: 12829423]
- Zoeller RT, Butnariu OV, Fletcher DL, Riley EP. Limited postnatal ethanol exposure permanently alters the expression of mRNAs encoding myelin basic protein and myelin-associated glycoprotein in cerebellum. *Alcohol Clin Exp Res* 1994;18:909–916. [PubMed: 7526726]

Figure 1. Prenatal ethanol-exposed mice display increased learned helplessness

Learned helplessness behavior was assessed as described in the Methods section. Data are graphed as a function of trial number; saccharin control (SAC) and ethanol-exposed (fetal alcohol spectrum disorder, FASD) mice ($n=8$) were used in this study. *Left:* Percentage of animals that escaped. The data were fit to the equation of a hyperbola using GraphPad Prism (version 4.0); the best-fit curves are shown. The curves fitting the saccharin control and prenatal ethanol-exposed data sets were significantly different from each other [$F(2,52)=44.0$, $p<0.0001$]. *Right:* Mean $\pm$ SEM escape latencies. Comparisons of different models [polynomial first order (straight line), one phase exponential decay curve, two phase exponential decay curve, hyperbola (one site binding)] using GraphPad Prism (version 4.0) found that both data sets were best fit by a polynomial first order (straight line). Linear regression analyses of the two data sets revealed that the slopes of the two lines were not different but the intercepts were significantly different [$F(1,445)=33.0$, $p<0.01$].

Figure 2. Prenatal ethanol-exposed mice display increased immobility in the forced swim test
Time spent in immobile behaviors, (floating, twitching, kicking) in seconds for saccharin (SAC) control (n=8) and prenatal ethanol-exposed (fetal alcohol spectrum disorder, FASD) (n=8) mice during the last two minutes of the three minutes of the swim task. $t(14)=8.41$, * indicates $p<0.01$.

Figure 3. Prenatal ethanol-exposed and saccharin control mice do not differ in locomotor activity

Locomotor activity, measured as counts (beam interruptions), over a 15-min period was measured in saccharin (SAC) control (n=8) and prenatal ethanol-exposed (fetal alcohol spectrum disorder, FASD) (n=9) mice, as described in the Materials and Methods.

Figure 4. Prenatal ethanol exposure is associated with reduced BDNF levels in the medial frontal cortex, but not hippocampal formation, of mice

Homogenates of the medial frontal cortex (MFC) and hippocampal formation (HPF) were prepared from adult saccharin (SAC) (n=8) and prenatal ethanol-exposed (fetal alcohol spectrum disorder, FASD) (n=7) mice and BDNF levels were quantified by ELISA. Data were expressed relative to the wet weight of the tissue sample. Compared to BDNF levels present in the medial frontal cortex of saccharin control mice, BDNF levels were significantly reduced in the medial frontal cortex of prenatal ethanol-exposed mice: $t(13)=2.7$, * indicates $p<0.02$.

Figure 5. Prenatal ethanol exposure is associated with reduced proBDNF levels in the medial frontal cortex, but not the hippocampal formation

Postnuclear (1,000 g_{max} , 6 min) fractions were prepared from the medial frontal cortex (MFC) and hippocampal formation (HPF) of adult saccharin (SAC; $n=6$ for HPF; $n=8$ for MFC) and prenatal ethanol-exposed (fetal alcohol spectrum disorder, FASD) ($n=6$ for HPF; $n=11$ for MFC) mice. proBDNF levels were analyzed by immunoblotting (see Methods). Data are presented as arbitrary units (defined as the adjusted volume measurement obtained from the immunoblot images divided by the amount of protein loaded onto the gel). Compared to proBDNF levels in the medial frontal cortex of saccharin control mice, proBDNF levels were significantly reduced in the medial frontal cortex of prenatal ethanol-exposed mice: $t(17)=2.44$, * indicates $p<0.02$.

Figure 6. Expression of housekeeping genes, cyclophilin A, HPRT1, 18S rRNA, and β-actin, in saccharin control and prenatal ethanol-exposed adult mouse hippocampal formation and medial frontal cortex

Levels of mRNA transcripts for cyclophilin A, HPRT1, 18S rRNA, and β-actin were determined as described in the Methods. Cycle threshold (Ct) values were determined in triplicate for prenatal ethanol-exposed (fetal alcohol spectrum disorder, FASD) (n=8) and saccharin (SAC) control (n=8) samples in both the hippocampal formation (HPF) and medial frontal cortex (MFC). Statistical significance was determined using a two-tailed t-test: cyclophilin A SAC vs. FASD in the MFC, $t(14) = 5.16$; 18S rRNA SAC vs. FASD in the HPF, $t(14) = 3.64$. ** designates $p < 0.01$.

Figure 7.

A. Mouse BDNF gene structure and alternative transcripts. The upper figure, modified from Aid et al. (2007), shows the exon/intron structure of the BDNF gene with boxes representing exons and lines representing introns. The lengths (number of nucleotides) of the exons and introns are not drawn to scale. Each of the 5' exons can be spliced together with exon IX, which contains the open reading frame (shaded) encoding proBDNF. Below the gene structure are schematic representations of the splice variants analyzed in Figure 7B. *B. Levels of BDNF total (exon IX-containing) and specific mRNA transcripts in the medial frontal cortex (MFC) and hippocampal formation (HPF) of saccharin control (SAC) and prenatal ethanol-exposed (fetal alcohol spectrum disorder, FASD) mice.* Real Time PCR was performed and data were calculated as described in the Methods. Compared to SAC control (n=8), prenatal ethanol-exposed mice (n=8) mice had significantly reduced levels of exons III [MFC, $t(14)=2.8$; HPF, $t(14)=4.2$], IV [MFC, $t(14)=8.1$; HPF, $t(14)=6.2$], VI [MFC, $t(14)=8.6$; HPF, $t(14)=3.6$], and IX [MFC, $t(14)=3.7$; HPF, $t(14)=3.9$]. * indicates $p<0.02$; ** indicates $p<0.01$.

Table 1

Sequences (5'→3') of the forward and reverse primer pairs used for PCR amplification.

Target	Forward primer	Reverse primer
BDNF exon I	CCTGCATCTGTTGGGGAG	GCCTTGTCCTGGGACGTTT
BDNF exon II	GGGTTGGTATACTGGGTT	GGATGGTCATCACTCTTCTC
BDNF exon III	GGGCCGGATGCTT	CCGTGGACGTTTACTTC
BDNF exon IV	CAGAGCAGCTGCCTTGAT	GCCTTGTCCTGGGACG
BDNF exon VI	CTGCCACCACTGCCTTGG	GGATGGTCATCACTCTTC
BDNF exon IX	CAGGTGAGAAGAGTGATGACC	ATTACGCTCTCCAGAGTCCC
β-actin	CCATCATGAAGTGTGACGTGG	GTCCGCCTAGAAGCATTTGCG
Cyclophilin A	TGCTGGACCAACACAAACG	GCCTTCTTTCACCTTCCAAA
HPRT1	AAGACTTGCTCGAGATGTCATGAA	ATCCAGCAGGTCAGCAAAGAA
18S rRNA	CGGCTACCACATCCAAGGAA	CCTGTATTGTTATTTTCGTCACCT

Table 2**Maternal weights and litter sizes for saccharin (SAC) control and ethanol-consuming mice**

Maternal weights were determined at the beginning of the drinking paradigm, as well as throughout drinking, and on postpartum day 6–7. Litter sizes were determined on Day 6–7 after delivery. The data are from six separate breeding rounds and are expressed as mean $\pm$ SEM; SAC (n=47) and ethanol (n=50) dams.

Prenatal diet/exposure	Maternal weight at beginning of drinking (g)	Maternal weight Day 6–7 after birth (g)	Litter size
SAC	20.97 ($\pm$ 2.63)	29.91 ($\pm$ 2.22)	5.9 ($\pm$ 2.29)
ethanol	21.50 ($\pm$ 3.23)	29.20 ($\pm$ 3.23)	6.3 ($\pm$ 1.99)