Published as: *J Mol Biol.* 2009 October 9; 392(5): 1117–1124.

The Structure and conformation of Lys-63 linked tetraubiquitin

Ajit B. Datta[†], Greg L. Hura[#], and Cynthia Wolberger^{†,*}

[†] Department of Biophysics and Biophysical Chemistry & the Howard Hughes Medical Institute, Johns Hopkins University School of Medicine, Baltimore, MD 21205

[#] Physical Biosciences Division, Advanced Light Source, Lawrence Berkeley National Laboratory, Berkeley, CA 94720, USA

Summary

Ubiquitination involves the covalent attachment of the ubiquitin C-terminus to the lysine sidechain of a substrate protein by an isopeptide bond. The modification can comprise a single ubiquitin moiety or a chain of ubiquitin molecules joined by isopeptide bonds between the C-terminus of one ubiquitin with one of the seven lysine residues in the next ubiquitin. Modification of substrate proteins with Lys63-linked polyubiquitin plays a key non-degradative signaling role in many biological processes including DNA repair and NF- κ B activation, whereas substrates modified by lysine-48 (Lys48) linked chains are targeted to the proteasome for degradation. The distinct signaling properties of alternatively linked ubiquitin chains presumably stems from structural differences that can be distinguished by effector proteins. We have determined the crystal structure of Lys63 tetraubiquitin at a resolution of 1.96 Å and performed Small Angle X-ray scattering (SAXS) experiments and molecular dynamics (MD) simulations to probe the conformation of Lys63 tetraubiquitin in solution. The chain adopts a highly extended conformation in the crystal, in contrast with the compact globular fold of Lys48 Ub₄. Small Angle X-ray scattering (SAXS) experiments show that the tetraubiquitin chain is dynamic in solution, adopting an ensemble of conformations that are more compact than the extended form in the crystal. The results of these studies provide a basis for understanding the differences in the behavior and recognition of Lys63 polyubiquitin chains.

Protein modification by covalent conjugation of one or more ubiquitin molecules, referred as ubiquitination (or ubiquitylation), has been shown to play important regulatory roles in many cellular processes including protein degradation, transcription, cell-cycle progression, viral infection and the immune response^{1,2}. This conjugation involves a cascade of enzymatic steps that results in the covalent attachment of the C-terminal glycine residue of ubiquitin to the ϵ -NH₂ of a lysine sidechain in the substrate protein³. Ubiquitin itself can also be ubiquitinated at any of its seven lysine residues⁴, or at its amino-terminus⁵, giving rise to different poly-ubiquitin chains.

Polyubiquitin chains of different linkage types, as distinguished by the lysine residue to which each successive ubiquitin C-terminus is attached, have distinct biological functions. The best understood among these are the Lys48-linked ubiquitin chains, which target proteins for degradation by the 26S proteasome⁶. Lys63-linked ubiquitin chains, on the other hand, play

*To whom correspondence should be addressed. cwolberg@jhmi.edu.

ACCESSION NUMBERS

Coordinates and structure factors have been deposited in the Protein Data Bank with accession number 3HM3.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

non-degradative roles in different signaling pathways⁷, notably NFκB transcription activation⁸ and the DNA damage response^{9, 10}. Other types of chain linkages have also been identified in cells¹¹, including branched and mixed-linkage chains^{12, 13}. Although the cellular roles of these “atypical” ubiquitin chains are yet to be studied in detail, they have been implicated in a few biological processes¹⁴. For example, K33/K29 mixed chains have been shown to regulate AMPK-related kinase activation while K29-linked poly-Ubs have been shown to play role in lysosomal degradation of proteins^{15, 16}.

The distinct roles played by different polyubiquitin chains indicate that effector molecules can distinguish among chains of differing linkage types. For example, the NZF domain of TAB2 binds specifically to Lys63-linked chains¹⁷, the ataxin3 UIM repeat preferentially binds to Lys48-linked chains¹⁸, and the UBAN motif of NEMO recognizes linear head-to-tail ubiquitin chains¹⁹. A thorough understanding of the molecular details of ubiquitin-mediated signaling therefore requires elucidation of the structures of different poly-ubiquitin chains and also their complexes with cognate partners.

The crystal structure of Lys48-linked tetraubiquitin determined at neutral pH showed it to adopt a compact globular conformation²⁰. That conformation could not be adopted by Lys63-linked tetraubiquitin as judged by the relation between the Lys63 side chains and the ubiquitin C-termini. Indeed, NMR studies^{21, 22} and an x-ray crystal structure¹⁷ of Lys63-linked diubiquitin show an open structure with limited contacts between ubiquitin monomers, as compared with the closed conformation of Lys48-linked diubiquitin.

In order to gain insight into the structure and topology adopted by longer Lys63-linked polyubiquitin chains, we determined the 1.96 Å resolution crystal structure of Lys63-linked tetraubiquitin. The crystals contain one tetra-ubiquitin chain per asymmetric unit and show a highly extended linear arrangement of ubiquitin monomers. The extended arrangement showed no inter-subunit interactions, in agreement with NMR results. The extended conformation of the Lys63 tetraubiquitin in the crystal suggests that the chain would likely adopt other conformations in solution. To further explore this conformational flexibility of the Lys63-linked tetraubiquitin chains in solution, we carried out small angle X-ray scattering measurements (SAXS) of Lys63 tetraubiquitin and used molecular dynamics simulations to deduce the ensemble of structures adopted by Lys63 tetraubiquitin in solution. We find that Lys63 tetraubiquitin adopts conformations in solution that are more compact than the extended form seen in the crystal, yet do not exhibit any specific inter-subunit interactions. These studies provide a detailed understanding of the structure and dynamics of Lys63 tetraubiquitin and how these chains could be recognized in the cell.

Experimental Details

Lys63-linked tetraubiquitin chain was synthesized using two ubiquitin mutants (K48R-K63R and Ub-D77) following Pickart and Raasi²³. All the enzymes required for this synthesis viz., E1 (human), Ubc13/MMS2 (Yeast) and Uuh1 (yeast) were also expressed and purified as recombinant proteins in *Escherichia coli*. After the chain of required length was synthesized, it was subjected to a final step of purification using size-exclusion chromatography in Superdex 75 prep-grade 26/60 column. The protein was then dialyzed in 10 mM Tris-HCl buffer (pH 7.6) containing 25 mM NaCl, overnight at 4°C. It was then concentrated using Vivaspin concentrators and stored at -80°C in small aliquots.

Crystals of Lys63 Ub4 were initially obtained at 20°C in hanging drop setups from sparse matrix screening with buffer 1 from Wizard II (Emerald Biosciences). The condition was then optimized to get fewer and better crystals. The best crystals were obtained using 4–6% PEG 3000 + 100 mM Na-acetate (pH 4.6–5.0) + 10 μM Zn acetate. However all the crystals were found to diffract poorly (best 3.5 Å) and also showed very high mosaicity and B-factor after

data processing. The diffraction improved substantially upon dehydration of the crystals by step-wise increase of PEG 3000 to 25% over a period of 4–5 days and a 1.95 Å dataset was collected from frozen crystals at 100 K at APS, Argonne, IL at the fixed wavelength beamline 14-BMC.

The data was processed in the space group $P2_13$ using the program HKL2000²⁴. The structure was solved by molecular replacement using the program MOLREP²⁵ from CCP4 Suite²⁶. The initial model was subjected to iterative building and refinement cycles using COOT²⁷ and REFMAC5²⁸. All the side chains were built first followed by the completion of the chain. Iso-peptide bonds were created towards the end with addition of water molecules.

SAXS data were collected at the SIBYLS beamline (beamline 12.3.1) at the Advanced Light Source in Lawrence Berkeley Labs. Data were collected from the sample at concentrations of 3, 6 and 9 mg/ml and no concentration dependence was observed. The sample buffer contained 10 mM Tris-HCl (pH 7.6), 50 mM NaCl and 0.5 mM EDTA at 10°C. Five and half second exposures were merged to create a final scattering profile. $P(r)$ functions were determined from the scattering profiles using GNOM²⁹. Scattering profiles were calculated from crystal structures using CRY SOL³⁰. The MES program was run as described³¹ with 3 independent runs per radius of gyration ranging from 20 to 60 Å at 1 Å steps.

Crystal Structure of Lys63 tetraubiquitin

Crystals of Lys63 tetraubiquitin formed in spacegroup $P2_13$ and contain one chain per asymmetric unit. Crystals obtained after initial rounds of optimization typically diffracted to 3.5–4.0 Å resolution. However, the diffraction improved significantly upon gradual dehydration of the crystals with step-wise increases of the precipitant concentration, yielding crystals diffracting to 1.96 Å resolution. The structure was solved by molecular replacement using the structure of monomeric ubiquitin³² as the search model. Data collection and refinement statistics are presented in Table 1.

The crystal structure shows a linear arrangement of the Lys63-linked ubiquitin monomers (Figure 1a & b) that is starkly different from the compact globular fold adopted by Lys48-linked tetra-ubiquitin (Figure 1c). Instead, the monomers in the Lys63-linked chain are joined by well-ordered linkages but do not contact each other. The absence of interactions between the individual ubiquitin monomers of the Lys63 Ub4 agrees with the previously reported NMR results, where no stable inter-subunit interactions could be observed in Lys63-linked di- and tetra-ubiquitin chains^{21, 22} as well as with the extended structure of crystallized di-ubiquitin¹⁷. The extended conformation of the tetraubiquitin chain and the absence of intra-chain contacts that could stabilize the observed conformation suggested that Lys63-linked tetraubiquitin in solution would likely be flexible and adopt other conformations in addition to that seen in the crystal.

We also note that the globular regions of each of the four ubiquitin monomers in the Lys63 tetraubiquitin chain alternate between two conformations that differ in the loop region comprising residues 6 through 12. Inclusion of this region during superposition of the globular region of the ubiquitin monomers (residues 1 through 70) increase the root mean square deviation to ~0.85 Å from ~0.4 Å. The reason for this difference remains unclear and could not be attributed to the differences in crystal packing environments of each monomer.

Recognition Surface of Lys63 Ubiquitin

Ubiquitin has a conserved hydrophobic patch comprising residues Leu8, Ile44 and Val70 that has been shown to be important for function³³. These residues mediate the interaction of ubiquitin with E2 enzymes³⁴ and also with ubiquitin binding domains (reviewed in³⁵). The

importance of these hydrophobic residues in the Lys48-linked ubiquitin chain recognition has been studied in detail^{36, 37, 38}. In addition, the hydrophobic patch has also been implicated in interactions with Lys63 linked chain interactions³⁹ and is shown to participate in the linear diubiquitin recognition by NEMO⁴⁰. In contrast with Lys48 tetraubiquitin, in which the hydrophobic patches from two adjacent monomers coalesce to form a single surface (Figure 1c), the hydrophobic patches in the crystal structure of Lys63 tetraubiquitin are exposed in all four ubiquitin monomers (Figure 1b). We also note that the hydrophobic patches of Lys63 monomers do not lie on the same surface of the molecule, but are rotated roughly 90 degrees relative to one another in successive monomers. This particular arrangement is probably favored in the crystal, where the molecules are conformationally “locked” in one particular orientation due to crystal packing. However, we note that one of the solution structures presented by Varadan et al.²² does also show this orthogonal positioning of the ubiquitin hydrophobic surface in successive monomers along with alternate arrangements. Given the presumed flexibility in the linkages that join the ubiquitin moieties in Lys63 chain, the individual monomers would probably be free to undergo rotational movements that would change the relative orientation of these patches.

Despite the strong tendency of ubiquitin to associate via the hydrophobic patches, we were surprised to find that these patches do not take part in crystal packing of the Lys63 Ub4 chains. Instead, the crystal contacts are formed by the interaction of an ubiquitin monomer in one chain with the isopeptide linkage of another chain in the crystal (Figure 2). This crystal packing interaction could potentially reflect a mode of interaction between polyubiquitin chains, or of ubiquitin-like domains with Lys-63-linked chains. Several proteins including IKK α and IKK β have been found to contain Ubiquitin like domains (UBLs). These UBL domains could possibly interact with Lys63 chains in a manner similar to that seen in the crystal. However, to our knowledge, no such interactions have been reported to exist in the literature.

Small angle X-ray scattering (SAXS) studies of Lys63 Ub4 in solution

Since the extended conformation of the Lys63 tetraubiquitin is clearly influenced by lattice contacts, we used small angle X-ray scattering (SAXS) combined with molecular dynamics (MD) simulations to explore the structure of Lys63 tetraubiquitin in solution. SAXS is a powerful tool to study the conformation of macromolecules in solution, particularly when a high resolution crystal structure is available⁴¹. We measured the x-ray scattering profile of Lys63 tetraubiquitin in solution (Fig. 3a) and derived the pair distribution function, $P(r)$, from the scattering profile to determine the distribution of interatomic vectors in the tetraubiquitin chain (Fig. 3b). As shown in Figure 3a, the measured x-ray scattering profile does not fit the theoretically calculated profiles from the crystal structures of either Lys63 tetraubiquitin (this study) or of Lys48 tetraubiquitin²⁰. This indicates that Lys63 tetraubiquitin adopts a distinct conformation in solution that is more compact than that seen in the crystal.

The interatomic vectors calculated in the pair distribution function (Fig. 3b) show that the maximum dimension of Lys63 tetraubiquitin in solution (D_{max}) is $118 \pm 2 \text{ \AA}$ with a radius of gyration (R_g) $32.3 \text{ \AA}$, which is considerably smaller than the D_{max} ($154 \text{ \AA}$) and R_g ($44 \text{ \AA}$) values calculated from the crystal structure. This apparent reduction of D_{max} and radius of gyration indicates that Lys63-linked tetraubiquitin is more compact in solution than in the crystal. At the same time, the peaks in the experimental $P(r)$ also show that Lys63 tetraubiquitin adopts a much more extended structure than the crystal structure of Lys48 tetraubiquitin (Figs. 3a & b). Interestingly, the second maximum in $P(r)$ of $\sim 41 \text{ \AA}$ corresponding to the average distance between any two consecutive ubiquitin monomers within the Lys63 polyubiquitin chain closely matches the calculated value from the crystal structure (Fig. 3b). However, the third maximum, which corresponds to the distances between first and third ubiquitin monomer or between the second and fourth, is considerably shorter in the observed profile ($63 \text{ \AA}$) as

compared to the one calculated from the crystal structure. The combination of a large Dmax but minor third and fourth maxima suggested that the Lys63 ubiquitin chains most likely exist as an ensemble of conformations at larger length scales.

Further analysis of the solution conformation of Lys63 tetraubiquitin was done using the program Minimum Ensemble Search (MES)³¹. This program finely samples the conformational space of the macromolecule using molecular dynamics (MD) simulations at high temperatures, where only covalent bonds and van der Waals interactions are taken into consideration. A genetic algorithm then identifies the minimal ensemble from all these conformations that best fits the data. The result shows that no single conformation from the MD simulations adequately explains the data. Instead, an optimal fit could be obtained with an ensemble of a minimum of three conformations (Figure 3c & 4). Inclusion of more conformations in the ensemble did not lead to any significant improvement in the fit, as measured by chi-squared values (data not shown). The first two dominant conformations (represented by 69% and 17% in the ensemble) (Figure 4a & b) preserve the overall linearity observed in the crystal structure but are notably more compact than the conformation in the crystal. The ensemble also contains a minor population (13% in the ensemble) (Figure 4c) that is highly compact as compared to the crystal structure.

Though the resolution of SAXS data cannot be used to determine the inter-subunit orientations accurately, we note that, in all of these MES conformations, the conserved hydrophobic interaction surface (colored red) of the ubiquitin monomers do not interact with each other and are either completely or partially exposed. This probably suggests that Lys63 chains are likely to be found in solution conformations that leave the hydrophobic patches exposed and would therefore be readily accessible for interaction with cognate proteins.

In summary, our results show that Lys63-linked polyubiquitin forms a relatively open conformation, in agreement with the existing biochemical and biophysical data. The elongated structures of the K63 tetraubiquitin, stabilized by a criss-crossed arrangement of chains within the crystal lattice, do not belong to any closed symmetry group. Thus, addition of more ubiquitin monomers to either the proximal or distal end of the chain can occur without requiring any rearrangement of the structure. Although the highly extended crystal structure of Lys63 tetraubiquitin agrees with predicted models based on diubiquitin structure¹⁷, the longer chain is notably more collapsed in solution.

Nevertheless, none of the representative solution conformations or the crystal structure buries the conserved hydrophobic patch of any of the ubiquitin monomers. This indicates that these patches remain available for interactions with other proteins.

The requirement of an ensemble instead of a single conformation to explain the SAXS data point towards the fact that Lys63 ubiquitin chains with four or more monomers do not form a single well-defined structure in solution. Instead, tetraubiquitin and presumably longer K63 chains exist in a dynamic equilibrium of many different conformations. Upon binding to a cognate protein, Lys63 chains would probably adopt one specific topology depending on the binding partner. This fluidity in the structure of Lys63 polyubiquitin may be important for its biology, where these chains are recognized by proteins with potentially different structures in different signaling pathways.

Moreover, the conformational space sampled by the free Lys63 polyubiquitin can be restricted upon attachment to substrate proteins, each of which could have a unique influence on the Lys63 polyubiquitin chain behavior. All these considered together point towards a highly plastic nature of Lys63-linked polyubiquitin mediated signaling mechanisms, which could account for the diverse roles played by this type of polyubiquitin modification in multiple signaling pathways.

Acknowledgments

We thank Michal Hammel from LBNL for help with the implementation of MES. Supported by the Howard Hughes Medical Institute (A.B.D and C.W.) and NCI SBDR grant # CA92584 (G.H.).

References

1. Haglund K, Dikic I. Ubiquitylation and cell signaling. *EMBO J* 2005;24:3353–3359. [PubMed: 16148945]
2. Mukhopadhyay D, Riezman H. Proteasome-independent functions of ubiquitin in endocytosis and signaling. *Science* 2007;315:201–205. [PubMed: 17218518]
3. Hershko A, Ciechanover A. The ubiquitin system. *Annu Rev Biochem* 1998;67:425–479. [PubMed: 9759494]
4. Pickart CM, Fushman D. Polyubiquitin chains: polymeric protein signals. *Curr Opin Chem Biol* 2004;8:610–616. [PubMed: 15556404]
5. Kirisako T, Kamei K, Murata S, Kato M, Fukumoto H, Kanie M, Sano S, Tokunaga F, Tanaka K, Iwai K. A ubiquitin ligase complex assembles linear polyubiquitin chains. *EMBO J* 2006;25:4877–4887. [PubMed: 17006537]
6. Pickart CM, Cohen RE. Proteasomes and their kin: proteases in the machine age. *Nat Rev Mol Cell Biol* 2004;5:177–187. [PubMed: 14990998]
7. Sun L, Chen ZJ. The novel functions of ubiquitination in signaling. *Curr Opin Cell Biol* 2004;16:119–126. [PubMed: 15196553]
8. Chen ZJ. Ubiquitin signalling in the NF-kappaB pathway. *Nat Cell Biol* 2005;7:758–765. [PubMed: 16056267]
9. Hofmann RM, Pickart CM. Noncanonical MMS2-encoded ubiquitin-conjugating enzyme functions in assembly of novel polyubiquitin chains for DNA repair. *Cell* 1999;96:645–653. [PubMed: 10089880]
10. Stelter P, Ulrich HD. Control of spontaneous and damage-induced mutagenesis by SUMO and ubiquitin conjugation. *Nature* 2003;425:188–191. [PubMed: 12968183]
11. Xu P, Duong DM, Seyfried NT, Cheng D, Xie Y, Robert J, Rush J, Hochstrasser M, Finley D, Peng J. Quantitative proteomics reveals the function of unconventional ubiquitin chains in proteasomal degradation. *Cell* 2009;137:133–145. [PubMed: 19345192]
12. Kim HT, Kim KP, Lledias F, Kisselev AF, Scaglione KM, Skowrya D, Gygi SP, Goldberg AL. Certain pairs of ubiquitin-conjugating enzymes (E2s) and ubiquitin-protein ligases (E3s) synthesize nondegradable forked ubiquitin chains containing all possible isopeptide linkages. *J Biol Chem* 2007;282:17375–17386. [PubMed: 17426036]
13. Li W, Ye Y. Polyubiquitin chains: functions, structures, and mechanisms. *Cell Mol Life Sci* 2008;65:2397–2406. [PubMed: 18438605]
14. Ikeda F, Dikic I. Atypical ubiquitin chains: new molecular signals. 'Protein Modifications: Beyond the Usual Suspects' review series. *EMBO Rep* 2008;9:536–42. [PubMed: 18516089]
15. Al-Hakim AK, Zagorska A, Chapman L, Deak M, Pegg M, Alessi DR. Control of AMPK-related kinases by USP9X and atypical Lys(29)/Lys(33)-linked polyubiquitin chains. *Biochem J* 2008;411:249–260. [PubMed: 18254724]
16. Chastagner P, Israël A, Brou C. Itch/AIP4 mediates Deltex degradation through the formation of K29-linked polyubiquitin chains. *EMBO Rep* 2006;7:1147–1153. [PubMed: 17028573]
17. Komander D, Reyes-Turcu F, Licchesi JDF, Odenwaelde P, Wilkinson KD, Barford D. Molecular discrimination of structurally equivalent Lys 63-linked and linear polyubiquitin chains. *EMBO Rep* 2009;10:466–473. [PubMed: 19373254]
18. Sims JJ, Cohen RE. Linkage-specific avidity defines the lysine 63-linked polyubiquitin-binding preference of rap80. *Mol Cell* 2009;33:775–783. [PubMed: 19328070]
19. Tokunaga F, Sakata S, Saeki Y, Satomi Y, Kirisako T, Kamei K, Nakagawa T, Kato M, Murata S, Yamaoka S, Yamamoto M, Akira S, Takao T, Tanaka K, Iwai K. Involvement of linear polyubiquitylation of NEMO in NF-kappaB activation. *Nat Cell Biol* 2009;11:123–132. [PubMed: 19136968]

20. Eddins MJ, Varadan R, Fushman D, Pickart CM, Wolberger C. Crystal structure and solution NMR studies of Lys48-linked tetraubiquitin at neutral pH. *J Mol Biol* 2007;367:204–11. [PubMed: 17240395]
21. Tenno T, Fujiwara K, Tochio H, Iwai K, Morita EH, Hayashi H, Murata S, Hiroaki H, Sato M, Tanaka K, Shirakawa M. Structural basis for distinct roles of Lys63- and Lys48-linked polyubiquitin chains. *Genes Cells* 2004;9:865–875. [PubMed: 15461659]
22. Varadan R, Assfalg M, Haririnia A, Raasi S, Pickart C, Fushman D. Solution conformation of Lys63-linked di-ubiquitin chain provides clues to functional diversity of polyubiquitin signaling. *J Biol Chem* 2004;279:7055–7063. [PubMed: 14645257]
23. Pickart CM, Raasi S. Controlled synthesis of polyubiquitin chains. *Meth Enzymol* 2005;399:21–36. [PubMed: 16338346]
24. Otwinowski Z, Minor W. Processing of X-ray Diffraction Data Collected in Oscillation Mode. *Methods in Enzymology* 1997;276:307–326.
25. Vagin A, Teplyakov A. MOLREP: an automated program for molecular replacement. *J Appl Cryst* 1997;30:1022.
26. CCP4. The CCP4 Suite: Programs for Protein Crystallography. *Acta Crystallographica Section D, Biological Crystallography* D50:760–763.
27. Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. *Acta Crystallogr D Biol Crystallogr* 2004;60:2126–2132. [PubMed: 15572765]
28. Murshudov GN, Vagin AA, Dodson EJ. Refinement of macromolecular structures by the maximum-likelihood method. *Acta Crystallogr D Biol Crystallogr* 1997;53:240–255. [PubMed: 15299926]
29. Konarev P, Volkov V, Sokolova A, Koch M, Svergun D. PRIMUS: a Windows PC-based system for small-angle scattering data analysis. *Journal of Applied Crystallography* 2003;36:1277–1282.
30. Svergun D, Barberato C, Koch M. CRY SOL - a Program to evaluate X-ray Solution Scattering of Biological Macromolecules from Atomic Coordinates. 1995;28:768–773.
31. Pelican M, Hura GL, Hammel M. Structure and Flexibility within proteins as identified through small angle X-ray scattering. *General Physiology and Biophysics* 2009;28:174–189. [PubMed: 19592714]
32. Vijay-Kumar S, Bugg CE, Cook WJ. Structure of ubiquitin refined at 1.8 Å resolution. *J Mol Biol* 1987;194:531–544. [PubMed: 3041007]
33. Thrower JS, Hoffman L, Rechsteiner M, Pickart CM. Recognition of the polyubiquitin proteolytic signal. *EMBO J* 2000;19:94–102. [PubMed: 10619848]
34. Eddins MJ, Carlile CM, Gomez KM, Pickart CM, Wolberger C. Mms2-Ubc13 covalently bound to ubiquitin reveals the structural basis of linkage-specific polyubiquitin chain formation. *Nat Struct Mol Biol* 2006;13:915–920. [PubMed: 16980971]
35. Hicke L, Schubert HL, Hill CP. Ubiquitin-binding domains. *Nat Rev Mol Cell Biol* 2005;6:610–621. [PubMed: 16064137]
36. Beal R, Deveraux Q, Xia G, Rechsteiner M, Pickart C. Surface hydrophobic residues of multiubiquitin chains essential for proteolytic targeting. *Proc Natl Acad Sci USA* 1996;93:861–866. [PubMed: 8570649]
37. Beal RE, Toscano-Cantaffa D, Young P, Rechsteiner M, Pickart CM. The hydrophobic effect contributes to polyubiquitin chain recognition. *Biochemistry* 1998;37:2925–2934. [PubMed: 9485444]
38. Raasi S, Orlov I, Fleming KG, Pickart CM. Binding of polyubiquitin chains to ubiquitin-associated (UBA) domains of HHR23A. *J Mol Biol* 2004;341:1367–1379. [PubMed: 15321727]
39. Ea C, Deng L, Xia Z, Pineda G, Chen ZJ. Activation of IKK by TNF α requires site-specific ubiquitination of RIP1 and polyubiquitin binding by NEMO. *Mol Cell* 2006;22:245–257. [PubMed: 16603398]
40. Rahighi S, Ikeda F, Kawasaki M, Akutsu M, Suzuki N, Kato R, Kensche T, Uejima T, Bloor S, Komander D, Randow F, Wakatsuki S, Dikic I. Specific recognition of linear ubiquitin chains by NEMO is important for NF- κ B activation. *Cell* 2009;136:1098–109. [PubMed: 19303852]
41. Putnam CD, Hammel M, Hura GL, Tainer JA. X-ray solution scattering (SAXS) combined with crystallography and computation: defining accurate macromolecular structures, conformations and assemblies in solution. *Q Rev Biophys* 2007;40:191–285. [PubMed: 18078545]

Figure 1. Crystal Structure of Lys63 linked tetraubiquitin

(a) Ribbon diagram and (b) Surface representation of Lys63 Ub4 crystal structure showing the highly extended chain conformation. The proximal ubiquitin is shown in cyan while the distal one is in yellow. In (b) the hydrophobic patches of individual ubiquitin monomers consisting of residues Leu8, Ile44 and Val70, which are important for interacting with cognate proteins are shown in red.

(c) Crystal structure of Lys48 tetraubiquitin (PDB ID 26OV) ²⁰. The structure of Lys48 is highly compact and globular. The hydrophobic patches from ubiquitin monomer can be seen to interact with each other partially burying the interaction surface. The coloring scheme has been kept identical to Lys63 tetraubiquitin.

Figure 2.

Crystal packing interactions of Lys63-linked tetraubiquitin showing two ubiquitin monomers from two different chains (surface representation in green and yellow) interacting at the isopeptide linkage of a third chain (cartoon representation). The isopeptide bond between the Lys63 of proximal ubiquitin (in red) and Gly76 of distal ubiquitin (in blue) are represented with sticks.

Figure 3. Small angle x-ray scattering (SAXS) of Lys63 tetraubiquitin in solution

(a) A comparison of the experimental scattering profile (black) with theoretically calculated profiles from the crystal structures of Lys63 Ub4 (green), Lys48 Ub4 (red) and the minimal ensemble (orange). The minimal ensemble (orange) of Lys63 tetraubiquitin was identified using the minimal ensemble search (MES) program which best fits the experimental data (black).

(b) The experimental electron pair distribution function ($P(r)$) (in black). The curve reveals a long maximum dimension with smeared 3rd and 4th Ub-Ub peaks suggesting an ensemble of distances on this length scale. In contrast, the theoretically calculated $P(r)$ for Lys48 tetramer (red) shows one broad peak indicating all Ub-Ub distances to be similar, while the $P(r)$ calculated Lys63 structure (in green) shows 4 major peaks that corresponds to all major Ub-Ub distances.

(c) Searching for the minimal ensemble using the Minimal Ensemble Search (MES) program³¹. The radius of gyration (R_g) of each model is plotted against the Chi^2 agreement to the experimental data. All the conformations (13,200) from which MES models were extracted are shown as grey circles. A single best model was identified (cyan) while the 3 conformations making up the MES model (orange squares) improves the Chi^2 agreement by 11% (orange line vs minimum of envelope formed by single conformations). The parameters for the two crystal structures are also shown (green and red circles).

Figure 4. Ensemble conformations Lys63 tetraubiquitin in solution obtained with minimal ensemble search (MES)

Three conformations of Lys63 tetraubiquitin (a), (b) and (c) that makes up the MES model are shown along with their respective percentages. Three hydrophobic residues (L8, I44 & V70) on the surface of the ubiquitins those are responsible for their interactions with cognate proteins are depicted in red.

Table 1**Data Collection & Refinement Statistics of K-63 Ub4 Crystals**

Data Collection Statistics	
Space Group	P2 ₁ 3
Cell dimensions	a= b=c= 106.95
Resolution (Å)	50-1.96 (2.01-1.96)
Rmerge (%)	8.1 (90)
Completeness (%)	99.6 (100)
Redundancy	42.9 (43.1)
I/sig(I)	54 (6)
Refinement	
R/R free (%)	19/24
No. of Atoms	
Protein	5208
Zn ²⁺	6
Water	238
r.m.s deviations	
Bond (Å)	0.027
Angles (degrees)	1.139

Values in the parenthesis denote the highest resolution shell