Episome-mediated Transfer of Drug Resistance in *Enterobacteriaceae*

X. Restriction and Modification of Phages by f_i^- R Factors

TSUTOMU WATANABE, TOSHIYA TAKANO, TOSHIHIKO ARAI,
HIROSHI NISHIDA, AND SACHIKO SATO

Department of Microbiology, Keio University School of Medicine, Tokyo, Japan

Received for publication 23 May 1966

ABSTRACT

WATANABE, TSUTOMU (Keio University, Tokyo, Japan), TOSHIYA TAKANO, TOSHIHIKO ARAI, HIROSHI NISHIDA, AND SACHIKO SATO. Episome-mediated transfer of drug resistance in *Enterobacteriaceae*. X. Restriction and modification of phages by f_i^- R factors. J. Bacteriol. 92:477-486. 1966.—An f_i^- R factor, which restricts phages λ , T1, and T7 without modifying them, was found to restrict and not to modify an F^- -specific phage, W-31, in *Escherichia coli* K-12, but not to restrict phage P-22 in *Salmonella typhimurium* LT-2, whereas other f_i^- R factors restricted and modified P-22 but not W-31; f_i^+ R factors did not restrict these phages. Transduction and lysogenization with phages λ and P-22 were reduced by these f_i^- R factors in K-12 and LT-2, respectively, and the transducing phages λ and P-22 were modified by these f_i^- R factors. Spontaneous as well as ultraviolet-induced production of phage P-22 and zygotic induction of phage λ were not significantly affected by any R factor. Injection of the nucleic acids of phages T1 and λ was not affected by R factors, but the injected phage nucleic acids were rapidly broken down in the bacteria carrying f_i^- R factors. The nucleic acids of the modified phages were not broken down in these bacteria. It was assumed from these results that the mechanism of restriction of phages by f_i^- R factors is due to the breakdown of the injected phage nucleic acids by a deoxyribonuclease(s), presumably located near the cell surface in the cells carrying f_i^- R factors. The deoxyribonuclease(s), formed in the cells carrying the nonmodifying f_i^- R factor, is considered to be different from that synthesized in the cells carrying the modifying f_i^- R factors. It was further shown that the average burst sizes of the unmodified as well as modified phages are slightly reduced by the presence of the f_i^- R factors.

We reported in a previous paper (31) that naturally occurring R factors can be classified into two types, f_i^+ and f_i^- . The term f_i is an abbreviation for fertility inhibition, and f_i^+ and f_i^- mean, respectively, the presence and absence of suppression of the function of the sex factor, F, of *Escherichia coli* K-12. The function of F that is suppressed by the f_i^+ R factors is possibly the formation of "F-pili" (7, 8, 21, 22, 30). We found that naturally occurring f_i^+ and f_i^- R factors belong to different groups in their superinfection immunity patterns, and also that some phages are restricted and modified by f_i^- R factors but not by f_i^+ R factors (31). The terms "restriction" and "modification" will be used as defined by Arber (3).

The results of our studies on the restriction and modification of phages by f_i^- R factors, reported

previously, may be summarized as follows. (i) The f_i^- R factors reduced the efficiency of plating (EOP) of phages λ and T1 in *E. coli* K-12, but f_i^+ R factors did not have this inhibitory action. One of the f_i^- R factors reduced the EOP of phage T7 as well. (ii) Phages λ and T1 underwent modifications in the hosts carrying some f_i^- R factors. (iii) At least two types of f_i^- R factors were recognized by their type of restriction and modification of these phages. (iv) Transduction of the ability to ferment galactose (gal^+) with Hft lysates of λ was reduced by the f_i^- R factors in the recipient. (v) Spontaneous production, as well as ultraviolet induction, of λ was not affected by any R factors. (vi) Adsorption of λ and T1 was not altered by the presence of any R factors.

We concluded from these results that the suppression of progeny formation of these phages by

f^- R factors is due to the inhibition of some step(s) after adsorption of the phages to the restrictive bacteria.

The above findings on the restriction and modification of phages by f^- R factors have been extended to an F^- -specific phage, W-31 (32), in *E. coli* K-12 and to phage P-22 in *Salmonella typhimurium* LT-2. In other systems of restriction of phages, it has been shown that the phage nucleic acids undergo breakdown shortly after injection into restrictive host cells (12, 14, 20, 28). We have also studied the injection and breakdown of the phage nucleic acids in bacteria carrying f^- R factors. In parallel with our studies on the restriction and modification of phages by f^- R factors, Yoshikawa, Miyake, and Wakabayashi (33) found that the formation of gal^+ recombinants in the cross of a gal^+ λ^+ Hfr strain of *E. coli* K-12 with a gal^- λ^- strain is not altered by the presence of f^- R factors in the F^- recipient, and concluded that zygotic induction of phage λ is not suppressed by the f^- R factors in the F^- recipient. We have also studied the zygotic induction of phage λ with an F^- recipient carrying various R factors. These results will be reported in the present paper.

MATERIALS AND METHODS

Culture media and drugs used were those described in previous papers (29, 31).

Strains W3102 (gal^- λ^- F^-), CSH-2 ($met F^-$), and Hfr H 3000 (B_1^- λ^+), all of which are substrains of *E. coli* K-12, were used. W3102 ($str-r$) and W3102 ($\lambda-r$ $str-r$), streptomycin-resistant and λ - and streptomycin-resistant derivatives of W3102, respectively, were used as well. *S. typhimurium* LT-2 and its derivatives *cysD-36* (cystine-requiring) and *tryD cysB-12 str-r* (tryptophan- and cystine-requiring and streptomycin-resistant) were also used.

Phage λ (wild type), phage W-31 (32), and phage P-22 (wild type) were employed.

The R factors listed in Table 1 are the same as those described in a previous paper (31).

General phage techniques followed were those described by Adams (1), except for those which will be specified in the text.

The method for transduction of gal^+ with Hft lysates of phage λ , described in a previous paper (31), was essentially similar to the original method of Morse, Lederberg, and Lederberg (24).

The method of transduction of chromosomal genes with phage P-22 in *S. typhimurium* LT-2 was the same as that described by Demerec and Hartman (11).

The conditions of ultraviolet induction of phage P-22 in P-22-lysogenic *S. typhimurium* LT-2 were the same as those used in the induction of phage λ in *E. coli* K-12 (31).

The methods of study of lysogenization frequencies were as follows. Penassay Broth (Difco) cultures of *E. coli* W3102 and *S. typhimurium* LT-2 were infected with phages λ and P-22, respectively, at various multi-

TABLE 1. *R* factors employed

Strain no. of R factor	Drug resistance markers ^a	f_i^b
N-1	<i>sul-r str-r tet-r</i>	—
N-3	<i>sul-r str-r tet-r</i>	—
R-15	<i>sul-r str-r</i>	—
222	<i>sul-r str-r chl-r tet-r</i>	+
N-6	<i>sul-r str-r tet-r</i>	+
N-9	<i>sul-r str-r tet-r</i>	+

^a The symbols *sul-r*, *str-r*, *chl-r*, and *tet-r* represent resistance to sulfonamide, streptomycin, chloramphenicol, and tetracycline, respectively.

^b Presence (+) and absence (—) of suppression of the function of *F* of *Escherichia coli* K-12; f_i is the abbreviation for fertility inhibition.

plicities of infection (MOI) at 37 C. The infected cultures were aerated at 37 C for 10 min, diluted 1:100 in ice-cold Penassay Broth, and washed three times with saline in a refrigerated centrifuge to remove free phage particles. The washed, infected bacteria were diluted with saline and plated on nutrient agar for colony isolation. After overnight incubation of the inoculated plates, the developed colonies were replica-plated onto soft nutrient agar layers seeded with indicators of the phages on the top of ordinary nutrient agar. After the replica plates were incubated overnight at 37 C, colonies with surrounding lytic zones were regarded as lysogenic and the others as nonlysogenic.

The method of study of zygotic induction of phage λ was the same as that described by Jacob and Wollman (16-18). The only difference was that we employed W3102 ($\lambda-r$ $str-r$) carrying various R factors as recipients in the crosses with Hfr H 3000 (λ^+).

The method of study of injection of phage nucleic acids was as follows. P^{32} -labeled phages were prepared mainly by the procedure of Hershey and Chase (15). W3102 was grown with aeration in radioactive H medium (containing P^{32} at a specific activity of 15 mc/mg of P) and infected with phage T1 at a MOI of 0.01. For preparing radioactive phage λ , CSH-2 was grown in a similar radioactive H medium and irradiated with ultraviolet light for phage induction. After almost complete clearing of the cultures, the lysates were centrifuged at 8,000 rev/min for 30 min. Pancreatic deoxyribonuclease (Sigma Chemical Co., St. Louis, Mo.) and pancreatic ribonuclease (Sigma) were added to the supernatant fractions (10 and 25 μ g/ml, respectively); the preparations were incubated at 37 C for 90 min, for digestion of the free nucleic acids, and were then centrifuged at 30,000 rev/min for 60 min. The sediment was suspended in tris(hydroxymethyl)amino-methane (Tris)-magnesium buffer (pH 7.5) containing 10 mM Tris-HCl and $MgCl_2$. (The volume of buffer used for suspension was 5% of the volume of the original lysates.) P^{32} -labeled phages in these suspensions were purified by sucrose density gradient centrifugation (5 to 20% sucrose in Tris-magnesium buffer) at 17,500 rev/min for 60 min (9, 25). Injection of phage nucleic acids was studied by the procedure of

Watanabe and Okada (32), a minor modification of the original method of Hershey and Chase (15).

The method used to study the breakdown of the injected phage nucleic acids was similar to that of Dussoix and Arber (12). Bacteria were grown in Lennox broth at 37 C with aeration, and the exponentially growing cells were centrifuged and resuspended in fresh Lennox broth to about 10^9 cells per milliliter. To 4.9 ml of this bacterial suspension was added 0.1 ml of the radioactive phage suspension with a titer of 10^9 plaque-forming units/ml. The phage-infected bacteria were aerated at 37 C, and 0.7-ml samples were taken periodically and added to equal volumes of 5% cold trichloroacetic acid. These samples were kept in an ice bath for 30 min, and were then centrifuged in the cold at 2,500 rev/min for 20 min. The radioactivity of heat-dried samples of the supernatant fluids was determined. The sediments were dissolved in 2 N NH_4OH , heat-dried, and tested for radioactivity. In parallel with these determinations, the samples taken at 3 min after infection were centrifuged at 8,000 rev/min, and the radioactivity of the supernatant fluids was determined.

RESULTS

Phage W-31, which was grown on W3102 (R^-), was plated on W3102 strains with and without R factors. The EOP and plaque size of W-31 were slightly reduced only on W3102 (N-1); if we take the titer of the phage on W3102 (R^-) as 1, the EOP of this phage on W3102 (N-1) was about 5×10^{-1} . Modification of this phage did not occur upon its growth on W3102 (N-1).

The frequency of lysogenization with phage λ was strongly reduced by the presence of f_i^- R factors in W3102 (Table 2). Surviving bacteria were about equal in number in all of the phage-infected W3102 strains with and without R factors.

Lft lysates of phage λ , obtained by ultraviolet irradiation of CSH-2, were used for transducing gal^+ into W3102 strains with and without R factors. The gal^+ Lft transductants obtained were further ultraviolet-irradiated, and the Hft lysates obtained were used for the transduction of gal^+ into W3102 strains with and without R factors. Phage λdg , like normal λ , was restricted by f_i^- R

factors; λdg was also modified in N-3 and R-15 by both of these f_i^- R factors, whereas it was not modified in N-1 (Table 3). Restriction and modification of λdg was previously reported to occur in P1-lysogenic bacteria (2). It may be worthwhile to point out that the suppression of gal^+ transduction by f_i^- R factors was not as severe as the reduction of EOP of normal λ (31).

Mating mixtures of Hfr H 3000 (λ^+) and W3102 (λ^- *str-r*) with and without R factors were interrupted at 60 min by blending, diluted in saline, and then plated in soft nutrient agar containing 1,000 $\mu\text{g}/\text{ml}$ of streptomycin with W3102 (*str-r*) as an indicator of phage λ . It was found that zygotic induction of phage λ is not significantly suppressed by any R factors in the F^- recipient.

Titration of phage P-22 propagated on LT-2 (R^-) was carried out with LT-2 strains with and without R factors as indicators. It was found that the EOP of P-22 was reduced by f_i^- R factors N-3 and R-15, but not by f_i^- R factor N-1 or by f_i^+ R factor 222 (Table 4). Plaques of P-22 on the

TABLE 2. Frequencies of lysogenization with phage λ of *Escherichia coli* W3102^a strains carrying various R factors

Strain	Frequency of lysogenized colonies ^b	
	No. of lysogenized colonies/no. studied	Per cent
W3102 (R^-)	20/127	15.7
W3120 (N-1)	8/298	2.7
W3102 (N-3)	1/328	0.3
W3102 (R-15)	17/450	3.8
W3102 (222)	28/127	22.0

^a W3102 is a $gal^- \lambda^- \text{F}^-$ mutant of *E. coli* K-12.

^b Each bacterial culture was infected with phage λ in Penassay Broth at a multiplicity of infection of 10. The infected culture was washed thoroughly and plated on nutrient agar for colony isolation. The resultant colonies were replica-plated onto soft nutrient agar seeded with W3102 (R^-) on top of ordinary nutrient agar.

TABLE 3. Host-induced modification of phage λdg on *Escherichia coli* W3102^a strains carrying various R factors

Hft lysate of λ prepared on	Frequency of transduction of ability to ferment galactose in recipient			
	W3102 (R^-)	W3102 (N-1)	W3102 (N-3)	W3102 (R-15)
W3102 (R^-)	2.3×10^{-3}	3.1×10^{-4}	3.4×10^{-4}	1.0×10^{-5}
W3102 (N-1)	3.1×10^{-3}	1.0×10^{-5}	4.3×10^{-5}	5.0×10^{-5}
W3102 (N-3)	2.0×10^{-3}	2.0×10^{-5}	1.5×10^{-3}	2.6×10^{-3}
W3102 (R-15)	2.7×10^{-3}	1.4×10^{-4}	6.7×10^{-3}	1.8×10^{-3}

^a W3102 is a $gal^- \lambda^- \text{F}^-$ mutant of *E. coli* K-12.

indicator strains carrying N-3 and R-15 were similar in shape but slightly smaller than those on the other indicator strains. P-22 was also modified by N-3 and R-15 (Table 4).

Phage P-22 grown on LT-2 (wild type) was used for transducing prototrophy into auxotrophic mutant strains of LT-2 with and without R factors. The frequencies of transductants were reduced by the presence of N-3 in the recipients (Table 5). R-15 gave comparable results, but no other R factor was inhibitory. Only complete transductants, and no abortive transductants (26), were counted in this experiment, because the frequencies of the latter were not reproducible. It may be worthwhile to emphasize that the reduction of transduction frequencies by N-3 and R-15 was less severe than the reduction of EOP by these R factors (Tables 4 and 5). When phage P-22, propagated on wild-type LT-2 carrying N-3, was used for transducing prototrophy into auxotrophic mutants of LT-2 carrying N-3, it was shown that transducing phage P-22 was also modified by N-3 (Table 5). Comparable results were again obtained with R-15.

The frequencies of lysogenization with phage P-22 were also markedly reduced by the presence of N-3 and R-15 in LT-2 (Table 6). The number of surviving bacteria were about equal in all of

the phage-infected LT-2 strains with and without R factors.

The effect of various R factors on the adsorption of phage P-22 was studied. LT-2 strains with and without R factors were grown in Penassay Broth at 37 C with aeration to about 5×10^8 cells per milliliter; 9 ml of each culture was mixed with 1 ml of phage P-22 grown on LT-2 (R⁻), 5×10^7 per milliliter, and aerated at 37 C. Samples (0.1 ml) were taken at various times, diluted with ice-cold Penassay Broth (1:100), and then centrifuged at 5,000 rev/min for 5 min in the cold. The supernatant fluids were treated with chloroform and assayed for free phage. There was no significant difference in the adsorption rates of P-22 to LT-2, with or without R factors.

Phage P-22 of lysogenic LT-2 strains with and without R factors was induced with ultraviolet light, and dose-response curves were obtained. No significant difference could be found in ultraviolet inducibility of phage P-22 in these strains. Spontaneous liberation of phage P-22 was also unaffected by any R factors.

Furthermore, no significant difference was found in the extent of injection of the nucleic acid of P³²-labeled phage T1 into W3102 strains with and without R factors (Table 7). Essentially

TABLE 4. Efficiencies of plating of phage P-22 on *Salmonella typhimurium* LT-2 strains carrying various R factors

P-22 grown on	Efficiency of plating on				
	LT-2 (R ⁻)	LT-2 (N-1)	LT-2 (N-3)	LT-2 (R-15)	LT-2 (222)
LT-2 (R ⁻)	1	9.5×10^{-1}	4.8×10^{-4}	2.7×10^{-4}	7.2×10^{-1}
LT-2 (N-3)	1	8.0×10^{-1}	1.2	1.1	8.0×10^{-1}
LT-2 (R-15)	1	7.3×10^{-1}	9.0×10^{-1}	1.0	1.0

TABLE 5. Frequencies of transduction with phage P-22 of prototrophy into auxotrophic mutants of *Salmonella typhimurium* LT-2 strains carrying various R factors

P-22 grown on	Recipient ^a	Selected marker ^b	Frequency of transductants (per plaque-forming unit)
LT-2 (R ⁻)	<i>cysD-36</i>	<i>cys</i> ⁺	6.5×10^{-6}
LT-2 (R ⁻)	<i>cysD-36</i> (N-3)	<i>cys</i> ⁺	6.7×10^{-7}
LT-2 (N-3)	<i>cysD-36</i>	<i>cys</i> ⁺	1.1×10^{-6}
LT-2 (N-3)	<i>cysD-36</i> (N-3)	<i>cys</i> ⁺	1.3×10^{-6}
LT-2 (R ⁻)	<i>tryD cysB-12 str-r</i>	<i>try</i> ⁺	1.7×10^{-6}
LT-2 (R ⁻)	<i>tryD cysB-12 str-r</i> (N-3)	<i>try</i> ⁺	3.2×10^{-6}
LT-2 (N-3)	<i>tryD cysB-12 str-r</i>	<i>try</i> ⁺	3.5×10^{-6}
LT-2 (N-3)	<i>tryD cysB-12 str-r</i> (N-3)	<i>try</i> ⁺	3.3×10^{-6}

^a Recipient strains *cysD-36* and *tryD cysB-12 str-r* are both substrains of *S. typhimurium* LT-2; *cysD-36* requires cystine, and *tryD cysB-12 str-r* requires tryptophan and cystine and is resistant to streptomycin.

^b The symbols *cys*⁺ and *try*⁺ represent nonrequirement of cystine and tryptophan, respectively.

similar results were obtained with P^{32} -labeled phage λ .

The fate of the injected phage nucleic acids was then studied. Figure 1A shows the kinetics of liberation of cold trichloroacetic acid-soluble P^{32} fraction after infection with P^{32} -labeled phage T1 of W3102 strains with and without R factors. The nucleic acid of phage T1 grown on W3102 (R^-) was rapidly broken down in the host bacteria carrying f_i^- R factors; more than 40% of the adsorbed phage nucleic acid was broken down within 10 min. In contrast, the nucleic acid of the phage was broken down less than 10% in W3102 (R^-) and W3102 (222). The nucleic acid of phage T1 grown on W3102 (N-3) and modified to both N-3 and R-15 was not broken down by more than 10% in W3102 (R^-), W3102 (N-3), and W3102 (R-15) but was broken down by about 40% in W3102 (N-1) as seen in (Fig. 1B). Comparable results were obtained with phage λ .

It was reported previously (31) that phage λ does not form visible plaques on W3102 (N-1) and that the plaques of phage T1 on this indicator

strain are much smaller than those on W3102 strains without R factors, with f_i^+ R factors, and with f_i^- R factors N-3 and R-15. No difference was found in the morphology of plaques of phages λ and T1 on W3102 strains without R factors, with f_i^+ R factors, and with f_i^- R factors N-3 and R-15. However, when the size of the plaques of phages T1 and λ was carefully re-examined in recent experiments, it was noted that the plaques of these phages on W3102 (N-3) and W3102 (R-15) were slightly but definitely smaller than those on W3102 strains without R factors and with f_i^+ R factors. Identical results were obtained also with phages λ and T1 modified to N-3 and R-15. These results suggested to us the possibility that the formation of progeny of these phages may be suppressed by the presence of these f_i^- R factors. Accordingly, the following experiments were undertaken with phage T1.

Since it was not possible to prepare a high-titer anti-T1 serum, as pointed out by Adams (1), we attempted in one-step growth experiments with T1 to remove free phage by washing with repeated centrifugation and resuspension. Penassay Broth cultures of W3102 strains with and without R factors were treated with 0.002 M KCN and then infected with phage T1 at a MOI of 0.1. In other experiments, the bacteria grown in Penassay Broth were washed and resuspended in T2 buffer (15), starved by aeration at 37 C for 30 min, and then infected with phage T1 at a similar MOI. The cultures infected under these conditions were washed four times with T2 buffer by repeated centrifugation and resuspension in the cold. The washed, infected bacteria were resuspended in cold Penassay Broth and then diluted in the same broth kept at 37 C to cell concentrations of 10^4 to 10^5 per milliliter. They were then aerated at 37 C, and 0.1-ml samples were periodically taken for determination of the infective centers. The results of such one-step growth experiments with phage T1 propagated on W3102 (R^-) are shown in Fig. 2A. The counts of bacterial infective centers of W3102, with and without R factors,

TABLE 6. Frequencies of lysogenization with phage P-22 of *Salmonella typhimurium* LT-2 strains carrying various R factors

Strain	Frequency of lysogenized colonies ^a	
	No. of lysogenized colonies/no. studied	Per cent
LT-2 (R^-)	534/1,088	49.1
LT-2 (N-1)	153/193	79.3
LT-2 (N-3)	1/1,796	0.06
LT-2 (R-15)	0/1,076	<0.09
LT-2 (222)	138/175	78.9

^a Each bacterial culture was infected with phage P-22 in Penassay Broth at a multiplicity of infection of 1. The infected culture was washed thoroughly and plated on nutrient agar for colony isolation. The resultant colonies were replica-plated onto a soft nutrient agar layer seeded with LT-2 (R^-) on the top of ordinary nutrient agar.

TABEL 7. Injection of nucleic acid of P^{32} -labeled phage T1 into *Escherichia coli* W3102 strains carrying various R factors

Strain	P^{32} adsorbed to bacteria ^a	P^{32} after blending ^a in		P^{32} in sediment after blending
		Supernatant fluid	Sediment	
W31-2 (R^-)	2.59×10^4	8.53×10^3	1.39×10^4	%
W3102 (N-1)	1.96×10^4	6.83×10^3	9.27×10^3	62.1
W3102 (N-3)	7.99×10^3	3.70×10^3	3.34×10^3	57.5
W3102 (222)	8.64×10^3	3.66×10^3	3.54×10^3	47.5
				49.2

^a Values (counts per minute) obtained from 1-ml samples after heat-drying.

FIG. 1. Breakdown of nucleic acid of ^{32}P -labeled T1 in *Escherichia coli* W3102 strains carrying various *R* factors. (A) Kinetics of formation of cold trichloroacetic acid-soluble ^{32}P fraction in the infection, with ^{32}P -labeled T1 grown on W3102 (R^-), of W3102 strains with and without *R* factors. (B) Similar kinetics in the infection, with ^{32}P -labeled T1 grown on W3102 (N-3), of W3102 strains with and without *R* factors.

before the liberation of progeny phage were in parallel with the EOP of the phage on these indicator strains. In addition, the latent period of T1 was not detectably altered by the presence of any *R* factors in the host bacteria. The average burst size of the phage, however, seemed to be slightly reduced in the hosts with fi^- *R* factors, especially with N-3 and R-15. The reduction in burst size in these strains was more marked when the host bacteria were starved before infection. One-step growth curves of T1 were similar when the phage grown on W3102 (N-3) and modified to N-3 and R-15 was used (Fig. 2B).

Single-burst experiments with phage T1 were then carried out in the hosts carrying various *R* factors. W3102 strains with and without *R* factors, grown at 37 C with aeration in Penassay Broth, were treated with 0.002 M KCN or starved by the procedures described above for one-step growth experiments. They were infected with phage T1 at a MOI of 0.1, and the infected bacteria were thoroughly washed with T2 buffer. They were then diluted with Penassay Broth so that 1 ml contained about 0.5 cell with a potentiality of liberating progeny phage. These bacterial suspensions were distributed into a series of small test tubes,

1 ml in each, and incubated at 37 C for 40 min. The contents of each tube were plated in 3 ml of soft nutrient agar with indicator strain W3102 for determining the free phage. The average burst sizes of T1 in single-burst experiments are shown in Table 8. It can be seen that the burst size of this phage was reduced by the presence of fi^- *R* factors, especially N-3. This tendency was more marked when the starved bacteria were infected as in one-step growth experiments. Comparable data were obtained also with phage T1 modified by N-3. Furthermore, it was shown that T1 was fully modified to N-3 by growing on W3102 (N-3) for a single cycle, as determined by plating the divided portions of the single-burst samples on both W3102 (R^-) and W3102 (N-3). Conversely, when the single-burst samples from W3102 (R^-) infected with T1, which had been modified by N-3, were plated on W3102 (R^-) and W3102 (N-3), it was found that the modification of T1 was completely lost during a single growth cycle on the R^- strain. These results clearly demonstrate, on the single infected-cell level, that the observed phenomena are really host-controlled modifications of the phage.

DISCUSSION

The results indicate that the formation of plaques by phage W-31 on *E. coli* K-12 and by phage P-22 on *S. typhimurium* LT-2 are reduced

FIG. 2. One-step growth curves of phage T1 in *Escherichia coli* W3102 (*gal*⁻ λ ⁻ F⁻) strains carrying various R factors. (A) One-step growth curves of T1, previously grown on W3102 (R⁻). (B) One-step growth curves of T1, previously grown on W3102 (N-3). The curves with solid symbols represent bacteria infected with T1 after treatment with 0.002 M KCN in Penassay Broth. The curves with open symbols represent bacteria infected with T1 after starvation in T2 buffer (15) for 30 min at 37 C.

by some *fi*⁻ R factors but not by *fi*⁺ R factors. Phage P-22 was also modified by the restricting *fi*⁻ R factors. The patterns of restriction and modification of these phages by R factors, reported here and in a previous paper (31), may be summarized as in Table 9. Not only was the formation of plaques of these phages suppressed by the *fi*⁻ R factors, but also the frequencies of lysogenization of K-12 with phage λ and of LT-2 with phage P-22 were reduced by the *fi*⁻ R factors. Furthermore, the frequencies of *gal*⁺ transduction with λ dg into a *gal*⁻ mutant of K-12 and of transduction with P-22 of prototrophy into auxotrophic mutants of LT-2 were reduced by the *fi*⁻ R factors in the recipients. It was also shown that

TABLE 8. Average burst sizes (*I*) of phage T1 in *Escherichia coli* W3102^a carrying various R factors studied by single-burst experiments

Phage T1 grown on	Condition of host bacteria at the time of phage infection	Average burst size on		
		W3102 (R ⁻)	W3102 (N-1)	W3102 (N-3)
W3102 (R ⁻)	Treated with KCN ^b	39.3	27.1	24.0
	Starved ^c	31.3	31.8	8.1
W3102 (N-3)	Treated with KCN	28.8	23.3	15.6
	Starved	28.3	21.4	9.0

^a W3102 is a *gal*⁻ λ ⁻ F⁻ mutant of *E. coli* K-12.

^b Bacteria in Penassay Broth were treated with 0.002 M KCN, and infected with T1 at a multiplicity of infection of 0.1. The infected bacteria were washed in the cold and resuspended in fresh Penassay Broth.

^c Bacteria were starved by aerating the T2 buffer suspension at 37 C for 30 min.

TABLE 9. Summary of restrictions and modifications of phages by R factors

Strain no. of R factor	<i>fi</i> ^a	Response of phage	Phage				
			λ	T1	T7	W-31	P-22
N-1.....	—	Restriction	+	+	+	+	—
		Modification	—	—	—	—	—
N-3.....	—	Restriction	+	+	—	—	+
		Modification ^b	+	+	—	—	+
R-15.....	—	Restriction	+	+	—	—	+
		Modification ^b	+	+	—	—	+
222, N-6, N-9..	+	Restriction	—	—	—	—	—
		Modification	—	—	—	—	—

^a Presence (+) and absence (—) of suppression of the function of F of *Escherichia coli* K-12; *fi* is the abbreviation for fertility inhibition.

^b Modification of phages by N-3 and R-15 occur simultaneously to both of these R factors.

the transducing phage λ dg and the transducing phage P-22 of prototrophy are modified by the f_i^- R factors N-3 and R-15 to both of them but not by N-1 like normal phages. It was pointed out that the suppression of these transductions by the f_i^- R factors was less severe than the reductions of EOP and lysogenization with these phages.

The adsorption of phage P-22 to LT-2 was not affected by any R factors. Similar findings have been obtained with phages λ and T1 in K-12 (31). It was demonstrated that the injection of the nucleic acids of phages T1 and λ is not detectably affected by the presence of any R factors.

It was further shown that the nucleic acids of phages T1 and λ grown on R^- cells of K-12 are broken down in bacteria carrying f_i^- R factors. These results of the injection of the phage nucleic acids and of their breakdown in restrictive bacteria indicate that the broken-down phage nucleic acids remain inside the cells if the bacteria are not treated with trichloroacetic acid. The nucleic acids of phages T1 and λ modified by N-3 were not broken down in the bacteria carrying N-3 and R-15, but underwent breakdown in bacteria carrying N-1. Less than 10% of the unmodified phage nucleic acids were broken down in cells carrying f_i^+ R factor 222 and in R^- cells, where no restriction is observed. About an equal extent of breakdown of nucleic acids was found with modified phages. It is not known whether this small fraction of breakdown of the phage nucleic acids is an artifact. Similar findings have been obtained by Dussoix and Arber (12) and Fukasawa (14) in their systems. Thus, the mechanism of suppression of formation of phage plaques by f_i^- R factors seems likely to be due to the breakdown of the injected phage nucleic acids, although there is an alternative possibility that this breakdown is a mere result of restriction (19). At any rate, the agent(s) responsible for the breakdown of the phage nucleic acids must be a deoxyribonuclease(s), because the phages listed above as restricted by the f_i^- R factors are all known to contain deoxyribonucleic acid (DNA) as their genetic material. Phage W-31 is not an exception (T. Miyake, *personal communication*). The modified phages may contain DNA with "host specificity" (3, 4), since they are not sensitive to this enzyme. The fact that the DNA of the phages, which were modified to N-3 and R-15, remain sensitive to breakdown in the cells with N-1 seems to suggest that these deoxyribonucleases are specifically formed by the presence of the f_i^- R factors and that the enzyme specified by N-1 is different from that specified by N-3 and R-15. There is an alternative possibility: the enzyme responsible for the breakdown of the phage DNA is a normal component of the cell and the injected

phage DNA undergoes some change which protects the DNA from the action of the enzyme, whereas this change does not occur in the restrictive bacteria (19). As in phage λ , spontaneous as well as ultraviolet-induced production of phage P-22 in lysogenic LT-2 was not affected by any R factors. One may suspect that this is due to the modification of prophages by f_i^- R factors. But this is not the case at least in N-1, because this R factor does not modify these phages. Furthermore, zygotic induction of phage λ was not affected by any R factors. Thus, the phage nucleic acids, which are present as prophages in the bacteria carrying f_i^- R factors, and which are transferred to the restrictive cells together with the host chromosome, are apparently able to replicate normally. Arber and Morse (5), Boyer (6), and Pittard (27) reported that the recombination frequencies and the linkage of chromosomal genes were reduced in the cross of Hfr strains with the restrictive F^- recipients and that these reductions are due to the breakdown of the transferred chromosome. We could not find any measurable reduction by f_i^- R factors of the recombination of chromosomal genes in the crosses of R^- Hfr with F^- R^+ strains (*unpublished data*). These results suggest that the enzyme responsible for the breakdown of the phage DNA in our systems may be located near the cell surface and that the chromosome transferred by conjugation and prophage λ attached to it may be somehow protected from this enzyme(s). In fact, Cordonnier and Bernardi (10) showed that endonuclease I is located near the cell wall, and Molholt and Fraser (23) found that the restriction of phages is removed upon conversion of restrictive *E. coli* to spheroplasts. We have recently shown, however, that endonuclease I is not involved with the mechanism of restriction of phages by f_i^- R factors by our finding that these phages are still restricted in an endonuclease I-less mutant of *E. coli* carrying f_i^- factors. These results will be reported in a separate paper. Our failure to show the restriction of bacterial chromosomes by f_i^- R factors might merely be due to the much weaker restriction compared to that by prophage P1.

The fact that transductions are less severely suppressed by f_i^- R factors than is the formation of phage plaques may be interpreted by assuming that the bacterial chromosomal genes in the transducing phage particles have a greater chance of surviving the inactivation by the nuclease. In other words, this difference may be due to the difference in target sizes between the DNA molecules of phages and chromosomal genes. Suppression of lysogenization with temperate phages by f_i^- R factors is easily and reasonably accounted for by the breakdown of the injected phage DNA.

N-3 and R-15 have so far exhibited identical patterns in the restriction and modification of various phages (Table 9), and also in their superinfection immunity (31). However, we have recently found that the restriction of the phages by R-15 but not by N-3 is temperature-sensitive. This finding is now under investigation and will be reported in a following paper.

It was further shown that the average burst size of phage T1 is reduced by the presence of f_i^- R factors, in one-step growth as well as in single-burst experiments. This point was not studied with other phages but, since the plaque sizes of phages λ and P-22, both modified and unmodified, on the indicators with N-3 and R-15 are slightly smaller than those on the indicators with f_i^+ R factors and without R factors, it is likely that the progeny formation of these phages is also suppressed slightly by the presence of f_i^- R factors. The reason for the small plaque size of the phages on the indicators with N-1 was previously interpreted as being due to the absence of modification of the phages (31). These results indicate that the formation of progeny phage by the phage DNA which escaped enzymatic breakdown after injection remains sensitive to some inhibitory mechanism in the cells with f_i^- R factors.

ACKNOWLEDGMENTS

We are indebted to D. Ushiba, H. Uetake, T. Fukasawa, and S. Toyama for helpful discussions.

This investigation was supported by Public Health Service research grant AI-04740 from the National Institute of Allergy and Infectious Diseases, and by a research grant from the Ministry of Education, Japan.

LITERATURE CITED

- ADAMS, M. H. 1950. Methods of study of bacterial viruses. *Methods Med. Res.* **2**:1-73.
- ARBER, W. 1964. Host specificity of DNA produced by *Escherichia coli*. III. Effects of transduction mediated by λ dg. *Virology* **23**:173-182.
- ARBER, W. 1965. Host-controlled modification of bacteriophage. *Ann. Rev. Microbiol.* **19**:365-378.
- ARBER, W., AND D. DUSOIX. 1962. Host specificity of DNA produced by *Escherichia coli*. I. Host controlled modification of bacteriophage λ . *J. Mol. Biol.* **5**:18-36.
- ARBER, W., AND M. L. MORSE. 1965. Host specificity of DNA produced by *Escherichia coli*. VI. Effects on bacterial conjugation. *Genetics* **51**:137-148.
- BOYER, H. 1964. Genetic control of restriction and modification in *Escherichia coli*. *J. Bacteriol.* **88**:1652-1660.
- BRINTON, C. C., JR. 1965. The structure, function, synthesis and genetic control of bacterial pili and a molecular model for DNA and RNA transport in gram negative bacteria. *Trans. N.Y. Acad. Sci. Ser. II* **27**:1003-1054.
- BRINTON, C. C., JR., P. GEMSKI, JR., AND C. CARNAHAN. 1964. A new type of bacterial pilus genetically controlled by the fertility factor of *E. coli* K12 and its role in chromosome transfer. *Proc. Natl. Acad. Sci. U.S.A.* **52**:776-783.
- BRITTEN, R. J., AND R. B. ROBERTS. 1960. High-resolution density gradient sedimentation analysis. *Science* **131**:32-33.
- CORDONNIER, C., AND G. BERNARDI. 1965. Localization of *E. coli* endonuclease I. *Biochem. Biophys. Res. Commun.* **20**:555-559.
- DEMEREK, M., AND Z. HARTMAN. 1956. Tryptophan mutants in *Salmonella typhimurium*. *Carnegie Inst. Wash. Publ.* **612**:5-17.
- DUSOIX, D., AND W. ARBER. 1962. Host specificity of DNA produced by *Escherichia coli*. II. Control over acceptance of DNA from infecting phage λ . *J. Mol. Biol.* **5**:37-49.
- DUSOIX, D., AND W. ARBER. 1965. Host specificity of DNA produced by *Escherichia coli*. IV. Host specificity of infectious DNA from bacteriophage λ . *J. Mol. Biol.* **11**:238-246.
- FUKASAWA, T. 1964. The course of infection with abnormal bacteriophage T4 containing non-glucosylated DNA on *Escherichia coli* strains. *J. Mol. Biol.* **9**:525-536.
- HERSHEY, A. D., AND M. CHASE. 1952. Independent functions of viral protein and nucleic acid in growth of bacteriophage. *J. Gen. Physiol.* **36**:39-56.
- JACOB, F., AND E. L. WOLLMAN. 1954. Induction spontanée du développement du bactériophage au cours de la recombinaison génétique chez *Escherichia coli*. *Compt. Rend.* **239**:455-456.
- JACOB, F., AND E. L. WOLLMAN. 1956. Sur les processus de conjugaison et de recombinaison génétique chez *E. coli*. I. L'induction par conjugaison ou induction zygotique. *Ann. Inst. Pasteur* **91**:486-510.
- JACOB, F., AND E. L. WOLLMAN. 1961. Sexuality and the genetics of bacteria, p.93-108. Academic Press, Inc., New York.
- LEDERBERG, S. 1965. Host-controlled restriction and modification of deoxyribonucleic acid in *Escherichia coli*. *Virology* **27**:378-387.
- LEDERBERG, S., AND M. MESELSOHN. 1964. Degradation of non-replicating bacteriophage DNA in non-accepting cells. *J. Mol. Biol.* **8**:623-628.
- MEYNELL, E., AND N. DATTA. 1965. Functional homology of the sex-factor and resistance transfer factors. *Nature* **207**:884-885.
- MEYNELL, E., AND N. DATTA. 1966. The relation of resistance transfer factors to the F factor (sex-factor) of *Escherichia coli* K12. *Genet. Res.* **7**:134-140.
- MOLHOLT, B., AND D. FRASER. 1965. Reversal of restriction for host modified T2 and T4 DNA upon conversion of non-permissive *Escherichia coli* to spheroplasts. *Biochem. Biophys. Res. Commun.* **19**:571-575.
- MORSE, M. L., E. M. LEDERBERG, AND J. LEDERBERG. 1956. Transduction in *Escherichia coli* K-12. *Genetics* **41**:142-156.
- NOMURA, M., B. D. HALL, AND S. SPIEGELMAN.

1960. Characterization of RNA synthesized in *Escherichia coli* after bacteriophage T2 infection. *J. Mol. Biol.* **2**:306-326.
26. OZEKI, H. 1956. Abortive transduction in purine-requiring mutants of *Salmonella typhimurium*. *Carnegie Inst. Wash. Publ.* **612**:97-106.
27. PITTARD, J. 1964. Effect of phage-controlled restriction on genetic linkage in bacterial crosses. *J. Bacteriol.* **87**:1256-1257.
28. UETAKE, H., S. TOYAMA, AND S. HAGIWARA. 1964. On the mechanism of host-induced modification. Multiplicity activation and thermolabile factor responsible for phage growth restriction. *Virology* **22**:202-213.
29. WATANABE, T., AND T. FUKASAWA. 1961. Episome-mediated transfer of drug resistance in *Enterobacteriaceae*. I. Transfer of resistance factors by conjugation. *J. Bacteriol.* **81**:669-678.
30. WATANABE, T., T. FUKASAWA, AND T. TAKANO. 1962. Conversion of male bacteria of *Escherichia coli* K-12 to resistance to f phages by infection with the episome "resistance transfer factor." *Virology* **17**:218-219.
31. WATANABE, T., H. NISHIDA, C. OGATA, T. ARAI, AND S. SATO. 1964. Episome-mediated transfer of drug resistance in *Enterobacteriaceae*. VII. Two types of naturally occurring R factors. *J. Bacteriol.* **88**:716-726.
32. WATANABE, T., AND M. OKADA. 1964. New type of sex factor-specific bacteriophage of *Escherichia coli*. *J. Bacteriol.* **87**:727-736.
33. YOSHIKAWA, M., K. MIYAKE, AND T. WAKABAYASHI. 1964. Studies on the mechanism of transfer of drug resistance in bacteria. 37. The effect of R factor on the process of UV induction and zygotic induction of prophage λ . *Med. Biol. (Tokyo)* **69**:344-347.