Published in final edited form as:

Hum Mutat. 2007 May ; 28(5): 469–476. doi:10.1002/humu.20471.

Signal Peptide Variants that Impair Secretion of Pancreatic Secretory Trypsin Inhibitor (SPINK1) Cause Autosomal Dominant Hereditary Pancreatitis

Orsolya Király^{1,#}, Arnaud Boulling^{2,3,#}, Heiko Witt^{4,#}, Cédric Le Maréchal^{2,3,5,6}, Jian-Min Chen^{2,3}, Jonas Rosendahl^{4,7}, Cinzia Battaggia⁸, Thomas Wartmann⁹, Miklós Sahin-Tóth^{1,*}, and Claude Férec^{2,3,5,6,*}

¹Department of Molecular and Cell Biology, Boston University, Goldman School of Dental Medicine, Boston, MA, 02118

²INSERM, U613, 29220 Brest, France

³Etablissement Français du Sang – Bretagne, 29220 Brest, France

⁴Department of Hepatology and Gastroenterology, Charité, 13353 Berlin, Germany

⁵Université de Bretagne Occidentale, Faculté de Médecine de Brest et des Sciences de la Santé, 29238 Brest, France

⁶Centre Hospitalier Universitaire (CHU) Brest, Hôpital Morvan, Laboratoire de Génétique Moléculaire et d'Histocompatibilité, 29220 Brest, France

⁷Department of Gastroenterology and Hepatology, University of Leipzig, 04103 Leipzig, Germany

⁸Department of Human and Animal Biology, Section of Anthropology, University of Rome "La Sapienza", 00133 Rome, Italy

⁹Division of Experimental Surgery; Department of Surgery, Otto-von-Guericke University, 39120 Magdeburg, Germany.

Abstract

Variants of the *SPINK1* gene encoding pancreatic secretory trypsin inhibitor have been described in association with chronic pancreatitis. These alterations are believed to cause a loss of function by either impairing the trypsin inhibitory activity or reducing expression. Here we report two novel *SPINK1* variants in exon 1 that affect the secretory signal peptide. The disease-associated c.41T>G (p.L14R) alteration was found in two European families with autosomal dominant hereditary pancreatitis, whereas the c.36G>C (p.L12F) variant was identified as a frequent alteration in subjects of African descent. The functional effects of both alterations and the previously reported c.41T>C (p.L14P) variant were characterized by activity assays and Western blots of wild-type and mutant *SPINK1* expressed in human embryonic kidney 293T and Chinese hamster ovary cells. Alterations p.L14R and p.L14P destined the inhibitor for rapid intracellular degradation and thereby abolished *SPINK1* secretion, whereas the p.L12F variant showed no detrimental effect. The results provide the first clear experimental demonstration that alterations which markedly reduce *SPINK1* expression

*Correspondence to Miklós Sahin-Tóth, 715 Albany Street, Evans-433; Boston, MA 02118; Tel: (617) 414-1070; Fax: (617) 414-1041; E-mail: miklos@bu.edu; or Claude Férec, INSERM, U613, Etablissement Français du Sang – Bretagne, 46 rue Félix Le Dantec, 29220 Brest, France. E-mail: claud.ferec@univ-brest.fr.

#These authors contributed equally to the project.

DATABASES. *SPINK1* reference sequence, NM_003122.2. Variant L12F, DQ790099. Variant L14R, DQ790100.

are associated with classic hereditary pancreatitis. Therefore, these variants should be classified as severe and regarded as disease-causing rather than disease-modifiers.

Keywords

pancreatic secretory trypsin inhibitor; PSTI; chronic pancreatitis; signal-peptide alteration; hereditary pancreatitis; human cationic trypsin; SPINK1

INTRODUCTION

The pancreatic secretory trypsin inhibitor is a 6.5 kDa protein synthesized, stored and secreted by the acinar cells of the exocrine pancreas. In humans, the inhibitor is encoded by the *SPINK1* (Serine Protease Inhibitor, Kazal type 1) gene on chromosome 5q32 (OMIM *167790). As a typical secretory protein, human *SPINK1* contains a 23 amino-acid signal peptide (Fig 1), which directs the protein to the endoplasmic reticulum, where, after translocation to the lumen and removal of the signal peptide, *SPINK1* achieves its mature fold. It has long been assumed that the physiological role of *SPINK1* is to protect the pancreas against premature trypsinogen activation by rapidly complexing free trypsin (Rinderknecht, 1986).

The strongest indication that *SPINK1* plays an essential protective role in pancreatic physiology came from relatively recent genetic studies, describing an association between the *SPINK1* c.101A>G (p.N34S, Chen et al., 2000) variant and chronic pancreatitis (Witt et al., 2000). Data taken from 8 larger studies in Europe and US indicate that 12.6 % of patients with chronic pancreatitis were heterozygous for N34S, and 3.6 % were homozygous (Witt et al., 2000; Pfützer et al., 2000; Plendl et al., 2001; Chen et al., 2001; Noone et al., 2001; Drenth et al., 2002; Threadgold et al., 2002; Truninger et al., 2002). In contrast, homozygous individuals were never identified among controls, whereas heterozygosity was detected at 1.9 % on average. Thus, the large majority of carriers never develop pancreatitis. In addition to these studies, N34S was also recognized as a major genetic risk factor in tropical pancreatitis in the Indian subcontinent (Chandak et al., 2002; Hassan et al., 2002; Bhatia et al., 2002; Schneider et al., 2002) and as a minor susceptibility factor in alcoholic chronic pancreatitis (Witt et al., 2001b). The mode of inheritance of the N34S alteration in chronic pancreatitis has been debated in the initial reports. The first study suggested that N34S may cause pancreatitis by an autosomal recessive mode with low penetrance (~25 %) or in an oligogenic fashion (Witt et al., 2000). Subsequently, based on negative genetic linkage studies, the lack of difference in age of onset between heterozygous and homozygous individuals and the observation that heterozygosity for N34S is common in the population, it was proposed that the N34S alteration is a “disease modifier”, which does not cause pancreatitis, but modifies the effect of other disease-causing genes (Pfützer et al., 2000). More recently, the N34S allele has been viewed as a relatively frequent genetic susceptibility factor to chronic pancreatitis, which exerts its effect in the context of other genetic and environmental factors, resulting in the highly variable penetrance and expression of the disease. Because *SPINK1* is supposed to protect the pancreas and the N34S variant seems to increase the risk of pancreatitis, the straightforward conclusion was drawn that the N34S alteration causes a loss of inhibitor function (Witt et al., 2000). However, such a functional defect has not been demonstrated so far, as a recent study indicated that recombinant N34S *SPINK1* retains full inhibitory activity (Kuwata et al., 2002).

In addition to the N34S alteration, several other *SPINK1* variations have been described in pancreatitis patients as well as healthy controls (see <http://www.uni-leipzig.de/pancreasmutation>). The large majority of alterations found in pancreatitis patients represent rare variants, therefore, disease-association and mode of inheritance has not been determined conclusively. Nevertheless, in some cases, the nature of

the alteration indicates an obligatory loss of SPINK1 expression, either due to splicing problems (c.87+1G>A; c.194+2T>C), shift of reading frame with premature termination (c.98dupA; c.27delC), loss of the initiator methionine codon (c.2T>C; p.M1?) or large genomic deletions in the *SPINK1* gene (Witt et al., 2000; Gaia et al., 2002; Le Maréchal, 2004; Masson et al. 2006).

Here we report that *SPINK1* alterations in the signal peptide are associated with a severe pancreatitis phenotype and these alterations result in decreased secretion of the mature SPINK1 peptide due to intracellular degradation.

MATERIALS AND METHODS

Plasmid construction, transfection, and Western blot analysis were executed independently in the Brest and Boston laboratories using somewhat different methods. Unless indicated otherwise, the Material and Methods section refers to the experiments performed in Boston.

Study subjects

The study was approved by the local ethical review committees. All patients gave their informed consent for genetic analysis. The study included 574 patients with hereditary or idiopathic chronic pancreatitis originating from France (n=320) and Germany (n=254). All 254 chronic pancreatitis patients originating from Germany (144 female, 110 male; mean age $\pm$ SD: 11.9 ± 6.9 years) were of German ethnicity. Of these subjects, 65 patients (42 female, 23 male; mean age $\pm$ SD: 12.1 ± 7.9 years) were classified as having hereditary pancreatitis and 189 patients (102 female, 87 male; mean age $\pm$ SD: 11.8 ± 6.5 years) were classified as having idiopathic chronic pancreatitis. The clinical diagnosis of CP was based on two or more of the following criteria: presence of a typical history of recurrent pancreatitis, radiological findings such as pancreatic calcifications and/or pancreatic ductal irregularities revealed by endoscopic retrograde pancreaticography or magnetic resonance imaging of the pancreas and/or pathological sonographic findings. Hereditary CP was diagnosed when two first-degree relatives or three or more second-degree relatives suffered from recurrent acute or chronic pancreatitis without any apparent precipitating factor. Patients were classified as having idiopathic CP when precipitating factors such as alcohol abuse, trauma, medication, infection, metabolic disorders and/or a positive family history were absent. Altogether 589 control subjects were included. The French controls consisted of 98 unrelated healthy bone marrow donors from Brest recruited between 1998 and 2000. The 340 German control subjects (181 female, 159 male; mean age $\pm$ SD: 56.9 ± 29.2 years) consisted of 162 medical students and staff recruited in Berlin between 1996 and 1998 and 178 healthy individuals recruited in Berlin for the Berlin Aging Study (BASE) between 1993 and 1995. To determine the frequency of *SPINK1* variants in non-Caucasians, we studied 151 individuals of African descent originating from Cameroon which were recruited for genetic population studies between 1986 and 1990 (132 female, 19 male; mean age $\pm$ SD: 33.8 ± 6.6 years).

Numbering of genetic variants is based on the cDNA sequence, indicated by a "c." before the number (den Dunnen and Paalman, 2003). Position +1 corresponds to the A of the ATG translation initiation codon located at nucleotide 121 in the NM_003122.2 *SPINK1* mRNA reference sequence.

Plasmid Construction

The *SPINK1* cDNA was amplified from IMAGE clone #1679809 (GenBank AA844948) using the oligonucleotide primers SPINK1_XhoI_sense 5'-CGG TGC TCG AGT CAA CTG ACC TCT GGA CGC AGA ACT TC-3' and SPINK1_BamHI_antisense 5'-ATA CAG GAT CCC AAC AAT AAG GCC AGT CAG GCC TCG CGG-3. The PCR product was digested with

XhoI and *BamHI* and ligated into the expression vector pcDNA3.1(−) (Invitrogen) digested with the same enzymes. Signal peptide mutants and the Glu-Glu-tagged SPINK1 constructs were generated by overlap extension PCR mutagenesis and ligated into pcDNA3.1(−). The Glu-Glu tag sequence (EYMPME) is derived from the polyoma virus medium T antigen (Grussenmeyer et al., 1985).

Cell Culture and Transfection

Human embryonic kidney (HEK) 293T cells were cultured in D-MEM supplemented with 10% fetal bovine serum, 4 mM glutamine and 1% penicillin/streptomycin solution at 37 °C in a humidified atmosphere containing 5% CO₂. A day before transfection, approximately 10⁶ cells were plated in 6-well tissue culture plates in culture medium without antibiotics. Transfections were performed using 4 µg pcDNA3.1(−)_SPINK1 plasmid, 2 µg pSV-β-galactosidase control plasmid (Promega), and 10 µL Lipofectamine 2000 (Invitrogen) in 2 mL OptiMEM medium supplemented with 2 mM glutamine. After 5 h incubation at 37 °C, 2 mL of feed medium (D-MEM with 20% fetal bovine serum, 4 mM glutamine) was added to each well and cells were incubated for an additional 24 h. Cells were then washed with 1 mL OptiMEM, 2 mM glutamine and the transfection medium was replaced with 2 mL OptiMEM, 2 mM glutamine. Time-courses of expression were measured starting from this medium change and were followed for 48 h. At the indicated times, 200 µL conditioned medium (i.e. 10 % of total volume) was removed from each well for activity assays and replaced with 200 µL fresh medium. At the conclusion of the transfection experiments, conditioned media and cells were harvested for Western blot analysis and β-galactosidase assays.

Trypsin Inhibitory Activity Determinations

Conditioned media from transfected cells were centrifuged for 5 min and 40 µL of the supernatant was mixed with 4 mg/mL bovine serum albumin and 80 nM recombinant human cationic trypsin (final concentrations) in a final volume of 50 µL and incubated for 1 min at room temperature. The chromogenic substrate CBZ-Gly-Pro-Arg-p-nitroanilide (150 µL) was then added to 0.14 mM concentration and 1 min rates of p-nitroaniline release were measured at 405 nm. Residual trypsin activity was expressed as percentage of the uninhibited maximal activity determined in the presence of 40 µL fresh medium devoid of SPINK1. Residual trypsin activities were converted to inhibitor concentrations using a calibration curve obtained by assaying serial dilutions of purified recombinant SPINK1.

β-Galactosidase Activity Assays to Determine Transfection Efficiency

Cells were co-transfected with 2 µg pSV-β-galactosidase plasmid. Forty-eight hours after transfection, the growth medium was removed and the cells were washed twice with PBS. Reporter Lysis Buffer (Promega; 200 µL) was added to each well and the plate was incubated for 15 min at room temperature. Cells were then scraped from the wells, the lysates were briefly vortexed and centrifuged for 5 min at 4 °C. Supernatants were assayed for β-galactosidase activity by measuring rates of o-nitrophenyl β-D-galactopyranoside hydrolysis using the β-galactosidase Enzyme Assay System (Promega). The resulting β-galactosidase activity rates were used to normalize SPINK1 expression to variations in transfection efficiency. SPINK1 concentrations were divided by the corresponding β-galactosidase activity values and multiplied by the average of β-galactosidase activities from all transfections in the experiment.

Total RNA Isolation and RT-PCR

RNA was isolated from cells transfected with wild type and mutant *SPINK1* constructs using the RNeasy kit (Qiagen) and 0.75 µg RNA was reverse-transcribed with the SMART PCR cDNA Synthesis Kit (Clontech). *SPINK1* cDNA was measured by semi-quantitative PCR using the gene-specific primer pair described under "*Plasmid Construction*". As a reference

control, the cDNA of the glyceraldehyde-3-phosphate dehydrogenase (GAPD) gene was also amplified using the pseudogene-free amplification conditions described by Harper et al. (2003).

Western Blotting

Samples of conditioned medium or cell extracts were run on a 21% Tris-glycine gel under reducing conditions and electrophoretically transferred onto an Immobilon-P membrane (Millipore) at 300 mA for 3 h. To detect the Glu-Glu-tag, the membrane was blocked with 5% milk powder solution overnight and incubated with an HRP-conjugated goat polyclonal antibody raised against the Glu-Glu tag (Abcam) at a concentration of 0.1 µg/mL for 1 h at room temperature. When untagged proteins were analyzed, the membrane was blocked in 5% milk/3% BSA overnight and hybridized with anti-human SPINK1 rabbit serum at a dilution of 1:5000 for 1 h at room temperature. The secondary antibody (ImmunoPure peroxidase conjugated goat anti-rabbit IgG; Pierce) was applied at a dilution of 1:50,000 for 1 h. HRP was detected using the SuperSignal West Pico Chemiluminescent Substrate (Pierce). Western blot analysis of conditioned media of Chinese hamster ovary (CHO) cells was performed using the PSKAN2 monoclonal mouse IgG antibody (Mobitec; Göttingen, Germany) at a 1:250 dilution, followed by peroxidase conjugated anti-mouse IgG (Amersham) at 1:10,000 dilution.

RESULTS

SPINK1 Signal Peptide Variants in Patients with Chronic Pancreatitis

Screening a previously described cohort of patients with familial or hereditary idiopathic pancreatitis from France for alterations in the *SPINK1* gene by denaturing high-performance liquid chromatography (Le Maréchal et al., 2004) identified a heterozygous T to G transversion at position c.41 (Supplementary Fig S1A) in a family of Bulgarian origin (Fig 2). This c.41T>G alteration resulted in the replacement of the leucine at amino acid position 14 with an arginine (p.L14R, Fig 1). The 20-year-old index patient was diagnosed at age 10. His father died from an acute attack of pancreatitis, and because the mother tested negative, he was an obligate carrier of the alteration. The father's mother was diagnosed with chronic pancreatitis at age 59 and was also heterozygous for L14R. Thus, the alteration segregated with pancreatitis over three generations, with complete penetrance. We also analyzed 254 German patients with chronic pancreatitis by direct DNA sequencing (Witt et al., 2000) and detected the L14R alteration in a 9-year old male patient from the county of Niedersachsen (Lower Saxony) (Fig 2). The alteration was present in the unaffected father and the affected grandmother. The greatgrandmother also suffers from chronic pancreatitis, but was not available for genetic testing. L14R was absent in 98 unrelated French healthy bone marrow donors, and in 340 German control subjects. No other pathogenic *PRSSI*, *CFTR* or other *SPINK1* alterations were detected in the two families with the L14R alteration.

The c.41T>C (p.L14P) alteration was identified in 2000 in a 5-year old German pediatric patient without a family history (Witt et al., 2000). The mother of the index patient carried the alteration, but had no pancreatitis. In the meantime, however, the mother also developed pancreatitis at age 41. This family was also negative for *PRSSI*, *CFTR* or other *SPINK1* alterations. L14P affects the same amino acid within the hydrophobic core of the signal peptide as the newly found L14R alteration. The strong association with pancreatitis indicates that both the L14R and L14P alterations can be regarded as relatively severe susceptibility factors for chronic pancreatitis.

We found another novel *SPINK1* variant that altered the signal peptide, c.36G>C (p.L12F; Supplementary Fig S1B), in 15 African subjects with idiopathic pancreatitis living in France. However, this variant turned out to be a common alteration in Africans, with an allele frequency

of 11 % (33/302 alleles). The L12F variation was not detected in Caucasian patients and controls from France or in 323 pancreatitis patients and 340 controls from Germany.

Signal Peptide Variants do not Affect mRNA Levels

Functional characterization of the effects of the signal peptide variants was performed by transiently transfecting HEK 293T cells with cDNA constructs encoding wild-type, L14P, L14R or L12F SPINK1 variants. To measure transfection efficiency, cells were routinely co-transfected with a plasmid encoding β -galactosidase. Semi-quantitative PCR analysis of cDNA preparations from transfected cells yielded products of equal intensity, indicating that mRNA levels expressed from the wild-type and the L14P, L14R or L12F constructs are comparable (Fig 3).

Pancreatitis-Associated Variations L14P and L14R Abolish SPINK1 Secretion

Secretion of SPINK1 protein by transfected cells was monitored by measuring the inhibitory activity of conditioned media against human cationic trypsin (Fig 4). After transfection with wild-type or L12F *SPINK1* cDNA, trypsin inhibitory activity in the medium of HEK 293T cells steadily increased during the 2-day time-course followed, whereas media from cells transfected with the L14R and L14P mutants showed minimal or no trypsin inhibitory activity, respectively. Residual trypsin activities were converted to actual SPINK1 concentrations using a calibration curve obtained by assaying serial dilutions of purified recombinant SPINK1. The resulting concentration values were then normalized to transfection efficiency using the β -galactosidase activity of the transfected cells. Time-courses of SPINK1 secretion expressed as normalized SPINK1 concentrations are presented in Fig 5. Clearly, pancreatitis-associated alterations L14R and L14P abolish secretion, whereas the L12F variant has no effect.

To demonstrate that the lack of trypsin inhibitory activity was the result of impaired secretion, aliquots of conditioned media were analyzed by SDS-PAGE and Western blotting. First, a rabbit polyclonal antibody raised against recombinant human SPINK1 was used. As shown in Fig 6A, wild-type and L12F SPINK1 proteins were visualized at comparable levels, whereas the L14P variant was not detected, and the L14R variant exhibited markedly reduced expression. Interestingly, the faint band observed for the L14R variant migrated slower than wild-type SPINK1, indicating that signal peptide processing probably occurred at an anomalous site. Indeed, analysis of the SPINK1 signal peptide sequence with the SignalP program (<http://www.cbs.dtu.dk/services/SignalP>) revealed that the L14R variant might be processed at the Ser18-Gly19 peptide-bond resulting in a mature peptide with an additional 5 amino acids (GNTGA) at the N terminus.

Because the L14R and L14P alterations occur in a heterozygous form in patients, we have also performed co-transfection experiments with wild-type and variant *SPINK1* cDNAs. Consistently, we found a slight to moderate increase in the secreted SPINK1 activity, which appeared to be due to increased secretion of the wild-type form, as judged by immunoblot analysis of differentially tagged SPINK1 proteins (not shown). These observations may indicate that in the presence of an inactive allele, compensatory mechanisms increase expression of the wild-type allele. On the other hand, co-transfection experiments do not appropriately model chromosomal bi-allelic expression and other interpretations of the results (e.g. homologous recombination between transfected plasmids) are also possible.

To investigate the intracellular fate of SPINK1 variants, epitope tagged versions of SPINK1 constructs were expressed and analyzed. Epitope tagging became necessary, because the rabbit polyclonal antibody gave strong non-specific and/or cross-reaction signals when tested on cell lysates. We placed the so-called Glu-Glu tag (EYMPME) at the C terminus of SPINK1. The epitope tagged SPINK1 construct was as efficiently secreted as untagged wild-type SPINK1

and the trypsin inhibitory activity was also uncompromised. In activity assays, conditioned medium from cells expressing epitope tagged versions of L14P and L14R SPINK1 exhibited no trypsin inhibition, whereas medium with epitope tagged L12F and wild-type SPINK1 inhibited trypsin comparably (not shown). On Western blots, the wild type and L12F constructs gave a comparably strong signal, while equal amounts of L14P and L14R media yielded no detectable SPINK1 (Fig 6B). Similarly, analysis of L14P and L14R cell lysates failed to reveal immunoreactive protein, indicating rapid intracellular degradation of the SPINK1 variants. As expected, wild-type and L12F SPINK1 proteins were clearly detected from cell lysates. Finally, to investigate whether the observed effects are dependent on the cell line used, CHO cells were also transfected with SPINK1 cDNAs. Western-blot analysis of conditioned media revealed the same secretion defect for the L14P and L14R variants, whereas the L12F variant was secreted as well as wild-type SPINK1 (Fig 6C).

DISCUSSION

Here we report that SPINK1 alterations which cause a complete block in secretion are associated with hereditary pancreatitis in heterozygotes. This is the first clear genetic-biochemical demonstration that severe loss-of-function *SPINK1* alterations are associated with a high risk of chronic pancreatitis, therefore, these alterations should be regarded as disease-causing rather than disease-modifiers. Potential examples of other severe *SPINK1* alterations in association with hereditary pancreatitis have been described previously (Witt et al., 2000; Le Maréchal et al., 2004; Masson et al., 2006). The results presented here support the notion that this group of SPINK1 alterations should be classified as “severe” and should be differentiated from “mild” alterations, particularly from the relatively frequent, archetypal N34S variant (Witt et al., 2001a; Le Maréchal et al., 2004; Masson et al., 2006). Mild alterations confer a smaller risk of pancreatitis, are typically found in association with idiopathic pancreatitis, and additional genetic and environmental risk factors are believed to influence their expression to a significant degree.

The L14R alteration is a novel variant, which we identified in two unrelated families, a Bulgarian family living in France, and a German family originating from the northern part of Germany. In both families, the alteration is inherited through several generations, thus establishing the diagnosis of autosomal dominant hereditary pancreatitis. The penetrance of the L14R alteration is high but incomplete, which is in agreement with the typical penetrance of dominant *PRSS1* (cationic trypsinogen) alterations associated with classic hereditary pancreatitis (Whitcomb et al., 1996; Le Maréchal et al., 2006). The newly described L14R alteration affects the same amino acid as the previously reported L14P alteration (Witt et al., 2000). The L14P alteration was originally identified in a 5-year old boy without a family history of pancreatitis. Interestingly, since that time, the mother, who also carried the alteration, developed pancreatitis. In addition to the two disease-causing alterations within the SPINK1 signal peptide, we also found a frequent polymorphic variant, L12F, in pancreatitis patients of African origin as well as in African controls, but not in white subjects. This variant served as an important control, demonstrating that not all signal peptide alterations are associated with pancreatitis and a direct relationship exists between the functional effects of the alterations and disease-association.

Functional analysis using transient expression in HEK 293T and CHO cells demonstrated that although mRNA expression was unaffected, both L14P and L14R abolished SPINK1 secretion, whereas L12F was secreted normally. The SPINK1 signal peptide conforms to the general paradigm of secretory signal peptides and comprises three functionally important regions; a positively charged N-terminal domain (5 amino acids), a hydrophobic central core (12 amino acids) and a polar C-terminal region (6 amino acids) including the signal-peptidase cleavage site (von Heijne G, 1990). The alterations studied here all disrupt the hydrophobic core region

and thus are expected to block translocation of preSPINK1 into the lumen of the endoplasmic reticulum. Alterations that affect the hydrophobic core of the signal peptide have been described and characterized in association with various human diseases. As shown in Table 1, the Leu→Pro change is a relatively frequent occurrence and a Leu→Arg alteration has also been detected. Functional analyses invariably demonstrated that these variations block secretion by inhibiting translocation of the preprotein into the endoplasmic reticulum. Therefore, signal peptide alterations that disrupt the hydrophobic core result in a loss of protein expression and consequently a loss of function.

It is arguable whether the constitutive secretory pathway of HEK or CHO cells used in our experiments adequately model the regulated secretory pathway of the pancreatic acinar cells. Unfortunately, there are no human acinar cell lines available which we could have used to characterize regulated secretion. The widely used rat acinar cell line AR42J (Jessop and Hay, 1980) exhibits diminished secretory function and its utility as a model to study acinus-related secretory phenomena is questionable. Given the difficulty in testing the SPINK1 variants in authentic acinar cells, we utilized epithelial cell lines that carry out constitutive secretion. In these cell lines, the alterations clearly impair secretion of SPINK1, which we interpret as a defect in targeting to the endoplasmic reticulum. Because targeting and translocation of secretory proteins to the endoplasmic reticulum is a universally conserved mechanism, we posit that the same defect is expected to occur in pancreatic acinar cells, eventually leading to decreased SPINK1 secretion through the regulated pathway as well. Direct measurements of SPINK1 levels from pancreatic juice sample of L14R or L14P carriers would provide strong corroborating evidence for our findings. However, juice samples from the patients described in this study were not available for analysis.

In summary, the present study demonstrates that the p.L14R and p.L14P signal peptide alterations result in essentially complete loss of SPINK1 secretion. Clinically, these loss-of-function alterations are associated with autosomal dominant hereditary pancreatitis, confirming the notion that severe *SPINK1* alterations represent strong risk factors for chronic pancreatitis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

ACKNOWLEDGMENTS

We thank the patients who participated in this study. We are grateful to Dr. A. Morali (Hôpital d'Enfants, Nancy, France) for contributing DNA samples.

Support for this study was received from the NIH (grant # DK058088 to M.S.-T.), the INSERM (Institut National de la Santé et de la Recherche Médicale to C.F.), the DFG (Wi 2036/2-1 to H.W.) and the Sonnenfeld-Stiftung, Berlin, Germany (to H.W.).

REFERENCES

- Arnold A, Horst SA, Gardella TJ, Baba H, Levine MA, Kronenberg HM. Mutation of the signal peptide-encoding region of the preproparathyroid hormone gene in familial isolated hypoparathyroidism. *J Clin Invest* 1990;86:1084–1087. [PubMed: 2212001]
- Bhatia E, Choudhuri G, Sikora SS, Landt O, Kage A, Becker M, Witt H. Tropical calcific pancreatitis: strong association with SPINK1 trypsin inhibitor mutations. *Gastroenterology* 2002;123:1020–1025. [PubMed: 12360463]
- Chandak GR, Idris MM, Reddy DN, Bhaskar S, Sriram PV, Singh L. Mutations in the pancreatic secretory trypsin inhibitor gene (PSTI/SPINK1) rather than the cationic trypsinogen gene (PRSS1) are significantly associated with tropical calcific pancreatitis. *J Med Genet* 2002;39:347–351. [PubMed: 12011155]

- Chen JM, Mercier B, Audrezet MP, Ferec C. Mutational analysis of the human pancreatic secretory trypsin inhibitor (PSTI) gene in hereditary and sporadic chronic pancreatitis. *J Med Genet* 2000;37:67–69. [PubMed: 10691414]
- Chen JM, Mercier B, Audrezet MP, Raguénés O, Quéré I, Férec C. Mutations of the pancreatic secretory trypsin inhibitor (PSTI) gene in idiopathic chronic pancreatitis. *Gastroenterology* 2001;120:1061–1064. [PubMed: 11265669]
- Drenth JP, te Morsche R, Jansen JB. Mutations in serine protease inhibitor Kazal type 1 are strongly associated with chronic pancreatitis. *Gut* 2002;50:687–692. [PubMed: 11950817]
- den Dunnen JT, Paalman MH. Standardizing mutation nomenclature: why bother? *Hum Mutat* 2003;22:181–182. [PubMed: 12938082]
- Gaia E, Salacone P, Gallo M, Promis GG, Brusco A, Bancone C, Carlo A. Germline mutations in CFTR and PSTI genes in chronic pancreatitis patients. *Dig Dis Sci* 2002;47:2416–2421. [PubMed: 12452372]
- Green PM, Mitchell VE, McGraw A, Goldman E, Giannelli F. Haemophilia B caused by a missense mutation in the prepeptide sequence of factor IX. *Hum Mutat* 1993;2:103–107. [PubMed: 8318985]
- Grussenmeyer T, Scheidtmann KH, Hutchinson MA, Eckhart W, Walter G. Complexes of polyoma virus medium T antigen and cellular proteins. *Proc Natl Acad Sci USA* 1985;82:7952–7954. [PubMed: 2415976]
- Harper LV, Hilton AC, Jones AF. RT-PCR for the pseudogene-free amplification of the glyceraldehyde-3-phosphate dehydrogenase gene (gapd). *Mol Cell Probes* 2003;17:261–265. [PubMed: 14580401]
- Hassan Z, Mohan V, Ali L, Allotey R, Barakat K, Faruque MO, Deepa R, McDermott MF, Jackson AE, Cassell P, Curtis D, Gelding SV, Vijayaravaghan S, Gyr N, Whitcomb DC, Khan AK, Hitman GA. SPINK1 is a susceptibility gene for fibrocalculous pancreatic diabetes in subjects from the Indian subcontinent. *Am J Hum Genet* 2002;71:964–968. [PubMed: 12187509]
- Jessop NW, Hay RJ. Characteristics of two rat pancreatic exocrine cell lines derived from transplantable tumors. *In Vitro* 1980;16:212.
- Kato M, Nanba E, Akaboshi S, Shiihara T, Ito A, Honma T, Tsuburaya K, Hayasaka K. Sonic hedgehog signal peptide mutation in a patient with holoprosencephaly. *Ann Neurol* 2000;47:514–516. [PubMed: 10762164]
- Kuwata K, Hirota M, Shimizu H, Nakae M, Nishihara S, Takimoto A, Mitsushima K, Kikuchi N, Endo K, Inoue M, Ogawa M. Functional analysis of recombinant pancreatic secretory trypsin inhibitor protein with amino-acid substitution. *J Gastroenterol* 2002;37:928–934. [PubMed: 12483248]
- Launz F, De La Salle C, Baas MJ, Schwartz A, Boval B, Cazenave JP, Caen JP. A Leu7Pro mutation in the signal peptide of platelet glycoprotein (GP)IX in a case of Bernard-Soulier syndrome abolishes surface expression of the GPIb-V-IX complex. *Br J Haematol* 2002;118:260–266. [PubMed: 12100158]
- Le Maréchal C, Chen JM, Le Gall C, Plessis G, Chipponi J, Chuzhanova NA, Raguénés O, Férec C. Two novel severe mutations in the pancreatic secretory trypsin inhibitor gene (SPINK1) cause familial and/or hereditary pancreatitis. *Hum Mutat* 2004;23:205.
- Le Maréchal C, Masson E, Chen JM, Morel F, Ruzsniowski P, Levy P, Férec C. Hereditary pancreatitis caused by triplication of the trypsinogen locus. *Nat Genet* 2006;38:1372–1374. [PubMed: 17072318]
- Masson E, Le Maréchal C, Chen JM, Frebourg T, Lerebours E, Férec C. Detection of a large genomic deletion in the pancreatic secretory trypsin inhibitor (SPINK1) gene. *Eur J Hum Genet* 2006;14:1204–1208. [PubMed: 16823394]
- Noone PG, Zhou Z, Silverman LM, Jowell PS, Knowles MR, Cohn JA. Cystic fibrosis gene mutations and pancreatitis risk: relation to epithelial ion transport and trypsin inhibitor gene mutations. *Gastroenterology* 2001;121:1310–1319. [PubMed: 11729110]
- Ozawa T, Takikawa Y, Niiya K, Ejiri N, Suzuki K, Sato S, Sakuragawa N. Factor VII Morioka (FVII L-26P): a homozygous missense mutation in the signal sequence identified in a patient with factor VII deficiency. *Br J Haematol* 1998;101:47–49. [PubMed: 9576180]
- Pidasheva S, Canaff L, Simonds WF, Marx SJ, Hendy GN. Impaired cotranslational processing of the calcium-sensing receptor due to signal peptide missense mutations in familial hypocalciuric hypercalcemia. *Hum Mol Genet* 2005;14:1679–1690. [PubMed: 15879434]

- Pfützer RH, Barmada MM, Brunskill AP, Finch R, Hart PS, Neoptolemos J, Furey WF, Whitcomb DC. PSTI/SPINK1 polymorphisms act as disease modifiers in familial and idiopathic chronic pancreatitis. *Gastroenterology* 2000;119:615–623. [PubMed: 10982753]
- Plendl H, Siebert R, Steinemann D, Grote W. High frequency of the N34S mutation in the SPINK1 gene in chronic pancreatitis detected by a new PCR-RFLP assay. *Am J Med Genet* 2001;100:252–253. [PubMed: 11343313]
- Rajpar MH, Koch MJ, Davies RM, Mellody KT, Kielty CM, Dixon MJ. Mutation of the signal peptide region of the bicistronic gene DSPP affects translocation to the endoplasmic reticulum and results in defective dentine biomineralization. *Hum Mol Genet* 2002;11:2559–2565. [PubMed: 12354781]
- Rinderknecht H. Activation of pancreatic zymogens. Normal activation, premature intrapancreatic activation, protective mechanisms against inappropriate activation. *Dig Dis Sci* 1986;31:314–321. [PubMed: 2936587]
- Schneider A, Suman A, Rossi L, Barmada MM, Beglinger C, Parvin S, Sattar S, Ali L, Khan AK, Gyr N, Whitcomb DC. SPINK1/PSTI mutations are associated with tropical pancreatitis and type II diabetes mellitus in Bangladesh. *Gastroenterology* 2002;123:1026–1030. [PubMed: 12360464]
- Seppen J, Steenken E, Lindhout D, Bosma PJ, Elferink RP. A mutation which disrupts the hydrophobic core of the signal peptide of bilirubin UDP-glucuronosyltransferase, an endoplasmic reticulum membrane protein, causes Crigler-Najjar type II. *FEBS Lett* 1996;390:294–298. [PubMed: 8706880]
- Threadgold J, Greenhalf W, Ellis I, Howes N, Lerch MM, Simon P, Jansen J, Charnley R, Laugier R, Frulloni L, Olah A, Delhaye M, Ihse I, Schaffalitzky de Muckadell OB, Andren-Sandberg A, Imrie CW, Martinek J, Gress TM, Mountford R, Whitcomb D, Neoptolemos JP. The N34S mutation of SPINK1 (PSTI) is associated with a familial pattern of idiopathic chronic pancreatitis but does not cause the disease. *Gut* 2002;50:675–681. [PubMed: 11950815]
- Truninger K, Witt H, Kock J, Kage A, Seifert B, Ammann RW, Blum HE, Becker M. Mutations of the serine protease inhibitor, Kazal type 1 gene, in patients with idiopathic chronic pancreatitis. *Am J Gastroenterol* 2002;97:1133–1137. [PubMed: 12014716]
- von Heijne G. The signal peptide. *J Membr Biol* 1990;115:195–201. [PubMed: 2197415]
- Whitcomb DC, Gorry MC, Preston RA, Furey W, Sossenheimer MJ, Ulrich CD, Martin SP, Amann ST, Gates LK Jr, Toskes PP, Liddle R, McGrath K, Uomo G, Post JC, Ehrlich GD. Hereditary pancreatitis is caused by a mutation in the cationic trypsinogen gene. *Nat Genet* 1996;14:141–145. [PubMed: 8841182]
- Witt H, Luck W, Hennies HC, Classen M, Kage A, Lass U, Landt O, Becker M. Mutations in the gene encoding the serine protease inhibitor, Kazal type 1 are associated with chronic pancreatitis. *Nat Genet* 2000;25:213–216. [PubMed: 10835640]
- Witt H, Hennies HC, Becker M. SPINK1 mutations in chronic pancreatitis. *Gastroenterology* 2001a; 120:1060–1061. [PubMed: 11265668]
- Witt H, Luck W, Becker M, Böhmig M, Kage A, Truninger K, Ammann RW, O'Reilly D, Kingsnorth A, Schulz HU, Halangk W, Kielstein V, Knoefel WT, Teich N, Keim V. Mutation in the SPINK1 trypsin inhibitor gene, alcohol use, and chronic pancreatitis. *JAMA* 2001b;285:2716–2717. [PubMed: 11386926]

Figure 1. Primary structure of human pancreatic secretory trypsin inhibitor (SPINK1). The amino acid sequence and alterations of the secretory signal peptide are indicated in the single-letter code.

Figure 2.

Identification of a new *SPINK1* missense alteration p.L14R, in two families with hereditary pancreatitis. Solid symbols represent patients with pancreatitis; open symbols indicate individuals without pancreatitis; the arrow points to the index patient. The numbers next to the symbols denote the current age and the age of disease onset (in parentheses). The genotypes of family members available for genetic analysis are also indicated. **Upper panel** Pedigree of the Bulgarian family living in France. **Lower panel** Pedigree of the German family.

Figure 3.

Effect of signal peptide alterations on *SPINK1* mRNA levels. **A.** Wild-type (wt) and L12F, L14R and L14P mutant *SPINK1* were transiently expressed in HEK 293T cells and total RNA was prepared and reverse-transcribed as described in *Materials and Methods*. The cDNA for *SPINK1* (**A**) and glyceraldehyde-3-phosphate dehydrogenase (**B**) were PCR-amplified and the semi-quantitative PCR-reactions were analyzed by agarose gel electrophoresis and ethidium bromide staining.

Figure 4.

Trypsin inhibitory activity in conditioned media of HEK 293T cells transfected with SPINK1 mutants. At the designated times, the capacity of conditioned media to inhibit human cationic trypsin was determined, as described in *Materials and Methods*. Residual trypsin activity is expressed as percentage of uninhibited trypsin activity. Data shown represent the mean $\pm$ SD of two parallel transfections. Solid squares, wild type SPINK1; open circles, mutant SPINK1.

Figure 5.

Secretion of wild type and mutant SPINK1 from HEK 293T cells. SPINK1 inhibitory activity was determined against human cationic trypsin as described in *Materials and Methods* and shown in Fig 4. Inhibitory activities were converted to SPINK1 concentrations using a calibration curve prepared with purified, recombinant SPINK1 and human cationic trypsin. For each transfection, a plasmid encoding β -galactosidase was co-transfected and at 48 hours β -galactosidase activity was determined from the lysed cells, as a measure of transfection efficiency. SPINK1 concentrations were then normalized to β -galactosidase activity. Data represent the mean $\pm$ SD of two experiments with two parallels each. Solid squares, wild type SPINK1; open circles, mutant SPINK1.

Figure 6.

Western blot analysis of SPINK1 expression in conditioned media and cell lysates of transfected HEK 293T cells. Aliquots of conditioned media (5 μ L) and cell lysates (10 μ L; ~3.5 mg/mL protein concentration) harvested 48 h post-transfection were subjected to polyacrylamide gel electrophoresis, transferred to Immobilon P membranes, and immunodetection was performed as described in *Materials and Methods*. **A.** SPINK1 in conditioned media was detected with a polyclonal rabbit antibody raised against recombinant human SPINK1. **B.** Expression of epitope-tagged SPINK1 proteins in conditioned media and cell lysates was visualized with a commercial anti-Glu-Glu tag antibody (Abcam). **C.** SPINK1

in the conditioned media of CHO cells was detected with the PSKAN2 monoclonal mouse IgG antibody against human SPINK1 (Mobitec; Gottingen, Germany).

Table 1

Mutations that affect the hydrophobic core of the signal peptide in human diseases. The OMIM database numbers and literature references are also indicated.

Protein	Mutation	Signal peptide	Disease association
prepro-parathyroid hormone (preproPTH) MIM * 168450	C18R	MIPAKDMAKVMIVMLAICFLTKSDG	familial isolated hypoparathyroidism MIM #146200; Arnold et al., 1990
coagulation factor IX (Christmas factor)	I17N	MQRVNMIMAEPSLITICLLGYLLSA	hemophilia B MIM #306900; Green et al., 1993
bilirubin UDP-glucuronosyltransferase MIM*191740	L15R	MAVESQGGRLVLGLLLCVLGPVSHA	Crigler-Najjar disease type II MIM #606785; Seppen et al., 1996
coagulation factor VII (Morioka)	L13P	MVSOALRLLCLLLGLOGCLA	hereditary factor VII deficiency MIM #227500; Ozawa et al., 1998
human sonic hedgehog (SHH) MIM *600725	L17P	MLLLARCLLLVLVSSLVCSGLA	holoprosencephaly 3 MIM #142945; Kato et al., 2000
platelet glycoprotein IX (GPIX) MIM*173515	L7P	MPAWGALFLLWATAEA	Bernard-Soulier syndrome MIM #231200; Lanza et al., 2002
dentine sialophosphoprotein (DSPP) MIM* 125485	Y6D	MKIITYFCIWAWAWA	dentine dysplasia type II MIM #125420; Rajpar et al., 2002
calcium-sensing receptor (CaSR) MIM +601199	L1IS L13P T14A	MAFYSCCWVLLALTWHTSA	hypocalciuric hypercalcemia. MIM #145980; Pidasheva et al., 2005