Published in final edited form as:

J Comp Neurol. 2008 August 10; 509(5): 526–550. doi:10.1002/cne.21762.

The glial investment of the adult and developing antennal lobe of *Drosophila*

Lynne A. Oland¹, John P. Biebelhausen¹, and Leslie P. Tolbert¹

¹ Arizona Research Laboratories Division of Neurobiology, University of Arizona, Tucson, AZ 85721, USA

Abstract

In recent years, the *Drosophila* olfactory system, with its unparalleled opportunities for genetic dissection of development and functional organization, has been used to study the development of central olfactory neurons and the molecular basis of olfactory coding. The results of these studies have been interpreted in the absence of a detailed understanding of the steps in maturation of glial cells in the antennal lobe. Here, we present a high-resolution study of the glia associated with olfactory glomeruli in adult and developing antennal lobes. The study provides a basis for comparison of findings in *Drosophila* with those in the moth *Manduca sexta* that indicate a critical role for glia in antennal lobe development. Using flies expressing GFP under a *Nervana2* driver to visualize glia for confocal microscopy, and probing at higher resolution with the electron microscope, we find that glial development in *Drosophila* differs markedly from that in moths: glial cell bodies remain in a ring around the glomerular neuropil; glial processes ensheath axon bundles in the nerve layer but likely contribute little to axonal sorting; their processes insinuate between glomeruli only very late and then form only a sparse, open network around each glomerulus; and glial processes invade the synaptic neuropil. Taking our results in the context of previous studies, we conclude that glial cells in the developing *Drosophila* antennal lobe are unlikely to play a strong role in either axonal sorting or glomerulus stabilization and that in the adult, glial processes do not electrically isolate glomeruli from their neighbors.

Keywords

glial cells; olfactory lobe; olfactory bulb; glomeruli

INTRODUCTION

Interactions between the neuronal and glial subsystems of the brain now have well recognized roles in numerous developmental processes, and in the mature brain operate in a dynamic partnership. Molecular genetic studies of the *Drosophila* nervous system have revealed it to be an excellent model in which to dissect the molecular basis of normal and disordered development and mature structure and function. The power of this approach has recently been extended to neuron-glial interactions (Chotard and Salecker, 2004; Freeman, 2006; Parker and Auld, 2006). In particular, the olfactory lobe of *Drosophila* is rapidly becoming a model system in which to study a variety of disorders for which the underlying mechanism may be a disruption in such interactions (Read et al., 2005; MacDonald et al., 2006). Although the general pattern of the glial investment of the adult antennal lobe has

Corresponding author: Lynne A. Oland, PhD, University of Arizona, Arizona, Research Laboratories Division of Neurobiology, PO Box 210077, Tucson, AZ 85721, ph: 520-621-6671, fax: 520-621-8282, email: lao@neurobio.arizona.edu.
Associate Editor: Ian Meinertzhagen

been noted (Jhaveri et al., 2000; Jhaveri and Rodrigues, 2002; Ang et al., 2003), no detailed study of glial structure and organization is available.

Considerable effort has been devoted to understanding the sequence of cellular changes that occurs during development of the *Drosophila* olfactory (antennal) lobe (Stocker et al., 1995; Jhaveri and Rodrigues, 2000, 2002; Jhaveri et al., 2000; Jefferis et al., 2004; Komiyama et al., 2007), but mainly from a neuronal perspective. A few studies have included results that briefly describe the state of glial investment at several times during metamorphic development (Jhaveri et al., 2000; Jefferis et al., 2004) when the antennal lobe develops its mature structure, but the sum of the data so far provides only an overview of the lobe's glial investment during development.

The goals of this study were to combine confocal and electron microscopy to determine the timing and extent of glial investment of the developing and adult antennal lobe of *Drosophila*, to place the developmental events in the context of known neuronal development in this system, and to compare the glial investment of the *Drosophila* antennal lobe with that of vertebrate and invertebrate olfactory systems in which the glial investment of the first-order olfactory neuropil has been well studied (e.g., Tolbert et al., 2004; Treloar et al., 1999; Valverde 1999).

Part of this work has been presented as an abstract (Oland et al., 2007).

MATERIALS AND METHODS

Animals

Stocks of *Drosophila melanogaster* were raised on standard cornmeal with yeast diet (Tucson *Drosophila* Stockcenter; stockcenter.arl.arizona.edu) at 25°C. Flies from the *Nrv2*-GAL4, UAS-GFP line (gift from P. Salvaterra) were used to visualize the cell bodies and processes of glial cells in the *Drosophila* central nervous system. The *Nervana2* (*Nrv2*) gene encodes a Na⁺, K⁺-ATPase β -subunit (Sun et al., 1999). Fluorescence is observed throughout the central nervous system in neurons and neuronal processes in embryos, but after embryogenesis, GFP expression is mainly restricted to the nuclei and processes of most, but not all, glial cells (Sun et al., 2001).

Pupae were staged during development by marking wandering third instar larvae in vials when they ceased wandering (denoted as 0 hours after puparium formation, APF). Animals were dissected at selected time points throughout adult metamorphic development and processed as described below.

Immunocytochemistry

Adult flies were anesthetized with carbon dioxide before dissection. Both adult and pupal brains (n=3–5 for each stage for each type of experiment) were dissected into Tris-buffered saline (20 mM Tris-HCl, 150 mM NaCl, pH 7.2) or phosphate-buffered saline (7.8 mM Na₂HPO₄·7H₂O, 2 mM NaH₂PO₄·H₂O, 150 mM NaCl, pH 7.4) and fixed overnight at 4°C in 4% paraformaldehyde in Tris or phosphate-buffered saline (TBS, PBS). In some cases, the initial fixation step was done in a research-grade microwave (Pelco) at 18°C (Power 2: 2 min on, 2 min off, 2 min on, 2 min off; then power 6: 10 sec on, 20 sec off, 10 sec on), after which the brains are placed in fresh fixative solution overnight. The brains were washed 3–5 × for 5–10 min each in buffered saline containing Triton X-100 and 0.1% sodium azide (TBSAT), blocked for 1–2 hrs, incubated in primary antibody ON at RT on a shaker, washed in buffered saline 3–5 × for 5–10 min each, incubated in secondary antibody on a shaker for 2 hrs at RT or overnight at 4°C, and finally washed 1 × 5 min in buffered saline, 2 × 5 min in 0.1M Tris or phosphate buffer, 1 × 15 min in 50:50 glycerol:distilled H₂O, and

80:20 glycerol:distilled H₂O. Brains were mounted in 80:20 glycerol:distilled H₂O between 2 coverslips separated by 3 strips of coverslip glued in position to form a small rectangular chamber with one open side.

For all antibodies, non-specific binding was blocked by placing brains in TBS or PBS with 0.1–0.2% Triton-X 100 and with 1% IgG-free BSA for 1 hr, and then incubating in primary antibody. Mouse anti-Synapsin 1 (3C11, Developmental Studies Hybridoma Bank (DSHB, Iowa City, IA) was used at 1:10 dilution in TBSA with 0.2–0.5% Triton, mouse anti-Repo antibody (8D12, DSHB) was used at 1:10 in PBS with 0.2% Triton, and mouse anti-Neuroglian (BP104, DSHB) was used at 1:50 in PBS with 0.2% Triton. All antibodies were raised against *Drosophila* epitopes. The secondary antibody was goat anti-mouse Cy3-conjugated antibody (Jackson ImmunoResearch, West Grove, PA) at 1:250 in buffered saline with 0.1% Triton. Control brains for nonspecific immunolabeling by the secondary antibody were prepared as described above, but with the primary antibody omitted, and imaged using microscope settings identical to those used for labeled brains.

Antibody characterization—C311 (anti-Synapsin, or anti-SYNORF1) is an affinity-purified IgG2b antibody raised in mouse against a glutathione-S-transferase Synapsin fusion protein. It recognizes nucleotide sequence 630–1970 of a cDNA clone expressed in *E. coli* BL21 (Klagges et al., 1996; Godenschwege et al., 2004). In synapsin mutants, Godenschwege et al. (2004) confirmed that in the absence of the protein, SYNORF1 labeling also is absent.

BP104 (anti-Neuroglian) is an IgG1 antibody raised in mouse against the nervous-system specific 180 kD form *Drosophila* Neuroglian (Hortsch et al., 1990). In affinity columns, the 180 kD form of Neuroglian could be purified to homogeneity and the derived amino acid sequence was identical to the sequence for the amino terminus of the 167 kD form (Bieber et al., 1989; Hortsch et al., 1990). The antibody recognizes an epitope in the cytoplasmic extension of the long form of the protein (Hortsch et al., 1990). Null mutations in the *nrg* gene are embryonic lethal mutations but two hypomorphic mutations with greatly reduced expression have been described Hall and Bieber (1996). Neuroglian protein expression was assessed using BP-104 in *nrg*⁻³ mutants, which are temperature-sensitive. When assessed during development at the restrictive temperature, labeling was eliminated in normally Neuroglian-positive neuronal processes and restricted to cell bodies (Hall and Bieber, 1996), and a Western Blot developmental series of wild-type *Drosophila* embryos indicated a gradually increasing expression of Neuroglian (Hortsch et al., 1990).

8D12 (anti-Repo) is an IgG1 antibody raised in mouse against amino acids 218–612 of *Drosophila* repo protein fused to a 6-histidine-tag and expressed in *E. coli*. (Campbell et al., 1994; Alfonso and Jones, 2002). *repo*, which is a putative target for *gcm* (glial cells missing), the primary regulator of glial cell fate in *Drosophila*, is a homeo-domain protein required for the differentiation and maintenance of most glial cells in the *Drosophila* nervous system, the midline glia excepted (Xiong et al., 1994; Halter et al., 1995).

Confocal microscopy

Whole-mount brains were viewed using either a Nikon PCM 2000 or a Zeiss 510 Meta laser scanning confocal system equipped with argon, green Helium Neon, and red Helium Neon lasers and appropriate filter sets. Confocal Assistant (copyright Todd Brelje; distributed by Bio-Rad), Zeiss LSM Image Browser, Corel Photopaint, and Corel Draw were used to create projections of confocal image stacks, to adjust image contrast and brightness, and to pseudocolor images. 3D images were generated using Amira's (version 4.1) Voltex2 module (2D hardware accelerated implementation for direct volume rendering) for the single channel (GFP) scalar data.

The number of neuropil glial cells in the adult antennal lobe was determined in 4 lobes from *Nrv2* animals that also were labeled with the Repo antibody. For each one, confocal images through the entire antennal lobe were collected at 40x, with a zoom of 1.5 and z interval of 0.5 μm , then projected into 2D substacks that were 10 μm in depth. The positions of glial cell bodies in each image were marked on an acetate sheet that was carried forward through the entire series of images to finally include a mark for every glial cell found in the images. When a glial cell body appeared in more than one image, it was counted in the image in which its profile was the largest.

Electron microscopy

Dissections were carried out in modified HL-3 medium (70 mM NaCl, 5 mM KCl, 1.5 mM CaCl_2 , 4 mM MgCl_2 , 10 mM NaHCO_3 , 5 mM trehalose, 115 mM sucrose, 5 mM Hepes, pH 7.2). For pupal animals, the head cuticle was opened and fat cells overlying the brain gently blown off with a stream of medium from a broken-tipped micropipette. For young pupal animals the head cuticle was only partially removed, leaving the cuticle supporting the eyes intact; for older pupae and adult animals, the brain was removed entirely from the head. Brains were transferred into 600 μl fresh fixative solution in small baskets constructed from beem capsules and Teflon mesh. These were placed in polyethylene Petri dishes containing the fixative solution and processed in the microwave as described above. The fixative solution was then replaced with fresh fixative solution and the brains were stored for up to 3 days at 4°C. The brains were washed in 0.1 M cacodylate solution with 7 mM CaCl_2 , then post-fixed in 2% osmium tetroxide in modified cacodylate buffer (0.1 M cacodylate, 0.02% CaCl_2 , and 2.25% sucrose) for 2 hrs, washed in modified cacodylate buffer 2×10 min, and processed in the microwave oven, first through a graded ethanol series (Temp: 18°C. Power 2 for 45 sec each), and then through a graded Epon/Araldite in 100% ethanol series (2 parts 100% EtOH: 1 part Epon/Araldite with DMP-30, 1:1, 1:2, 100% Epon/Araldite with DMP-30 $\times 2$) (Temp: 18°C. Power 2 for 3 min each). Finally, the brains were embedded in Epon/Araldite with DMP-30 and hardened in a 60°C oven ON.

The embedded brains were thin-sectioned on an RMC PowerTome ultra-microtome, picked up on Formvar-coated slot grids, and viewed in a JEOL 1200X electron microscope equipped with a Gatan digital camera. Some of the sections were lightly stained with lead citrate. Images collected with Gatan Digital Micrograph software were adjusted in CorelDraw for contrast and brightness. Montages comprising a half to an entire cross-section of the antennal lobe at each stage examined were constructed in Adobe Photoshop or Corel Draw and printed on a large format printer.

Fixative solution for electron microscopy included 2.5% glutaraldehyde, 2.5% paraformaldehyde, and 7mM CaCl_2 in 0.1 M cacodylate buffer, pH 7.3 (Meinertzhagen and O'Neil, 1991). Modified HL-3 solution consisted of 70 mM NaCl, 5mM KCl, 20 mM MgCl_2 , 2.2 mM NaOH, 10 mM NaHCO_3 , and 5 mM BES, pH 7.2.

RESULTS

In *Drosophila*, as in other insects, glial cell populations typically are distinguished from one another based on their positions in the CNS as well as the size, shape, and staining characteristics of their cell bodies and nuclei (Strausfeld, 1976; Ito et al., 1995; Edwards and Tolbert, 1998). In this study, we focused primarily on the neuropil-associated glial cells.

Glial investment of the neuropil of the adult antennal lobe: confocal microscopy

The glial-cell investment of the neuropil of the adult antennal lobe was examined using a *Nervana2-Gal4*, UAS-GFP transgenic line in which glial cell bodies and processes in the

Drosophila central nervous system express GFP (Sun et al., 1999, 2001). Both cell body (cortical) glia and neuropil glia were consistently GFP-positive in the antennal lobes of *Nrv2* metamorphosing animals, but in adults, neuropil glia always were GFP-positive while the cell body glia were GFP-positive in most, but not all, preparations. Although not all neuropil glial cells throughout the brain express GFP at the stages examined in the *Nrv2* animals, double-labeling experiments with anti-Repo (Fig. 1), which labels glial cell nuclei, showed that 95% of the Repo-positive neuropil glial cells in the *Nrv2* antennal lobe also were GFP-positive. In the 4 adult antennal lobes in which glial cells were counted, only 2 cells were found that were GFP-positive, but Repo-negative. The high level of congruence in the antennal lobe between GFP and Repo-positive labels indicates that the use of the *Nrv2* line provides an accurate view not only of the positioning of glial cell bodies but also of the investment of the neuropil by glial processes. The number of glial cells in the adult antennal lobe, including both the Repo-positive and Repo-negative cells, was ~ 80 (76, 78, 80, and 85). No consistent spatial relationship was found between the position of glial cell bodies and individual glomeruli that would suggest the possibility of a glia-based positioning signal for establishing glomerular territories. Images of sections of the antennal lobe in the *Nrv2* strain double-labeled with an antibody against Synapsin highlight the thin glial net cast around each glomerulus (Fig. 2A,B). The GFP-labeled glial processes around the glomeruli are often velate (Fig. 2C,E), sometimes branched, and the network of glial processes is usually perceived as a border only when several sections are stacked together; in single optical sections, it is often difficult to locate the boundaries between glomeruli. Glial processes also invade glomerular territory, appearing throughout the body of each glomerulus, but at different densities in different glomeruli, as well as within different regions of the same glomerulus (Fig. 2A–C). Sometimes the processes seen within the neuropil of glomeruli form tiny circles when viewed at the light-microscopic level (Fig. 2C; 12B); in electron micrographs they are seen to be encircling large dendritic processes or bundles of smaller dendrites (Fig. 5B,E).

Figure 3 shows a series of images of 5- μ m-thick stacks of optical sections viewed from the most anterior surface of the antennal lobe to the posterior region adjacent to the protocerebral neuropil, permitting examination of the glial investment through the entire extent of the antennal lobe neuropil. The glomerular envelopes, which are not robust anywhere in the lobe, and indeed the glomerular structure of the neuropil, are often more difficult to resolve in the posterior region of the antennal lobe (Fig. 3: compare sections 1–90 with section 91–120, glomerular borders less conspicuous in the latter).

Glial processes are most abundant in association with primary neurite tracts entering the antennal lobe neuropil from the neuronal cell body clusters, especially the lateral cluster (Fig. 1B, 2B, 3: sections 41–70), and it is only along these tracts that glial cell bodies may be seen within the neuropil; in all other cases they reside at the perimeter of the neuropil. A small cluster of glial cell bodies (typically 2–4 cells) encircles the entrance of the lateral-cluster primary neurite tract into the neuropil; these glial cells extend long, parallel processes along that tract.

The neuropil glial cells also extend processes throughout the coarse central neuropil of the lobe, where they form a tracery of processes (Fig. 3, sections 61–100); images from the electron microscope reveal that these processes typically ensheath or partially ensheath the large dendrites (Fig. 5E). They also ensheath the axonal bundles that extend out of the lobe to higher order centers in the brain (Fig. 4A,B). The axons of olfactory receptor neurons form a layer around the antennal lobe neuropil and this nerve layer is invested with glial processes that arise, at least partially, from the same glial cells that invest the neuropil (Fig. 2C,D). It has been reported in larval *Drosophila* brain that cortex (cell body) glia also contribute to the sheath surrounding neuropil regions (Pereanu et al., 2005).

The antenno-cerebral tract carrying the axons of projection neurons to higher order brain centers is enveloped by glial cell processes that are present as the tract forms within the coarse central neuropil of the lobe (Fig. 3: sections 91–120, 4A). Glia also surround the antennal commissure, as noted by Yao et al. (2007), and ensheath separate tracts within it. Small clusters of 4–6 glial cell bodies gate either side of the commissure (Fig. 4B).

3D projections were made through portions of 3 adult *Nrv2* antennal lobes that were labeled with an antibody against *Drosophila* neuroglial. This antibody labels a substantial sub-population of ORN axons. Figure 4C shows a frame from one of these projections (for movie showing rotation of the image stack, see Supplementary material 1). The image clearly illustrates the porous or net-like character of the glomerular envelopes as well as the variability in infiltration of the glomerular neuropil by glial processes. Some of the glial processes are velate; most are fine and sparsely branching. The arrow marks the position of the primary neurite tract that arises from the lateral cluster of neuronal cell bodies.

Glial investment of the neuropil of the adult antennal lobe: electron microscopy

Because the glial processes investing the *Drosophila* antennal lobe are often extremely fine, and because the Repo-labeling of the antennal lobe showed that ~ 5% of the neuropil glial cells were not GFP-positive, we used the electron microscope to ensure that the picture of the glial investment that emerged from the confocal images of *Nrv2* antennal lobes was a representative one. At the EM level, glial cell processes were identified by the presence of ribosomes and glycogen particles and by their typically darkly staining cytoplasm (Figs. 5,6; Tolbert and Hildebrand, 1981).

Ultra-thin sections were cut at the level of the lateral-cluster primary neurite tract and/or the antennal nerve entry site into the antennal lobes of 5 adult *Nrv2* brains. Electron micrographs were made across the entire antennal lobe neuropil, the individual images were assembled into large montages, and the glial processes highlighted. Based on these initial images, higher power electron micrographs were made of selected regions.

The most striking feature of these EM cross-sections was the lack of an obvious visual signature that distinguished individual glomeruli from each other; scanning across the entire neuropil revealed neither large dendritic branches, as appears in cockroach (Ernst et al., 1977), nor a continuous glial border, as appears in the moth (Tolbert and Hildebrand, 1981), as indicators that the edge of a glomerulus has been crossed. Consistent with the light-level data, the glial investment of the neuropil of the adult lobe was not extensive, and differed markedly in distribution compared to that of the moth (Tolbert and Hildebrand, 1981; Oland et al., 1999), rodent or opossum (Pinching and Powell, 1971; Valverde and Lopez-Mascaraque, 2004).

Glial processes could be followed short distances from the perimeter of the neuropil toward the center of the lobe (Fig. 5B) but nowhere was there a complete glial border around glomerular neuropil, which was defined by the presence of a high density of synapses in regions that had a high proportion of fine processes and that were located beneath the border of glial cells surrounding the entire neuropil. Glial processes also could be found within glomerular neuropil (Fig. 5D). Although serial sections were not examined, the lack of any recurring pattern of spatial relationship with pre- or post-synaptic processes at the thousands of synapses present in 10 complete cross-sections of the adult antennal lobe neuropil suggests that these glial processes are not engaged at synaptic sites in modulation of synaptic activity. Large axonal and dendritic profiles, whether in glomerular neuropil (Fig. 5B,D), in the coarse central neuropil (Fig. 5E), or entering the neuropil in the primary neurite tracts (Fig. 5C) were ensheathed by thin glial processes. Those in the glomerular or coarse neuropil typically were individually ensheathed, either partially or fully, while those

traveling in tracts were typically ensheathed as bundles. The glial processes at the outer edges of the primary neurite tract often were multi-layered (Fig. 5C), and here scalariform-like junctions could be found linking glial processes. No clear evidence of any synapse-like profiles or other junctions between neuronal and glial processes was found.

Examination of the base of the adult antennal nerve close to the place at which it enters the lobe showed glial processes arranged in parallel with the incoming olfactory receptor axons, ensheathing them until they enter the lobe with no consistent evidence of any characteristic glial structure or junctional type to mark the transition from peripheral nerve to CNS (not shown). As described by Stocker et al. (1979) for more distal portions of the nerve, the proximal nerve contains bundles of small or large axons and most of the large axons are ensheathed by glia while the smaller ones are ensheathed as bundles. In the nerve layer of the antennal lobe, glial processes ensheathed small bundles of axons (Fig. 5A). Some regions of the glial border around the neuropil did not include axonal bundles, instead comprising only a thin border that separated neuronal cell bodies from the glomerular synaptic neuropil (Fig. 6A).

In one preparation, ultra-thin sections were cut at 10- μ m intervals in the horizontal plane from the dorsal to the ventral edge of the antennal lobe neuropil with the goal of verifying the finding at the light-microscopic level that the glial investment of the lobe varied from the anterior to posterior regions of the lobe. At each level, a montage of the entire neuropil was made and the glial processes highlighted. The glial investment of the neuropil was more sparse and glomerular borders were most diffuse in the posterior region (data not shown), confirming the light-level observations described above (Fig. 3). The poor glomerular definition in the posterior region is consistent with the findings of others (Laissue et al., 1999). In addition to the processes of neuropil glial cells, the antennal lobe is also invested lightly by tracheoles (Fig. 5A,B). Confirming data reported by Pereanu et al. (2007) for the *Drosophila* brain in general, the few tracheoles found in the antennal lobe neuropil were always found to be associated with glial cell processes (Fig. 5B). A similar pattern was described for the moth antennal lobe, where most tracheoles, though not all, are found within the glial envelopes surrounding glomeruli or associated with glial processes in the coarse central neuropil (Oland and Evans, 2000).

Development of the glial investment of the antennal lobe: confocal microscopy

Previous studies of the developing *Drosophila* antennal lobe neuropil by others (Jhaveri et al., 2000; Jhaveri and Rodrigues, 2002; Jefferis et al., 2004) have provided an overview of the development of its glial investment, which indicated that the investment of the antennal lobe neuropil developed too late for glial cells to be involved in the process of odotopic map development or the subsequent process of glomerulus construction. The apparent lack of a role for glial cells in these processes stands in stark contrast to their roles in glomerulus development in the moth, and likely in marsupials and mammals. Because we wished to know whether glial cells associated with the neuropil might be in position to interact with neurons in developmentally important ways during any phase of development of the primary olfactory pathway in *Drosophila*, and because the finest glial processes are likely to be undetectable at the light level, we completed a developmental series both to confirm the general results of the previous studies and to provide a finer grain view of glial changes as receptor axons grow into the lobe and as glomeruli mature.

Previous work had demonstrated that the antennal lobe neuropil is demarcated by glia at 12% of development, prior to the arrival of ORN axons at ~18% of development (Jefferis et al., 2004; Jhaveri et al., 2000; Jhaveri and Rodrigues, 2002). That work further suggested that glial processes begin to extend into the neuropil at around 50% of development (Jhaveri et al., 2000; Jefferis et al., 2004). Based upon this work and time points significant for the

development of the other cell classes in the lobe (see Figs. 15, 16), the following stages of pupal development were selected for study: 25, 30, 33, 40, 45, 50, 60, 67, and 75%.

At 25% of development, proliferation of glial cells surrounding the neuropil has ceased and ORN axons have arrived at the antennal lobe (Stocker et al., 1995; Jhaveri et al., 2000; Jhaveri and Rodrigues, 2002; Jefferis et al., 2004). The glial border is almost unbroken around the circumference of the neuropil although it may only a few microns thick in some places. As the axons enter, they course within and below the glial layer, encountering along the way mainly unbranched glial processes extending from the glial cell bodies toward the neuropil (Fig. 7A,B, L). During the period through 50% of development (Fig. 7C–F,L), glial processes gradually surround bundles of axons and some glial processes begin to form a boundary between the inner face of the nerve layer and the neuropil. In sections where the apical regions of glomeruli are visible, thick bundles of ORN axons that have become associated with a particular glomerulus can be seen in cross-section (Fig. 7F, 2D inset showing similar cross-section through a mature nerve). During subsequent stages, axonal fascicles in the nerve layer become further subdivided and the sheaths around them become more robust (Fig. 7G–L). Consistent with the EM data (see below), fortuitous sections show axons traveling long distances in the nerve layer without obvious changes in fascicle association.

Turning to the development of the glial investment of the neuropil, Figure 8A shows that at 25% of development there was no extension of glial processes into the neuropil even as the branches of ORN axons begin to invade the neuropil. By 30% of development, the first glial process ingrowth was seen within the neuropil in the form of a short fringe of processes projecting into the neuropil for a few microns (Fig. 8B) along sections of the glial border. This phase of ingrowth continues from 30% through 40%, as the glial fringe expands into the neuropil (Fig. 8C,D), but there is no evidence in the antennal lobes of *Nrv2* animals that glial cells have extended processes deep into the neuropil at this time. Glomerulus formation has already begun, however, with projection neurons already forming tufted arbors and ORN axons beginning to grow into the territory occupied by the dendrites of projection neurons (Jhaveri et al., 2000; Jhaveri and Rodrigues, 2002; Jefferis 2004; Hummel and Zipursky, 2004).

By 45% of development, the glial fringe around the neuropil has further thickened, particularly in the region of the lateral primary neurite tract as well as in the medial portion of the lobe (Fig. 8E, 11A). A few distinct glial processes are evident within the neuropil by 50% when glomeruli first become consistently detectable with synaptic markers (Fig. 8F, 11A–B; also Jefferis et al., 2004). At this point, glial processes extend toward the center of the antennal lobe, although the amount of ingrowth along the entirety of the glial border is quite modest. The greatest ingrowth of glial processes remained in the region of the primary neurite tract arising from the lateral cluster of neuron cell bodies (Fig. 8F; 11A,B).

As development continues through 60%, there is continued ingrowth of glial processes from the glial border (Figure 9A,B; 11C). The number of glial processes has increased, but the boundaries of glomeruli, though detectable with synaptic markers, could not be distinguished based upon their glial surrounds. A few glial processes could be followed from their originating cell body to near the center of the neuropil, and by 67% of development some processes had begun to branch and contribute to the formation of the glial net that eventually surrounds glomeruli (Figs. 9D–F; 11C,D) and to the investment of large processes in the coarse central neuropil (Fig. 9C). Only at this stage was it possible to begin to discern an organized glial structure bounding glomeruli (Figs. 9D, 11D), though the mature neuronal architecture of the glomeruli already has been well established (Hummel et al., 2003). Though now developed enough to be a rudimentary net, the structure is

recognizable as a boundary only when several optical sections were stacked together. Hereafter the net slowly becomes more refined, but never becomes a complete border. The extensive investment and further thickening of the glial processes investing the primary neurite tracts are readily apparent by 75% of development (Fig. 10A, 11D,E). Furthermore, fine glial processes now widely invest the synaptic neuropil of the glomeruli (Fig. 10B, 11E). At the time resolution we used, we were unable to determine with certainty differences in the timing of establishment of the glomerular borders and invasion of the glomerular neuropil by glial processes. We did, however, have some preparations in which we could identify glomerular borders, but saw few glial processes in the glomerular neuropil, suggesting that glial invasion of the glomerular neuropil likely lags slightly behind establishment of the glial net around the glomerular surface.

Figure 11 shows a developmental series through roughly the middle of the antennal lobe in which the developing glomerular neuropil in *Nrv2* antennal lobes is labeled with anti-Synapsin. In this context, the gradual invasion of glial processes into the neuropil is clearly shown to occur after glomerular structure is well established.

Development of the glial investment of the antennal lobe: electron microscopy

Electron microscopy of preparations at each of the stages examined with the confocal microscope was carried out to ensure that the finest glial processes were detected. Before approximately 50% of development, only a few thin glial processes extend into the neuropil, typically in irregular patches along the perimeter of the neuropil except in the region of the primary neurite tracts, where glial processes consistently appear intercalated among the large neurites by 50% of development (Fig. 12A) and also ensheath the tract as a whole. By 60–67%, the primary neurite tract of the lateral cluster of neuronal cell bodies is extensively invested, confirming the data acquired with the confocal microscope (Figs. 9–11). Glial processes also ensheath large dendrites within the glomerular neuropil, either partially or fully (Fig. 12B). The network of processes that in the confocal images was seen to be investing the new glomeruli was visible in the EM only as small, discontinuous profiles of processes scattered in the neuropil (Fig. 12C), consistent with what would be expected when taking 60-nm-thick sections through a network of processes.

In the nerve layer, bundles of olfactory receptor axons are well ensheathed by glial processes by 50%–67% of development (Fig. 12D,E). Though the sizes of the fascicles varied widely, there was little evidence of a local region of sorting either in the nerve or within the nerve layer, such as loosening of fascicles, axonal profiles with different orientations in the same region of the nerve, or sudden differences in axonal morphology as the axons turned toward the neuropil from the nerve layer (Oland et al., 1998; Au et al., 2002; Treloar et al., 2002). The overall investment pattern seen in the adult gradually develops over the remaining stages of metamorphosis.

Although the region in which the antennal nerve enters the neuropil in the adult is not organized in a manner that is markedly different from the pattern seen in more distal parts of the nerve, i.e. distinguished either by a varied orientation of axonal bundles as seen in the EM or by the pattern of glial cells seen in the confocal microscope, we asked whether the entry region of the antennal nerve is distinctive in the developing animals, especially during the early period of axon ingrowth (Fig. 13; see Supplementary material 2 for rotations), as it is in *Manduca* (Rössler et al., 1999). Bundles of receptor axons, which were labeled in *Nrv2* animals with antibodies against the IgG superfamily member Neuroglian (Hall and Bieber, 1997), show no evidence of extensive re-sorting (Figs. 7G; 13A,B,D) in any discrete region of the antennal nerve equivalent to the glia-rich sorting zone seen in the moth (Rössler et al., 1999) or in the nerve layer within the lobe (Figs. 7E,I; 13D) equivalent to the inner nerve layer of the vertebrate olfactory bulb (Au et al., 2000). Rather, small bundles of axons in the

nerve gradually shift position between larger fascicles as they travel to the lobe; the positions of the larger fascicles also change gradually relative to each other, much like the fiber bundles in a rope (Fig. 13A). The entry to the lobe is not marked consistently by any obvious glial structure at early developmental times, although a single large glial cell body was present adjacent to the entry region in many preparations at 40% of development and younger (Fig. 13C). The entry region is invested lightly by glial processes at least as early as 25% of development. Through 50% of development, glial cell bodies investing the nerve generally remain just distal to the region in which the olfactory and mechanosensory branches of the nerve split, leaving the olfactory entry region invested only lightly with glial processes. Receptor axons crossing this region (a distance of about 20 μ m) change direction rather abruptly to enter the nerve layer that surrounds the neuropil (7G; Fig. 13B,D), and then separate into smaller fascicles that spread widely and course in the nerve layer in loose tracts to their glomerular targets (Fig. 13D, see Supplemental material 2 for rotations).

DISCUSSION

The primary olfactory pathways of insects in general have an organizational plan similar to that of mammals, including humans, and thus provide useful model systems in which to study cellular and molecular interactions fundamental to brain development. We have provided here a detailed study of the glial investment of the neuropil of the adult and developing antennal lobe of the brain of *Drosophila melanogaster*.

The adult antennal lobe

Three striking features are characteristic of the adult antennal lobe in *Drosophila* (Fig 13): (1) glomeruli are surrounded by only a sparse network of glial processes, clearly distinct from the continuous multi-layered envelopes of the moth (Tolbert and Hildebrand, 1981) and the complex glial envelopes that also incorporate periglomerular cells in mice, rats, cats, and opossums (Pinching and Powell, 1971; Willey, 1973; Tolbert and Hildebrand, 1981; Raisman, 1985; Valverde and Lopez-Mascaraque, 1991; Valverde et al., 1992; Bailey and Shipley, 1993); (2) large dendrites and axons or axon bundles are almost always ensheathed to some extent, whether they are in the nerve layer, the body of a glomerulus, the primary neurite tracts, or the coarse central neuropil; and (3) glial processes are scattered throughout the synaptic neuropil of the glomeruli, unlike in the moth (Tolbert, 1989; Oland et al., 1999) and more like in the vertebrates (Chiu and Greer, 1996).

Across a wide variety of both vertebrate and invertebrate species, glomeruli are characteristic organizational features of the first-order olfactory neuropil. In numerous mammals, a marsupial, and the moth *Manduca sexta*, these glomeruli are surrounded and separated by more or less complete envelopes of glial cell bodies and processes (Pinching and Powell, 1971; Willey, 1973; Tolbert and Hildebrand, 1981; Raisman, 1985; Valverde and Lopez-Mascaraque, 1991; Oland et al. 1999). In contrast, in many insects, including honeybee (Gascuel and Masson, 1991; Hähnelin and Bicker, 1996; Schröter and Malun, 2000; Rössler et al., 2002), cockroach (Ernst et al., 1977), and *Drosophila melanogaster* (reported here), glial envelopment of glomeruli is present but far less robust. And in *Xenopus* (Byrd and Burd, 1991; Nezhlin et al., 2003), glial borders appear to be lacking altogether in the adult; instead, large dendrites of antennal lobe neurons cup and define the borders of the glomeruli. In bee, ant, and *Drosophila*, glial cell bodies surround the antennal lobe neuropil or large, multi-glomerular neuropil compartments within the antennal lobe (Schröter and Malun, 2000; Rössler et al., 2002), and in bee, glial cell bodies also can be found in the coarse central neuropil. In bee, glial processes do form a cap around the apical and lateral portions of each glomerulus (Hähnelin and Bicker, 1996; Brown, 2001), but whether the glia form a net of processes or a sheet-like envelope, was not addressed by the authors. The variability in completeness of the glial envelope is thus wide, and our current

results, indicating that in *Drosophila* each glomerulus has a surrounding net rather than a more complete envelope, place *Drosophila* at the sparse end of the spectrum in insects.

The absence of a continuous glial border around glomeruli in *Drosophila* and some other insects and the absence of any glial border in frogs raise the question, under what circumstances are complete or robust glial borders required? Is glomerular size an important determinant?

A review of the extant comparative data suggests that larger glomeruli tend to have more robust envelopes. Vertebrate and marsupial glomeruli range in size from 50 to 200 μm in diameter and are well surrounded by glia; amphibian glomeruli are smaller, ranging from 20 to 50 μm (Hildebrand and Shepherd, 1997), and lack glial borders. In *Manduca*, glomeruli range in size from 50 to 100 μm (Hildebrand and Montague, 1986), while those of *Heliothis*, another moth are 25–50 μm (Vickers et al., 2003); the glial borders of the former are robust, while those of the latter appear in sectioned material to be similar, but no detailed information is available. The glomeruli of bees range from 30 to 50 μm in diameter and have a partial glial surround (Hahnlein and Bicker, 1996) and those in *Drosophila* are about 20 μm in diameter, with a net-like glial surround. Roach glomeruli are about 40 μm in diameter and are bounded mainly by thick dendrites and perhaps some glial processes (Ernst and Boeckh, 1983). While the sizes of glomeruli in many other insects are known (e.g., Ignell et al., 2001; Rössler et al., 2002), their glial investment is as yet not characterized. In short, there may be a rough size correlation.

Is there a functional difference that might arise from the dramatically different glial surrounds of *Manduca* and *Drosophila* glomeruli? A single glomerulus in the *Manduca* antennal lobe receives, on average, input from approximately 5,000 receptor neurons, 5–6 uniglomerular projection neurons, and roughly 360 interneurons as well as some 500 multi-glomerular projection neurons contribute to the set of glomeruli in patterns that are not yet well understood (Homberg et al., 1989). Mathematical modeling of the multi-layered glial envelope in *Manduca* (Goriely et al., 2002) suggested that this glial envelope may provide a significant physical barrier that would block diffusion of potassium ions from an activated glomerulus to neighboring glomeruli on a time scale that would limit activity to the intra-glomerular circuits of the active glomerulus. In addition, the extracellular $[\text{K}^+]$ would increase with activity, possibly influencing neuronal activity within the active glomerulus for many seconds.

In contrast, a *Drosophila* glomerulus receives input from fewer than 50 receptor neurons, up to 3 projection neurons, and an as yet unknown number of local interneurons (Stocker et al., 1990). The porous, net-like structure of the glial surround for each glomerulus in *Drosophila* clearly cannot constitute a physically insulating barrier between glomeruli, yet individual glomeruli have been shown by imaging responses to different odorants to be functionally distinct (Wang et al., 2003), indicating that a physical barrier is not necessary to enable glomeruli to maintain their discrete odotopic representation. *Drosophila* must rely essentially entirely on neuronal circuitry to isolate and sharpen the glomerular response to particular odorants (e.g. Silbering and Galizia, 2007). Perhaps there is no need for a glial barrier to prevent effects on neighboring glomeruli when so few axons are simultaneously activated in a glomerulus, or perhaps, in systems like *Drosophila* in which some glomeruli are more broadly tuned (Fishilevich and Vosshall, 2005), a robust glial border is unneeded. Examination of glomerular responses to olfactory stimuli after selective removal of glial cells in a species with large glomeruli would shed light on whether this is indeed a likely explanation for the lack of glial envelopes for glomeruli in *Drosophila*.

The prevalence of glial processes *within* the synaptic neuropil of the glomeruli also is widely variable among species. In mammals, astrocytes with several distinct morphologies can be found within the glomerular neuropil (Pinching and Powell, 1971; Bailey and Shipley, 1993; Chiu and Greer, 1996), each associated with a single glomerulus. Astrocytic processes define subcompartments of the glomeruli and are unevenly distributed within these subcompartments. For example, the region in rat in which receptor axons synapse with terminal dendritic branches is mainly devoid of astrocytic processes while more central regions display typical tri-partite synaptic arrangements with subsets of central neurons (Chao et al., 1997). Nevertheless, infiltration of glial processes into glomerular neuropil is generally less than is found in other synaptic neuropil areas of the brain (Chao et al., 1997), and glial processes are not found in particularly close association with synaptic complexes as they are in other neuropils (Chao et al., 1997; Kasowski et al., 1999).

In the moth, as in the bee and *Drosophila*, glial cell bodies are never found within the synaptic glomerular neuropil (Oland et al., 1999; Hähnlein and Bicker, 1996). Processes of the 75–100 simple neuropil glia that form the multi-layered glial envelope for each glomerulus in *Manduca* extend no more than a few microns into the glomerular neuropil, and the processes of an additional 4–5 “complex” neuropil glial cells accompany fascicles of olfactory receptor axons into the apical portion of the glomeruli, a few of their processes extending deeply into the body of the glomerulus and branching (Fig. 14A; Oland et al., 1999). The paucity of glial processes in the central third of the glomeruli, the glomerular region most dense in synapses, suggests, however, that glial cells in this system are not modulating synaptic function locally. Glial processes also extensively ensheath large dendrites entering the base of the glomeruli from the central neuropil of the lobe, presumably serving to insulate the processes electrically in a manner analogous to astrocytic enwrapping of dendrites and neuronal cell bodies in the vertebrate nervous system (Halassa et al., 2007).

Drosophila glomerular neuropil, in contrast, is infiltrated by many glial processes (Fig. 14B). The majority of these processes are directed toward ensheathment of large dendritic processes, but many glial processes also generally invest the glomerular neuropil. They do not display any preferential close association with synapses. Perhaps they sit in position not to regulate synapses directly but to regulate the general neuronal microenvironment during periods of neuronal activity in the manner of astrocytes and insect glial cells associated with other neuropils (Kuffler et al. 1966; Coles, 1989; Kofuji and Newman, 2004; Tsacopoulos et al., 1988; Tsacopoulos et al., 1994; Tsacopoulos and Magistretti, 1996; Shepherd and Tyrer, 1985; Seal et al., 1998; Borycz et al., 2002; Soustelle et al., 2002; Kofuji and Newman, 2004; Allen and Barres, 2005; Lohr & Deitmer, 1999; Freeman et al., 2003; Fellin and Carnignoto, 2004; Heil et al., 2007).

It should be noted that the results of the current study in *Drosophila* show that neuropil glial cells in the antennal lobe are not simply ensheathing axons. In mammals, both oligodendrocytes and astrocytes ensheath dendrites (Peters et al., 1991) and also contribute to the glial envelope around glomeruli (Valverde and Lopez-Mascaraque, 1991), but astrocytes also invade synaptic glomerular neuropil, a function that falls outside of the oligodendrocytes' repertoire. Neuropil glial cells in *Drosophila* not only ensheath axons and dendrites and form the glial net around glomeruli, they also invade glomerular neuropil, suggesting that neuropil glial cells should be not be considered as functionally homologous only to oligodendrocytes (Edenfield et al., 2005; Pereanu et al., 2005; Freeman and Doherty, 2006), but rather as sharing some of the functional capacities of both astrocytes and oligodendrocytes.

The developing antennal lobe

In moth, rodent, and opossum, neuron-glia interactions have been implicated in several key steps in development of the olfactory pathway. In the moth, a complex picture has evolved in which neuron-glia interactions and certain cell signaling molecules appear to be critical both for olfactory receptor axon sorting and targeting and for stabilization of newly forming glomeruli (Tolbert et al., 2004; Oland and Tolbert 2003; Gibson et al., 2001; Kaneko and Nighorn, 2003; Gibson and Tolbert, 2006). First, ingrowing axons encounter a glial cell-rich sorting zone region at the junction of the antennal nerve and antennal lobe and, as they course through it, dramatically reroute and sort into fascicles destined to terminate in specific glomeruli or groups of glomeruli (Oland et al., 1998), with the sorted axons taking a rather direct route to their final targets. In developing moths deficient in glia, receptor axons fail to fasciculate and properly target glomeruli, with many axons completely bypassing the antennal lobe (Rössler et al., 1999). In the mouse, sorting occurs in the olfactory nerve layer of the bulb, which can be subdivided into outer and inner nerve layers. Axonal morphology and fasciculation differ in the two layers; the axons travel in bundles around the lobe while in the outer layer, but once they enter the inner layer, which has a characteristic glial population different from that in the outer layer, they become less tightly bundled and sort into fascicles targeting specific glomeruli (Au et al., 2002; Treloar et al., 2002). Thus in this system, too, axonal interaction with a particular set of glial cells sets the stage for final sorting and targeting.

Our observations of the *Drosophila* antennal nerve at both the confocal and EM levels indicate that there is no discrete sorting region in the *Drosophila* nerve. This is consistent with the findings of Rodrigues and her colleagues (Bhalerao et al., 2003; Sen et al., 2005), who showed using OR22a and OR47b lines that the axons of each class neither travel together in the nerve nor approach their glomeruli together, although axons belonging to individual olfactory receptor neuron classes do course through the nerve layer in stereotypic patterns to reach their target glomerulus (Ang et al., 2003). We found no evidence in our study of the development of the nerve layer at the confocal level or in the electron microscope for extensive sorting within the nerve layer in the antennal lobe.

Taken together, the data suggest that axonal sorting in *Drosophila* depends mainly on: 1) rough segregation of OR classes on the antenna (DeBruyne et al., 2001; Couto et al., 2005); 2) on cues expressed by the axons (Hummel and Zipursky, 2004), some of which may be lineage-specific (Endo et al., 2007), that allow some sorting along the nerve; and 3) on cues intrinsic to the antennal lobe (Ang et al., 2003; Komiyama et al., 2007) that guide axons to their specific targets within the neuropil. Glial cells in *Drosophila* thus appear to have no significant role in the sorting of olfactory receptor axons at the base of the antennal nerve or in the nerve layer.

While the mechanisms of sorting and targeting of receptor axons are only partially understood in any olfactory pathway, the subsequent establishment and stabilization of glomeruli involves glial cells in several well-studied systems. During normal development in *Manduca* (Tolbert et al., 2004), receptor axons establish axonal protoglomeruli that evolve into the mature glomerular architecture seen in the adult antennal lobe. Importantly, glia activated by ingrowing receptor axons surround protoglomeruli and are required to stabilize them during the subsequent ingrowth of, first, projection neuron dendrites and later, local interneuron dendrites (Fig. 15A). By the end of normal development, each glomerulus in *Manduca* is nearly enveloped by a multilayered glial sheath (Figs. 14A, 15A). If development proceeds with an insufficient number of stabilizing glia, however, the axonal protoglomerular template quickly dissolves (Baumann et al., 1996), resulting in an aglomerular neuropil in the adult.

A similar sequence of glial changes occurs in the developing vertebrate olfactory pathways although there the requirement for glial cells in glomerulus construction has not been tested directly. As receptor axons grow into the olfactory bulb, they encounter a plexus of glial processes at the level of the presumptive glomerular layer that comprise the terminal branches of radial glia (Bailey et al., 1999). The processes of these glial cells branch to form “glial glomeruli” and as they do so, the axons organize their terminal branches within the territories of the glial glomeruli to form protoglomeruli. Astrocytic processes, along with a few oligodendrocytic processes, extend to surround the newly forming glomeruli; the olfactory ensheathing cells that accompany the ingrowing axons may interleave with the astrocytes that form the glial borders around glomeruli but do not extend processes into the glomerular neuropil (Valverde and Lopez-Mascaraque, 1991; Treloar et al., 1999).

In *Drosophila*, the first ORN axons to arrive at the antennal lobe (at ~18% of development) are associated with a small group of glial cells on the periphery of the lobe and remain associated with these glia during a short waiting period prior to their entry into the antennal lobe neuropil (Jhaveri et al., 2000). Projection neurons already have targeted specific regions of the developing antennal lobe neuropil prior to arrival of the receptor axons (Jefferis et al., 2004) and the projection neuron map comes into rough register with the later developing receptor neuron protoglomerular array (Hummel and Zipursky, 2004), by mechanisms not yet well understood, to begin the process of glomerulus development. By a variety of measures (Jhaveri et al., 2000; Jhaveri et al., 2002; Hummel and Zipursky, 2004; Jefferis et al., 2004), glomerular architecture is well established by 50% of development (Fig 15B). While we have found that the glial investment of the neuropil begins with the formation of a fringe of glial processes at 30–40% of development, somewhat earlier than previously reported, the growth of glial processes into the lobe is minimal as late as 50% of development, confirming previous studies (Jhaveri et al., 2000; Jefferis et al., 2004), and even then is largely restricted at first to the region of the primary neurite tracts with little evidence of any rapid extension to surround developing glomeruli.

Through the remainder of metamorphic development, the number of glial processes extending into the neuropil gradually increases, as does the degree to which they branch, the end result being the net of glial processes around each glomerulus (Fig. 14B). The pattern of branching is regulated in some way by interaction with sensory neurons as neuropil glia branch aberrantly in atonal mutants, which lack a subset of olfactory receptor axons (Jhaveri and Rodrigues, 2002). The timecourse over which the glial investment of the *Drosophila* antennal lobe is established relative to that of the other cellular elements contributing to glomeruli differs markedly from that seen in *Manduca* (Figs. 15–16; Tolbert et al., 2004) and several vertebrates (Bailey et al., 1999; Valverde and Lopez-Mascaraque, 1991; Treloar et al., 1999). It precludes the possibility that neuropil glia are involved directly in the stabilization of glomeruli, though leaves open several other roles in the maturation of glomeruli, including synapse maturation and dendritic pruning, as occurs in other neuropils such as the mushroom bodies (Marin et al., 2005). Also, during the early phases of ORN axon ingrowth, neuropil glial cells may interact with ingrowing axons in the nerve layer or in the midline-located transient interhemispheric fibrous ring (TIRF) in ways that indirectly affect the organization of the glomerular array. Indeed, a recent study has implicated Derailed-expressing glia, especially those in the TIRF, in modulating axonal Wnt5 signaling; lack of either results in midline crossing and targeting defects in ORN axons (Yao, 2007).

Figure 16 provides a chart of the timing and sequence of known neuronal and glial events in the development of *Drosophila melanogaster* antennal lobes.

Summary

The glomeruli of the *Drosophila* antennal lobe share with most other species, vertebrate and invertebrate alike, the presence of glial processes around each glomerulus. This glial structure is minimal in *Drosophila*, appearing in the form of an overlying net of processes rather than more extensive arrangement of processes that together form a continuous envelope. Glial processes extend into the glomerular neuropil, but do not form morphological associations with synapses. In the developing olfactory pathway, glial cells do not form a discrete axonal sorting zone either in the antennal nerve or in the nerve layer, and they extend processes around glomerular neuropil too late to be involved in the process of glomerulus stabilization, though they extend into the neuropil during the period of synaptogenesis and may contribute to that process. These findings suggest that in small olfactory neuropils, with relatively few neuronal components, glial cells are unlikely to be necessary for sorting of receptor axons, for development of glomeruli (although they may affect axon targeting), or for isolation of odor activity within discrete glomeruli.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge the contributions of: Patricia Jansma, M.S., who manages the Arizona Research Laboratories Division of Neurobiology (ARLDN) Imaging Facility and provided expert technical help with the confocal and electron microscopy; Marvin Landis, from the University of Arizona Campus Computing and Information Technology Department, who generated the 3D movies; Jennifer Crane, now at Stower's Institute for Medical Research, Kansas City MO, who generated much of Figure 15 when she was a graduate student at the University of Arizona; Andrea Wellington from the laboratory of our ARLDN colleague Konrad Zinsmaier, who provided advice on *Drosophila* rearing; Jane Lim, an undergraduate student in our group who provided the images of the antennal nerve labeled with antibodies to neuroglian (Fig. 12); and members of our laboratory group with whom we had many constructive discussions about this work. Funding for J Biebelhausen was provided by HHMI funding (52003749) via the University of Arizona's Undergraduate Biology Research Program overseen by Carol Bender, M.S. Funding for L Oland and L Tolbert was provided by NIH R01 DC004598 to L Tolbert and by NIH P01 NS28495 to L Oland (Project 2).

The primary antibodies used in this study were obtained from the Developmental Studies Hybridoma Bank developed under the auspices of the NICHD and maintained by the University of Iowa.

References

- Alfonso TB, Jones BW. Gcm2 promotes glial cell differentiation and is required with glial cells missing for macrophage development in *Drosophila*. *Dev Biol.* 2002; 248:369–83. [PubMed: 12167411]
- Allen NJ, Barres BA. Signaling between glia and neurons: focus on synaptic plasticity. *Curr Opin Neurobiol.* 2005; 15:542–48. [PubMed: 16144764]
- Ang LH, Kim J, Stepensky V, Hing H. Dock and pak regulate olfactory axon pathfinding in *Drosophila*. *Development.* 2003; 130:1307–16. [PubMed: 12588847]
- Au WW, Treloar HB, Greer CA. Sublaminar organization of the mouse olfactory bulb nerve layer. *J Comp Neurol.* 2002; 446:68–80. [PubMed: 11920721]
- Bailey MS, Puche AC, Shipley MT. Development of the olfactory bulb: evidence for glia-neuron interactions in glomerular formation. *J Comp Neurol.* 1999; 415:423–48. [PubMed: 10570454]
- Bailey MS, Shipley MT. Astrocyte subtypes in the rat olfactory bulb: morphological heterogeneity and differential laminar distribution. *J Comp Neurol.* 1993; 328:501–26. [PubMed: 8429132]
- Baumann PM, Oland LA, Tolbert LP. Glial cells stabilize axonal protoglomeruli in the developing olfactory lobe of the moth *Manduca sexta*. *J Comp Neurol.* 1996; 373:118–28. [PubMed: 8876467]

- Bhalerao S, Sen A, Stocker R, Rodrigues V. Olfactory neurons expressing identified receptor genes project to subsets of glomeruli within the antennal lobe of *Drosophila melanogaster*. *J Neurobio*. 2003; 54:577–92.
- Bieber AJ, Snow PM, Hortsch M, Patel NH, Jacobs JR, Traquina ZR, Schilling J, Goodman CS. *Drosophila* neuroglian: A member of the immunoglobulin superfamily with extensive homology to the vertebrate neural adhesion molecule L1. *Cell*. 1989; 59:447–60. [PubMed: 2805067]
- Borycz J, Borycz JA, Loubani M, Meinertzhagen IA. tan and ebony genes regulate a novel pathway for transmitter metabolism at fly photoreceptor terminals. *J Neurosci*. 2002; 22:10549–57. [PubMed: 12486147]
- Brown, S. Ultrastructural analysis of activity-dependent volume changes in the antennal lobe of the worker honey bee *Apis Mellifera*. Dissertation. University of Otago; Dunedin, New Zealand: 2001.
- de Bruyne M, Foster K, Carlson JR. Odor coding in the *Drosophila* antenna. *Neuron*. 2001; 30:537–52. [PubMed: 11395013]
- Byrd CA, Burd GD. Development of the olfactory bulb in the clawed frog, *Xenopus laevis*: a morphological and quantitative analysis. *J Comp Neurol*. 1991; 314:79–90. [PubMed: 1797876]
- Campbell G, Göring H, Lin T, Spana E, Andersson S, Doe C, Tomlinson A. RK2, a glial-specific homeodomain protein required for embryonic nerve cord condensation and viability in *Drosophila*. *Development*. 1994; 120:2957–66. [PubMed: 7607085]
- Chao TI, Kasa P, Wolff JR. Distribution of astroglia in glomeruli of the rat main olfactory bulb: Exclusion from the sensory subcompartment of neuropil. *J Comp Neurol*. 1997; 388:191–210. [PubMed: 9368837]
- Chiu K, Greer CA. Immunocytochemical analyses of astrocyte development in the olfactory bulb. *Brain Res Dev Brain Res*. 1996; 95:28–37.
- Chotard C, Salecker I. Neurons and glia: team players in axon guidance. *Trends Neurosci*. 2004; 27:655–61. [PubMed: 15474165]
- Clyne PJ, Warr CG, Freeman MR, Lessing D, Kim J, Carlson JR. A novel family of divergent seven-transmembrane proteins: candidate odorant receptors in *Drosophila*. *Neuron*. 1999; 22:327–38. [PubMed: 10069338]
- Coles JA. Functions of glial cells in the retina of the honeybee drone. *Glia*. 1989; 2:1–9. [PubMed: 2523335]
- Couto A, Alenius M, Dickson BJ. Molecular, anatomical, functional organization of the *Drosophila* olfactory system. *Curr Biol*. 2005; 15:1535–47. [PubMed: 16139208]
- Devaud J-M, Acebes A, Ramaswami M, Ferrus A. Structural and functional changes in the olfactory pathway of adult *Drosophila* take place at a critical age. *J Neurobiol*. 2003; 56:13–23. [PubMed: 12767029]
- Dubin AE, Harris GL. Voltage-activated and odor-modulated conductances in olfactory neurons of *Drosophila melanogaster*. *J Neurobiol*. 1997; 32:123–37. [PubMed: 8989668]
- Edenfield G, Stork T, Klämbt C. Neuron-glia interaction in the insect nervous system. *Curr Opin Neurobio*. 2005; 15:34–39.
- Edwards, JS.; Tolbert, LP. Insect neuroglia. In: Harrison, FW.; Locke, M., editors. *Microscopic anatomy of invertebrates*, Vol 11B: Insecta. New York: Wiley-Liss; 1998. p. 449-66.
- Endo K, Aoki T, Yoda Y, Kimura K, Hama C. Notch signal organizes the *Drosophila* olfactory circuitry by diversifying the sensory neuronal lineages. *Nat Neurosci*. 2007; 10:153–59. [PubMed: 17220884]
- Ernst KD, Boeckh J, Boeckh V. A neuroanatomical study on the organization of the central antennal pathways in insects. *Cell Tissue Res*. 1977; 176:285–306. [PubMed: 832298]
- Ernst KD, Boeckh J. A neuroanatomical study on the organization of the central antennal pathways in insects. III. Neuroanatomical characterization of physiologically defined response types of deutocerebral neurons in *Periplaneta americana*. *Cell Tissue Res*. 1983; 229:1–22. [PubMed: 6831538]
- Fellin T, Carnignoto G. Neurone-to-astrocyte signalling in the brain represents a distinct multifunction unit. *J Physiol*. 2004; 559:3–15. [PubMed: 15218071]
- Fishilevich E, Vosshall LB. Genetic and functional subdivision of the *Drosophila* antennal lobe. *Curr Biol*. 2005; 15:1548–53. [PubMed: 16139209]

- Freeman MR. Sculpting the nervous system: glial control of neuronal development. *Curr Opin Neurobiol.* 2006; 16:119–25. [PubMed: 16387489]
- Freeman MR, Delrow J, Kim J, Johnson E, Doe CQ. Unwrapping glial biology: Gcm target genes regulating glial development, diversification, function. *Neuron.* 2003; 38:567–80. [PubMed: 12765609]
- Freeman MR, Doherty J. Glial cell biology in *Drosophila* and vertebrates. *Trends Neurosci.* 2006; 29:82–90. [PubMed: 16377000]
- Gascuel J, Masson C. Developmental study of afferented and deafferented bee antennal lobes. *J Neurobiol.* 1991; 22:795–810. [PubMed: 1779223]
- Gibson NJ, Tolbert LP. Activation of epidermal growth factor receptor mediates receptor axon sorting and extension in the developing olfactory system of the moth *Manduca sexta*. *J Comp Neurol.* 2006; 495:554–52. [PubMed: 16498681]
- Godenschwege TA, Reisch D, Diegelmann S, Elberle K, Funk N, Heisenberg M, Hoppe V, Hoppe J, Klagges BRE, Martin J-R, Nikitina EA, Putz G, Reifegerste R, Reisch N, Rister J, Schaupp M, Scholz H, Schwaerzel M, Wernet U, Zars TD, Buchner S, Buchner E. Flies lacking all synapsins are unexpectedly healthy but are impaired in complex behaviour. *Eur J Neurosci.* 2004; 20:611–22. [PubMed: 15255973]
- Goriely AR, Secomb TW, Tolbert LP. Effect of the glial envelope on extracellular K(+) diffusion in olfactory glomeruli. *J Neurophysiol.* 2002; 87:1712–22. [PubMed: 11929893]
- Hähnlein I, Bicker G. Morphology of neuroglia in the antennal lobes and mushroom bodies of the brain of the honeybee. *J Comp Neurol.* 1996; 367:235–45. [PubMed: 8708007]
- Halassa MM, Fellin T, Haydon PG. The tripartite synapse: roles for gliotransmission in health and disease. *Trends in Mol Med.* 2007; 13:54–63. [PubMed: 17207662]
- Hall SG, Bieber AJ. Mutations in the *Drosophila* neuroglial cell adhesion molecule affect motor neuron pathfinding and peripheral nervous system patterning. *J Neurobiol.* 1997; 32:325–40. [PubMed: 9058324]
- Halter DA, Urban J, Rickert C, Ner SS, Ito K, Travers AA, Technau GM. The homeobox gene *repo* is required for the differentiation and maintenance of glia function in the embryonic nervous system of *Drosophila melanogaster*. *Development.* 1995; 121:317–32. [PubMed: 7768175]
- Heil JE, Oland LA, Lohr C. Acetylcholine-mediated axon-glia signaling in the developing insect olfactory system. *Eur J Neurosci.* 2007; 26:1227–41. [PubMed: 17767501]
- Hildebrand, JG.; Montague, RA. Functional organization of olfactory pathways in the central nervous system of *Manduca sexta*. In: Payne, TL.; Birch, MC.; Kennedy, CEJ., editors. *Mechanisms in insect olfaction*. Oxford University Press; Oxford: 1986. p. 279-85.
- Hildebrand JG, Shepherd GM. Mechanisms of olfactory discrimination: converging evidence for common principles across phyla. *Annu Rev Neurosci.* 1997; 20:595–631. [PubMed: 9056726]
- Homberg U, Christensen TA, Hildebrand JG. Structure and Function of the deutocerebrum in insects. *Ann Rev Ent.* 1989; 34:477–501.
- Hortsch M, Bieber AJ, Patel NH, Goodman CS. Differential splicing generates a nervous system-specific form of *Drosophila* neuroglial. *Neuron.* 1990; 4:697–709. [PubMed: 1693086]
- Hummel T, Vasconcelos ML, Clemens JC, Fishilevich Y, Vosshall LB, Zipursky SL. Axonal targeting of olfactory receptor neurons in *Drosophila* is controlled by Dscam. *Neuron.* 2003; 37:221–31. [PubMed: 12546818]
- Hummel T, Zipursky SL. Afferent induction of olfactory glomeruli requires N-Cadherin. *Neuron.* 2004; 42:77–88. [PubMed: 15066266]
- Ignell R, Anton S, Hansson BS. The antennal lobe of orthoptera – anatomy and evolution. *Brain Behav Evol.* 2001; 57:1–17. [PubMed: 11359044]
- Ito K, Urban J, Technau GM. Distribution, classification and development of *Drosophila* glial cells in the late embryonic and early larval ventral nerve cord. *Roux's Arch Dev Biol.* 1995; 204:284–307.
- Jefferis GS, Marin EC, Stocker RF, Luo L. Target neuron prespecification in the olfactory map of *Drosophila*. *Nature.* 2001; 414:204–08. [PubMed: 11719930]
- Jefferis GS, Vyas RM, Berdnik D, Ramaekers A, Stocker RF, Tanaka NK, Ito K, Luo L. Developmental origin of wiring specificity in the olfactory system of *Drosophila*. *Development.* 2004; 131:117–30. [PubMed: 14645123]

- Jhaveri D, Rodrigues V. Sensory neurons of the Atonal lineage pioneer the formation of glomeruli within the adult *Drosophila* olfactory lobe. *Development*. 2002; 129:1251–60. [PubMed: 11874920]
- Jhaveri D, Sen A, Rodrigues V. Mechanisms underlying olfactory neuronal connectivity in *Drosophila* - the atonal lineage organizes the periphery while sensory neurons and glia pattern the olfactory lobe. *Dev Biol*. 2000; 226:73–87. [PubMed: 10993675]
- Kaneko M, Nighorn A. Interaxonal Eph-ephrin signaling may mediate sorting of olfactory sensory axons in *Manduca sexta*. *J Neurosci*. 2003; 23:11523–38. [PubMed: 14684856]
- Kasowski HJ, Kim H, Greer CA. Compartmental organization of the olfactory bulb glomerulus. *J Comp Neurol*. 1999; 407:261–74. [PubMed: 10213094]
- Klagges BR, Heimbeck G, Godenschwege TA, Hofbauer A, Pfulgfelder GO, Reifegerste R, Reisch D, Schaupp M, Buchner S, Buchner E. Invertebrate synapsins: a single gene codes for several isoforms in *Drosophila*. *J Neurosci*. 1996; 16:3154–65. [PubMed: 8627354]
- Kofuji P, Newman EA. Potassium buffering in the central nervous system. *Neuroscience*. 2004; 129:1045–56. [PubMed: 15561419]
- Komiyama T, Sweeney LB, Schuldiner O, Garcia KC, Luo L. Graded expression of semaphorin-1a cell-autonomously directs dendritic targeting of olfactory projection neurons. *Cell*. 2007; 128:399–410. [PubMed: 17254975]
- Kuffler SW, Nicholls JG, Orkand RK. Physiological properties of glial cells in the central nervous system of amphibia. *J Neurophys*. 1966; 29:768–87.
- Laissue PP, Reiter C, Hiesinger PR, Halter S, Fischbach KF, Stocker RF. Three-dimensional reconstruction of the antennal lobe in *Drosophila melanogaster*. *J Comp Neurol*. 1999; 405:543–552. [PubMed: 10098944]
- Lienhard MC, Stocker RF. The development of the sensory neuron pattern in the antennal disc of wild-type and mutant (*Iz3, ss**) *Drosophila melanogaster*. *Dev*. 1991; 112:1063–75.
- Lohr C, Deitmer JW. Dendritic calcium transients in the leech giant glial cell in situ. *Glia*. 1999; 26:109–18. [PubMed: 10384876]
- MacDonald JM, Beach MG, Porpiglia E, Sheehan AE, Watts RJ, Freeman MR. The *Drosophila* cell corpse engulfment receptor Draper mediates glial clearance of severed axons. *Neuron*. 2006; 50:869–81. [PubMed: 16772169]
- Marin EC, Jefferis GS, Komiyama T, Zhu H, Luo L. Representation of the glomerular olfactory map in the *Drosophila* brain. *Cell*. 2002; 109:243–55. [PubMed: 12007410]
- Marin EC, Watts RJ, Tanaka NK, Ito K, Luo L. Developmentally programmed remodeling of the *Drosophila* olfactory circuit. *Dev*. 2005; 132:725–37.
- Meinertzhagen IA, O'Neil SD. Synaptic organization of columnar elements in the lamina of the wild type in *Drosophila melanogaster*. *J Comp Neurol*. 1991; 305:232–63. [PubMed: 1902848]
- Nezlin LP, Heermann S, Schild D, Rossler W. Organization of glomeruli in the main olfactory bulb of *Xenopus laevis* tadpoles. *J Comp Neurol*. 2003; 464:257–68. [PubMed: 12900923]
- Oland LA, Evans S. The tracheal system of the developing primary olfactory pathway of *Manduca sexta*: trachea do not play a guidance or targeting role for ingrowing receptor axons. *Arthropod Struct Dev*. 2000; 29:185–96. [PubMed: 18088926]
- Oland LA, Tolbert LP. Key interactions between neurons and glial cells during neural development in insects. *Annu Rev Entomol*. 2003; 48:89–110. [PubMed: 12194908]
- Oland LA, Biebelhausen JP, Tolbert LP. Development of the glial investment of glomeruli in the *Drosophila* olfactory lobe. *Chemical Senses*. 2007 in press.
- Oland LA, Marrero HG, Burger I. Glial cells in the developing and adult olfactory lobe of the moth *Manduca sexta*. *Cell Tissue Res*. 1999; 297:527–45. [PubMed: 10460499]
- Oland LA, Pott WM, Higgins MR, Tolbert LP. Targeted ingrowth and glial relationships of olfactory receptor axons in the primary olfactory pathway of an insect. *J Comp Neurol*. 1998; 398:119–38. [PubMed: 9703031]
- Parker RJ, Auld VJ. Roles of glia in the *Drosophila* nervous system. *Semin Cell Dev Biol*. 2006; 17:66–77. [PubMed: 16420983]

- Pereanu W, Shy D, Hartenstein V. Morphogenesis and proliferation of the larval brain glia in *Drosophila*. *Dev Biol*. 2005; 283:191–203. [PubMed: 15907832]
- Pereanu W, Spindler S, Cruz L, Hartenstein V. Tracheal development in the *Drosophila* brain is constrained by glial cells. *Dev Biol*. 2007; 302:169–80. [PubMed: 17046740]
- Peters, A.; Palay, SL.; Webster, H deF. The fine structure of the nervous system. New York: Oxford University Press; 1991.
- Pinching AJ, Powell TP. The neuropil of the glomeruli of the olfactory bulb. *J Cell Sci*. 1971; 9:347–77. [PubMed: 4108057]
- Raisman G. Specialized neuroglial arrangement may explain the capacity of vomeronasal axons to reinnervate central neurons. *Neuroscience*. 1985; 14:237–54. [PubMed: 3974880]
- Ray K, Rodrigues V. Cellular events during development of the olfactory sense organs in *Drosophila melanogaster*. *Dev Biol*. 1995; 167:426–38. [PubMed: 7875369]
- Read RD, Goodfellow PJ, Mardis ER, Novak N, Armstrong JR, Cagan RL. A *Drosophila* model of multiple endocrine neoplasia type 2. *Genetics*. 2005; 171:1057–81. [PubMed: 15965261]
- Rössler W, Kuduz J, Schürmann FW, Schild D. Aggregation of f-actin in olfactory glomeruli: a common feature of glomeruli across phyla. *Chem Senses*. 2002; 27:803–10. [PubMed: 12438205]
- Rössler W, Oland LA, Higgins MR, Hildebrand JG, Tolbert LP. Development of a glia-rich axon-sorting zone in the olfactory pathway of the moth *Manduca sexta*. *J Neurosci*. 1999; 19:9865–77. [PubMed: 10559396]
- Roy B, Singh AP, Shetty C, Chaudhary V, North A, Landgraf M, VijayRaghavan K, Rodrigues V. Metamorphosis of an identified serotonergic neuron in the *Drosophila* olfactory system. *Neural Dev*. 2007; 2:20. [PubMed: 17958902]
- Schröter U, Malun D. Formation of antennal lobe and mushroom body neuropils during metamorphosis in the honeybee, *Apis mellifera*. *J Comp Neurol*. 2000; 422:229–245. [PubMed: 10842229]
- Seal RP, Daniels GM, Wolfgang WJ, Forte MA, Amara SG. Identification and characterization of a cDNA encoding a neuronal glutamate transporter from *Drosophila melanogaster*. *Receptors Channels*. 1998; 6:51–64. [PubMed: 9664622]
- Sen A, Shetty C, Jhaveri D, Rodrigues V. Distinct types of glial cells populate the *Drosophila* antenna. *BMC Dev Biol*. 2005; 5:25. [PubMed: 16281986]
- Shepherd D, Tyrer NM. Inhibition of GABA uptake potentiates the effects of exogenous GABA on locust skeletal muscle. *Comp Biochem Physiol C*. 1985; 82:315–21. [PubMed: 2866905]
- Silbering AF, Galizia CG. Processing of odor mixtures in the *Drosophila* antennal lobe reveals both global inhibition and glomerulus-specific interactions. *J Neurosci*. 2007; 27:11966–77. [PubMed: 17978037]
- Soustelle L, Besson MT, Rival T, Birman S. Terminal glial differentiation involves regulated expression of the excitatory amino acid transporters in the *Drosophila* embryonic CNS. *Dev Biol*. 2002; 248:294–306. [PubMed: 12167405]
- Stocker RF. Fine structural comparison of the antennal nerve in the homeotic mutant *Antennapedia* with the wild-type antennal and second leg nerves of *Drosophila melanogaster*. *J Morphol*. 1979; 160:209–22. [PubMed: 110943]
- Stocker RF, Lienhard MC, Borst A, Fischbach KF. Neuronal architecture of the antennal lobe in *Drosophila melanogaster*. *Cell Tissue Res*. 1990; 262:9–34. [PubMed: 2124174]
- Stocker RF, Tissot M, Gendre N. Morphogenesis and cellular proliferation pattern in the developing antennal lobe of *Drosophila melanogaster*. *Roux's Arch Dev Biol*. 1995; 205:62–72.
- Strausfeld, NJ. Atlas of an Insect Brain. New York: Springer-Verlag; 1976.
- Sun B, Xu P, Salvaterra PM. Dynamic visualization of nervous system in live *Drosophila*. *Proc Natl Acad Sci U S A*. 1999; 96:10438–43. [PubMed: 10468627]
- Sun B, Xu P, Wang W, Salvaterra PM. In vivo modification of Na(+), K(+)-ATPase activity in *Drosophila*. *Comp Biochem Physiol B Biochem Mol Biol*. 2001; 130:521–36. [PubMed: 11691629]

- Tissot M, Gendre N, Hawken A, Störtkuhl KF, Stocker RF. Larval chemosensory projections and invasion of adult afferents in the antennal lobe of *Drosophila*. *J Neurobiol*. 1997; 32:281–97. [PubMed: 9058321]
- Tolbert LP. Afferent axons from the antenna influence the number and placement of intrinsic synapses in the antennal lobes of *Manduca sexta*. *Synapse*. 1989; 3:83–95. [PubMed: 2919370]
- Tolbert LP, Hildebrand JG. Organization and synaptic ultrastructure of glomeruli in the antennal lobes of the moth *Manduca sexta*: A study using thin sections and freeze-fracture. *Proc Roy Soc Lond B*. 1981; 213:279–301.
- Tolbert LP, Oland LA, Tucker ES, Gibson NJ, Higgins MR, Lipscomb BW. Bidirectional influences between neurons and glial cells in the developing olfactory system. *Prog Neurobiol*. 2004; 73:73–105. [PubMed: 15201035]
- Treloar HB, Purcell AL, Greer CA. Glomerular formation in the developing rat olfactory bulb. *J Comp Neurol*. 1999; 413:289–304. [PubMed: 10524340]
- Treloar HB, Feinstein P, Mombaerts P, Greer CA. Specificity of glomerular targeting by olfactory sensory axons. *J Neurosci*. 2002; 22:2469–77. [PubMed: 11923411]
- Tsacopoulos M, Evequoz-Mercier V, Perrottet P, Buchner E. Honeybee retinal glial cells transform glucose and supply the neurons with metabolic substrate. *PNAS*. 1988; 85:8727–31. [PubMed: 3186756]
- Tsacopoulos M, Magistretti PJ. Metabolic coupling between glia and neurons. *J Neurosci*. 1996; 16:877–85. [PubMed: 8558256]
- Tsacopoulos M, Veuthey AL, Saravelos SG, Perrottet P, Tsoupras G. Glial cells transform glucose to alanine, which fuels the neurons in the honeybee retina. *J Neurosci*. 1994; 14:1339–51. [PubMed: 8120629]
- Valverde F. Building an olfactory glomerulus. *J Comp Neurol*. 1999; 415:419–22. [PubMed: 10570453]
- Valverde F, Lopez-Mascaraque L. Neuroglial arrangements in the olfactory glomeruli of the hedgehog. *J Comp Neurol*. 1991; 307:658–74. [PubMed: 1714466]
- Valverde F, Santacana M, Heredia M. Formation of an olfactory glomerulus: morphological aspects of development and organization. *Neurosci*. 1992; 49:255–75.
- Vickers NJ, Poole K, Linn CE Jr. Consequences of interspecies antennal imaginal disc transplantation on organization of olfactory glomeruli and pheromone blend discrimination. *J Comp Neurol*. 2003; 466:377–88. [PubMed: 14556295]
- Wang JW, Wong AM, Flores J, Vosshall LB, Axel R. Two-photon calcium imaging reveals an odor-evoked map of activity in the fly brain. *Cell*. 2003; 112:271–82. [PubMed: 12553914]
- Wiley TJ. The ultrastructure of the cat olfactory bulb. *J Comp Neurol*. 1973; 152:211–32. [PubMed: 4775753]
- Xiong W-C, Okano H, Patel NH, Blendy JA, Montell C. *repo* encodes a glial-specific homeo-domain protein required in the *Drosophila* nervous system. *Genes and Development*. 1994; 8:981–994. [PubMed: 7926782]
- Yao Y, Wu Y, Yin C, Ozawa R, Aigaki T, Wouda RR, Noordermeer JN, Fradkin LG, Hing H. Antagonistic roles of Wnt5 and the Drl receptor in patterning the *Drosophila* antennal lobe. *Nat Neurosci*. 2007; 10:1423–32. [PubMed: 17934456]

Fig. 1.

Nuclei of Repo-positive (magenta) glial cells co-localize with GFP-labeled (green) cell bodies of glial cells in adult *Nrv2* antennal lobes. **A.** Nearly all of the neuropil glial cell nuclei are white, indicating that most of the glial processes present in the neuropil of *Nrv2* lobes are visible. Glial processes form a lacey network around glomeruli (*/**) and are especially prominent in the coarse central neuropil (c). Note difference in density of glial processes in * vs ** glomeruli. The large Repo-only cell bodies (upper left) belong to glial cells positioned within the neuronal cell body cluster; the medially located Repo-positive cell body (arrowhead) lies just under the perineurial sheath. **B.** A small group of glial cells (arrow) mark the entry of the primary neurite tract extending from the lateral cluster of neuron cell bodies. Stack z dimensions: A = 10 μm , B = 4 μm .

Fig. 2.

Glial processes in the primary olfactory pathway. **A–E.** Adult antennal lobe. **A,B.** Antibody to Synapsin I (magenta) labels glomerular neuropil in an anterior plane (**A**) and a more posterior plane (**B**) in a *Nrv2* GFP-expressing (green) fly. Fine glial processes extend between glomeruli (arrows). Arrowhead: One of the cortical glial cells contributing to glial network around neuronal cell bodies (ncb). an: antennal nerve. **C–E.** Higher power images of the glial investment of the adult glomeruli (*) and the coarse central neuropil (c). Neuropil glial cells contribute to the ensheathment of axon bundles (arrows in C,D) in the nerve layer and dendritic processes (arrowheads) in the neuropil. Two glomeruli are indicated by fine dotted lines. Note the variability in density of glial investment of these as well as the wide variation in glial process dimensions. g: glial cell body. **D.** Glial ensheathment (arrow) of unlabeled axonal fascicles at the apical edge of a glomerulus; inset shows neuroglial-labeled axon bundles in a similar region. **E.** Example of velate glial processes forming a sheath, by position likely to be a small tract of fibers. Stack z dimensions: A = 7.5 μm , B = 2 μm ; C = 4 μm ; D = 3 μm ; E = 2 μm .

Fig. 3.

Frontal sections showing glial investment through the depth of the adult antennal lobe beginning at the anterior surface. Images of the glial border around neuropil were deliberately saturated to allow fine glial processes in the neuropil to be visible. Note clear delineation of glomeruli by glial processes visible in more anterior sections (sections 1–100) compared to those in the most posterior sections (sections 101–120). an: antennal nerve; c: coarse central neuropil; ncb: neuronal cell bodies; nl: nerve layer; large arrow in sxs 41–70: lateral primary neurite tract; arrow in sxs 111–120: antennal commissure.

Fig. 4.

Glia-ensheathed tracts from the adult antennal lobe (AL, borders indicated with dotted line) and characteristics of the glial network around *Drosophila* glomeruli. **A.** Antenno-cerebral tract (arrow) extends posteriorly into the protocerebrum, the entire tract as well as fascicles within the tract ensheathed by glial processes. Neuropil labeled with antibody against Synapsin I (magenta). **B.** Tracts within the antennal commissure (ac) also are ensheathed and the entry/exit regions of the tract marked by a loose cluster of glial cell bodies (arrows). act: antenno-cerebral tract. Stack z dimensions: A,B = 2 μ m. **C.** Frame from a rotation series of a volume-rendered projection of glial cells and their processes in the neuropil of the adult antennal lobe, *Nrv2* strain. Arrow indicates position of the primary neurite tract for the lateral cluster of neuronal cell bodies. Stack z dimension: 26 μ m.

Fig. 5.

Glial investment of the adult antennal lobe: electron micrographs. **A.** Glial processes surround axon fascicles in the nerve layer and **(B)** extend into the neuropil from the glial border (g and arrowheads). ncb: neuronal cell bodies in an overlying cluster; t: tracheole. **C.** A portion of the lateral primary neurite tract showing ensheathing glial processes (arrowheads), sometimes in layers. Scalariform-like junctions (arrows) form between glial layers. **D.** Glial processes (arrowheads) within the synaptic neuropil (several synapses are circled) and **(E)** ensheathing large processes in the coarse central neuropil. Schematic diagram showing approximate region from which each panel was taken appears in Figure 6.

Fig. 6. Glial investment of the adult antennal lobe: electron micrographs (continued). **A.** In some regions, the synaptic neuropil directly abuts a very thin glial border around the neuropil (arrowheads). Several synapses are circled. pn: primary neurites traveling toward the tract that will carry them into the neuropil. **B.** Schematic diagram of a cross-section of the AL at a level typical of that from which the EM images in this figure were obtained. Boxes indicate regions similar to those used for each of the panels shown.

Fig 7. Developmental series showing the nerve layer. Olfactory receptor axons (labeled with *Drosophila* anti-neuroglial, magenta) initially course under and through the glial layer surrounding the nascent neuropil (arrowheads in **A,B**). Views of the nerve layer in **C,E,G,I** show regions in which the axons are mainly seen in longitudinal section; views in **D,F,H,J,K** show regions with axons mainly seen in cross-section. Arrows in **F,J,K** point to apical region of glomeruli where axon bundles have turned into the glomeruli, with fascicles ensheathed by glia. **A,B**. The first axons entering, and forming, the nerve layer are not ensheathed by glia and are only loosely traveling together. **C–K**. Axons in the nerve layer gradually become ensheathed by glial processes and travel long distances without changing to another fascicle. **G**. Clear view of the entry region of the AN, where axon bundles abruptly change course to enter the nerve layer. Stack z dimensions = 1–1.5 μ m. Scale bar in **I** applies to **A–K**. **L**. Schematic diagram summarizing nerve layer development. Each stage is a 3D representation of a piece of the nerve layer taken from a slab through the AL, as indicated by the turquoise section in the upper diagram. The nerve layer thickens with time

(and more axons). The glial investment, initially only on the outside, gradually extends processes toward the center of the lobe and gradually ensheaths fascicles. Note that although glial processes that extend from the glial cells ensheath axon bundles in the nerve layer *and* extend into the neuropil, processes extending into the neuropil have been left out of these diagrams in order to keep the focus processes occurring in the nerve layer.

Fig. 8.

Glial-cell investment of the antennal lobe neuropil at 25–50% development, as shown by *Nrv2* labeling. Each image in this and Figures 9 and 10 is a 2-D stack of 10 optical sections ($z = 5 \mu\text{m}$) through approximately the same region of the lobe, unless noted otherwise. **A.** At 25%, a border of glial cells surrounds the AL neuropil and clearly demarcates its boundaries. No glial process ingrowth is yet evident. **B.** At 30%, a small fringe of glial-cell process ingrowth into the neuropil is now visible (arrow) along the inside of portions of the glial border. **C.** By 33%, there is a thickening of the glial fringe, with more processes extending further into the neuropil (arrow). The greatest extension of glial processes from the developing fringe is seen in the region of primary neurite ingrowth (*). **D.** Thickening of the fringe and its development in other portions of the glial border surrounding the neuropil is seen through 40% of development (arrows). **E.** At 45%, there is modest process ingrowth toward the middle of the lobe, mainly confined to the region of primary neurite ingrowth that occurs at a more or less stereotypical position along the lateral portion of the glial border. Processes also extend early from a portion of the medial border (arrow). **F.** By 50%, distinct glial processes are seen growing into the lobe toward its center (arrows). an, antennal nerve.

Fig. 9.

Glial-cell investment of the antennal lobe neuropil at 60% and 67% of development as shown by *Nrv2* labeling. **A,B.** 60%. Thickening of the glial border around the lobe along with increased process growth into the antennal lobe neuropil (arrows); some processes are already velate. Individual glial processes have reached the center of the lobe. Glomeruli cannot yet be distinguished by their glial process surrounds. Arrowhead: primary neurite tract. **C–F.** 67%. **C.** Glial processes have begun to form loose networks around a few glomeruli (*) and the glial investment of the coarse central neuropil (c) becomes obvious (arrows). **D.** Glial cell processes heavily invest the primary neurite tract (†) and extend deep into the neuropil (arrows). **E,F.** Higher power views showing branched and often velate glial processes. an, antennal nerve. Stack z dimensions for E,F = 5 μ m.

Fig. 10.

Glial-cell investment of the AL neuropil at 75% development. **A.** Within the neuropil there are now many glial processes (arrow) projecting from the glial border around the neuropil. A single glial cell body appears to have migrated a short distance along the primary neurite tract into the neuropil (arrowhead). **B.** At higher magnification, glial processes (arrows) are seen within the synaptic neuropil of glomeruli. an, antennal nerve. Stack z dimension for A = 5 μm ; B = 2 μm .

Fig. 11.

Developmental series showing the gradually increasing glial investment of the neuropil in the context of Synapsin labeling. Left column: Glial cells and their processes (*Nrv2*); middle column: synapsin labeling (magenta); right column: combined images. Each image is a 1–1.5 μm stack taken through the region of the AL in which the tract of primary neurites from lateral cluster neurons (white arrows in left column) enters the neuropil. Mature glomeruli at this level, roughly in the middle of the AL, are less discrete in appearance than they are more anteriorly (see Fig. 3). Arrows in **E** point to fine glial processes within the glomerular neuropil. an: antennal nerve. Scale bar in **A** applies to all images.

Fig. 12.

Electron micrographs from antennal lobes at 50% (**A,D**) and 67% (**B,C,E**) of development. **A.** Glial processes beginning to enwrap large neurites of the lateral primary neurite tract. **B.** Large dendrites within a glomerulus partially ensheathed by glial processes (arrowheads). **C.** Fine glial processes extending into the glomerular neuropil, with a synapse circled. **D.** Small fascicle (asterisk) of olfactory receptor axons embedded in the glial border of the neuropil (g, arrowhead). **E.** Receptor axon fascicles (asterisk) at the apical edge of a glomerulus. g: glial processes; gcb: glial cell body.

Fig. 13.

Antennal nerve at entry to antennal lobe. 40% of metamorphic development. **A.** Neuroglial-positive axons (magenta) traveling in parallel fascicles along the antennal nerve (AN), gradually changing position within the nerve. Small fascicles move between larger ones along the length of the nerve (white arrows show regions of the nerve in which examples appear). **B.** Entry region of the nerve (bracket, ~20 μm) showing small, mostly aligned, neuroglial-positive axon fascicles that abruptly change direction as they enter the antennal lobe nerve layer. **C.** Entry region of nerve (bracket) showing the relatively small glial investment of this region compared to that in the more distal nerve. AN outlined in magenta. Arrow: large glial cell body that often appears in the entry region. **D.** 2-D projection of neuroglial-labeled receptor axons as they approach, enter and grow in tracts through the nerve layer of the antennal lobe, with no obvious sorting along that trajectory. Stack z dimensions: **A,B** = 1 μm ; **C** = 50 μm ; **D** = 33 μm .

A *Manduca* **B *Drosophila*** **Fig. 14.**

Comparison of the glial investment of adult *Manduca* and *Drosophila* glomeruli. Schematic diagrams on left show surface views of the extensive glomerular sheath formed by glial cells and their processes in *Manduca* and the network of processes surrounding a *Drosophila* glomerulus. Diagrams on the right, which represent very low power electron microscopic views through the body of one glomerulus and a portion of another, show 1) the easily recognized multi-layered glial border in *Manduca* compared to the lack of an obvious border in *Drosophila*; 2) glial sheaths (circular structures) enwrapping axon fascicles entering the glomeruli from the nerve layer (NL); 3) the sparseness of glial processes from the central region of a *Manduca* glomerulus compared to the relatively homogeneous distribution of glial processes throughout a *Drosophila* glomerulus. Inset: A *Drosophila* glomerulus drawn to the same scale as the *Manduca* glomerulus in **A**. GL: glomerular layer. CN: central coarse neuropil.

Fig. 15.

Schematic diagram comparing sequence of events in development of *Manduca* (A) and *Drosophila* (B) glomeruli. Green: glial cells; magenta: ORN axons; black: projection neurons; blue: local interneurons.

Fig. 16.

Chart placing developmental changes in glial cell investment of the neuropil in the context of development of the neuronal elements of the *Drosophila* olfactory pathway. (1) Lienhard and Stocker (1991); (2) Ray and Rodrigues (1995); (3) Stocker et al. (1995); (4) Dubin and Harris (1997); (5) Tissot et al. (1997); (6) Clyne et al. (1999); (7) Jhaveri et al. (2000); (8) Jhaveri and Rodrigues (2002); (9) Devaud et al. (2003); (10) Jefferis et al. (2004); (11) Hummel and Zipursky (2004); (12) Sen et al. (2005); (13) Marin et al. (2005); (14) Roy et al. (2007); (15) Yao et al. (2007); (*) this paper.