Published in final edited form as:

Auton Neurosci. 2008 December 15; 144(1-2): 61–75. doi:10.1016/j.autneu.2008.09.006.

The Development of Nicotinic Receptors in the Human Medulla Oblongata: Inter-relationship with the Serotonergic System

Jhodie R. Duncan, David S. Paterson, and Hannah C. Kinney

Department of Pathology, Children's Hospital Boston and Harvard Medical School, Boston, Massachusetts, 02115, USA

Abstract

Maternal cigarette smoking during pregnancy adversely affects fetal development and increases the risk for the sudden infant death syndrome (SIDS). In SIDS we have reported abnormalities in the medullary serotonergic (5-HT) system, which is vital for homeostatic control. In this study we analyzed the inter-relationship between nicotinic receptors (nAChRs), to which nicotine in cigarette smoke bind, and the medullary 5-HT system in the human fetus and infant as a step towards determining the mechanisms whereby smoking increases SIDS risk in infants with 5-HT defects. Immunohistochemistry for the $\alpha 4$ nAChR subunit and 5-HT neurons was applied in fetal and infant medullae (15–92 postconceptional weeks, $n=9$). The distribution of different nAChRs was determined from 39–82 postconceptional weeks ($n=5$) using tissue autoradiography for ^3H -nicotine, ^3H -epibatidine, ^3H -cytisine, and ^{125}I -bungarotoxin; the findings were compared to laboratory 5-HT_{1A} and 5-HT transporter binding data, and 5-HT neuronal density. Alpha4 immunoreactivity was ubiquitously expressed in medullary nuclei related to homeostatic functions from 15 weeks on, including rhombic lip germinal cells. At all ages, $\alpha 4$ co-localized with 5-HT neurons, indicating a potential site of interaction whereby exogenous nicotine may adversely affect 5-HT neuronal development and function. Binding for heteromeric nAChRs was highest in the inferior olive, and for homomeric nAChRs, in the vagal complex. In the paragigantocellularis lateralis, 5-HT_{1A} receptor binding simultaneously increased as $\alpha 7$ binding decreased across infancy. This study indicates parallel dynamic and complex changes in the medullary nicotinic and 5-HT systems throughout early life, i.e., the period of risk for SIDS.

Keywords

Acetylcholine; Autoradiography; Arcuate nucleus; Cigarette smoking; Paragigantocellularis lateralis; Sudden infant death syndrome

Introduction

Maternal cigarette smoking during pregnancy is a major public health problem world-wide despite extensive warnings about its harmful effects upon the fetus (Chiolero et al., 2005; Aliyu et al., 2007; Raatikainen et al., 2007). Indeed, 20–25% of women in the general population in the United States smoke during pregnancy (Chiolero et al., 2005; Raatikainen et al., 2007),

Correspondence: Jhodie R. Duncan, PhD, Department of Pathology, Enders 1109, Children's Hospital Boston, 300 Longwood Avenue, Boston, MA., 02115, jhodie.duncan@childrens.harvard.edu, Telephone: 617 919-4525, Fax: 617 730-0168.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

with up to 75% of women in high risk populations for infant mortality and morbidity e.g., American Indians in the Northern Plains (Iyasu et al., 2002) and Australian aborigines (Gilchrist et al., 2004). Cigarette smoking during pregnancy is associated in the offspring with prematurity (Chiolero et al., 2005; Raatikainen et al., 2007), low birth weight (Chiolero et al., 2005; Raatikainen et al., 2007), and cognitive/attention deficits (Julvez et al., 2007). In addition, the risk for the sudden infant death syndrome (SIDS), the leading cause of postneonatal infant mortality in the United States, is increased five-fold or more in infants born to mothers who smoke during pregnancy compared to those whose mothers do not smoke (Blair et al., 1996). SIDS is the sudden death of an infant what remains unexplained after a complete autopsy and death scene investigation (Willinger et al., 1991); typically a seemingly healthy infant is found dead after a sleep period. Reports of impairment of cardiorespiratory function and/or arousal in infants exposed to cigarette smoke during gestation suggest that such exposure harms the fetal development of cardiorespiratory and arousal pathways, the majority of which are located in the brainstem (Kahn et al., 1994; Lewis and Bosque, 1995; Parslow et al., 2004). Brainstem neurotransmitter abnormalities are found in SIDS infants, particularly in the serotonergic (5-HT) system in the medulla oblongata which is involved in the modulation of cardiorespiration and arousal under homeostatic stress (Panigrahy et al., 2000; Kinney et al., 2001; Kinney et al., 2003; Kinney, 2005; Paterson et al., 2006b) and thus these abnormalities in the 5-HT system are likely to play an important role in the failure to respond to a life-threatening event after birth. SIDS infants exposed to cigarette smoke during fetal development also demonstrate abnormal binding to nicotinic receptors (nAChRs) in mesopontine regions related to arousal in combination with the medullary 5-HT abnormalities (Duncan et al., 2008). Given that nicotine is a major neurotoxic component of cigarette smoke and is a potent agonist to the brain's endogenous nAChRs, the SIDS data suggest that the increased risk for SIDS due to maternal smoking during pregnancy may be related to rostral nAChRs abnormalities induced by exogenous nicotine in association with the caudal 5-HT abnormalities. Critical to determining the inter-relationships of nAChR and the medullary 5-HT system and the pathogenesis of SIDS is a basic understanding of the development of nAChRs relative to medullary 5-HT neurons, receptors, and transporter (5-HTT) in the fetal period and infancy. Yet, virtually nothing is known about the relevant chemical anatomy in the developing human medulla, and this we undertook the following baseline study to help bridge this gap in knowledge.

Nicotinic receptors are pentameric, ligand-gated ion channels comprised of heteromeric or homomeric subunits encoded by 9α ($\alpha 2-10$) and 3β ($\beta 2-4$) genes (Gotti et al., 2006), with $\alpha 7$ being the only subunit known to form homomeric receptors in the human brain. These subunits are located at pre-, post- and/or peri-synaptic sites (Lucas-Meunier et al., 2003) and each nAChR subtype displays unique physiological and pharmacological properties (Cimino et al., 1995; Hellstrom-Lindh and Court, 2000; Gotti et al., 2006), as well as distinct neuroanatomical distributions at different developmental time-points (Cimino et al., 1995; Adams et al., 2002; Tribollet et al., 2004). Presynaptic nAChRs regulate neurotransmitter release, while postsynaptic nAChRs activate intracellular signaling and gene transcription (McKay et al., 2007). Nicotine exposure is known to have multiple effects on the 5-HT system (Mihailescu et al., 1998; Kenny et al., 2001; Slotkin et al., 2007b), and thus the expression of nAChRs by 5-HT neurons may play an important role in the 5-HT abnormalities reported in SIDS. In the human infant, the medullary 5-HT system is comprised of 5-HT neuronal cell bodies in the raphé (raphé obscurus and raphé pallidus), extra-raphé (paragigantocellularis lateralis [PGCL], gigantocellularis, intermediate reticular zone, and subtrigeminal nucleus), and ventral surface (embedded within the arcuate nucleus) (Kinney et al., 2007). In a series of normative studies of the human fetal and infant medulla, we have reported the temporal and topographic profiles of 5-HT neuronal cell bodies (Kinney et al., 2007), 5-HT receptors (Zec et al., 1996; Paterson et al., 2004a), and 5-HT transporter (5-HTT) (Paterson et al., 2004a).

In the following study, we expanded upon this valuable 5-HT-related database by delineating the spatial and temporal sequence of nAChR receptor development relative to maturational changes in the medullary 5-HT system. We tested the following hypotheses: 1) nAChRs are concentrated in regions related to medullary sites that contain 5-HT cell bodies and/or receive 5-HT terminals and are involved in cardiorespiratory control and/or arousal; 2) nAChRs are expressed by 5-HT cell bodies, indicating a potential site whereby exogenous nicotine in cigarette smoke interacts directly with 5-HT neurons to alter 5-HT development and/or function; 3) nAChR subtypes are expressed in the fetal medulla in sites related to cardiorespiration and/or arousal, indicating a site of interaction with exogenous nicotine during prenatal development that may alter the developmental program of these sites and thereby alter cardiorespiratory patterning and arousal responses postnatally; and 4) rapid changes are occurring in the development of 5-HT-related markers and nAChRs in parallel, suggesting that the fetal and infant time-frame, i.e. the risk period for SIDS, is a critical period in the development of medullary 5-HT system relevant to nAChRs. To test these hypotheses, we used single- and double-labeling immunocytochemistry and tissue section autoradiography, and we integrated the new nAChR-related findings with the previous 5-HT related findings (Kinney et al., 2001; Kinney et al., 2003; Paterson et al., 2006b) in our ever-expanding database of human brainstem chemical anatomy in early life. Given the extraordinary difficulty in collecting normative brainstems from human fetuses and infants, we believe this combined analysis is justified and necessary.

Materials and Methods

Clinical Database

Brainstems were obtained from autopsies in the Departments of Pathology, Children's Hospital Boston and Brigham and Women's Hospital, Boston, MA, and the office of the Chief Medical Examiner, San Diego, CA with approval of the Committee on Clinical Investigation. Fetal age was determined by the standard method of foot-length measurements. All ages (in weeks) were expressed as postconceptional age which is the sum of gestational plus postnatal age.

Single-label Immunocytochemistry for the Developmental Profile of nAChRs

Immunocytochemistry was performed with antibodies to $\alpha 4$ and 5-HT cells (Table 1). Medullae were fixed in 4% paraformaldehyde, cut into 3 mm blocks, embedded in paraffin, serially or stepped sectioned at 4 μ m-thick in the coronal plane, and mounted upon glass microscopic slides. The paraffin sections were de-waxed and re-hydrated through to water. Antigen retrieval was preformed using 10% citrate buffer at 195 °F for 10 minutes, and following phosphate buffered saline (PBS) washes, sections were treated with peroxidase blocking reagent (Dako Cytomation, Carpinteria, CA) for 20 minutes. Tissue sections were blocked for 1 hour in 4% normal goat serum and 0.1% triton X following incubation with the primary antibody (Table 1). Following PBS washes, tissues were incubated with secondary antibodies followed by ABC elite kit (1hr, Vector, Burlingame, CA) and reacted with di-amino denzadine substrate (Dako Cytomation, Carpinteria, CA). All sections for the same antibody were stained in the same run and in all cases staining failed to occur with omission of the primary antibody. In this study the Chemicon and Covance $\alpha 4$ nAChR antibodies produced identical immunostaining patterns in the paraffin-embedded human medulla which was comparative to that of positive control tissues (rat brain) and to those reported previously in rodents (Cucchiaro and Commons, 2003) and rabbits ((Cucchiaro and Commons, 2003; Centeno et al., 2004). Specificity for human brain tissues using western blot characteristics for the Covance antibody has been previously reported (Whiting and Lindstrom, 1988) with immunolabelling being eliminated in brain sections from $\alpha 4$ knock out mice (Whiteaker et al., 2006). For analysis optimum results for $\alpha 4$ were obtained using the Covance antibody for DAB staining and the Chemicon antibody for immunofluorescence.

Each case had a minimum of three stained sections for analysis. The medullary nuclei were identified anatomically in reference to the human brainstem atlas of Olszewski and Baxter (1982), with the exception of the raphé nuclei, which were classified according to the atlas of Paxinos and Huang (1995). The levels and nuclei at which analysis was performed are presented with their atlas number (Olszewski and Baxter, 1982) in parentheses: 1) mid-medulla, level of nucleus of Roller (Plate X), for analysis of the nucleus of the solitary tract (NTS), dorsal motor nucleus of vagus (DMX), hypoglossal nucleus, principal inferior olive (ION), medial accessory olive, and arcuate nucleus; 2) rostral medulla (Plate XII), for analysis of the raphé obscurus, paragigantocellularis lateralis (PGCL), gigantocellularis, intermediate reticular zone, dorsal accessory olive, ION, and arcuate nucleus. As defined by us, the medullary 5-HT system consists of neurons with their cell bodies located in the raphé (raphé obscurus and raphé magnus), extra-raphé (PGCL, gigantocellularis, intermediate reticular zone, and subtrigeminal nucleus) and ventral positions, including the arcuate nucleus, and projection sites that mediate cardiovascular and respiratory functions, e.g., NTS, DMX, and hypoglossal nucleus (Kinney et al., 2007).

Immunostained sections were visualized using a Olympus BX51 microscope (Olympus America Inc., Melville, NY) and representative images captured using an Olympus DP71 camera (Olympus America Inc., Melville, NY) and Microsuite Biological Suite 5 computer software. Images were saved as .tiff files and image size and contrast adjusted in Adobe Photoshop Software (Adobe systems, San Jose, CA). Figures were constructed in CorelDraw Graphics Suite 12 (Corel Corporation, Ottawa, CA) including the addition of text and scale bars.

Nicotinic Receptor Binding in the Infant Medulla with Tissue Autoradiography

We investigated nAChR binding in newly acquired specimens with 4 different radioligands, each with different affinities to different subunits of the nAChRs (Table 2). Blocks of fresh infant medullae, stored at -70°C until use, were serially sectioned at $20\text{ }\mu\text{m}$ on a motorized Leitz cryostat and mounted on glass microscopic slides. For each radioligand, the general methods of tissue autoradiography were identical (Kinney et al., 1993); specific details for each radioligand are provided in Table 2. The unfixed sections were preincubated in buffer for 30–60 minutes, followed by incubation with the selected radioligand diluted in the same buffer. Non-specific binding was determined in a subset of adjacent sections that were incubated with the radioligand and excess “cold” displacer (Table 2). To remove unbound ligand, the sections were washed in a series of changes of buffer followed by distilled water. The sections were then dried under a stream of anhydrous air and exposed to film or phosphorimage plates (Table 2). A set of ^3H - or ^{125}I -standards (Amersham, Piscataway, NJ) was included in each X-ray cassette to permit the conversion of the optical densities of silver grains in autoradiograms to specific activity of tissue-bound ligand to femtomole per milligram (fmol/mg) of tissue. Receptor binding density (expressed as the specific activity of tissue-bound ligand) was analyzed in 11 nuclei of each medulla (see above). The one exception was ^3H -nicotine binding in which we used for reference and comparison the values previously obtained by us in 5 nuclei (Kinney et al., 1993; Nachmanoff et al., 1998). The selected nuclei were analyzed at 2 standardized brainstem levels (2 specific and 1 nonspecific autoradiograms for each nucleus, 4 measurements for each paired nucleus), according to standard sampling procedures established in our laboratory (Kinney et al., 2001). As there was no significant difference in binding measurements between the right and left (paired) nuclei, binding densities were combined and expressed as a mean $\pm$ standard error of the mean.

Generation of Brainstem Autoradiograms

The MCID Elite 6 imaging system (Imaging Research Inc., Ontario, CA) was used to perform quantitative densitometry of autoradiographs. Optical densities were converted to specific

activities of tissue bound ligand in fmol/mg tissue with ^3H or ^{125}I -standards. Receptor binding density was determined in each brainstem nucleus by digitizing the boundaries of nuclei upon the color image of the autoradiogram displayed on the computer monitor. Calculations to equate for nonspecific binding were made if needed. Specific activity data are presented as computer-generated mosaics with an increasing linear color step scale for each ligand. The quantitative data for comparison of binding levels across different radioligands are presented with a “normalized scale” of the percentage of binding relative to a nucleus with high binding, thus the ION (100%) for nAChR markers or raphé obscurus (100%) for 5-HT markers. Binding levels are then compared by the following scale: 0: no binding; 1+: low binding (1–25%); 2+: moderate binding (26–50%); 3+: high binding (51–75%); 4+: very high binding (76–100%); 5+: intense binding (>101%).

Serotonergic Markers in the Infant Medulla

For neurochemical analysis of 5-HT_{1A} receptors and 5-HTT binding, and immunocytochemistry analysis for quantification of 5-HT cell number as markers of the 5-HT system, we combined nAChR observations in recently acquired tissue specimens with 5-HT related observations based upon the established database in our laboratory (Zec et al., 1996; Paterson et al., 2006b; Kinney et al., 2007). Our methods and distribution patterns for 5-HT_{1A} receptors, 5-HTT binding, and 5-HT neuronal density have been published previously (Paterson et al., 2006b; Kinney et al., 2007).

Double-Label Immunofluorescence for the Co-localization of Nicotinic Receptors and 5-HT Neurons

Double-label immunofluorescence was performed to determine co-localization and distribution of $\alpha 4$ nAChR (Chemicon) and 5-HT neurons (Table 1) using alternating sections on the paraffin-embedded cases above (n=5). The same methods described above were employed with the exception of 20 minutes of citrate buffer retrieval and no peroxidase blocking reagent treatment before the cocktail mixture of antibodies was applied to the tissue ON at 4°C after which the appropriate Alexa-Fluor secondary antibodies (Invetrogen Molecular Probes, Eugene, Oregon, USA) were applied (Table 1, 1 hr). Omission of the primary antibody resulted in no staining. Slides were viewed using either FITC or TRITC filters using a Olympus BX51 microscope (Olympus America Inc., Melville, NY). Images were captured using a Photometrics Cool Snap Fx camera (Photometrics, Tucson, CA) and MICD Elite 6.0 program (Imaging Research Inc., Ontario, Canada). Images were pseudo-colored and merged using Adobe Photoshop 6.0.

Statistical Analysis

All data are presented as the mean $\pm$ standard error of the mean. Linear regression analysis was used to determine the effects of postconceptional age and postmortem interval on receptor binding. P-values <0.05 were considered significant.

Results

Immunocytochemistry

Clinical Database—For immunocytochemistry, 9 medullae fixed in 4% paraformaldehyde and embedded in paraffin were examined. The ages of the cases in postconceptional weeks were: 15, 23, 30, 37, 38, 42, 52 (3 postnatal months), 80 (10 postnatal months), and 92 (approximately 1 year) weeks. The mean age was 45.4 ± 8.5 postconceptional weeks. The causes of death were: spontaneous abortion (n=2); cardiopulmonary disease (n=3); Fryns Syndrome (n=1); neuroblastoma (n=1), and mild hypoxic-ischemic encephalopathy (n=2). There were no histopathologic brainstem changes in any case upon standard neuropathologic examination.

Postmortem intervals ranged from 11–48 hours (median 12 hours), with no visually obvious effect of postmortem interval upon immunostaining.

The Developmental Distribution of the $\alpha 4$ nAChR Subunit—The earliest time-point examined by us of $\alpha 4$ immunostaining was 15 gestational weeks, i.e., early in the second trimester, and the latest time-point, 13 postnatal months, i.e., the end of infancy. At 15 weeks, we previously reported that early differentiating 5-HT neurons are present in raphé obscurus, raphé pallidus, gigantocellularis, and PGCL, as well as scattered within the arcuate nucleus along the ventral medullary surface (Kinney et al., 2007). At this same age, we now report that $\alpha 4$ was expressed by differentiating (5-HT and non-5-HT) neurons in virtually all nuclei throughout the medulla, but with variable intensity of immunostaining (Figure 1). Notably, moderate immunostaining was present upon neuronal cell bodies in the raphé obscurus (Figure 1A) and extra-raphé (Figure 1E), i.e., sites of 5-HT cell bodies. Intense immunoreactivity was also present in sites involved in cardiorespiratory control and which are known to receive heavy 5-HT projections, i.e., hypoglossal nucleus (upper airway patency) (Figure 1A) and DMX (preganglionic outflow of the parasympathetic system), with a mild degree of immunostaining in the nucleus ambiguus (innervation of larynx and pharynx). The undifferentiated neurons of the primordia of the ION expressed mild immunostaining (Figure 1A). Of note, $\alpha 4$ immunostaining was detected in the early differentiating cytoplasm of scattered germinal cells of the rhombic lip (Figure 1B); these immunopositive cells were immature with a small amount of cytoplasm rimming the large nucleus. Furthermore, $\alpha 4$ immunopositive neurons with trailing processes were seen adjacent to the rhombic lip at the ventrolateral medullary surface (Figure 1C). Intense punctate immunoreactivity was also present throughout the neuropil (Figure 1E, F). In the 8 cases ranging in age from 23 gestational weeks (midgestation) to 13 postnatal months, the pattern of $\alpha 4$ immunostaining was essentially the same as that at 15 gestational weeks with no obvious developmental differences in the intensity or topography (Figure 2). The two exceptions were found in known derivatives of the rhombic lip, i.e., arcuate nucleus and ION (Figure 3). In both of these regions, there was an obvious visual increase in the intensity of immunostaining from 23 to 44 gestational weeks (Figure 3); thereafter staining intensity did not visually change.

Relationship of $\alpha 4$ to 5-HT Neuronal Cell Bodies—With the application of double-label immunocytochemistry, we found that as early as 15 gestational weeks, 5-HT (and non 5-HT) neurons in the raphé obscurus (Figure 4A–C) and extra-raphé (data not shown) co-localized with $\alpha 4$. This same pattern of co-localization was observed at term (Figure 4D–F), 3 postnatal months, and 10 postnatal months (Figure 5). In addition, 5-HT neurons in the PGCL (Figure 6C), and arcuate nucleus (Figure 6F) co-localized with $\alpha 4$ at these ages. The $\alpha 4$ immunostaining was discrete and punctuate upon the 5-HT cell bodies (Figures 4 and 6), consistent with postsynaptic receptor localization (Figure 6C).

The Quantitative Distribution of nAChR Binding in the Human Infant Medulla Autoradiography Clinical Database—We compared the binding patterns of three specific nAChR radioligands in newly accrued tissues to each other and to binding to the broad radioligand ^3H -nicotine (the later determined in our archival dataset) (Kinney et al., 1993; Nachmanoff et al., 1998). In the current (non-published cases), nAChR receptor binding to ^3H -epibatidine, ^3H -cytisine, and ^{125}I -bgt was analyzed in 5 infant cases at 39 and 41 weeks, and 4, 6 and 9 postnatal months (mean: 55.5 ± 6.9 postconceptional weeks). The causes of death were: congenital heart disease, $n=3$; acute pneumonia, $n=1$; and accidental asphyxia, $n=1$; there was no primary brainstem histopathology in any case. The mean postmortem interval was 14.6 ± 2.0 hours. There was no effect of postmortem interval upon nAChR binding, as previously reported by us for ^3H -nicotine (Nachmanoff et al., 1998; Duncan et al., 2008). We also compared the quantitative distribution of nAChR binding with each of the three radioligands

under study (exception ^3H -cytisine due to insufficient data) to 5-HT receptor and transporter binding, and 5-HT neuronal density in our archival database (Paterson et al., 2006b; Kinney et al., 2007).

General observations—The relative distribution of nAChR binding with ^3H -nicotine, ^3H -epibatidine, ^3H -cytisine and, ^{125}I -bgt varied across medullary nuclei depending on the radioligand used (Table 3, 4) (Figure 7), as expected given that differences in binding patterns are based on the affinity of each ligand to specific nAChR subunits (Table 2). For all radioligands except ^{125}I -bgt, the highest binding was striking in the ION (Table 3) (Figure 7). Accordingly, we arbitrarily defined the binding level in the ION as the “internal standard” at 100% for each nAChR radioligand and calculated the binding in each nucleus for each radioligand as a percent of this standard (Table 4). In this way, we were able to compare the relative binding levels among the radioligands to each other. While several developmental changes in nAChR binding levels were observed with different radioligands we found that the major postnatal changes in binding occurred in the DMX because they involved changes in more than one radioligand, i.e., ^3H -epibatidine and ^{125}I -bgt (Table 5) (Figure 8).

^3H -Nicotine Binding—From analysis in this study and the archival dataset, we observed that ^3H -nicotine binding which demonstrates affinity to all nAChRs was the highest in the ION (Table 3), with 50–80% higher levels than that in any other nucleus ($p < 0.001$) (Figure 7A) (Kinney et al., 1993). The lowest detectable binding was in the raphé obscurus, no binding was detectable in the arcuate nucleus at the ventral surface (Table 3). There were no significant changes in ^3H -nicotine binding in any nucleus analyzed over the age-range studied (38–66 postconceptional weeks) (Table 5).

^3H -Epibatidine Binding—Tritiated-epibatidine binding (affinity to the $\alpha 2-4$, $\beta 2$, and $\beta 4$) was visually prominent in all medullary nuclei, in contrast to the overall low binding in these nuclei to ^3H -nicotine (with the obvious exception of the ION). As with ^3H -nicotine binding, ^3H -epibatidine binding was highest in the ION, with binding levels approximately 80% higher than those of the raphé obscurus and arcuate nucleus and 43% higher than the NTS and DMX (Table 3) (Figure 7B). Notably, binding in the arcuate nucleus was visually distinct in contrast to its absence with ^3H -nicotine binding (Figure 7). Of all the radioligands analyzed, the pattern and intensity of ^3H -epibatidine binding most closely resembled that of the $\alpha 4$ immunostaining. There was a significant decrease in ^3H -epibatidine binding with increasing postconceptional age in two nuclei, the DMX ($p = 0.01$) (Table 5) (Figure 8A) and dorsal accessory olive ($p = 0.02$) (Table 5).

^3H -Cytisine Binding—In the 2 available cases (39 and 41 postconceptional weeks), ^3H -cytisine binding to $\alpha 4\beta 2$ was highest in the ION, as well as in the dorsal accessory olive (Table 3). The binding in all tegmental nuclei, however, was very low, even more so than that for ^3H -nicotine (Table 4) (Figure 7C). Interestingly, binding in the medial accessory olive was approximately 80% lower than that in the other components of the olivary complex, i.e., ION and dorsal accessory nucleus. Evaluation of the developmental profile in infancy was precluded by the small sample size.

^{125}I -Bungarotoxin Binding—The pattern of binding to homomeric nAChRs with ^{125}I -bgt was distinctly different from that of ^3H -nicotine, ^3H -epibatidine, and ^3H -cysteine, i.e., all heteromeric nAChRs. In contrast to the highest binding with all of the heteromeric radioligands in the ION, binding with ^{125}I -bgt to $\alpha 7$ was highest in the DMX and NTS (Table 3, 4), being three-fold higher than that in the ION in the DMX, and one-fold higher in the NTS (Table 3) (Figure 7D). The lowest binding was present in the arcuate nucleus, although it was still visually distinct. There were significant decreases in ^{125}I -bgt binding with increasing postconceptional

age in four nuclei: NTS ($p=0.02$), DMX ($p=0.03$) (Figure 8B), PGCL ($p=0.01$), and arcuate nucleus ($p=0.002$) (Table 5). These changes are in contrast to the findings above of changes occurring in two nuclei with ^3H -epibatine and in none with ^3H -nicotine.

Interactions between nAChR binding and Markers of the 5-HT System during Infancy

—Using the unique archival normative data concerning 5-HT markers in our laboratory, we compared the developmental profiles of nAChR expression and 5-HT neuronal density, 5-HT receptor binding, and 5-HTT binding of the medullary 5-HT system. This analysis revealed that in only one nucleus, the PGCL, were there simultaneous changes in both nicotinic and 5-HT-related markers, i.e., there was an increase in 5-HT_{1A} receptor binding and decrease in $\alpha 7$ receptor binding with increasing age in early infancy (Table 5) (Figure 9), with the “cross-over” time-point at 2–3 months. While we found decreasing nAChR binding with increasing postnatal age in the vagal complex (DMX [^3H -epibatine and ^{125}I -bgt] and NTS [^{125}I -bgt]), dorsal accessory olive (^3H -epibatine), and arcuate nucleus (^{125}I -bgt), there were no concomitant changes in these nuclei in any 5-HT marker (Table 5). In the hypoglossal nucleus, 5-HT_{1A} receptor binding increased with increasing postconceptional age, yet there were no changes in nAChR binding over the same time-period. We found no developmental changes for either nicotinic or 5-HT markers in the raphe obscurus, intermediate reticular zone, or gigantocellularis nucleus i.e., in all sites of 5-HT neurons (Kinney et al., 2007), ION, or medial accessory olive (Table 5). In addition, 5-HTT binding and 5-HT neuronal densities were unchanged in any nuclei over infancy (Table 5). In essence, we found that dynamic changes in either the nAChR and/or 5-HT markers occurred in 6 of the 11 (55%) nuclei sampled throughout infancy indicating complex developmental profiles simultaneously in the nicotinic and 5-HT systems in the medulla (Figure 10).

Discussion

The agonist nicotine in maternal cigarette smoke crosses the placenta into the fetal circulation and then crosses the fetal blood-brain-barrier to bind to brain nAChRs, with the potential to alter the development and function of the cholinergic, as well as other neurotransmitter, systems (Luck et al., 1985; Slotkin et al., 2005; Slotkin et al., 2007a; Dwyer et al., 2008). In this study, we defined the developmental topography of cholinergic nAChRs relative to the medullary 5-HT system in the human fetus and infant as a first step towards deciphering the underlying basis of the effect of exogenous nicotine in maternal cigarette smoke upon the developing human 5-HT system, i.e., a system of special interest in the link between SIDS and maternal cigarette smoking during pregnancy. Our major finding is that the maturational profiles of nAChRs and markers of 5-HT neuronal cell bodies, receptors, and transporter are complex and dynamic relative to one another in early life, suggesting a vulnerable period in their simultaneous development. Moreover, nAChR expression is present in virtually all medullary nuclei, including those that are comprised by 5-HT cell bodies or that receive 5-HT projections. The expression of nAChRs in cardiovascular, respiratory, and motor nuclei in the human fetal and infant medulla is supported by extensive animal data (Swanson et al., 1987; Hellstrom-Lindahl et al., 1998; Ferreira et al., 2000; Cucchiari and Commons, 2003; Kubin and Fenik, 2004; Padley et al., 2007). Rodent studies suggest that the cholinergic projections to medullary nuclei are primarily from the pedunculopontine tegmental nucleus and laterodorsal tegmental nucleus in the mesopons that are critical for arousal and the generation of REM sleep, suggesting that nAChRs also play an important role in these components of state regulation (Frank et al., 2001; Lena et al., 2004; Padley et al., 2007; Rukhadze and Kubin, 2007). Of note, prenatal nicotine exposure interrupts normal sleep/wake cycling by reducing periods of REM sleep in the neonatal rat (Frank et al., 2001), and $\beta 2$ knockout mice experience increased periods of REM (Lena et al., 2004) and impaired respiration which is enhanced upon nicotine exposure (Cohen et al., 2002). In the following discussion, we highlight insights from our data into the potential neuroanatomic substrate of the adverse effects of maternal cigarette smoke (nicotine)

on the development and function of cardiovascular and respiratory 5-HT related networks in the human medulla.

Fetal Expression of nAChRs in the Human Medulla

In the immunocytochemical analysis of nAChR expression, we focused upon the $\alpha 4$ subunit due to its high affinity to nicotine (Benwell et al., 1988; Flores et al., 1992) and its known role in the medullary control of homeostasis (Fonck et al., 2005; Hatori et al., 2006; Shao et al., 2008). We found $\alpha 4$ expression at 15 gestational weeks by a subpopulation of germinal cells in the rhombic lip, the origin of cerebellar-relay nuclei in particular in the medulla and pons (Rakic and Sidman, 1982; Landsberg et al., 2005). This observation suggests that $\alpha 4$ plays a trophic role upon at least a subset of rhombic lip precursors. In embryonic and fetal development, acetylcholine and nicotine via nAChRs experimentally influence neuronal cell proliferation, migration, survival, and differentiation (Opanashuk et al., 2001; Torrao and Britto, 2002). Functional nAChRs are present on stem and progenitor cells in the cerebral cortex of the mouse as early as embryonic day 10 (Schneider et al., 2002). The trophic role of acetylcholine during early life is underscored by experiments in which ablation of basal forebrain cholinergic neurons results in reduced laminar thickness, neuronal size, and dendritic tree field in the cortical projection targets (Hohmann et al., 1988; Robertson et al., 1998). In fetal monkey brains from offspring of mothers exposed to nicotine, regional neuronal cell loss, decrease in cell size, and abnormal neurite outgrowth are found (Slotkin et al., 2005), suggesting that nAChR-mediated trophic effects are adversely altered by exogenous nicotine exposure; importantly, the prenatal effects persist postnatally (Good et al., 2006; Slotkin et al., 2007b). Of direct relevance to the rhombic lip, multiple nAChR subtypes are expressed by cultured rat granule neurons, derivatives of the rhombic lip in the cerebellar cortex (Didier et al., 1995), as well as by the human external granular layer (Court et al., 1995). Interestingly, the mouse external granule cells express $\alpha 3$, not $\alpha 4$, nAChR subunits, a potential species-, age- and/or region-specific difference of $\alpha 4$ expression in the human rhombic lip (Opanashuk et al., 2001). In addition to the rhombic lip, we observed nAChR expression in multiple post-mitotic neuronal subtypes in medullary nuclei as early as the second trimester, the earliest time-point studied by us. Parenthetically, nAChRs have been reported in human fetal brainstem homogenates as early as 4–6 gestational weeks (Hellstrom-Lindahl and Court, 2000).

Nicotinic Receptors and Their Relationship to the Medullary 5-HT System

There is considerable functional evidence for important nicotinic-5-HT interactions in animal models. Nicotine injected into rat brains, for example, inhibits 5-HT neuronal firing, an affect blocked by the nAChR antagonist mecamylamine (Touiki et al., 2007). Nicotine exposure also increases 5-HT release and alters 5-HT neuronal firing in rat hippocampal and mesencephalic slices in a dose-dependent manner (Mihailescu et al., 1998; Kenny et al., 2001). In rats, prenatal nicotine via maternal osmotic pumps from gestational day 4 to term suppresses 5-HT_{1A} receptor binding and upregulates 5-HT₂ receptor binding at 6 postnatal months (Slotkin et al., 2007b). In our study, the $\alpha 4$ subunit is expressed by the soma and processes of 5-HT (and non-5-HT) neurons at 15 gestational weeks (and thereafter) in the raphé (raphé obscurus), extraraphé (gigantocellularis and PGCL) and ventral surface (arcuate nucleus). Thus, exogenous nicotine in maternal cigarette smoke potentially binds to these receptors to adversely influence 5-HT neuronal development and/or function via receptor-mediated interactions. In the infant, nicotine from second hand smoke could act at these sites as well.

Of the 11 nuclei in the medulla that we analyzed both with nicotinic and 5-HT markers, only the PGCL demonstrates changes in the two systems simultaneously, whereas one nucleus (hypoglossal nucleus) demonstrates changes only in 5-HT markers (5-HT_{1A} receptor binding), and four nuclei (NTS, DMX, dorsal accessory nucleus, and arcuate nucleus), changes in nAChR binding only. Over this same time-period, 5-HTT binding and 5-HT neuronal cell

density do not change. This analysis was based upon the premise that receptor and transporter binding levels are reasonable “markers” of transmitter function and that developmental changes within them are indicators of how the transmitter system as a whole changes with maturation. While we do not propose that a change in one system is driving a change in the other, the simultaneous occurrence of changes in the two systems suggests an exceptionally “active” (susceptible) time for these two systems in the PGCL relative to each other, notably during the peak age for SIDS. Thus, it seems appropriate to focus in the future upon the PGCL as a “model” medullary site to probe nicotinic-5-HT interactions both at the physiological and neuropathologic levels. The basis of the rise in 5-HT_{1A} receptor binding and concomitant decrease in $\alpha 7$ receptor binding during early infancy is unknown, begging the question of the type of underlying changes in neuronal number, terminal projections, dendritic arborization, and/or spine maturation that “account” for changes in receptor number and/or affinity. Nevertheless, receptor changes for each system may be independent of other structural changes, given receptor binding increases for one ligand and decreases for the other, but this speculation remains to be proven. The preponderance of nuclei sampled with changes in nAChR receptor binding changes (five of the eleven nuclei), as opposed to those with 5-HT_{1A} binding (two of the ten nuclei), suggests enhanced developmental plasticity in nicotinic as opposed to 5-HT receptors in early infancy, but again, considerably more information is needed for certainty.

The Localization of nAChRs in the Sites Relevant to Cardiopulmonary Function and Arousal in the Human Fetal and Infant Medulla

Paragigantocellularis lateralis—The PGCL, at least in part, is the human homologue of the preBötzinger complex in rodents, i.e., the putative site of respiratory rhythm generation (Paterson et al., 2004b). Within the preBötzinger complex in mouse models, nicotine alters the respiratory pattern by exciting inspiratory neurons via $\alpha 4\beta 2$ and not $\alpha 7$ receptors (Shao and Feldman, 2002). Of major interest in regards to the effect of maternal cigarette smoke and nicotine upon arousal in the off-spring is also evidence that the PGCL plays a role in sleep/wake patterning (Darnall et al., 2005). Physiological studies in human infants exposed to cigarette smoke during gestation demonstrate abnormal arousals (Lewis and Bosque, 1995; Franco et al., 1999; Horne et al., 2002; Blair et al., 2006); moreover, prospective monitoring studies indicate deficits in cortical and subcortical arousals in SIDS infants prior to death (Kahn, 1992; Schechtman et al., 1992; Kato et al., 2003). Thus, the question arises as to the site of action of prenatal nicotine upon brainstem arousal systems, given that arousal is mediated by descending, as well as ascending, pathways that influence subcortical, as opposed to cortical, arousal (Siegel, 2002; Sinton and McCarley, 2004; Saper et al., 2005). Descending (subcortical) pathways include 5-HT neurons in the raphe obscurus (Brown et al., 2008) and PGCL (Bradley et al., 2002; Darnall et al., 2005), and their disruption results in sleep fragmentation and the virtual loss of REM sleep. We speculate that acetylcholine via the mesopontine source nuclei constitutively affects descending arousal through influencing 5-HT neuronal firing in these medullary sites, an idea for testing in animal models with transmitter-specific neuroanatomic and neurophysiologic methods.

The vagal complex—We found that the DMX (preganglionic parasympathetic outflow) and NTS (visceral sensory input) are the nuclei with the highest homomeric ($\alpha 7$) receptor binding (¹²⁵I-bgt) in the infant medulla. Dissociated preganglionic neurons of the rat DMX express functional nAChRs (Nabekura et al., 1995), and application of nicotine to medullary slices causes depolarization of DMX neurons (Ito et al., 1989). Acetylcholine excites cardiac vagal neurons that control heart rate and cardiac function and that are located in the DMX and nucleus ambiguus in part via the activation of direct ligand-gated postsynaptic nAChRs (Wang et al., 2001). In human infants, $\alpha 7$ nAChR binding is three-fold higher in the DMX and one-fold higher in the NTS than the ION, the site with the highest binding density for heteromeric nAChRs. This observation potentially suggests a greater role for $\alpha 7$ compared to other nAChR

subunits in the function(s) of the vagal complex. Noteworthy is the observation that the most dramatic changes in nAChR binding occur in the DMX and NTS across human infancy.

Arcuate nucleus—This nucleus, comprised of neurons of heterogeneous transmitter phenotypes clustered over the ventral surface of the human medulla, is postulated to be homologous to the respiratory chemosensitive fields in cats (Filiano et al., 1990; Paterson et al., 2006a; Benarroch et al., 2007). Chemosensitivity in the arcuate nucleus has largely been attributed to 5-HT neurons embedded in this region (Paterson et al., 2006a; Kinney et al., 2007), as 5-HT neurons on the ventral surface of the rat have been shown to be chemosensitive (Richerson et al., 2001). Structural and neurotransmitter abnormalities have been reported in the arcuate nucleus in SIDS cases by us and others (Panigrahy et al., 1997; Mallard et al., 1999; Panigrahy et al., 2000; Kinney et al., 2003; Paterson et al., 2006b), including reduced muscarinic receptor binding (Kinney et al., 1995) but not altered nAChR binding (Nachmanoff et al., 1998; Duncan et al., 2008). In the present study, we found that 5-HT neurons (as well as non-5-HT neurons) in the human arcuate nucleus express $\alpha 4$, suggesting a site of interaction whereby acetylcholine or nicotine influences the central response to carbon dioxide. Indeed, low doses of nicotine applied to the ventral medullary surface of cats elevate blood pressure and respiration, and high doses decrease heart rate and increase respiration (Dev and Loeschke, 1979). Using the radioligand ^3H -nicotine, we previously reported that there are no binding abnormalities to nAChRs in the arcuate nucleus in SIDS cases compared to controls (Nachmanoff et al., 1998; Duncan et al., 2008). Yet, the present study suggests that the use of the ^3H -nicotine may have masked a specific nAChR subunit binding abnormality because it was too broad and did not visually highlight this region in autoradiographs. The presence of visually detectable binding with ^3H -epibatidine and ^{125}I -bgt that we observed in this study underscores the need in future studies to apply these specific radioligands in order to detect potential nAChR binding differences between SIDS and controls in the arcuate nucleus.

Principle inferior olive—In this study, the ION was the single medullary region with the highest binding for all radioligands analyzed for heteromeric (but not homomeric) nAChRs in the infant period; indeed, binding density in this site was 5–8-fold higher than that of any other nucleus, including the dorsal and medial accessory olives, subdivisions of the human inferior olivary complex. The ION, the sole source of climbing fibers to the Purkinje cells, is known to receive cholinergic input from mesopontine ACh neurons (Woolf and Butcher, 1989), and is critical for somatic motor, respiratory, and autonomic coordination (Urbano et al., 2006; Zhuang et al., 2008). The specific role of nAChRs in the human infant ION is unknown, but their heavy concentration at this site likely increases the susceptibility of this nucleus to nicotine neurotoxicity at least through infancy.

Potential Limitations of the Study

A potential limitation of this study is the small sample size of fetal and infant medullae; yet normative tissues from human cases in early life are extraordinarily difficult to accrue, given that autopsies are extremely rare in this age-group. Despite the small sample size, we found robust developmental changes that underscore their potential validity. Obviously, ongoing accrual of cases is needed for confirmation and is in progress. While results from knockout mice studies on the specificity of currently available antibodies to nAChRs are unclear (Whiteaker et al., 2006; Moser et al., 2007), extrapolation of these findings to other species, including humans, should be made with caution (Jones and Wonnacott, 2005). However, the authors acknowledge that until specificity can be validated, interpretation of the results obtained with the use of the $\alpha 4$ nAChR antibody in the human medulla should be made in combination with the distribution patterns of ligand binding. Furthermore the absence of a reliable $\alpha 7$ antibody for use in human postmortem tissues and thus the inability to assess its protein distribution with Immunocytochemistry is another limitation of this study. In rats,

however, $\alpha 7$ protein expression is present in the NTS, DMX, hypoglossal nucleus and extra-raphe (a region containing 5-HT neurons) (Dominguez del Toro et al., 1994). Thus, $\alpha 7$ binding expression seen in the human infant medulla in this study appears similar to $\alpha 7$ protein expression determined by immunohistochemistry in the rat.

Conclusions

Cholinergic nAChRs are ubiquitous in the human fetal and infant medulla, including in the medullary 5-HT system. Importantly, 5-HT neurons express nAChRs as early as the second trimester, providing the substrate whereby nicotine neurotoxicity is triggered by exposure to maternal cigarette smoke during pregnancy and environmental smoke thereafter. This study provides plausibility for the biological hazards of maternal cigarette smoking upon the developing brainstem and SIDS risk, and underscores the public health message against smoking during pregnancy to reduce this risk.

Acknowledgments

The authors thank Dr. Kathryn G. Commons for assistance in the use of nAChR immunocytochemistry, Dr. Felicia L. Trachtenberg for statistical advice, and Mr. Richard A. Belliveau for technical assistance in this study. We are grateful to Dr. Henry F. Krous, Ms. Elizabeth Haas, and the Office of the Chief Medical Examiner, San Diego, CA, for assistance in the accrual of infant brainstem specimens.

Support: Supported by the First Candle/SIDS Alliance, CJ Martin Fellowship (Nation Health and Medical Research Council of Australia), National Institute of Child Health and Development (R37-HD20991 and PO1-HD045991), and Developmental Disabilities Research Center, Children's Hospital Boston (P30-HD18655).

References

- Adams CE, Broide RS, Chen Y, Winzer-Serhan UH, Henderson TA, Leslie FM, Freedman R. Development of the $\alpha 7$ nicotinic cholinergic receptor in rat hippocampal formation. *Brain Res Dev Brain Res* 2002;139(2):175–187.
- Aliyu MH, Salihu HM, Wilson RE, Kirby RS. Prenatal smoking and risk of intrapartum stillbirth. *Arch Environ Occup Health* 2007;62(2):87–92. [PubMed: 18316266]
- Baker KG, Halliday GM, Hornung JP, Geffen LB, Cotton RG, Tork I. Distribution, morphology and number of monoamine-synthesizing and substance P-containing neurons in the human dorsal raphe nucleus. *Neuroscience* 1991;42(3):757–775. [PubMed: 1720227]
- Benarroch EE, Schmeichel AM, Low PA, Parisi JE. Depletion of putative chemosensitive respiratory neurons in the ventral medullary surface in multiple system atrophy. *Brain* 2007;130(Pt 2):469–475. [PubMed: 17235127]
- Benwell ME, Balfour DJ, Anderson JM. Evidence that tobacco smoking increases the density of (-)-[3H] nicotine binding sites in human brain. *J Neurochem* 1988;50(4):1243–1247. [PubMed: 3346676]
- Blair PS, Fleming PJ, Bensley D, Smith I, Bacon C, Taylor E, Berry J, Golding J, Tripp J. Smoking and the sudden infant death syndrome: results from 1993–5 case-control study for confidential inquiry into stillbirths and deaths in infancy. Confidential Enquiry into Stillbirths and Deaths Regional Coordinators and Researchers. *BMJ* 1996;313(7051):195–198. [PubMed: 8696194]
- Blair PS, Sidebotham P, Berry PJ, Evans M, Fleming PJ. Major epidemiological changes in sudden infant death syndrome: a 20-year population-based study in the UK. *Lancet* 2006;367(9507):314–319. [PubMed: 16443038]
- Bradley SR, Pieribone VA, Wang W, Severson CA, Jacobs RA, Richerson GB. Chemosensitive serotonergic neurons are closely associated with large medullary arteries. *Nat Neurosci* 2002;5(5):401–402. [PubMed: 11967547]
- Brown JW, Sirlin EA, Benoit AM, Hoffman JM, Darnall RA. Activation of 5-HT_{1A} receptors in medullary raphe disrupts sleep and decreases shivering during cooling in the conscious piglet. *Am J Physiol Regul Integr Comp Physiol* 2008;294(3):R884–894. [PubMed: 18094064]
- Centeno ML, Luo J, Lindstrom JM, Caba M, Pau KY. Expression of $\alpha 4$ and $\alpha 7$ nicotinic receptors in the brainstem of female rabbits after coitus. *Brain Res* 2004;1012(1–2):1–12. [PubMed: 15158155]

- Chiolero A, Bovet P, Paccaud F. Association between maternal smoking and low birth weight in Switzerland: the EDEN study. *Swiss Med Wkly* 2005;135(35–36):525–530. [PubMed: 16323070]
- Cimino M, Marini P, Colombo S, Andena M, Cattabeni F, Fornasari D, Clementi F. Expression of neuronal acetylcholine nicotinic receptor alpha 4 and beta 2 subunits during postnatal development of the rat brain. *J Neural Transm Gen Sect* 1995;100(2):77–92. [PubMed: 8962687]
- Cohen G, Han ZY, Grailhe R, Gallego J, Gaultier C, Changeux JP, Lagercrantz H. beta 2 nicotinic acetylcholine receptor subunit modulates protective responses to stress: A receptor basis for sleep-disordered breathing after nicotine exposure. *Proc Natl Acad Sci USA* 2002;99(20):13272–13277. [PubMed: 12228730]
- Court JA, Perry EK, Spurdin D, Griffiths M, Kerwin JM, Morris CM, Johnson M, Oakley AE, Birdsall NJ, Clementi F, Perry RH. The role of the cholinergic system in the development of the human cerebellum. *Brain Res Dev Brain Res* 1995;90(1–2):159–167.
- Cucchiario G, Commons KG. Alpha 4 nicotinic acetylcholine receptor subunit links cholinergic to brainstem monoaminergic neurotransmission. *Synapse* 2003;49(3):195–205. [PubMed: 12774304]
- Darnall RA, Harris MB, Gill WH, Hoffman JM, Brown JW, Niblock MM. Inhibition of serotonergic neurons in the nucleus paragigantocellularis lateralis fragments sleep and decreases rapid eye movement sleep in the piglet: implications for sudden infant death syndrome. *J Neurosci* 2005;25(36):8322–8332. [PubMed: 16148240]
- Dev NB, Loeschcke HH. A cholinergic mechanism involved in the respiratory chemosensitivity of the medulla oblongata in the cat. *Pflugers Arch* 1979;379(1):29–36. [PubMed: 571102]
- Didier M, Berman SA, Lindstrom J, Bursztajn S. Characterization of nicotinic acetylcholine receptors expressed in primary cultures of cerebellar granule cells. *Brain Res Mol Brain Res* 1995;30(1):17–28. [PubMed: 7609640]
- Dominguez del Toro E, Juiz JM, Peng X, Lindstrom J, Criado M. Immunocytochemical localization of the alpha 7 subunit of the nicotinic acetylcholine receptor in the rat central nervous system. *J Comp Neurol* 1994;349(3):325–342. [PubMed: 7852628]
- Duncan JR, Randall LL, Belliveau RA, Trachtenberg FL, Randall B, Habbe D, Mandell F, Welty TK, Iyasu S, Kinney HC. The effect of maternal smoking and drinking during pregnancy upon (3)h-nicotine receptor brainstem binding in infants dying of the sudden infant death syndrome: initial observations in a high risk population. *Brain Pathol* 2008;18(1):21–31. [PubMed: 17924983]
- Dwyer JB, Broide RS, Leslie FM. Nicotine and brain development. *Birth Defects Res C Embryo Today* 2008;84(1):30–44. [PubMed: 18383130]
- Ferreira M, Singh A, Dretchen KL, Kellar KJ, Gillis RA. Brainstem nicotinic receptor subtypes that influence intragastric and arterial blood pressures. *J Pharmacol Exp Ther* 2000;294(1):230–238. [PubMed: 10871317]
- Filiano JJ, Choi JC, Kinney HC. Candidate cell populations for respiratory chemosensitive fields in the human infant medulla. *J Comp Neurol* 1990;293(3):448–465. [PubMed: 2324324]
- Flores CM, Rogers SW, Pabreza LA, Wolfe BB, Kellar KJ. A subtype of nicotinic cholinergic receptor in rat brain is composed of alpha 4 and beta 2 subunits and is up-regulated by chronic nicotine treatment. *Mol Pharmacol* 1992;41(1):31–37. [PubMed: 1732720]
- Fonck C, Cohen BN, Nashmi R, Whiteaker P, Wagenaar DA, Rodrigues-Pinguet N, Deshpande P, McKinney S, Kwok S, Munoz J, Labarca C, Collins AC, Marks MJ, Lester HA. Novel seizure phenotype and sleep disruptions in knock-in mice with hypersensitive alpha 4* nicotinic receptors. *J Neurosci* 2005;25(49):11396–11411. [PubMed: 16339034]
- Franco P, Groswasser J, Hassid S, Lanquart JP, Scaillet S, Kahn A. Prenatal exposure to cigarette smoking is associated with a decrease in arousal in infants. *J Pediatr* 1999;135(1):34–38. [PubMed: 10393601]
- Frank MG, Srere H, Ledezma C, O'Hara B, Heller HC. Prenatal nicotine alters vigilance states and AchR gene expression in the neonatal rat: implications for SIDS. *Am J Physiol Regul Integr Comp Physiol* 2001;280(4):R1134–1140. [PubMed: 11247836]
- Gilchrist D, Woods B, Binns CW, Scott JA, Gracey M, Smith H. Aboriginal mothers, breastfeeding and smoking. *Aust NZ J Public Health* 2004;28(3):225–228.
- Good CH, Bay KD, Buchanan RA, McKeon KA, Skinner RD, Garcia-Rill E. Prenatal exposure to cigarette smoke affects the physiology of pedunculopontine nucleus (PPN) neurons in development. *Neurotoxicol Teratol* 2006;28(2):210–219. [PubMed: 16469482]

- Gotti C, Zoli M, Clementi F. Brain nicotinic acetylcholine receptors: native subtypes and their relevance. *Trends Pharmacol Sci* 2006;27(9):482–491. [PubMed: 16876883]
- Haan EA, Jennings IG, Cuello AC, Nakata H, Fujisawa H, Chow CW, Kushinsky R, Brittingham J, Cotton RG. Identification of serotonergic neurons in human brain by a monoclonal antibody binding to all three aromatic amino acid hydroxylases. *Brain Res* 1987;426(1):19–27. [PubMed: 2891407]
- Hatori E, Sakuraba S, Kashiwagi M, Kuribayashi J, Tsujita M, Hosokawa Y, Takeda J, Kuwana S. Association of nicotinic acetylcholine receptors with central respiratory control in isolated brainstem-spinal cord preparation of neonatal rats. *Biol Res* 2006;39(2):321–330. [PubMed: 16874407]
- Hellstrom-Lindahl E, Court JA. Nicotinic acetylcholine receptors during prenatal development and brain pathology in human aging. *Behav Brain Res* 2000;113(1–2):159–168. [PubMed: 10942042]
- Hellstrom-Lindahl E, Gorbounova O, Seiger A, Mousavi M, Nordberg A. Regional distribution of nicotinic receptors during prenatal development of human brain and spinal cord. *Brain Res Dev Brain Res* 1998;108(1–2):147–160.
- Hohmann CF, Brooks AR, Coyle JT. Neonatal lesions of the basal forebrain cholinergic neurons result in abnormal cortical development. *Brain Res* 1988;470(2):253–264. [PubMed: 3219585]
- Horne RS, Ferens D, Watts AM, Vitkovic J, Lacey B, Andrew S, Cranage SM, Chau B, Greaves R, Adamson TM. Effects of maternal tobacco smoking, sleeping position, and sleep state on arousal in healthy term infants. *Arch Dis Child Fetal Neonatal Ed* 2002;87(2):F100–105. [PubMed: 12193515]
- Ito C, Fukuda A, Nabekura J, Oomura Y. Acetylcholine causes nicotinic depolarization in rat dorsal motor nucleus of the vagus, in vitro. *Brain Res* 1989;503(1):44–48. [PubMed: 2575433]
- Iyasu S, Randall LL, Welty TK, Hsia J, Kinney HC, Mandell F, McClain M, Randall B, Habbe D, Wilson H, Willinger M. Risk factors for sudden infant death syndrome among northern plains Indians. *JAMA* 2002;288(21):2717–2723. [PubMed: 12460095]
- Jensen AA, Frolund B, Liljefors T, Krogsgaard-Larsen P. Neuronal nicotinic acetylcholine receptors: structural revelations, target identifications, and therapeutic inspirations. *J Med Chem* 2005;48(15):4705–4745. [PubMed: 16033252]
- Jones IW, Wonnacott S. Why doesn't nicotinic ACh receptor immunoreactivity knock out? *Trends Neurosci* 2005;28(7):343–345. [PubMed: 15979499]
- Julvez J, Ribas-Fito N, Torrent M, Fornis M, Garcia-Esteban R, Sunyer J. Maternal smoking habits and cognitive development of children at age 4 years in a population-based birth cohort. *Int J Epidemiol* 2007;36(4):825–832. [PubMed: 17550944]
- Kahn A. Infant sudden death. *J Pharm Belg* 1992;47(1):33–37. [PubMed: 1625161]
- Kahn A, Groswasser J, Sottiaux M, Kelmanson I, Rebuffat E, Franco P, Dramaix M, Wayenberg JL. Prenatal exposure to cigarettes in infants with obstructive sleep apneas. *Pediatrics* 1994;93(5):778–783. [PubMed: 8165078]
- Kato I, Franco P, Groswasser J, Scaillet S, Kelmanson I, Togari H, Kahn A. Incomplete arousal processes in infants who were victims of sudden death. *Am J Respir Crit Care Med* 2003;168(11):1298–1303. [PubMed: 12917226]
- Kenny PJ, File SE, Rattray M. Nicotine regulates 5-HT(1A) receptor gene expression in the cerebral cortex and dorsal hippocampus. *Eur J Neurosci* 2001;13(6):1267–1271. [PubMed: 11285026]
- Kinney HC. Abnormalities of the brainstem serotonergic system in the sudden infant death syndrome: a review. *Pediatr Dev Pathol* 2005;8(5):507–524. [PubMed: 16222475]
- Kinney HC, Belliveau RA, Trachtenberg FL, Rava LA, Paterson DS. The development of the medullary serotonergic system in early human life. *Auton Neurosci* 2007;132(1–2):81–102. [PubMed: 17236817]
- Kinney HC, Filiano JJ, Sleeper LA, Mandell F, Valdes-Dapena M, White WF. Decreased muscarinic receptor binding in the arcuate nucleus in sudden infant death syndrome. *Science* 1995;269(5229):1446–1450. [PubMed: 7660131]
- Kinney HC, Filiano JJ, White WF. Medullary serotonergic network deficiency in the sudden infant death syndrome: review of a 15-year study of a single dataset. *J Neuropathol Exp Neurol* 2001;60(3):228–247. [PubMed: 11245208]
- Kinney HC, O'Donnell TJ, Kriger P, White WF. Early developmental changes in [3H]nicotine binding in the human brainstem. *Neuroscience* 1993;55(4):1127–1138. [PubMed: 8232902]

- Kinney HC, Randall LL, Sleeper LA, Willinger M, Belliveau RA, Zec N, Rava LA, Dominici L, Iyasu S, Randall B, Habbe D, Wilson H, Mandell F, McClain M, Welty TK. Serotonergic brainstem abnormalities in Northern Plains Indians with the sudden infant death syndrome. *J Neuropathol Exp Neurol* 2003;62(11):1178–1191. [PubMed: 14656075]
- Kubin L, Fenik V. Pontine cholinergic mechanisms and their impact on respiratory regulation. *Respir Physiol Neurobiol* 2004;143(2–3):235–249. [PubMed: 15519558]
- Landsberg RL, Awatramani RB, Hunter NL, Farago AF, DiPietrantonio HJ, Rodriguez CI, Dymecki SM. Hindbrain rhombic lip is comprised of discrete progenitor cell populations allocated by Pax6. *Neuron* 2005;48(6):933–947. [PubMed: 16364898]
- Lena C, Popa D, Grailhe R, Escourrou P, Changeux JP, Adrien J. Beta2-containing nicotinic receptors contribute to the organization of sleep and regulate putative micro-arousals in mice. *J Neurosci* 2004;24(25):5711–5718. [PubMed: 15215293]
- Lewis KW, Bosque EM. Deficient hypoxia awakening response in infants of smoking mothers: possible relationship to sudden infant death syndrome. *J Pediatr* 1995;127(5):691–699. [PubMed: 7472818]
- Lucas-Meunier E, Fossier P, Baux G, Amar M. Cholinergic modulation of the cortical neuronal network. *Pflugers Arch* 2003;446(1):17–29. [PubMed: 12690458]
- Luck W, Nau H, Hansen R, Steldinger R. Extent of nicotine and cotinine transfer to the human fetus, placenta and amniotic fluid of smoking mothers. *Dev Pharmacol Ther* 1985;8(6):384–395. [PubMed: 4075937]
- Mallard C, Tolcos M, Leditschke J, Campbell P, Rees S. Reduction in choline acetyltransferase immunoreactivity but not muscarinic-m2 receptor immunoreactivity in the brainstem of SIDS infants. *J Neuropathol Exp Neurol* 1999;58(3):255–264. [PubMed: 10197817]
- McKay BE, Placzek AN, Dani JA. Regulation of synaptic transmission and plasticity by neuronal nicotinic acetylcholine receptors. *Biochem Pharmacol* 2007;74(8):1120–1133. [PubMed: 17689497]
- Mihailescu S, Palomero-Rivero M, Meade-Huerta P, Maza-Flores A, Drucker-Colin R. Effects of nicotine and mecamylamine on rat dorsal raphe neurons. *Eur J Pharmacol* 1998;360(1):31–36. [PubMed: 9845269]
- Moser N, Mechawar N, Jones I, Gochberg-Sarver A, Orr-Urtreger A, Plomann M, Salas R, Molles B, Marubio L, Roth U, Maskos U, Winzer-Serhan U, Bourgeois JP, Le Sourd AM, De Biasi M, Schroder H, Lindstrom J, Maelicke A, Changeux JP, Wevers A. Evaluating the suitability of nicotinic acetylcholine receptor antibodies for standard immunodetection procedures. *J Neurochem* 2007;102(2):479–492. [PubMed: 17419810]
- Nabekura J, Ueno S, Ogawa T, Akaike N. Colocalization of ATP and nicotinic ACh receptors in the identified vagal preganglionic neurone of rat. *J Physiol* 1995;489(Pt 2):519–527. [PubMed: 8847644]
- Nachmanoff DB, Panigrahy A, Filiano JJ, Mandell F, Sleeper LA, Valdes-Dapena M, Krous HF, White WF, Kinney HC. Brainstem 3H-nicotine receptor binding in the sudden infant death syndrome. *J Neuropathol Exp Neurol* 1998;57(11):1018–1025. [PubMed: 9825938]
- Olszewski, J.; Baxter, D. *Atlas of the human brainstem*. Karger; Basel: 1982.
- Opanashuk LA, Pauly JR, Hauser KF. Effect of nicotine on cerebellar granule neuron development. *Eur J Neurosci* 2001;13(1):48–56. [PubMed: 11135003]
- Padley JR, Kumar NN, Li Q, Nguyen TB, Pilowsky PM, Goodchild AK. Central command regulation of circulatory function mediated by descending pontine cholinergic inputs to sympathoexcitatory rostral ventrolateral medulla neurons. *Circ Res* 2007;100(2):284–291. [PubMed: 17204655]
- Panigrahy A, Filiano J, Sleeper LA, Mandell F, Valdes-Dapena M, Krous HF, Rava LA, Foley E, White WF, Kinney HC. Decreased serotonergic receptor binding in rhombic lip-derived regions of the medulla oblongata in the sudden infant death syndrome. *J Neuropathol Exp Neurol* 2000;59(5):377–384. [PubMed: 10888367]
- Panigrahy A, Filiano JJ, Sleeper LA, Mandell F, Valdes-Dapena M, Krous HF, Rava LA, White WF, Kinney HC. Decreased kainate receptor binding in the arcuate nucleus of the sudden infant death syndrome. *J Neuropathol Exp Neurol* 1997;56(11):1253–1261. [PubMed: 9370236]
- Parslow PM, Cranage SM, Adamson TM, Harding R, Horne RS. Arousal and ventilatory responses to hypoxia in sleeping infants: effects of maternal smoking. *Respir Physiol Neurobiol* 2004;140(1):77–87. [PubMed: 15109930]

- Paterson DS, Belliveau RA, Trachtenberg F, Kinney HC. Differential development of 5-HT receptor and the serotonin transporter binding in the human infant medulla. *J Comp Neurol* 2004a;472(2):221–231. [PubMed: 15048689]
- Paterson DS, Thompson EG, Kinney HC. Serotonergic and glutamatergic neurons at the ventral medullary surface of the human infant: Observations relevant to central chemosensitivity in early human life. *Auton Neurosci* 2006a;124(1–2):112–124. [PubMed: 16458076]
- Paterson DS, Trachtenberg FL, Thompson EG, Belliveau RA, Beggs AH, Darnall R, Chadwick AE, Krous HF, Kinney HC. Multiple serotonergic brainstem abnormalities in sudden infant death syndrome. *JAMA* 2006b;296(17):2124–2132. [PubMed: 17077377]
- Paterson, S.; Thompson, E.; Kinney, H. Neurons In the rostral ventrolateral medulla of the human infant express neurochemical markers consistent with preBötzinger complex neurons. Society for Neuroscience Annual Meeting; October 23–27; San Diego CA. 2004b.
- Paxinos, G.; Huang, X-F. Atlas of the human brainstem. San Diego: Academic Press, Inc; 1995.
- Raatikainen K, Huurinen P, Heinonen S. Smoking in early gestation or through pregnancy: a decision crucial to pregnancy outcome. *Prev Med* 2007;44(1):59–63. [PubMed: 16959307]
- Rakic, P.; Sidman, RL. Development of the human central nervous system. In: Haymaker, W.; Adams, RD., editors. *Histology and histopathology of the nervous system*. Springfield, Illinois: Charles C. Thomas; 1982. p. 3–145.
- Richerson GB, Wang W, Tiwari J, Bradley SR. Chemosensitivity of serotonergic neurons in the rostral ventral medulla. *Respir Physiol* 2001;129(1–2):175–189. [PubMed: 11738653]
- Robertson RT, Gallardo KA, Claytor KJ, Ha DH, Ku KH, Yu BP, Lauterborn JC, Wiley RG, Yu J, Gall CM, Leslie FM. Neonatal treatment with 192 IgG-saporin produces long-term forebrain cholinergic deficits and reduces dendritic branching and spine density of neocortical pyramidal neurons. *Cereb Cortex* 1998;8(2):142–155. [PubMed: 9542893]
- Rukhadze I, Kubin L. Mesopontine cholinergic projections to the hypoglossal motor nucleus. *Neurosci Lett* 2007;413(2):121–125. [PubMed: 17174027]
- Saper CB, Scammell TE, Lu J. Hypothalamic regulation of sleep and circadian rhythms. *Nature* 2005;437(7063):1257–1263. [PubMed: 16251950]
- Schechtman VL, Harper RM, Wilson AJ, Southall DP. Sleep state organization in normal infants and victims of the sudden infant death syndrome. *Pediatrics* 1992;89(5 Pt 1):865–870. [PubMed: 1579396]
- Schneider AS, Atluri P, Shen Q, Barnes W, Mah SJ, Stadfelt D, Goderie SK, Temple S, Fleck MW. Functional nicotinic acetylcholine receptor expression on stem and progenitor cells of the early embryonic nervous system. *Ann NY Acad Sci* 2002;971:135–138. [PubMed: 12438103]
- Shao XM, Feldman JL. Pharmacology of nicotinic receptors in preBotzinger complex that mediate modulation of respiratory pattern. *J Neurophysiol* 2002;88(4):1851–1858. [PubMed: 12364511]
- Shao XM, Tan W, Xiu J, Puskar N, Fonck C, Lester HA, Feldman JL. Alpha4* nicotinic receptors in preBotzinger complex mediate cholinergic/nicotinic modulation of respiratory rhythm. *J Neurosci* 2008;28(2):519–528. [PubMed: 18184794]
- Siegel, J. *The Neural Control of Sleep and Waking*. New York: Springer-Verlag; 2002.
- Sinton CM, McCarley RW. Neurophysiological mechanisms of sleep and wakefulness: a question of balance. *Semin Neurol* 2004;24(3):211–223. [PubMed: 15449215]
- Slotkin TA, MacKillop EA, Rudder CL, Ryde IT, Tate CA, Seidler FJ. Permanent, sex-selective effects of prenatal or adolescent nicotine exposure, separately or sequentially, in rat brain regions: indices of cholinergic and serotonergic synaptic function, cell signaling, and neural cell number and size at 6 months of age. *Neuropsychopharm* 2007a;32(5):1082–1097.
- Slotkin TA, Ryde IT, Tate CA, Seidler FJ. Lasting effects of nicotine treatment and withdrawal on serotonergic systems and cell signaling in rat brain regions: separate or sequential exposure during fetal development and adulthood. *Brain Res Bull* 2007b;73(4–6):259–272. [PubMed: 17562392]
- Slotkin TA, Seidler FJ, Qiao D, Aldridge JE, Tate CA, Cousins MM, Proskocil BJ, Sekhon HS, Clark JA, Lupo SL, Spindel ER. Effects of prenatal nicotine exposure on primate brain development and attempted amelioration with supplemental choline or vitamin C: neurotransmitter receptors, cell signaling and cell development biomarkers in fetal brain regions of rhesus monkeys. *Neuropsychopharmacology* 2005;30(1):129–144. [PubMed: 15316571]

- Swanson LW, Simmons DM, Whiting PJ, Lindstrom J. Immunohistochemical localization of neuronal nicotinic receptors in the rodent central nervous system. *J Neurosci* 1987;7(10):3334–3342. [PubMed: 2822866]
- Torraso AS, Britto LR. Neurotransmitter regulation of neural development: acetylcholine and nicotinic receptors. *An Acad Bras Cienc* 2002;74(3):453–461. [PubMed: 12378313]
- Touiki K, Rat P, Molimard R, Chait A, de Beaurepaire R. Effects of tobacco and cigarette smoke extracts on serotonergic raphe neurons in the rat. *Neuroreport* 2007;18(9):925–929. [PubMed: 17515803]
- Tribollet E, Bertrand D, Marguerat A, Raggenbass M. Comparative distribution of nicotinic receptor subtypes during development, adulthood and aging: an autoradiographic study in the rat brain. *Neuroscience* 2004;124(2):405–420. [PubMed: 14980390]
- Urbano FJ, Simpson JI, Llinas RR. Somatomotor and oculomotor inferior olivary neurons have distinct electrophysiological phenotypes. *Proc Natl Acad Sci U S A* 2006;103(44):16550–16555. [PubMed: 17050678]
- Wada E, Wada K, Boulter J, Deneris E, Heinemann S, Patrick J, Swanson LW. Distribution of alpha 2, alpha 3, alpha 4, and beta 2 neuronal nicotinic receptor subunit mRNAs in the central nervous system: a hybridization histochemical study in the rat. *J Comp Neurol* 1989;284(2):314–335. [PubMed: 2754038]
- Wang J, Irnaten M, Neff RA, Venkatesan P, Evans C, Loewy AD, Mettenleiter TC, Mendelowitz D. Synaptic and neurotransmitter activation of cardiac vagal neurons in the nucleus ambiguus. *Ann NY Acad Sci* 2001;940:237–246. [PubMed: 11458681]
- Whiteaker P, Cooper JF, Salminen O, Marks MJ, McClure-Begley TD, Brown RW, Collins AC, Lindstrom JM. Immunolabeling demonstrates the interdependence of mouse brain alpha4 and beta2 nicotinic acetylcholine receptor subunit expression. *J Comp Neurol* 2006;499(6):1016–1038. [PubMed: 17072836]
- Whiting PJ, Lindstrom JM. Characterization of bovine and human neuronal nicotinic acetylcholine receptors using monoclonal antibodies. *J Neurosci* 1988;8(9):3395–3404. [PubMed: 3171681]
- Willinger M, James LS, Catz C. Defining the sudden infant death syndrome (SIDS): deliberations of an expert panel convened by the National Institute of Child Health and Human Development. *Pediatr Pathol* 1991;11(5):677–684. [PubMed: 1745639]
- Woolf NJ, Butcher LL. Cholinergic systems in the rat brain: IV. Descending projections of the pontomesencephalic tegmentum. *Brain Res Bull* 1989;23(6):519–540. [PubMed: 2611694]
- Zec N, Filiano JJ, Panigrahy A, White WF, Kinney HC. Developmental changes in [3H]lysergic acid diethylamide ([3H]LSD) binding to serotonin receptors in the human brainstem. *J Neuropathol Exp Neurol* 1996;55(1):114–126. [PubMed: 8558166]
- Zhuang J, Xu F, Frazier DT. Hyperventilation evoked by activation of the vicinity of the caudal inferior olivary nucleus depends on the fastigial nucleus in anesthetized rats. *J Appl Physiol* 2008;104(5):1351–1358. [PubMed: 18356483]

Figure 1. $\alpha 4$ immunoreactivity in the medulla at 15 gestational weeks

(A) Immunostaining for the $\alpha 4$ nAChR showed ubiquitous distribution. (B) A subpopulation of rhombic lip cells with differentiating cytoplasm were intensely immunoreactive for $\alpha 4$, as well as (C), cells with trailing processes migrating away from this region. (D) Diagrammatic representation of the human medulla at 15 weeks gestation. (E, F) $\alpha 4$ immunopositive punctuate fibers/axons in the neuropil. HG, hypoglossal nucleus; ION, inferior olive; PCA, postconceptional age; RO, raphé obscurus. Scales A: 255 μm ; B: 41 μm ; C 6.2 μm , E: 26 μm ; F: 12 μm .

Figure 2. $\alpha 4$ immunoreactivity in the medulla at term and infancy

Nicotinic receptor immunoreactivity for $\alpha 4$ was ubiquitously expressed in all nuclei, including the (A) raphé obscurus; (C) dorsal motor nucleus of vagus (DMX), hypoglossal nucleus (HG); (D) arcuate nucleus and (E) inferior olive (ION). (B) A diagram of the human medulla at term is provided for orientation. PCA, postconceptional age. Scales A: 30 μm ; C: 65 μm ; D: 93 μm ; E: 103 μm .

Figure 3. Comparisons of $\alpha 4$ nAChR immunoreactivity in the medulla at 23 and 44 weeks post conception
 $\alpha 4$ immunoreactive cells at 23 weeks (A, C) and 44 weeks (B, D) in the arcuate nucleus (A, B) and inferior olive (ION) (C, D). In both nuclei there is a visually obvious increase in immunoreactivity in the cell soma over the last half of gestation. PCA, postconceptional age. Scales A, B: 93 μ m; C: 104 μ m; D: 87 μ m.

Figure 4. Colocalization of $\alpha 4$ and 5-HT neurons in the medulla

(A, D) $\alpha 4$ immunopositive cells (red) and (B, E) PH8 immunopositive (5-HT) neurons (green) in the raphe obscurus at 15 weeks (A-C) and at term (D-F). (C, F) Colocalization of expression is demonstrated by merging of images (yellow). A subpopulation of neurons expressing $\alpha 4$ which were not serotonergic, yet are observed in association to 5-HT cells (F, arrow). PCA, postconceptional age. Scales: A-C: 12 μm ; D-F: 20 μm .

Figure 5. Colocalization of $\alpha 4$ and 5-HT neurons in the raphe obscurus

(A) $\alpha 4$ immunopositive cells (red) and (B) PH8 immunopositive (5-HT) neurons (green) in the raphe obscurus at 10 postnatal months. (C) Colocalization of all visible 5-HT neurons is demonstrated by merging of the image (yellow). A subset of non-5-HT neurons also express $\alpha 4$ in this region. PCA, postconceptional age. Scales: A-C: 100 μ m.

Figure 6. Colocalization of $\alpha 4$ and 5-HT neurons in the medulla

(A, D) $\alpha 4$ immunopositive neurons (red) and (B, E) PH8 immunopositive (5-HT) neurons (green) in the paragigantocellularis lateralis (PGCL) at 10 months postnatal (A-C) and arcuate nucleus at term (D-F), with colocalization demonstrated by merging of the images (yellow) (C, F). A subpopulation of neurons expressing $\alpha 4$ were not serotonergic, yet were in close proximity to 5-HT cells (C). At high magnification the $\alpha 4$ immunostaining is punctate, consistent with receptors (C). Scales: D-F: 8.4 μm ; A-C: 42 μm .

Figure 7. nAChR binding in the infant medulla

Pseudocolored autoradiographic images of ³H-nicotine (A), ³H-epibatidine (B), ³H-cytisine (C) and ¹²⁵I-bgt (D) binding in the infant medulla (mid-level) with specific activity scales and graphs of binding in select nuclei for each radioligand where binding levels are presented for each ligand as a percent of binding in the inferior olive nucleus (ION) arbitrarily defined as 100%. Differences in binding patterns are based on the affinity of each ligand to specific nAChR subunits. The highest density of binding with ³H-epibatidine was seen in the inferior olive ION (B). The highest binding density for ¹²⁵I-bgt was in the dorsal motor nucleus of vagus (DMX) (D). ARC, arcuate nucleus; HG, hypoglossal nucleus; NTS, nucleus of the solitary tract; PGCL, paragigantocellularis lateralis; RO, raphe obscurus.

Figure 8. Development profiles of nAChR binding in the dorsal motor nucleus in the infant medulla Receptor binding for both ^3H -epibatidine (A) and ^{125}I -bgt (B) significantly decreased ($p<0.05$) over infancy in the dorsal motor nucleus (DMX). PCA, postconceptional age.

Figure 9. Developmental profile of nAChR and 5-HT receptor binding in the paragigantiocellulus lateralis

There was a significant reduction ($p < 0.05$) in $\alpha 7$ ($^{125}\text{I-bgt}$) and a significant increase in 5-HT_{1A} ($^3\text{H-8-OH-DPAT}$) receptor binding over infancy in the paragigantiocellulus lateralis (PGCL). PCA, postconceptional age.

Figure 10. Comparative development sequences of nAChRs and 5-HT markers in the human medulla

Profiles for the nAChR and 5-HT systems in the (A) paragigantocellularis lateralis; (B) arcuate nucleus; and (C) nucleus of the solitary tract show that dynamic changes occur in different nuclei and across development. – no change; ▸ decrease; ◀ increase. * fetal 5-HT neuronal densities from Kinney et al., 2007. NA; data not available for receptor and transporter binding in the fetal period. This diagram highlights the complexity of the developmental changes in the nicotinic and 5-HT systems in the human medulla relative to one another.

Table 1
Antibodies used to assess the developmental profile of nAChRs in the medulla oblongata of the human fetus and infant

Antibody	Host	Source/cat number/lot	Immunogen	Dilution	Peptide sequence	Labels
$\alpha 4$ nAChR	Guinea pig polyclonal IgG	Chemicon/AB5590/0608038451	Synthetic peptide corresponding to AA 568–588 of the rat nAChR	1:200	AA 568–588 of rat nAChR	Nerve fibers and cell bodies corresponding to <i>in situ</i> pattern using rat nAChR probes (Wada et al., 1989)
$\alpha 4$ nAChR (MAB-299)	Rat monoclonal IgG1	Covance/MRT-613R/147939002	Purified rat brain nAChR	1:500	* Characterized by Whiting and Lindstrom, 1988 * Common epitope of tryptophan hydroxylase (TPOH), tyrosine hydroxylase and phenylalanine hydroxylase.	Binds to the extracellular surface of $\alpha 4$ nAChR subunits (Whiting and Lindstrom, 1988)
PH8 for tryptophan hydroxylase, key biosynthetic enzyme	Murine monoclonal IgG	Chemicon/MAB5278/16401065	Immunoglobulin fraction from murine ascites fluid	1:1000		In fixed human tissue PH8 binds specifically to TPOH due to antigenic changes induced by fixation (Haan et al., 1987; Baker et al., 1991; Kinney et al., 2007)

Table 2

Autoradiographic procedures for nAChRs

Ligand	Preferential Binding	Supplier/ Catalogue/Lot	Concentration	Incubation time/temp	Displacer	Detection Method
³ H-Nicotine (Reference for comparison radioligand Kinney et al., 1993; Nachmanoff et al., 1998; Duncan et al., 2008)	All nAChR subunits (Jensen et al., 2005)	PerkinElmer/NE T827/3529280	3.5 nM (79.5 Ci/mmol)	25 min/RT	10 µM L-nicotine hydrogen	³ H sensitive film (LKB Ultrafilm): 12 weeks
³ H-Epipibatidine	α2–4, β2, β4 (Jensen et al., 2005)	PerkinElmer/NE T1102/3557074	0.5 nM (56 Ci/mmol)	60 min/RT	300 µM L-nicotine hydrogen	phosphoimage plates, BAS TR2025 system: 3 weeks
³ H-Cytisine	α4β2 subtype (Jensen et al., 2005)	PerkinElmer/NE T1504/3559630	2nM (27 Ci/mmol)	60 min/RT	100 µM L-nicotine hydrogen	phosphoimage plates, BAS TR2025 system: 6 weeks
¹²⁵ I-Bungarotoxin	α7 (Jensen et al., 2005)	PerkinElmer/NE X126/BE31670	1.5 nM (120 Ci/mmol)	120 min/4°C	1 mM L-nicotine hydrogen	¹²⁵ I-sensitive film (Kodak BioMax MR): 5 days

Table 3

Nicotinic receptor binding in the human infant medulla (fmol/mg tissue)

Nuclei	³ H-nicotine (all nAChRs) (n=5)	³ H-epibatidine ($\alpha 2-4$, $\beta 2, \beta 4$) (n=5)	³ H-cytisine ($\alpha 4 \beta 2$) (n=2)	¹²⁵ I-bgt ($\alpha 7$) (n=5)
HG	9.90 $\pm$ 1.00 *	25.65 $\pm$ 4.35	5.20	2.25 $\pm$ 0.52
NTS	7.50 $\pm$ 1.00 *	43.22 $\pm$ 4.67	2.61	5.44 $\pm$ 0.46
DMX	9.27 $\pm$ 1.23 *	41.63 $\pm$ 4.91	2.58	10.79 $\pm$ 0.87
MAO	17.00 $\pm$ 1.50 *	35.14 $\pm$ 3.77	7.05	3.12 $\pm$ 0.69
DAO	15.2 $\pm$ 1.70 *	40.02 $\pm$ 6.70	10.65	4.51 $\pm$ 1.08
ION	23.98 $\pm$ 1.56	75.08 $\pm$ 5.89	33.02	3.39 $\pm$ 1.00
IRZ	10.88 $\pm$ 1.31	26.63 $\pm$ 4.01	5.34	4.14 $\pm$ 1.30
GC	9.70 $\pm$ 1.11	25.57 $\pm$ 5.44	6.94	4.10 $\pm$ 1.21
PGCL	8.77 $\pm$ 1.08	24.56 $\pm$ 5.04	6.23	3.34 $\pm$ 1.25
RO	4.60 $\pm$ 0.69	14.25 $\pm$ 2.34	3.20	2.60 $\pm$ 0.84
ARC	No detectable binding *	16.21 $\pm$ 2.20	3.43	2.11 $\pm$ 0.26

Values are expressed as means $\pm$ SEM and are relative to their own radioligand.

* Data for certain nuclei for ³H-nicotine binding have been previously published (Nachmanoff et al., 1998). Due to tissue constraints, only 2 cases were available for ³H-cytisine analysis and are used for reference.

Abbreviations: HG, hypoglossal nucleus; NTS, nucleus of the solitary tract; DMX, dorsal motor nucleus; MAO, medial accessory olive; DAO, dorsal accessory olive; ION, inferior olive; IRZ, intermediate reticular zone; GC, gigantocellularis; PGCL, paragigantocellularis lateralis; RO, raphé obscurus; ARC, arcuate nucleus.

Summary of observations of nAChR and 5-HT markers in the human infant medulla

Table 4

Source	$\alpha 4$ protein expression: single label	Immunohistochemistry $\alpha 4$ nAChR & 5-HT co-localization: double label	^3H -nicotine (all nAChRs)	^3H -epibatidine ($\alpha 2\text{-}4$, $\beta 2$, $\beta 4$)	^3H -cytisine ($\alpha 4\beta 2$)	Receptor Binding ^{125}I -bgt ($\alpha 7$)	^3H -8-OH-DPAT (5-HT_{1A})	^{125}I -RTL-55 (5-HTT)
RO	Present	Present	+	+	+	+++	+++	+++
GC	Present	Present	++	++	+	+++++	++	+++
PGCL	Present	Present	++	++	+	++++	++	++
ARC	Present	Present	0	+	+	++++	+	+
HG	Present	No 5-HT cells	++	++	+	++++	+	+++
NTS	Present	No 5-HT cells	++	+++	+	++++	++	++
DMX	Present	No 5-HT cells	++	+++	+	++++	Not measured	Not measured
ION	Present	No 5-HT cells	+++	++++	++++	++++	+	Not measured

Key: Scale as a percent of the ION (nAChRs) or RO (5-HT markers [Paterson et al., 2004a; Paterson et al., 2006b]): 0: no binding; 1–25% low binding +; 26–50% moderate binding ++; 51–75% high binding +++; 76–100% very high binding ++++; above 101% intense binding +++++.

Abbreviations: RO, raphé obscurus; GC, gigantocellularis; PGCL, paragigantocellularis lateralis; ARC, arcuate nucleus; HG, hypoglossal nucleus; NTS, nucleus of the solitary tract; DMX, dorsal motor nucleus; ION, inferior olive.

Table 5

Comparison of developmental changes in binding to 5-HT_{1A}, 5-HTT, 5-HT neuronal density and nAChRs with increasing age in the human infant medulla

Nuclei	5-HT _{1A} *	5-HTT *	5-HT neuronal density *	Epibatidine	Bungarotoxin	Nicotine
PGCL			No change	No change	Decrease (0.01)	No change
HG	Increase (0.04)	No change	No neurons	No change	No change	* Not reported
NTS	Increase (0.003)	No change	No neurons	No change	Decrease (0.02)	* Not reported
ARC	No change	No change	No change	No change	Decrease (0.002)	No change
DAO	No change	No change	No neurons	Decrease (0.02)	No change	* Not reported
DMX	Not measured	Not measured	No neurons	Decrease (0.01)	Decrease (0.03)	No change
IRZ	No change	No change	No change	No change	No change	No change
GC	No change	No change	No change	No change	No change	No change
RO	No change	No change	No change	No change	No change	No change
ION	No change	Not measured	No neurons	No change	No change	No change
MAO	No change	No change	No neurons	No change	No change	* Not reported

* The 5-HT-related data have been previously published and are shown here for comparison (Paterson et al., 2004a; Paterson et al., 2006b), ³H-nicotine data have been previously published in select nuclei (Nachmanoff et al., 1998).

Abbreviations: PGCL, paraventricular nucleus; HG, hypothalamic nucleus; NTS, nucleus of the solitary tract; ARC, arcuate nucleus; DAO, dorsal accessory olive; DMX, dorsal motor nucleus; IRZ, intermediate reticular zone; GC, gigantocellularis; RO, raphe obscurus; ION, inferior olive; MAO, medial accessory olive.