Published in final edited form as:

J Electrocardiol. 2009 ; 42(6): 511–516. doi:10.1016/j.jelectrocard.2009.06.008.

The many faces of repolarization instability: which one is prognostic?

Vladimir Shusterman, MD, PhD^{¶, *}, Rachel Lampert, MD[§], and Barry London, MD, PhD[¶]

[¶] University of Pittsburgh, Pittsburgh, PA

^{*} PinMed, Inc., Pittsburgh, PA

[§] Yale University, New Haven, CT

Abstract

Instabilities of the STT segment's magnitude, and in particular the 0.5 beat/cycle oscillations (T-wave alternans, TWA), have been linked to the heightened risk of ventricular tachyarrhythmias (VTA) and sudden cardiac death (SCD). During the last decade theoretical, experimental and clinical research efforts have focused primarily on TWA, examining its mechanisms and predictive value using time-invariant cutoff values. However, recent evidence suggests that such a single-snapshot test of a single-frequency (TWA) oscillation using a constant cutoff value might be suboptimal for risk stratification, due to several reasons.

First, it is well-known that the risk of VTA/SCD evolves over time with changes in electrophysiological substrate, environmental and physiological triggers, and the impact of other physiological (e.g., circadian) rhythmicity. Hence, the outcome of TWA testing might depend on the time of day, as Holter-based TWA studies have demonstrated. Furthermore, currently employed single-snapshot testing with a binary cutoff value may not coincide with the periods of heightened risk for VTA/SCD and may not yield prognostic information, as a recent TWA sub-study of the SCD-HeFT trial has showed. Second, the analysis focused on TWA alone ignores the existence of multiple (alternating and nonalternating) forms of repolarization instability that have been shown to arise or increase prior to the onset of VTA/SCD.

Summarizing, recent studies have identified multiple forms of repolarization instabilities modulated by distinct mechanisms, which might have different prognostic values. Therefore, the assessment of TWA needs to be dynamic and personalized to take into account the time evolution of risk and individual history.

Keywords

Repolarization instability; T-wave alternans; ventricular tachyarrhythmias; sudden cardiac death

Introduction

Beat-to-beat alterations of the electrocardiographic repolarization segment have long been recognized as a marker of increased risk for ventricular tachyarrhythmias and sudden cardiac

Address for correspondence: Vladimir Shusterman, 200 Lothrop Street, Room B535, Pittsburgh, PA 15213, shustermanv@upmc.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

death.¹ During the last decade, remarkable progress has been made in this knowledge area. A number of theoretical, experimental, and clinical studies have provided a wealth of evidence linking the 0.5 cycle/beat oscillations in the amplitude of the STT segment, referred to as the T-wave alternans (TWA), to the heightened risk of VTA.^{2,3} Moreover, the repolarization instability has been shown to have a direct pathophysiological connection with the initiation of arrhythmias.⁴ As a result of these multidisciplinary research efforts, T-wave alternans testing has become a clinical tool for arrhythmia risk stratification.⁵ Recently, an increase in T-wave alternans has been reported preceding the imminent onset of VTA in several animal models⁶ and clinical studies conducted in patients with structural heart disease, suggesting that the utility of TWA might be further extended to serve as a short-term predictor of these life-threatening events.^{7,8}

However, recent evidence suggests that currently employed single-snapshot testing of a single-frequency (0.5 cycle/beat) oscillation using a one-size-fits-all cutoff value might be suboptimal for risk stratification, due to several reasons. First, the underlying assumption that both T-wave alternans and the risk of arrhythmias do not change over time, is oversimplified.^{9,10} T-wave alternans has been shown to exhibit circadian changes, peaking in the morning and declining at night.¹¹ Thus, the outcome of TWA testing is likely to be different if the testing is done at different times of day. Furthermore, the risk of arrhythmias and sudden death also evolves over longer time scales as well, following the changes in the underlying electrophysiological substrate. In particular, a recent SCD-HeFT trial has failed to show that T-wave alternans predicts arrhythmic events or mortality in patients with heart failure (n=490, ischemic heart disease in 49% of subjects, left ventricular ejection fraction of 25%), but showed that its prognostic significance gradually evolves over 15–30 month time scale with the progression of changes in the underlying heart disease.^{9,10} This provides further support for the notion that currently employed single-snapshot testing may not coincide with the periods of heightened risk for VTA/SCD and may not yield prognostic information. In addition, the analysis focused on TWA alone ignores the existence of multiple forms of repolarization instability that have been shown to arise or increase prior to the onset of VTA/SCD both in patients with cardiomyopathy⁷ and long-QT syndrome.¹² Analysis of such instabilities may provide complementary information that may help to improve the risk assessment with respect to arrhythmias and sudden death.

To clarify the predictive value of T-wave alternans and other forms of repolarization instability, one needs to determine the physiological covariates and their respective weights, modulating the relationship between the instability and arrhythmogenesis (Figure 1).

The restitution covariate

The restitution curve, i.e. the relationship between the cardiac electrical Action-Potential Duration (APD) and a preceding Diastolic Interval (DI) has been introduced by Nolasco and Dahlen.¹³ In their seminal study conducted in the frog ventricular muscle strips over 40 years ago, the researchers noticed striking similarities between the cardiac APD/DI relationship and a simple electronic amplifier with a negative feedback. In such an electronic system, if G is a gain function, I is an input, F is a fraction of an output O , and X is an independent signal, then the output is $O = G(I)$ and $I = X - F(O)$. Similarly, the transfer functions for the cardiac tissue paced at the cycle length (CL) can be expressed as: $APD_1 = f(DI_0)$ and $DI_1 = f(APD_1) = CL_1 - APD_1$.¹³

Assuming that an APD depends only on the preceding DI (a “no-memory” assumption), the relationship between the two variables can be expressed by a non-linear curve resembling a logarithmic function. If the pacing cycle length (CL) is kept constant, then $d(DI_1)/d(APD) = -1$, and a simple 1-dimensional, iterative map, referred to as the cobweb, can be used to examine

this nonlinear system's dynamics. It is not difficult to show that the slope of the curve defining the relationship between APD and DI determines the stability of this dynamical system.¹³ In particular, if the slope is between 0 and 1, the system is stable, and a small perturbation δ would produce damped oscillations (transients) that will return to the equilibrium (steady state). In contrast, if the slope is >1 , a small perturbation would lead to an ever-growing oscillation (under-damping). And if the slope of the APD curve is exactly equal to 1, then the system is bi-stable, producing sustained, period-2 oscillations (i.e. alternans).^{13,14}

This method provides an intuitive tool for tracking the nonlinear dynamics of the cardiac electrical activity at the expense of several important simplifications. First, it assumes that an APD depends only on a single DI, which is not realistic for cardiac myocytes. It has been shown by several investigators that an APD strongly depends on a much longer (at least 60-sec) history of CL, which can be better approximated by one or two exponential functions.^{15,16} Second, the simplified, restitution-only approach also ignores the inverse relationship between conduction velocity and cycle length, which plays a crucial role in cardiac physiology and may become compromised in heart failure.^{17,18} Finally, this method disregards the impact of Ca^{2+} handling and several other important covariates (discussed later in this article), and as expected, several recent studies have showed that this theoretical construct, taken alone, fails to predict T-wave alternans or arrhythmias in clinical settings.¹⁹ Nevertheless, the restitution-based approach has proved useful in several settings, where fast pacing protocols could be employed to achieve a relatively short CL (Figure 2A), while all other covariates are kept constant.^{13,20,21}

A recent analysis of the spatial distribution of the restitution curves in different regions of the human heart has showed a wide spectrum of results.²² This spatial heterogeneity suggests that the analysis needs to be extended from a single restitution curve to the entire spatial distribution of the curves in different regions of the heart. Furthermore, the restitution-based approach can also be extended from a “memoryless” APD, which depends on a single DI, to an APD, which has “memory” of a longer history of cardiac beats.²³ Yet, the recently proposed theoretical constructs incorporating such a memory still could not provide an accurate approximation of the real-life dynamical behavior of the cardiac electrophysiological parameters.²³ This implicitly suggests the importance of other physiological covariates that remain unaccounted for by the restitution-only based approach.

The calcium covariate

The Ca^{2+} -related mechanisms play a pivotal role in the excitation-contraction coupling, and L-type Ca^{2+} channels also determine a key ionic current that is active during the plateau phase of cardiac repolarization. In the course of the development of heart failure, calcium handling becomes compromised, including decreased peak systolic Ca^{2+} , elevated diastolic Ca^{2+} , and prolongation of the Ca^{2+} transient.¹⁸ In humans with heart failure, these changes are associated with the decreased expression of the sarco(endo)plasmic reticulum Ca^{2+} -ATPase (SERCA), decreases in phospholamban, and increased expression of the Na/Ca exchanger.²⁴ The resulting prolongation of the Ca^{2+} transient and intracellular Ca^{2+} overload can lead to premature Ca^{2+} release from the sarcoplasmic reticulum and delayed afterdepolarizations.¹⁸ As a result, the Ca^{2+} alternans can often emerge at near-normal heart rates, giving rise to APD-alternans and T-wave alternans (Figure 2B). In particular, our group has demonstrated pronounced alternans (alternation of up to 40% of the peak amplitude of the Ca^{2+} transient) in a TNF- α genetic mouse model of heart failure, using optical mapping of Ca^{2+} in isolated, Langendorff-perfused hearts.¹⁸ In this model, the Ca^{2+} -alternans was found at relatively long pacing cycle lengths. Furthermore, changes in the Ca^{2+} concentration affected arrhythmia inducibility in this genetic mouse model, confirming the essential role of abnormal calcium handling in the mechanism of arrhythmogenesis.¹⁸ Similarly, Wilson et al. has recently reported that Ca^{2+}

alternans was the driving force generating action-potential alternans at slow heart rates in a canine heart-failure wedge preparation.²⁵

Of note, repolarization alternans associated with Ca^{2+} alternans has been also observed in pharmacologically induced Long-QT syndrome in arterially perfused canine left-ventricular wedge preparations (Figure 2C).²⁶

The adrenergic covariate

It has long been recognized that sympathetic nervous system activity modulates the magnitude of repolarization instability. In a canine model of ischemia, Nearing and Verrier have shown that modifications of the stellate ganglion activity directly affect the level of T-wave alternans.²⁷ In contrast, beta-adrenergic blockade has been shown to suppress the magnitude of T-wave alternans in humans with ischemic cardiomyopathy and left ventricular ejection fraction of < 40%.²⁸ Further support for the role of autonomic activity in the emergence of T-wave alternans has been provided by the observations of circadian changes in the magnitude of T-wave alternans in Holter recordings obtained from patients with a prior myocardial infarction (Figure 2F).¹¹ Mental stress in humans with implantable devices has been shown to exacerbate the level of T-wave alternans, and the magnitude of the increase in T-wave alternans was strongly correlated with the level of norepinephrine and the incidence of arrhythmic events in a two-year follow-up (Figure 2E).^{29,30,31}

The mechanisms by which adrenergic stimulation promotes repolarization alternans may involve additional influx of Ca^{2+} into the myocytes, overloading the calcium-handling cascades, which are chronically impaired in heart failure.^{32,33} In particular, it is likely that Ca^{2+} -alternans and impaired Ca^{2+} handling are the driving forces behind the emergence of T-wave alternans with β -adrenergic stimulation in our studies of the TNF- α genetic mouse model of heart failure, since Ca^{2+} -alternans play a pivotal role in the mechanisms of repolarization instability and arrhythmias accompanying development of heart failure in these animals.³⁴

The conduction velocity covariate

The inverse relationship between the conduction velocity and basic cycle length is well documented.¹⁷ This relationship plays a particularly important role at short CL, and conduction-velocity alternans have been observed at faster heart rates, during various types of a narrow-QRS tachycardia.³⁵ Preferential conduction slowing at short pacing cycle lengths (CL < 80 ms) has been also observed in a TNF- α genetic mouse model of heart failure, suggesting that impaired adaptation of the conduction system to faster heart rates may contribute to the emergence of alternans in the setting of heart failure (Figure 2D).¹⁸

Other physiological covariates

In addition to the L-type Ca^{2+} channels, an incomplete recovery, temporal instability, or heterogeneous distribution of other ionic currents, which are active during repolarization, may also lead to the emergence of repolarization instability. Indeed, a number of intracellular processes, including modifications of repolarizing potassium currents (I_{Kr}) and changes in excitability, have been implicated in the mechanisms of repolarization instability.^{36,37,38} We note also that functional changes in the repolarizing ionic channels, including diminished repolarizing K^+ currents (I_{to} and I_{K1}), have been documented in heart failure, suggesting possible involvement of these channels (in addition to the Ca^{2+} -related mechanisms discussed earlier) in the emergence of instabilities in this setting.¹⁸

Adding to the complexity of the “repolarization-instability” conundrum, the alternans has been also reported in association with pharmacologically induced Brugada syndrome in the right

ventricular canine tissue (Figure 2G). In contrast to the other settings described above, the alternans occurred at slow pacing rates and disappeared when the pacing accelerated. The presumable mechanism of this instability includes the alternating I_{Na} causing alternations in I_{to} and L-type Ca^{2+} currents in the right ventricular epicardium.³⁹

Discerning the impact of physiological covariates in real-life data

Summarizing, the results of clinical, experimental, and theoretical studies have provided a wealth of evidence supporting statistical and mechanistic links between the occurrence of repolarization instability and arrhythmias. The notion that the measurement of repolarization instability may provide important diagnostic and prognostic information for the arrhythmia risk assessment has become well recognized.^{1,2,5,7,8,9,11,31} Several clinical trials have confirmed the predictive value of T-wave alternans,^{1,2,11,31} while other trials have failed to demonstrate a strong predictive value.¹⁰ The inconsistencies most likely arise from the differences in the underlying physiological mechanisms driving the instabilities in different settings, leading to a common end-point from a multitude of sub-cellular, intra-cellular, and multi-cellular processes (Figure 1). Thus, a better understanding of these physiological covariates, along with the measurement caveats,⁴⁰ is needed for improving the accuracy of arrhythmia risk assessment.

In a real-life setting, electrophysiologic data are usually collected at the time when at least some of the covariates exhibit simultaneous changes. Thus, the algorithms are needed to differentiate the respective impacts and weights of each participating covariate. Figure 2 shows one possible algorithm that allows, to a first approximation, determine the relative weights of the primary physiological covariates. We view the flow chart in Figure 2 as a working hypothesis guiding our ongoing studies. More experimental and clinical data will be needed to determine the quantitative weight of each covariate and refine the probabilities of transition from one mechanism to another. We note, however, that the flow chart might fail when all covariates have similar weights, making it difficult to discern predominant, major effects. The practical implementation and testing of this algorithm will require continuous or serial tracking of individual repolarization dynamics, along with other electrophysiological parameters, including changes in heart rate, QRS duration, circadian variability, physical activity and psychological status.^{11,31} The improvement can also be achieved by tracking the changes against the individual's baseline and using "personalized" (i.e. individually tailored) pattern recognition analysis.^{7,40,41}

Acknowledgments

The project was supported in part by National Institutes of Health National Heart, Lung, and Blood Institute grants R44HL077116 (Dr. Shusterman) and R01 HL62300 (Dr. London), and American Heart Association Established Investigator Award (Dr. London).

References

1. Rosenbaum DS, Jackson LE, Smith JM, Garan H, Ruskin JN, Cohen RJ. Electrical alternans and vulnerability to ventricular arrhythmias. *N Engl J Med* 1994;330:235–241. [PubMed: 8272084]
2. Rosenbaum DS, Albrecht P, Cohen RJ. Predicting sudden cardiac death from T wave alternans of the surface electrocardiogram: Promise and Pitfalls. *J Cardiovasc Electrophysiol* 1996;7:1095–1111. [PubMed: 8930743]
3. Jordan PN, Christini DJ. Characterizing the contribution of voltage- and calcium-dependent coupling to action potential stability: implications for repolarization alternans. *Am J Physiol Heart Circ Physiol* 2007;293:H2109–118. [PubMed: 17586611]
4. Armoundas AA, Tomaselli GF, Esperer HD. Pathophysiological basis and clinical application of T-wave alternans. *J Am Coll Cardiol* 2002;40:207–217. [PubMed: 12106921]

5. Costantini O, Hohnloser SH, Kirk MM, Lerman BB, Baker JH, Sethuraman B, Dettmer MM, Rosenbaum DS. ABCD Trial Investigators. The ABCD (Alternans Before Cardioverter Defibrillator) Trial: strategies using T-wave alternans to improve efficiency of sudden cardiac death prevention. *J Am Coll Cardiol* 2009;53(6):471–9. [PubMed: 19195603]
6. Nearing BD, Verrier RL. Progressive increases in complexity of T-wave oscillations herald ischemia-induced ventricular fibrillation. *Circ Res* 2002;91(8):727–32. [PubMed: 12386150]
7. Shusterman V, Goldberg A, London B. Upsurge in T-wave alternans and nonalternating repolarization instability precedes spontaneous initiation of ventricular tachyarrhythmias in humans. *Circulation* 2006;113(25):2880–7. [PubMed: 16785339]
8. Swerdlow CD, Zhou X, Voroshilovsky O, Abeyaratne A, Gillberg J. High amplitude T-wave alternans precedes spontaneous ventricular tachycardia or fibrillation in ICD electrograms. *Heart Rhythm* 2008;5(5):670–6. [PubMed: 18452868]
9. Rosenbaum DS. T-Wave Alternans in the Sudden Cardiac Death in Heart Failure Trial Population: Signal or Noise? *Circulation* 2008;118:2015–2018. [PubMed: 19001031]
10. Gold MR, Ip JH, Costantini O, Poole JE, McNulty S, Mark DB, Lee KL, Bardy GH. Role of Microvolt T-Wave Alternans in Assessment of Arrhythmia Vulnerability Among Patients With Heart Failure and Systolic Dysfunction: Primary Results From the T-Wave Alternans Sudden Cardiac Death in Heart Failure Trial Substudy. *Circulation* 2008;118:2022–2028. [PubMed: 18955671]
11. Verrier RL, Nearing BD, La Rovere MT, Pinna GD, Mittleman MA, Bigger JT Jr, Schwartz PJ. ATRAMI Investigators. Ambulatory electrocardiogram-based tracking of T wave alternans in postmyocardial infarction patients to assess risk of cardiac arrest or arrhythmic death. *J Cardiovasc Electrophysiol* 2003;14(7):705–11. [PubMed: 12930249]
12. Nemec J, Buncová M, Shusterman V, Winter B, Shen WK, Ackerman MJ. QT interval variability and adaptation to heart rate changes in patients with long QT syndrome. *Pacing Clin Electrophysiol* 2009;32(1):72–81. [PubMed: 19140916]
13. Nolasco JB, Dahlen RW. A graphic method for the study of alternation in cardiac action potentials. *J Appl Physiol* 1968;25:191–196. [PubMed: 5666097]
14. Weiss JN, Karma A, Shiferaw Y, Chen PS, Garfinkel AZ. Qu. From Pulsus to Pulseless: The Saga of Cardiac Alternans. *Circ Res* 2006;98:1244–1253. [PubMed: 16728670]
15. Elharrar V, Atarashi H, Surawicz B. Cycle length-dependent action potential duration in canine cardiac Purkinje fibers. *Am J Physiol* 1984;247(6 Pt 2):H936–45. [PubMed: 6507640]
16. Lux RL, Hilbel T, Brockmeier K. Electrocardiographic measures of repolarization revisited: why? what? how? *J Electrocardiol* 2001;34:259–264. [PubMed: 11781965]
17. Cherry EM, Fenton FH. Suppression of alternans and conduction blocks despite steep APD restitution: electrotonic, memory, and conduction velocity restitution effects. *Am J Physiol Heart Circ Physiol* 2004;286:2332–2341.
18. London B, Baker LC, Lee JS, Shusterman V, Choi BR, Kubota T, McTiernan CF, Feldman AM, Salama G. Calcium-dependent arrhythmias in transgenic mice with heart failure. *Am J Physiol Heart Circ Physiol* 2003;284(2):H431–41. [PubMed: 12388316]
19. Narayan SM, Franz MR, Lalani G, Kim J, Sastry A. T-wave alternans, restitution of human action potential duration, and outcome. *J Am Coll Cardiol* 2007;18;50(25):2385–92. [PubMed: 18154963]
20. Garfinkel A, Kim YH, Voroshilovsky O, Qu Z, Kil JR, Lee MH, Karagueuzian HS, Weiss JN, Chen PS. Preventing ventricular fibrillation by flattening cardiac restitution. *PNAS* 2000;97(11):6061–6066. [PubMed: 10811880]
21. Koller ML, Maier SKG, Gelzer AR, Bauer WR, Meesmann M, Gilmour RF. Altered dynamics of action potential restitution and alternans in humans with structural heart disease. *Circulation* 2005;112:1542–1548. [PubMed: 16157783]
22. Nash MP, Bradley CP, Sutton PM, Clayton RH, Kallis P, Hayward MP, Paterson DJ, Taggart P. Whole heart action potential duration restitution properties in cardiac patients: a combined clinical and modelling study. *Exp Physiol* 2006;91.2:339–354. [PubMed: 16452121]
23. Kalb SS, Dobrovolny HM, Tolkacheva EG, Idriss SF, Krassowska W, Gauthier DJ. The restitution portrait: a new method for investigating rate-dependent restitution. *J Cardiovasc Electrophysiol* 2004;15(6):698–709. [PubMed: 15175067]

24. Arai M, Alpert NR, MacLennan DH, Barton P, Periasamy M. Alterations in sarcoplasmic reticulum gene expression in human heart failure: a possible mechanism for alterations in systolic and diastolic properties of the failing myocardium. *Circ Res* 1993;72:463–469. [PubMed: 8418995]
25. Wilson LD, Jeyaraj D, Wan X, Hoeker GS, Said TH, Gittinger M, Laurita KR, Rosenbaum DS. Heart failure enhances susceptibility to arrhythmogenic cardiac alternans. *Heart Rhythm* 2009;6:251–259. [PubMed: 19187920]
26. Shimizu W, Antzelevitch C. Cellular and Ionic Basis for T-Wave Alternans Under Long-QT Conditions. *Circulation* 1999;99:1499–1507. [PubMed: 10086976]
27. Nearing BD, Huang AH, Verrier RL. Dynamic tracking of cardiac vulnerability by complex demodulation of the T wave. *Science* 1991;252:437–440. [PubMed: 2017682]
28. Rashba EJ, Cooklin M, MacMurdy K, Kavesh N, Kirk M, Sarang S, Peters RW, Shorofsky SR, Gold MR. Effects of Selective Autonomic Blockade on T-Wave Alternans in Humans. *Circulation* 2002;105:837–842. [PubMed: 11854124]
29. Kop WJ, Krantz DS, Nearing BD, Gottdiener JS, Quigley JF, O'Callahan M, DelNegro AA, Friehling TD, Karasik P, Suchday S, Levine J, Verrier RL. Effects of acute mental stress and exercise on t-wave alternans in patients with implantable cardioverter defibrillators and controls. *Circulation* 2004;109(15):1864–1869. [PubMed: 15037524]
30. Lampert R, Shusterman V, Burg MM, Lee FA, Earley C, Goldberg A, McPherson CA, Batsford WP, Soufer R. Effects of psychologic stress on repolarization and relationship to autonomic and hemodynamic factors. *J Cardiovasc Electrophysiol* 2005;16(4):372–7. [PubMed: 15828878]
31. Lampert R, Shusterman V, Burg M, McPherson C, Batsford W, Goldberg A, Soufer R. Anger-induced T-Wave alternans predicts future ventricular arrhythmias in patients with implantable cardioverter-defibrillators. *J Am Coll Cardiol* 2009;53:774–8. [PubMed: 19245968]
32. Jin YT, Hasebe N, Matsusaka T, Natori S, Ohta T, Tsuji S, Kikuchi K. Magnesium attenuates isoproterenol-induced acute cardiac dysfunction and β -adrenergic desensitization. *Am J Physiol Heart Circ Physiol* 2007;292:H1593–H1599. [PubMed: 17114241]
33. Chatelain P, Laruel A, Beaufort P, Meysmans L, Clinet M. Prevention of calcium overload and down-regulation of calcium channels in rat heart by SR 33557, a novel calcium entry blocker. *Cardioscience* 1992;3:117–123. [PubMed: 1322733]
34. Shusterman V, Goldberg A, London B. Adrenergic stimulation promotes T-wave alternans in a TNF- α genetic mouse model of congestive heart failure. *Heart Rhythm* 2005;2:S142–143.
35. Morady F, DiCarlo LA Jr, Baerman JM, de Buitelir M, Kou WH. Determinants of QRS alternans during narrow QRS tachycardia. *J Am Coll Cardiol* 1987 Mar;9(3):489–99. [PubMed: 3819196]
36. Hua F, Gilmour RF. Contribution of I_{K_r} to rate-dependent action potential dynamics in canine endocardium. *Circ Res* 2004;94:810–819. [PubMed: 14963001]
37. O'Rourke B, Ramza BM, Marban E. Oscillations of membrane current and excitability driven by metabolic oscillations in heart cells. *Science* 1994;265:962–966. [PubMed: 8052856]
38. Armondas AA, Tomaselli GF, Esperer HD. Pathophysiological basis and clinical application of T-wave alternans. *J Am Coll Cardiol* 2002;40:207–217. [PubMed: 12106921]
39. Morita H, Zipes DP, Lopshire J, Morita ST, Wu J. T wave alternans in an in vitro canine tissue model of Brugada syndrome. *Am J Physiol Heart Circ Physiol* 2006;291:421–428.
40. Shusterman V, Goldberg A. Tracking repolarization dynamics in real-life data. *J Electrocardiol* 2004;37 (Suppl):180–6. [PubMed: 15534838]
41. Shusterman V, Aysin B, Anderson KP, Beigel A. Multidimensional rhythm disturbances as a precursor of sustained ventricular tachyarrhythmias. *Circulation Research* 2001;88:705–712. [PubMed: 11304493]

Figure 1.
Factors affecting repolarization instability

Figure 2.
Block-diagram of an algorithm differentiating the origins of repolarization instability