Published in final edited form as:

Ann N Y Acad Sci. 2009 July ; 1170: 11–17. doi:10.1111/j.1749-6632.2009.04369.x.

WNT5a in tongue and fungiform papilla development: distinct roles compared with WNT10b and other morphogenetic proteins

Hong-Xiang Liu¹, Ann M. Staubach Grosse², Katherine D. Walton², Daniel A. Saims¹, Deborah L. Gumucio², and Charlotte M. Mistretta^{1,*}

¹School of Dentistry, University of Michigan, Ann Arbor, USA

²Medical School, University of Michigan, Ann Arbor, USA

Abstract

Fungiform papillae are complex taste organs that develop in a pattern on anterior tongue in rodent embryos. Several intrinsic secreted molecules are important for papilla development and patterning, including SHH, BMPs, Noggin, EGF and WNTs. Recent data about roles of WNTs in regulation of tongue and fungiform papilla development lead to new insights about the importance of tissue and timing contexts when studying effects of morphogenetic proteins. WNT/ β -catenin signaling is required for formation of fungiform papillae, but not for determining tongue size and shape. In contrast, WNT5a apparently is important for tongue outgrowth, but not papilla development. Preliminary data from WNT5a mutant mice separate genetic programs for papilla number from those for tongue shape and size.

Keywords

Tongue; taste papilla; development; WNT; SHH; BMP; Noggin; EGF

Fungiform papillae are taste organs with a patterned distribution on the anterior tongue and contain one taste bud each in rodents[1,2]. These papillae are highly complex sensory organs. At maturity they include: an epithelial covering with a taste bud in the apical papilla epithelium and a taste bud-free lateral papilla epithelium; a mesenchymal core of connective tissue underlying the epithelium; and, an innervation from the chorda tympani nerve (to taste bud) and the lingual branch of the trigeminal nerve (to epithelium surrounding the taste bud and papilla walls)[2,3]. Because of the complexity of these sensory organs, many tissue specific contexts must be considered when studying regulation of fungiform papilla development. In turn, tissues will be at different morphological stages during embryonic and postnatal development. To fully understand regulatory mechanisms and signaling factors in fungiform papillae, tissue and temporal variables must be well defined.

In rodents, the fungiform papilla develops from a homogeneous epithelium of the initial lingual swellings to a papilla placode or thickened epithelial cell cluster on the spatulate tongue, to a well defined papilla (Fig. 1A)[1,2]. There are many stage-specific factors that are prominent in fungiform papilla formation: 1) formation of the tongue; 2) development of lingual epithelium and mesenchyme; 3) placode formation and invagination; 4) formation of the early papilla; 5) differentiation of papilla epithelium; 6) mesenchymal core development; 7) differentiation of inter-papilla epithelium; and, 8) innervation to papilla and taste bud.

*Corresponding author: 1011 N. University Ave, School of Dentistry, University of Michigan, Ann Arbor, MI 48109-1078; Email: chmist@umich.edu; Tel: +1-734 647 3911; Fax: +1-734 647 2110..

Of the many tissue and temporal factors, it has been demonstrated that intact sensory innervation is not required for tongue and papilla development[4,5]. In organ culture that excludes intact sensory ganglia, the tongue forms from mandibular swellings; and, papilla placodes develop from a homogeneous tongue epithelium and progress to form papillae that mirror those in the embryo in number, temporal and spatial patterns and basic molecular markers[5,6,7]. Patterning and essential molecular phenotypes are, therefore, intrinsic to lingual tissues.

Intrinsic secreted factors in fungiform papilla development

In the past decade the *in vitro* tongue culture system[4] and loss- and gain-of-function mouse models have been increasingly used to contribute to our understanding of papilla development. Numerous secreted regulatory molecules have been identified to be important for regulating tongue and papilla development, e.g., sonic hedgehog (SHH)[5,7,8], bone morphogenetic proteins (BMPs)[9], the BMP antagonist Noggin[9], epidermal growth factor (EGF)[10] and WNTs[11,12,13].

Regulatory roles for these molecules are complex and are concentration- and stage-dependent, summarized in Fig. 1B. For example, in E14+2 day tongue cultures, exogenous SHH increases papillae at low concentration but reduces papillae at high concentration (Fig. 1B, SHH)[14]. BMP2, 4 and 7 each has dual stage-specific effects on fungiform papilla formation in tongue culture, i.e., increases (E13) or reduces (E14) papilla number with BMP added to culture medium or via beads (Fig. 1B, BMPs)[9]. The BMP antagonist, Noggin, increases papilla number at both E13 and E14 whereas exogenous Follistatin is ineffective in culture (Fig. 1B, Noggin, Follistatin)[9]. EGF reduces papilla number both at E13 and at E14 (Fig. 1B, EGF)[10]. In contrast to the secreted factors SHH, BMPs and EGF, that effect a decrease in fungiform papilla formation, activation of WNT canonical signaling with LiCl increases papillae at E12.5 in mouse (Fig. 1B, WNT/LiCl)[11,12,13], whereas signaling via WNT5a does not alter papilla number (Fig. 1B, WNT5a).

Organ cultures treated with antagonists or inhibitors illustrate the nature and extent of signaling effects for SHH, BMP and EGF (Fig. 2). Exogenous SHH, BMP and EGF each decreases papilla number compared to control cultures in standard medium. Blocking SHH signaling with cyclopamine (Fig. 2, SHH and CYCL) increases fungiform papilla number by more than double that in control organ culture and induces papilla formation on the papilla-free intermolar eminence[5,7]. When BMP signaling is antagonized with Noggin treatment (Fig. 2, BMP2, 4, 7 and Noggin), papilla number is increased by 60% compared to control cultures and large clustered papillae form[9]. Similarly, C56 (Fig. 2, EGF and C56), an EGF receptor (EGFR) inhibitor, leads to about 40% increase of fungiform papilla number compared to cultures with standard medium[10]. We know that the EGF protein is found throughout the E13-E18 tongue epithelium whereas EGFR undergoes a progressive restriction to the epithelium *between* fungiform papillae[10]. Thus, EGF regulation of papilla formation signals through EGFR expressed in inter-papilla epithelium and results in an increase in cell proliferation between papillae[10].

In short, to build this complex taste organ — the fungiform papilla — critical tissue and temporal contexts are imperative to the study of regulatory factors. To form and pattern the papillae, the inter-papilla lingual tissue also is an essential part of the puzzle. At any given time and lingual location, variations of cell fate determination, proliferation and differentiation will contribute to cellular responsiveness. Secreted regulatory factors do not act in isolation but signaling pathways interact. Recent data about roles of WNTs in regulation of fungiform papillae lead to new insights about the importance of tissue and timing contexts in tongue and fungiform papilla development.

WNT/ β -catenin signaling is required for formation of fungiform papillae, but not for tongue size and shape

Three groups have reported that canonical WNT signaling is required for fungiform papilla formation[11,12,13]. Essential components for canonical WNT signaling, TOPGAL [a β -galactosidase gene driven by a T cell factor (TCF) β -catenin responsive promoter] and the transcription factor Lef1, have restricted expression in developing fungiform papillae (Fig. 3). Deletion of either β -catenin[13] or Lef1[12] leads to a striking loss of papillae, demonstrating the requirement of canonical WNT signaling for fungiform papilla development (Fig. 3). To activate canonical WNT signaling pathways, WNT10b may be an important endogenous agonist. The expression of WNT10b is restricted in developing fungiform papillae in parallel with TOPGAL and Lef1[12]. In E14.5 WNT10b null mutants, fungiform papillae are reduced in size compared to wild type (Fig. 3). Thus, WNT10b is proposed to act via canonical signaling pathways to regulate fungiform papilla development[12].

It is noteworthy that in postnatal day 1 Lef1 $-/-$ tongue, where fungiform papillae are “atrophied” or missing, tissue positions for papillae are maintained (Fig. 3). This suggests that there are gene programs specifically for fungiform papilla formation, separate from that for other lingual tissue development. This is supported by new data on tongue and fungiform papilla development in Wnt5a null mutants.

WNT5a is important for tongue outgrowth, but not for papilla development

In contrast to WNT10b, WNT5a has a unique expression pattern in vertebrate embryos. Perinatal lethal WNT5a null mutant embryos have a striking phenotype with truncated face, limbs and tail demonstrating the importance of WNT5a in proximal-distal outgrowth in vertebrate embryos[15]. Recently, there is increased interest in roles of WNT5a in several aspects of embryonic development, e.g., cell polarity and directional movement[16], airway and vascular tubulogenesis[17], and mammary gland development[18].

Our laboratory has performed gene array analysis in E14+2 day tongue cultures. Multiple gene expression patterns were compared between anterior tongue where fungiform papillae are abundantly distributed and the intermolar eminence which is papilla-free. We found that WNT5a expression is increased by 8 fold in anterior tongue compared to intermolar eminence. In our preliminary data, Western blot analysis revealed that WNT5a is expressed in embryonic (E13-E16), but not in postnatal tongue.

To learn if WNT5a has a role in development of tongue and taste papillae, in preliminary experiments we have compared embryonic tongues in WNT5a $-/-$ mice with those of wild type littermates (Fig. 4). At E16, length of oral tongue in WNT5a $-/-$ mice was reduced to 65% of wild type length. Portions of the tongue that were anterior, or posterior, to the anterior-most border of the intermolar eminence were reduced by similar proportions. However, pharyngeal tongue in null mice was reduced to only 40% of wild-type. Whereas length of oral tongue was severely compromised in null mice, width was similar to that in wild type. Overall, anterior tongue area was substantially reduced and shape was radically altered. However, considering the much truncated anterior tongue, preliminary analysis indicates that numbers of fungiform papillae were not different on mutant tongues relative to wild type. The single circumvallate papilla on posterior tongue also was sustained without an obvious alteration in WNT5a mutant tongues at E16 (Fig. 4).

In summary, our preliminary data indicate that WNT5a is important for embryonic tongue outgrowth. This is distinct from roles for WNT10b and other regulatory factors in

development of fungiform papilla numbers and pattern. Even with a much reduced anterior tongue, numbers of fungiform papillae are not different on mutant tongues relative to wild type. Therefore, the WNT5a preliminary data support the novel idea that spatial constraints may not be a major factor in fungiform papilla formation. The data provide evidence of separate genetic programs for papilla number from those for tongue shape and size.

Acknowledgments

The research was supported by NIDCD, NIH Grant DC 000456 (CMM) and NIDDK, NIH Grant DK065850 (DLG).

References

1. Mistretta, CM. Topographical and histological study of the developing rat tongue, palate, and taste buds. In: Bosma, JF., editor. *The Third Symposium on Oral Sensation and Perception: The Mouth of the Infant*. Springfield: Thomas; 1972. p. 163-187.
2. Mistretta, CM. Developmental neurobiology of taste. In: Getchell, T.; Doty, R.; Bartoshuk, L.; Snow, J., editors. *Smell and Taste in Health and Disease*. NY: Raven Press; 1991. p. 35-64.
3. Mistretta CM. The role of innervation in induction and differentiation of taste organs: introduction and background. *Ann. NY Acad. Sci.* 1998; 855:1–13. [PubMed: 9929582]
4. Mbiene JP, MacCallum DK, Mistretta CM. Organ cultures of embryonic rat tongue support tongue and gustatory papilla morphogenesis in vitro without intact sensory ganglia. *J. Comp. Neurol.* 1997; 377:324–340. [PubMed: 8989649]
5. Liu HX, MacCallum DK, Edwards C, Gaffield W, Mistretta CM. Sonic hedgehog exerts distinct, stage-specific effects on tongue and taste papilla development. *Dev. Biol.* 2004; 276:280–300. [PubMed: 15581865]
6. Nosrat CA, MacCallum DK, Mistretta CM. Distinctive spatiotemporal expression patterns for neurotrophins develop in gustatory papillae and lingual tissues in embryonic tongue organ cultures. *Cell Tiss. Res.* 2001; 203:35–45.
7. Mistretta CM, Liu HX, Gaffield W, MacCallum DK. Cyclopamine and jervine in embryonic rat tongue cultures demonstrate a role for Shh signaling in taste papilla development and patterning: fungiform papillae double in number and form in novel locations in dorsal lingual epithelium. *Dev. Biol.* 2003; 253:450–468.
8. Hall JM, Bell ML, Finger TE. Disruption of sonic hedgehog signaling alters growth and patterning of lingual taste papillae. *Dev. Biol.* 2003; 255:263–277. [PubMed: 12648489]
9. Zhou YQ, Liu HX, Mistretta CM. Bone morphogenetic proteins and noggin: inhibiting and inducing fungiform papilla development. *Dev. Biol.* 2006; 297:198–213. [PubMed: 16828469]
10. Liu HX, Henson BS, Zhou YQ, D'Silva NJ, Mistretta CM. Fungiform papilla pattern: EGF regulates inter-papilla lingual epithelium and decreases papilla number via PI3K/Akt, MEK/ERK and p38 MAPK signaling. *Dev. Dyn.* 2008; 237:2378–2393. [PubMed: 18729215]
11. Okubo T, Pevny LH, Hogan BLM. Sox2 is required for development of taste bud sensory cells. *Genes Dev.* 2006; 20:2654–2659. [PubMed: 17015430]
12. Iwatsuki K, Liu HX, Grunder A, Singer MA, Lane TF, Grosschedl R, Mistretta CM, Margolske RF. Wnt signaling interacts with Shh to regulate taste papilla development. *Proc. Nat. Acad. Sci. (USA)*. 2007; 104:2253–2258. [PubMed: 17284610]
13. Liu F, Thirumangalathu S, Gallant NM, Yang SH, Stoick-Cooper CL, Reddy ST, Andl T, Taketo MM, Dlugosz AA, Moon RT, Barlow LA, Millar SE. Wnt-beta-catenin signaling initiates taste papilla development. *Nat. Genet.* 2007; 39:106–112. [PubMed: 17128274]
14. Liu, HX.; Mistretta, CM. Concentration-dependent effects of Shh and agonist on taste papilla formation. 29th Annual Meeting of Association on Chemoreception Sciences; 2007. [abstract].
15. Yamaguchi TP, Bradley A, McMahon AP, Jones SA. Wnt5a pathway underlies outgrowth of multiple structures in the vertebrate embryo. *Development*. 1999; 126:1211–1223. [PubMed: 10021340]

16. Witze ES, Litman ES, Argast GM, Moon RT, Ahn NG. Wnt5a control of cell polarity and directional movement by polarized redistribution of adhesion receptors. *Science*. 2008; 320:365–369. [PubMed: 18420933]
17. Loscertales M, Mikels AJ, Hu JK, Donahoe PK, Roberts DJ. Chick pulmonary Wnt5a directs airway and vascular tubulogenesis. *Development*. 2008; 135:1365–1376. [PubMed: 18305003]
18. Roarty K, Serra R. Wnt5a is required for proper mammary gland development and TGF-beta-mediated inhibition of ductal growth. *Development*. 2007; 134:3929–3939. [PubMed: 17898001]
19. Mistretta CM, Liu HX. Development of fungiform papillae: Patterned lingual gustatory organs. *Arch. Histol. Cytol.* 2006; 69:199–208. [PubMed: 17287575]

Figure 1.
A: Diagram for temporal progression of lingual tissue during fungiform papilla formation [adapted from 19]. In rodents, rat (R) and mouse (M), the fungiform papilla develops from a homogeneous epithelium of the initial lingual swellings at R E13/M E11.5, to a papilla placode or thickened epithelial cell cluster on the spatulate tongue (R E14/M E12.5), to a well defined papilla (R E15–16/M E13.5–14.5). **B:** Summary of effects of secreted molecules (SHH, BMPs, Noggin, Follistatin, EGF, WNT/LiCl, WNT5a) on fungiform papilla number in tongue cultures begun at R E13/M E11.5 or R E14/M E12.5 (↑ represents an increase of papilla number; ↓ decrease; – no change; ? indicates no published data; dashed lines indicate preliminary data only). SHH has dual, concentration-specific effects in E14+2 day rat tongue cultures, increasing papillae at low concentration and decreasing papillae at high[14]. LiCl (see WNT/LiCl) has been used to activate canonical WNT signaling in tongue cultures.

Figure 2.

Scanning electron photomicrographs of E14+2 day rat tongue cultures with addition of SHH, BMPs, EGF or antagonists in culture[9,10,14]. The number in the top right corner of each image represents average number of fungiform papillae for each treatment. STAND: standard medium as a control; CYCL: cyclopamine, an alkaloid that disrupts Shh signaling; Noggin: BMP antagonist; C56: compound 56, a specific inhibitor of EGFR. With SHH (high concentration), BMPs or EGF in the culture medium, fungiform papilla number is reduced. In contrast, with addition of antagonist to block signaling, papilla number is increased (CYCL, Noggin, C56). Formation of fungiform papillae in an atypical tongue region, the intermolar eminence, is observed with CYCL.

Figure 3.

Requirement of WNT canonical signaling in fungiform papilla formation. **First column:** photomicrographs of E14.5 tongue sections indicate the restricted expression of TOPGAL [a β -galactosidase gene driven by a T cell factor (TCF) β -catenin responsive promoter], Lef1, and WNT10b to developing fungiform papillae (arrows)[12]. **Middle and right columns:** scanning electron microscope images of tongues from wild type (WT, middle column) or null mutant (KO, right column). Arrows point to fungiform papillae. Asterisks mark missing or "atrophied" papillae. In null mutants of Postnatal Day 1 β -catenin[13], or Lef1[12], and E14.5 WNT10b[12], fungiform papillae are dramatically reduced. It is noteworthy that the tissue positions are "held" even when papillae are missing in Lef1 knockouts.

Figure 4.

A: Scanning photomicrographs of E16 mouse tongues from wild type (WT) and WNT5a null mutants (^{-/-}). The tongue is shorter and misshapen in WNT5a^{-/-} mutant compared to wild type. **B:** Scanning electron photomicrographs indicate that there is no apparent alteration of circumvallate papilla (arrows) in WNT5a^{-/-} mutant compared to wild type. **C:** Histograms of regional tongue length (oral, pharyngeal and total) and fungiform papilla number. Whereas the length of tongue is reduced in WNT5a^{-/-} mutant, papilla number is not different from wild type.