Published in final edited form as:

Invest Ophthalmol Vis Sci. 2009 September ; 50(9): 4173–4184. doi:10.1167/iov.08-3136.

Embryonic Corneal Schwann Cells Express Some Schwann Cell Marker mRNAs, but No Mature Schwann Cell Marker Proteins

Abigail H. Conrad, Michael Albrecht, Maya Pettit-Scott, and Gary W. Conrad

From the Division of Biology, Kansas State University, Manhattan, Kansas

Abstract

Purpose—Embryonic chick nerves encircle the cornea in pericorneal tissue until embryonic day (E)9, then penetrate the anterior corneal stroma, invade the epithelium, and branch over the corneal surface through E20. Adult corneal nerves, cut during transplantation or LASIK, never fully regenerate. Schwann cells (SCs) protect nerve fibers and augment nerve repair. This study evaluates SC differentiation in embryonic chick corneas.

Methods—Fertile chicken eggs were incubated from E0 at 38°C, 45% humidity. Dissected permeabilized corneas plus pericorneal tissue were immunostained for SC marker proteins. Other corneas were paraffin embedded, sectioned, and processed by in situ hybridization for corneal-, nerve-related, and SC marker gene expression. E9 to E20 corneas, dissected from pericorneal tissue, were assessed by real-time PCR (QPCR) for mRNA expression.

Results—QPCR revealed unchanging low to moderate *SLIT2/ROBO* and *NTN/UNC5* family, *BACE1*, and *CADM3/CADM4* expressions, but high *NEO1* expression. *EGR2* and *POU3F1* expressions never surpassed *PAX3* expression. *ITGNA6/IT-GNB4* expressions increased 20-fold; *ITGNB1* expression was high. SC marker *S100* and *MBP* expressions increased; *MAG*, *GFAP*, and *SCMP* expressions were very low. Antibodies against the MPZ, MAG, S100, and SCMP proteins immunostained along pericorneal nerves, but not along corneal nerves. In the cornea, *SLIT2* and *SOX10* mRNAs were expressed in anterior stroma and epithelium, whereas *PAX3*, *S100*, *MBP*, and *MPZL1* mRNAs were expressed only in corneal epithelium.

Conclusions—Embryonic chick corneas contain SCs, as defined by *SOX10* and *PAX3* transcription, which remain immature, at least in part because of stromal transcriptional and epithelial translational regulation of some SC marker gene expression.

The cornea is one of the most highly innervated tissues on the surface of the body. During chick development, corneal nerves, part of the peripheral nervous system, are derived from neural crest cells, which reside together with ectodermal placode– derived nerve cells in the ophthalmic lobe of the trigeminal ganglion.¹ Hamburger-Hamilton stage 13 to 15,² embryonic day (E)4, trigeminal ganglion ophthalmic lobe placode– derived postmitotic nerves express high levels of transmembrane axon-guidance EphA3 receptor kinase.³ These nerves extend from the trigeminal ganglion into the ophthalmic periocular head mesenchyme, which does not express ephrin-A2 and -A5.⁴ Trigeminal ganglion ophthalmic postmitotic neural crest– derived nerves follow the placode-derived nerves to the periocular mesenchyme, where, from E5 to E8, these periocular nerve bundles subdivide and extend dorsally and ventrally around the cornea, forming a limbal pericorneal nerve ring, but are repelled from entering the cornea.^{5,6} Soluble neurorepellent semaphorin 3A (SEMA 3A), produced by developing lens and diffusing through the cornea and into the adjacent periocular mesenchyme, is responsible for

Corresponding author: Abigail H. Conrad, Division of Biology, Kansas State University, Manhattan, KS 66506-4901; aconrad@ksu.edu.

Disclosure: A.H. Conrad, None; M. Albrecht, None; M. Pettit-Scott, None; G.W. Conrad, None

corneal exclusion of neuropilin receptor-expressing periocular nerves during this period.⁷ In addition, secreted neuroguidance SLIT ligands are synthesized by embryonic lens epithelia, and could interact with axonal transmembrane roundabout (ROBO) receptors expressed on outer surfaces of growing axons to repel corneal nerve growth.⁸ On E9 neural crest– derived limbal ring nerves defasciculate, and sensory nerves invade the anterior corneal stroma simultaneously from all around its perimeter,^{1,5,6,9} then branch and extend anterocentrally, penetrating the epithelium by E12 and reaching the cornea center by E14.⁶ It has been suggested⁶ that extracellular highly sulfated keratan sulfate proteoglycan (KSPG) accumulation, beginning in the posterior corneal stroma by E9,^{10,11} blocks the diffusion of lens SEMA 3A and probably also lens SLIT2, thus allowing the neural crest– derived sensory periocular nerves to grow into the anterior corneal stroma. Subsequent progressive accumulation of highly sulfated KSPG anteriorly across the stroma from E9 to E16¹⁰ guides the corneal nerves toward the epithelium as they seek to avoid highly sulfated KSPG.^{12,13}

Peripheral nervous system nerves are accompanied by neural crest– derived Schwann cells (SCs),¹⁴ which undergo three main developmental transitions: from migrating neural crest cells to SC precursors, SC precursors to immature SCs, and immature SCs to mature myelinating SCs or mature nonmyelinating SCs.¹⁵ The final transition requires intimate contact between nerves and SCs and is accompanied by withdrawal of SCs from the cell cycle.¹³ As SCs differentiate, they express SC-related proteins in transition-stage characteristic patterns, as summarized in Table 1. Transcription factors SOX10, PAX3, POU3F1, and EGR2 are expressed in SCs in a temporally orchestrated pattern: SOX10 is expressed in neural crest cells and all subsequent SC developmental stages including myelination¹⁶; PAX3 is expressed in neural crest cells, SC precursors, immature SCs, and mature nonmyelinating SCs, but not in mature myelinating SCs¹⁷; POU3F1 is expressed in immature SCs and induces the transition of immature SCs to mature myelinating SCs, but later is reduced in expression in mature myelinating SCs¹⁸; expressions of EGR2, myelin-associated glycoprotein (MAG), myelin basic protein (MBP), and P0 (MPZ) genes are strongly upregulated at the onset of myelination in mature myelinating SCs and remain high throughout myelination.¹⁵ MPZ, with expression directly regulated by Sox10,¹⁹ is expressed by SC precursors, immature SCs, and mature myelinating SCs; S100 protein (S100) is expressed by immature SCs and by mature myelinating and nonmyelinating SCs, but not by SC precursors; and glial fibrillary acid protein (GFAP) is expressed by immature SCs and mature nonmyelinating SCs, but not by mature myelinating SCs.¹⁵ In addition, integrin beta 1 (ITGNB1) is expressed only in mature nonmyelinating SCs.¹⁵

If nerves are damaged such that close contacts with SCs are lost, mature myelinating SCs dedifferentiate into immature SCs and actively contribute to nerve regrowth.²⁰ Neuregulin (NRG)-1 type III,²¹ (NRG1-III), a sensory and motor nerve-derived factor (SMDF) neuregulin isoform,²² is expressed by both peripheral nervous system axons²¹ and SCs²³ and determines both whether adjacent SCs will myelinate nerves or just ensheath them²⁴ and how thick the myelin sheath will be.²⁵ BACE1, a β -secretase that is expressed in peripheral nerves, processes SMDF neuregulin so that it can interact with its receptors; BACE1 is required for proper myelination in mice.²⁶ CADMs, cell adhesion molecules, mediate close membrane interactions between adjacent cells. The interaction of axon-expressed CADM3 and SC-expressed CADM4 has been shown to be critical for myelination.²⁷ Secreted netrins (NTNs) function as positive axon guidance factors when they bind nerve transmembrane deleted-in-colorectal-cancer (DCC)^{28,29} or neogenin (NEO)³⁰ receptors, as repulsive factors when they interact with transmembrane UNC5 receptors^{29,31} or with UNC5-DCC heterodimers,³² and as neuronal survival factors.³³ Alternatively, NTNs can be diverted from their role in axon guidance to regulate epithelial cell adhesion and migration by interacting with epithelial cell membrane $\alpha 6 \beta 4$ and $\alpha 3 \beta 1$ integrins.³⁴ SCs interact with extracellular matrix molecules to stimulate peripheral nerve regeneration.³⁵ Indeed, SCs accompanying stromal and subepithelial corneal

nerves have been hypothesized to facilitate stromal and subepithelial nerve regeneration after rabbit corneal injury.³⁶

When corneas are damaged during LASIK surgery, corneal transplantation, or accidental injury, nerve regeneration is slow and often incomplete.^{37,38} Evidence for the presence of SCs in the cornea rests on ultrastructural observations in adult mammalian corneas,^{39–41} but an attempt to demonstrate their presence by staining for SC marker GFAP and S100 proteins immunocytoologically was unsuccessful.³⁹ To examine in more detail the expression of SC-related genes in the cornea, we used quantitative real-time PCR (QPCR) and in situ hybridization to reveal expressions of SC marker mRNAs, and immunocytochemical staining to detect SC marker protein expressions. Our results show that around the periphery of the cornea mature myelinating SCs associate closely with and myelinate pericorneal nerves. In contrast, within the cornea proper, many SC marker-expressing cells are not closely associated with corneal nerves. Moreover, several SC marker mRNAs are expressed in the corneal epithelium, but only the basal SC marker *SOX10* is expressed in the corneal stroma, suggesting that corneal SCs are restricted to the immature SC stage and may be transcriptionally regulated in the stroma and translationally regulated in the epithelium.

MATERIALS AND METHODS

Embryo Culture and Cornea Isolation

Fertile White Leghorn chicken eggs were transferred to a 38°C incubator on E0 and incubated at 45% humidity. Corneas from embryos of the desired age were dissected in sterile saline G (137 mM NaCl, 2.7 mM KCl, 4.3 mM Na₂HPO₄, 1.4 mM KH₂PO₄, 6.1 mM glucose, 0.6 mM MgSO₄, and 0.1 mM CaCl₂ [pH 7.4]). For RNA isolation, the corneas were dissected from the pericorneal tissue, and then quick frozen in liquid nitrogen and stored at –70°C until used. These corneas include both central, mid-central, and lateral corneal tissue, as defined by Müller et al.⁴⁰

RNA Isolation and QPCR

RNA isolation, cDNA synthesis, and QPCR were performed as described previously.^{6,42} Sequences for genes of interest were obtained from GenBank (gi; <http://www.ncbi.nlm.nih.gov/Genbank>; provided in the public domain by the National Center for Biotechnology Information, Bethesda, MD). PCR primers for QPCR were designed (Designer 31 Molecular Beacons Design; Sigma-Aldrich, St. Louis, MO) to amplify fragments between 80 and 150 base pairs in length and are listed in Table 2. Gene names conform to guidelines established by the Second International Workshop on Poultry Genome Mapping, 1994 (described at <http://www.chicken-genome.org>). Each primer set generates only one amplified band with chick corneal cDNA. For QPCR, cDNA dilutions (X) were chosen so that the cycle threshold (Ct, the number of PCR cycles required for the fluorescent signal that measures product accumulation for a given primer set to exceed background levels) for housekeeping gene glyceraldehyde-3-phosphate dehydrogenase (*GAPDH*) was between 13 and 16. All comparative QPCR reaction series consisted of duplicates for 1× and ×/10 cDNA dilutions for each PCR primer pair. Primer pair efficiencies for the 10-fold cDNA dilution were between 90% and 110%. *GAPDH* expression was chosen for normalization of all gene expression. Because corneal *GAPDH* expression was greater than the expression of any of the other genes included in this study, graphic representation of the normalized data employs a negative y-axis.

Corneal Nerve and SC Immunohistochemical Staining

Whole cornea staining was performed as described previously.^{6,43} The corneas were dissected to include pericorneal tissue as well as peripheral and central corneal tissue. Each cornea was

slit radially through the peripheral two thirds of the tissue three times around the cornea to allow antibodies to diffuse laterally into the corneal stroma and epithelium.⁴³ Primary antibodies included: corneal nerves: anti-neuronal class III β -tubulin (TUJ1; CRP Inc., Denver, PA), diluted 1:1000; anti-GFAP (Abcam Inc., Cambridge, MA), diluted 1:500; anti-MAG (United States Biological, Swampscott, MA), diluted 1:500; anti-MPZ (1E8; Developmental Studies Hybridoma Bank, University of Iowa, Iowa City, IA), diluted 1:500; anti-S-100 (Vector Laboratories, Inc., Burlingame, CA), diluted 1:500; and anti-SCMP (anti-SMP; Developmental Studies Hybridoma Bank), diluted 1:500. The secondary antibody was horseradish peroxidase–conjugated goat anti-mouse IgG [H+L] (Invitrogen, Carlsbad, CA) diluted 1:400. Stained corneas were stored in 100% glycerol at 4°C. They were photographed with a dissecting microscope (model M5; Wild Heerbrugg, Heerbrugg, Switzerland) with a digital camera (Coolpix 995; Nikon, Tokyo, Japan) and transmitted light from a light source placed below the glass stage.

RNA Probe Synthesis

PCR primers used to generate cloned gene fragments for *Kera*, 833 bp; *Lum*, 944 bp; *Slit2*, 647 bp; *MPZL1*, 662 bp; *Sox10*, 1091 bp; *PAX3*, 810 bp; *S100*, 970 bp; and *MBP*, 909 bp probes are listed in Table 3. The cDNAs, obtained by reverse transcription PCR from E10 to 14 embryo corneal RNA, were cloned into pGEM-T vectors (Promega, Madison, WI), and sense and antisense riboprobes were synthesized with DIG RNA labeling kit (Roche, Indianapolis, IN) as described previously,^{44,45} digested with DNase I, precipitated, re-dissolved in hybridization buffer, quantitated by dot blot, and stored at –20°C.

In Situ Hybridization on Corneal Sections

E14 eye fronts were dissected into PBS (137 mM NaCl, 2.7 mM KCl, 4.3 mM Na₂HPO₄·7H₂O, 1.4 mM KH₂PO₄ [pH 7.3]) containing 3.7% formaldehyde, freed of adherent lens and retina, fixed overnight at 4°C in modified Carnoy's,⁴⁶ washed three times in PBS to remove all fixative, dehydrated through an EtOH series, washed two times in xylenes, and embedded in paraffin. Paraffin block serial sections of 10 μ m were mounted on slides (SuperFrost Plus; Fisher, Pittsburgh, PA), dried overnight at 40°C, and stored in desiccated boxes at –20°C until hybridization. Hybridization was performed as described previously,^{46,47} except that dewaxing was done with xylenes, prehybridization, and hybridization solutions were made according to Moorman et al.,⁴⁸ and incubations were conducted in a gasket-sealed metal tray (InSlide Out; Boekel, Feasterville, PA) containing wipes soaked in 50% formamide/50% one time SSC (Kimwipes; Kimberly Clark, Neenah, WI) to prevent dehydration. Controls were performed using the sense digoxigenin-labeled RNA transcript of the corresponding antisense RNA probe, and final probe concentrations were 1 ng labeled probe/ μ L hybridization buffer. Hybridization solution (15–30 μ L) was placed on each set of three sections; prehybridization and hybridization were conducted at 65°C. High-stringency washes were performed at 65°C. Antibody reaction and stain development were conducted in the same trays with 50 to 100 μ L solution per section set and 50% PBS/50% H₂O as the humidifying solution. Anti-digoxigenin-AP Fab fragments (Roche) were diluted 1:300 and probes were detected (BM Purple AP Substrate; Roche). Staining was stopped by washing in pH 5.5 PBS, and the slides were rinsed in pH 7.5 PBS, mounted in 70% glycerol/30% PBS, viewed with a microscope (Diaphot 300; Nikon), and photographed with a digital camera (Coolpix 995; Nikon).

Results

Corneal Expression of Genes Involved in Nerve Growth and SC Function by QPCR

Secreted NTNs are a complex family of nerve guidance molecules that can be attractive guidance factors when they interact with nerve transmembrane NEO receptors, or repulsive guidance factors when they interact with nerve transmembrane UNC5 receptors. In the

developing chick cornea, *NTN2L* expression was low and unchanging from E9 to E20, whereas *NTN1* and *NTN4* expressions were 10- and 200-fold lower, respectively, and increased only slightly from E9 to E20 (Fig. 1). Expressions of genes for repulsive NTN receptors *UNC5A-C* were marginally higher than *NTN2L* expression, with *UNC5A* and *UNC5B* expressions virtually equivalent to, and *UNC5C* expression consistently approximately 7.5-fold higher than, *NTN2L* expression (Fig. 1). In contrast, expression of the positive transmembrane NTN receptor *NEO1* was 100-fold higher than that of *NTN2L*, was 10-fold higher than the expression of the most highly expressed negative receptor, *UNC5C*, and remained constant from E9 to E20 (Fig. 1). The *SLIT/ROBO* neuroguidance family members, on the other hand, were more highly expressed than any of the NTNs or their negative receptors, especially at the earlier ages, when the nerves are just entering the cornea,⁶ with *SLIT2* expression 20-fold higher and *ROBO1* expression 100-fold higher than *NTN2L* expression from E9 to E14 (Fig. 1), but then expressions of both *SLIT2* and *ROBO1* declined 10-fold from E14 to E20, when nerves are continuing to grow in the corneal epithelium⁶ (Fig. 1). In contrast, expression of *ROBO2* remained constant at approximately 0.5-fold that of *SLIT2* from E9 to E14, did not decline from E14 to E20, and was approximately 7.5-fold higher than *SLIT2* and *ROBO1* by E20. Expression of *BACE1*, whose enzymatic protein product is essential for processing neuronal neuregulin SMDF before its interaction with SC receptors,²⁶ was moderate throughout corneal development, and did not change as the cornea became highly innervated (Fig. 1).

Expressions of the *CADMs* most closely associated with the axon–SC interaction necessary for myelination, *CADM3* and *–4*, were moderate for neuronally expressed *CADM3* and approximately 7.5-fold lower for SC-specific *CADM4* in the cornea throughout development, and rose less than 2-fold from E9 to E20 (Fig. 2). *CADM1* and *–2* were 7.5-fold more highly expressed than *CADM3* from E9 to E12, while nerves were penetrating the stroma and extending into the epithelium, but then declined in expression from E12 to E20 as *CADM3* expression rose, although they remained more highly expressed than *CADM4* (Fig. 2).

The expressions of the SC transcription factors *PAX3*, *POU3F1*, and *EGR2* are regulated independently of one another, as they in turn regulate migration and stage transition of SCs from SC precursors to mature myelinating SCs in the peripheral nervous system.^{17,18} In the developing chick cornea, *PAX3* was expressed at a moderate level, which did not change appreciably from E9 to E20 (Fig. 2). In contrast, *POU3F1* and *EGR2* expressions began 10-fold lower than that of *PAX3*, and rose 2-fold from E9 to E14. Then from E14 to E20, *POU3F1* expression declined to 7.5-fold lower than *PAX3*, whereas *EGR2* expression rose to 0.5-fold that of *PAX3* (Fig. 2). However, *PAX3* expression remained greater than that of both *POU3F1* and *EGR2* throughout corneal development.

In addition to their functions as nerve guidance factors, extracellular NTNs have been found in the basement membranes of epithelial tissues where they bind integrins $\alpha3\beta1$ and $\alpha6\beta4$ and become adhesive/migratory regulators for epithelial cells.³⁴ In the developing chick cornea, the most highly expressed integrin was *ITGNB1*, with expression constant from E9 through E14, and then declining slightly from E14 to E20 (Fig. 3). In contrast, expression of *ITGNA3* began 100-fold lower than *ITGNB1* expression, rose approximately 7.5-fold from E9 to E14, and then declined back to the E9 level by E20 (Fig. 3). Although their ratios remained the same during E9 to E20, the significant difference in their expression levels suggests that *ITGNB1* and *ITGNA3* may not always function together in the developing chick cornea. On the other hand, *ITGNB4* and *ITGNA6* expressions were almost 100-fold lower than that of *ITGNB1* at E9, but rose coordinately 20-fold to a level similar to that of *ITGNB1* by E20. Their more equivalent expressions suggest that *ITGNA6* and *ITGNB4* function together in the developing chick cornea.

Various marker proteins are made by SCs as they differentiate from SC precursors to mature myelinating or nonmyelinating SCs¹⁵ (Table 1). *MBP* mRNA and protein expressions were strongly upregulated as the SCs transitioned from immature SCs to mature myelinating SCs.¹⁵ In the embryonic chick cornea, *MBP* mRNA expression was moderate on E9, increased approximately twofold from E9 to E12, but then did not increase beyond that level from E12 to E20 (Fig. 4). Similarly, the expression of *S100*, observed in all SCs in the peripheral nervous system,¹⁵ began at the same level as that of *MBP* in the cornea, declined slightly by E12, and then rose approximately 7.5-fold from E12 to E20, having equivalent expression with *MBP* on E14 and E20 (Fig. 4). In contrast, expressions of *GFAP*, expressed in mature nonmyelinating SCs,¹⁵ and of *SCMP* and *MAG*, upregulated in mature myelinating SCs,¹⁵ were 50- to 100-fold lower than the expression of *MBP* and *S100* on E9, and did not increase from E9 to E20 (Fig. 4). No *MPZ* gene has been cloned in chicks, but genes for P0-like protein 1 (*MPZL1*) and P0-like protein 3 (*MPZL3*), which share significant homology with *MPZ* in their extracellular and transmembrane epitopes, have been identified.^{49,50} In the chick cornea, *MPZL1* expression was 10-fold higher than *MBP* expression at E9 and did not change from E9 to E20 (Fig. 4). Expression of *MPZL3* was the same as that of *MBP* and *S100* on E9, but then increased 10-fold from E9 to E20, such that the expressions of *MPZL1* and *MPZL3* were almost equivalent by E20 (Fig. 4).

Corneal Expressions of Genes Involved in Nerve Growth or Cell Adhesion by In Situ Hybridization

In the E14 chick cornea, in situ hybridization revealed that the gene for one of the corneal KSPG core proteins, keratocan (*KERA*; Figs. 5A–C), was expressed by stromal keratocytes and by endothelial cells lining the posterior surface of the cornea, but was not expressed in cells of the epithelium that cover the corneal anterior surface. *SLIT2* (Figs. 5D–F), whose protein product is a secreted molecule that interacts with ROBO receptors on nerve growth cones,⁸ was also expressed by anterior stromal cells. Unlike *KERA*, however, *SLIT2* was also expressed by the epithelial cells that cover the anterior surface of the E14 cornea and was not expressed by posterior corneal stroma cells or corneal endothelium. A third expression pattern was displayed by *MPZL1*, a transmembrane cell adhesion glycoprotein with an intracellular domain very different from that of SC-specific *MPZ*.^{49,50} *MPZL1* was expressed only in the corneal epithelium (Figs. 5G–I), where its expression appeared stronger in the peripheral and mid-corneal epithelium than in the central corneal epithelium (Fig. 5G, arrowheads). *MPZL1* was expressed primarily in the basal layer of the corneal epithelial cells (Fig. 5I).

Pericorneal Expressions of Proteins Characteristic of Nerves and SCs

From E14 through E21, pericorneal nerve bundles, as visualized by neuronal-specific class III β -tubulin immunostaining, formed a ring around the cornea (Fig. 6A, white arrow), defasciculated and extended through the limbus to the peripheral cornea (Fig. 6A, white arrowhead), and continued extension and significant branching radially toward the center of the cornea (Fig. 6A, black arrowhead). In contrast, immunostaining revealed that the SC marker proteins *MPZ*, *S100*, *MAG*, and *SCMP*, products of genes known to be upregulated in mature myelinating SCs,¹⁵ accumulated in the presumptive SCs accompanying the limbal pericorneal ring of nerve bundles (Figs. 6B–E, white arrows) and along defasciculating nerve trunks as they branched away from the limbal ring toward the peripheral cornea (Figs. 6B–E, white arrowheads), but then ceased to be detected from the peripheral to the central cornea (Figs. 6B–E, black arrowheads) where corneal nerves continued to extend (Figs. 6A, black arrowhead). This pericornea-restricted staining pattern was identical for all these proteins characteristic of mature myelinating SCs, even though *MPZ* and *S100* are also known to be expressed in immature SCs,¹⁵ and there was a 100-fold greater expression of *S100* mRNA in the developing cornea than of *MAG* and *SCMP* (see Fig. 4). Immunostaining for *GFAP* in the chick pericorneal tissue and peripheral and central cornea was not detected (data not shown).

Pericorneal and Corneal Expressions of Genes Involved in SC Function by In Situ Hybridization

The expressions of corneal keratocyte mRNA and SC marker mRNAs in pericorneal nerve bundles are shown in Figure 7. An overview of pericorneal nerve bundle location relative to the cornea is shown in Figures 8B, 8E, 8H, 8K, 8N, and 8Q. Stromal KSPG core protein lumican *LUM* mRNA was not expressed in the pericorneal nerve bundles (Fig. 7A), whereas immunocytochemically detected nerve-specific class III β -tubulin protein was heavily expressed in the many individual nerve fibers that composed the pericorneal nerve bundles (Fig. 7B). Transcription factor *SOX10* mRNA, expressed by all peripheral nervous system neural-crest–derived SCs at all stages of differentiation,¹⁵ was highly expressed by many cells within this pericorneal nerve bundle (Fig. 7C), as was *MBP* mRNA, an SC marker known to be significantly upregulated in myelinating SCs¹⁵ (Fig. 7E). In contrast, *S100* mRNA (Fig. 7F), which is expressed in immature, mature myelinating, and mature nonmyelinating SCs,¹⁵ was not as highly expressed by as many cells within the pericorneal nerve bundle, and transcription factor *PAX3* mRNA (Fig. 7D), which is not expressed in mature myelinating SCs and has been shown to repress MBP expression in primary mouse SC cultures,¹⁷ was expressed by few cells within the pericorneal nerve bundle.

In the E14 cornea itself, mRNA expression of the KSPG core protein *LUM*, like that of *KERA*, was high in keratocytes across the entire corneal stroma and along the anterior side of the iris (Fig. 8A), but ceased at the cornea–limbus border (Fig. 8B), and did not occur in the corneal epithelial cell layer (Fig. 8C). In contrast, nerve-specific class III β -tubulin protein immunocytochemical staining occurred in the anterior corneal stroma (Figs. 8D, 8F, black arrowheads), throughout the iris (Fig. 8D), in limbal pericorneal nerve bundles (Fig. 8E, ring with arrow) and other defasciculated nerve segments (Fig. 8E), in nerve segments penetrating from the corneal stroma into the epithelial cell layer (Fig. 8F, white arrowhead), and in both basal and anterior epithelial layers (Fig. 8F). The SC transcription factor *SOX10* mRNA expression pattern was similar to nerve-specific protein expression, occurring in peripheral and central anterior corneal stroma (Figs. 8G, 8I, black arrowheads), throughout the iris (Fig. 8G), in the limbus in both pericorneal nerve ring bundles (Fig. 8H, ring with arrow) and individual limbal cells (Fig. 8H), in SCs penetrating from the anterior stroma into the corneal epithelial layer (Fig. 8I, white arrowheads), and in many cells throughout the basal and anterior corneal epithelium (Fig. 8I). SC transcription factor *PAX3* mRNA (Figs. 8J–L) expression patterns were similar to those of *SOX10*, except that *PAX3* mRNA was not expressed in the corneal stroma, was expressed in few SC cells within the limbal pericorneal nerve bundle (Fig. 8K, ring with arrow), and was more heavily expressed in anterior epithelial cells than in basal epithelial cells (Fig. 8L). Similarly, immature and mature myelinating and nonmyelinating SC marker *S100* mRNA (Figs. 8M–O) was expressed most strongly in the corneal anterior epithelium, throughout the iris, and throughout the limbus, but less strongly within the pericorneal nerve bundle than *SOX10*, and not at all in the stroma. Finally, the expression pattern of myelinating SC marker *MBP* mRNA (Fig. 8P–R) was similar to *PAX3* and *SOX10* patterns, except that *MBP* expression was high in the pericorneal nerve ring bundle (Fig. 8Q, ring with arrow), but less robust in other limbal tissue and the iris, and largely restricted to the anterior corneal epithelium (Fig. 8R).

To examine the spatial relationships between SC marker mRNA-expressing cells and nerve class III β -tubulin protein-expressing nerve segments, we developed sections of E14 corneas, first by in situ hybridization techniques to reveal the SC markers (blue) and then by double staining with antibody to reveal neuronal tubulin (red). *LUM* mRNA expression was restricted to the anterior iris (Fig. 9A), whereas neuronal tubulin protein was expressed throughout the iris (Figs. 9A, 9C, 9E, 9G, 9I). *LUM* mRNA expression was so intense in the corneal stroma (Fig. 9B) that neuronal tubulin staining was difficult to record, but corneal epithelial nerve

segments were seen in both the basal and anterior stroma (Fig. 9B). There were many *SOX10* mRNA-expressing cells in the iris, some of which (Fig. 9C, small white arrows) were immediately adjacent to neuronal tubulin–stained cell segments (Fig. 9C, large white arrows), whereas other *SOX10*-expressing cells did not appear to be intimately associated with neuronal tubulin protein-stained cell segments (Fig. 9C, black arrows). The same relationships occurred between *SOX10* mRNA-expressing SCs and neuronal tubulin–stained nerve segments in the anterior corneal stroma and corneal epithelium (Fig. 9D). Similarly, for iris *PAX3* (Fig. 9E), *S100* (Fig. 9G), and *MBP* (Fig. 9I) expressions, some SC marker–expressing cells were in immediate contact with neuronal tubulin–stained nerve segments (Figs. 9E, 9G, 9I, small and large white arrows), whereas others were not (Figs. 9E, 9G, 9I, black arrows). These same relationships between *PAX3*, *S100*, and *MBP* SC marker mRNA-expressing cells and neuronal tubulin–stained nerve segments occurred in the corneal epithelial layer (Figs. 9F, 9H, 9J). Notably, in the iris, *PAX3*, *S100*, and *MBP* mRNA-expressing cells occupied the anterior iris, where *LUM* mRNA-expressing cells also reside, whereas in the corneal stroma these SC marker mRNAs were not expressed in tissue where *LUM* mRNA was expressed. In addition, in all tissues shown, neuronal class III β -tubulin–stained nerve segments were visible that had no apparent contact with any SC marker–expressing cells.

DISCUSSION

Evidence of SCs in the cornea has previously been based on microscopic descriptions of cells flanking corneal nerves in adult mammalian corneas, but has not demonstrated any SC marker gene expression in the cornea.^{39–41} In this study, QPCR showed that genes for transcription factors characteristic of peripheral nervous system SCs, *SOX10*, *PAX3*, *POU3F1*, and *EGR2*, were expressed in the embryonic chick cornea. *PAX3* expression remained higher than both *POU3F1* and *EGR2* expressions throughout development. In mouse SC development, expression of *PAX3* declines after SC transition from immature to mature myelinating SCs,¹⁸ but forced elevated expression of *PAX3* suppresses myelinating SC expression of *MBP* and returns SCs to a nonmyelinating state.¹⁷ Thus, it is possible that chick corneal SCs do not transition to mature myelinating SCs, because *PAX3* expression keeps *MBP* and other myelin-related gene expression repressed.¹⁷ QPCR expression profiles also revealed moderate expression of *MBP* and *S100*, and low expression of *GFAP*, upregulated in mature nonmyelinating SCs,⁵¹ and *SCMP* and *MAG*, upregulated in mature myelinating SCs.¹⁵ These SC transcription factor and marker gene expression profiles support the conclusion that chick embryonic corneal SCs are in an immature SC stage throughout embryonic corneal development (Table 1). Whereas *ITGNB1*, a marker for mature nonmyelinating SCs,⁵¹ was highly expressed in the embryonic cornea, we currently have no evidence that it is expressed specifically in corneal SCs.

The in situ expression profiles of *SOX10*, *PAX3*, *S100*, and *MBP* coupled with immunochemical profiles of neuronal-specific class III β -tubulin in embryonic chick pericorneal nerve bundles revealed that many cells in pericorneal nerve bundles express *SOX10* mRNA, characteristic of all SCs; and/or *MBP* mRNA, characteristic of mature myelinating SCs.¹⁵ Some cells express *S100*, characteristic of immature as well as mature myelinating and nonmyelinating SCs,¹⁷ and few cells express *PAX3*, characteristic of immature or mature nonmyelinating SCs.

Colocalization of these SC marker mRNAs in a pericorneal nerve bundle in patterns where many myelinating and a few immature SCs are expected supports the conclusion that our *SOX10*, *PAX3*, *S100*, and *MBP* probes detected SCs in corneal and pericorneal tissues. These four mRNAs and neuronal tubulin also colocalized in the corneal epithelium, although often not with such close approximation between SC marker-expressing cells and neuronal tubulin–identified nerve segments as was seen in pericorneal nerve bundles. Outside the central nervous system, *SOX10* expression is restricted to neural crest–derived SCs and melanocytes,⁵² and there are no melanocytes in the cornea. During embryogenesis, *PAX3* expression is found in

trunk skeletal muscle progenitor cells,⁵³ ectodermal placode cells that form trigeminal ganglion ophthalmic lobe ectodermal placode– derived nerves,⁵⁴ optic cup retinal pigment epithelium and optic stalk cells,⁵⁵ and melanocyte stem cells,⁵⁵ as well as in craniofacial nonmyelinating SCs cells.¹⁷ Again, there are no melanocytes or skeletal muscle cells in the cornea, and ectodermal placode– derived nerves do not contribute to corneal nerves.¹ Thus *SOX10*- and *PAX3*-expressing cells in the cornea could only be SCs.

On the other hand, in addition to their expression in SCs, *S100B* is expressed in developing skeletal muscle cells,⁵⁶ chondrocytes,⁵⁷ adipocytes,⁵⁸ melanocytes,⁵⁹ some brain lymphocytes,⁵⁹ and Langerhans cells,⁶⁰ and *MBP* is expressed in thymocytes⁶¹ and other hemopoietic progenitors⁶² in Golli forms that are splice variants of the same gene that encodes myelin-specific MBP mRNA.⁶³ Golli mRNAs contain three exons unique to Golli-MBP, spliced in-frame to exon 1b plus part of intron 1 of the myelin-MBP sequence (HMBP), or to exons 1b-7 of the myelin-MBP sequence (MBP2). No complete chick Golli-MBP protein coding sequences are currently in the GenBank database, but there is an EST (expressed sequence tag) containing a Golli-specific sequence and a few 3' bases identical with the most 5' bases of chick myelin-MBP 5' UTR. If chick Golli-MBPs are similar to mouse Golli-MBPs, our chick MBP QPCR- and in situ primers exclude all HMBP sequences, but could be detecting MBP2 mRNA, since mouse MBP2 exon 1b-7 mRNA is alternatively spliced in ways identical with mouse myelin-MBP mRNA,^{62,63} and shares the same 3' UTR region. Whereas skeletal muscle cells, chondrocytes, adipocytes, and melanocytes are not present in the cornea, the human cornea has recently been shown to have some Langerhans cells in its epithelium, some potential antigen-presenting dendritic cells in its anterior stroma, and some macrophages in its posterior stroma.⁶⁴ Our in situ studies showed no *S100* or *MBP* expression in the anterior or posterior stroma, but did show expression of *S100* and *MBP* mRNAs in the epithelial cell layer where Langerhans cells have been identified. Thus, it is possible that in the corneal epithelium some *S100*- and *MBP2*-expressing cells could be Langerhans cells. Notably, however, another lymphoid cell– expressed gene observed in this study, *MPZL1*, was expressed primarily in the basal epithelium (Fig. 5I), whereas *S100* and *MBP* mRNAs, as well as *SOX10* and *PAX3* mRNAs, were expressed strongly in the anterior epithelium (Figs. 8I, 8L, 8O, 8R), suggesting that *S100* and *MBP* probes also detect SCs in the corneal epithelium.

S100, *SCMP*, and *MAG* mRNAs were expressed in the cornea, but were not translated into enough protein to be detected immunocytochemically in the central cornea, whereas their protein products were detected in the pericorneal nerves that defasciculate into corneal nerves. These observations suggest that there may be translational regulation of the expression of these mRNAs, as well as that of *MBP*, in embryonic corneal epithelial SCs, where *S100* and *MBP* mRNAs have been localized. Mechanisms for SC posttranslational regulation have yet to be delineated. Intriguingly, *S100*-, *MBP*-, and *PAX3*-expressing cells were identified in the iris, too, where nerves become myelinated and contain MBP protein.⁶⁵ Notably, *SOX10*-, *PAX3*-, *S100*-, and *MBP*-expressing cells cohabited in anterior iris tissue where *LUM* mRNA was expressed, but where lumican core protein is not posttranslationally modified by addition of sulfated KS chains.¹⁰ In contrast, SCs within the sulfated KS-rich E14 corneal stroma expressed *SOX10*, but not *PAX3*, *S100*, or *MBP* mRNAs. It is possible that extracellular matrix sulfated KS chains somehow repress downstream SC marker transcription in *SOX10*-expressing SCs in the corneal stroma. Mechanisms for glycosaminoglycan regulation of SC transcription remain to be resolved, but chondroitin/dermatan sulfate⁶⁶ and heparan sulfate⁶⁷ proteoglycans have been shown to influence SC migration, myelination, and peripheral sensory axon regeneration.

In both the E14 iris and the E14 cornea, some SC marker expressing cells appeared to be in contact with nerve segments, whereas others appeared not to be, although it is possible that the nerves they contacted were just not exposed in the corneal section viewed. Nevertheless, in the

iris, close nerve-SC interaction developed sufficiently to allow nerves to become myelinated, whereas in the corneal stroma and epithelium nerve myelination did not occur. It is possible that something in the cornea prevents the necessary close adhesion between SCs and nerves that myelination demands. CADMs are a family of cell-adhesion molecules.⁶⁸ Typically *CADM1* and -3 are expressed by nerve cells, *CADM1* and -4 are expressed by SCs, *CADM4* expression is upregulated in SCs when they begin to myelinate, and heterophilic interaction between nerve axon-*CADM3* and SC-*CADM4* mediates an axon-SC interaction that is essential for peripheral nervous system myelination.^{27,69} In the developing cornea, *CADM4* expression remained the lowest of the *CADM* expressions throughout corneal development, apparently not high enough to support myelination. Similarly, in the developing chick cornea moderate *BACE1* expression did not increase during development, supporting further the conclusion that developing chick corneal SCs remain in an immature SC stage.

SLIT2 expression in embryonic chick anterior corneal stroma and epithelium places it in the region of the cornea where nerves extend and branch during chick embryonic development.^{5,6} Expression of *SLIT2* has also been reported in mouse embryo corneal epithelium.⁷⁰ Although classically thought to act as chemorepellant axon guidance factors, in vertebrates SLIT/ROBO interactions are also positive regulators for peripheral nervous system sensory axon extension and branching, especially in trigeminal ganglion axon branching in the embryonic head,⁷¹ depending on which extracellular matrix component binds them.⁷² In *Drosophila*, SLITs bound by heparan sulfate sidechain-bearing core protein Glypican-1 are transformed into nerve elongation/branching-stimulating forms.⁷³ *SLIT2* and KSPG core protein mRNAs⁴² are highly expressed in E9 to E14 anterior embryonic chick corneal stromal cells, before accumulation of highly sulfated KS in the extracellular matrix advances from the posterior into the anterior corneal stroma on E14 to E16.¹⁰ It is attractive to imagine that KSPG core proteins act as positive corneal nerve extension/branching chemoregulators during early corneal nerve development by binding with *SLIT2* protein, since each corneal sensory nerve that enters the anterior cornea supports from 200 to 3000 epithelial nerve endings.⁵⁶ Further studies on the binding properties of corneal KSPG core proteins with SLITs will be necessary to better understand SLIT function in corneal nerve branching.

Corneal expression of the NTN chemoattractant receptor gene *NEO1*, the most ubiquitously expressed NTN receptor outside the central nervous system,⁷⁴ was high throughout corneal development, consistent with the possibility that corneal NTNs also may be acting through the *NEO1* receptor to attract corneal nerve extension. Alternatively, NTNs are expressed in developing epithelia,³⁴ where they function in epithelial cell adhesion and migration by interacting with the integrins $\alpha6\beta4$ and $\alpha3\beta1$.³⁴ In the developing chick cornea, expressions of *ITGNA6* and *ITGNB4* rose coordinately from E9 to E20, suggesting that $\alpha6\beta4$ is present in sufficient concentration to interact with corneal NTNs to play a role in corneal epithelial cell adhesion and migration. Although *ITGNB1* expression was approximately 100-fold higher than *ITGNA3* expression, their expression patterns also paralleled one another during corneal development, suggesting that $\alpha3\beta1$ could also function in corneal epithelial cell adhesion and migration, with the additional *ITGNB1* participating in other as yet undefined corneal roles. In addition, adhesion/migration of lymphocytes or epithelial cells in corneal epithelium may also be facilitated by *MPZL1*, expressed in mesenchymal and hemopoietic cells in mice,⁴⁹ and *MPZL3*, expressed in epidermal keratinocytes in mice,⁵⁰ as these genes have not been shown to be expressed by SCs. *MPZL1*, the more highly expressed in embryonic chick cornea, was expressed chiefly in the peripheral corneal basal epithelial layer, where it could facilitate cell migration on the basement membrane underlying the epithelial layer.⁴⁹

SCs are thought to stimulate nerve regeneration by positive signals that are independent of providing a tract for the nerves to regrow along.^{15,36} Human stromal nerves are ensheathed by thin SC extensions suggestive of immature SCs,⁷⁵ and chick embryonic SCs show gene

expression profiles characteristic of immature SCs, suggesting that they are in an appropriate developmental stage to stimulate corneal nerve regeneration after hatching. Mechanisms to temporarily stimulate SC development in posttransplantation anterior corneas or post-LASIK flaps may improve posttransplantation or post-LASIK nerve regeneration.

Acknowledgments

Supported by National Institutes of Health EY000952 (GWC); the Howard Hughes Medical Institute Biological Sciences Education Program HHMI 52003734 (MA); and the Terry C. Johnson Center for Basic Cancer Research at Kansas State University (MA, MP-S).

References

1. Lwigale PY. Embryonic origin of avian corneal sensory nerves. *Dev Biol* 2001;239:323–337. [PubMed: 11784038]
2. Hamburger V, Hamilton HL. A series of normal stages in the development of the chick embryo. *J Morphol* 1951;88:49–94.
3. Flanagan JG, Vanderhaeghen P. The ephrins and eph receptors in neural development. *Ann Rev Neurosci* 1998;21:309–345. [PubMed: 9530499]
4. Jayasena CS, Flood WD, Koblar SA. High EphA3 expressing ophthalmic trigeminal sensory axons are sensitive to ephrin-A5-Fc: implications for lobe specific axon guidance. *Neuroscience* 2005;135:97–109. [PubMed: 16054765]
5. Bee JA. The development and pattern of innervation of the avian cornea. *Dev Biol* 1982;92:5–15. [PubMed: 6179807]
6. Conrad AH, Straffuss JM, Wittman MD, Conway S, Conrad GW. Thyroxine increases the rate but does not alter the pattern of innervation during embryonic chick corneal development. *Invest Ophthalmol Vis Sci* 2008;49:139–153. [PubMed: 18172086]
7. Lwigale PY, Bronner-Fraser M. Lens-derived Semaphorin3A regulates sensory innervation of the cornea. *Dev Biol* 2007;306:750–759. [PubMed: 17499699]
8. Niclou SP, Jia L, Raper JA. Slit2 is a repellent for retinal ganglion cell axons. *J Neurosci* 2000;20:4962–4974. [PubMed: 10864954]
9. Riley NC, Lwigale PY, Conrad GW. Specificity of corneal nerve positions during embryogenesis. *Mol Vis* 2001;7:297–304. [PubMed: 11754335]
10. Funderburgh JL, Caterson B, Conrad GW. Distribution of proteoglycans antigenically related to corneal keratan sulfate proteoglycans. *J Biol Chem* 1987;262:11634–11640. [PubMed: 2957372]
11. Zhang Y, Conrad AH, Tasheva ES, et al. Detection and quantification of sulfated disaccharides from keratan sulfate and chondroitin/dermatan sulfate during chick corneal development by ESI-MS/MS. *Invest Ophthalmol Vis Sci* 2005;46:1604–1614. [PubMed: 15851558]
12. Perris R, Perissinotto D, Pettway Z, Bronner-Fraser M, Morgelin M, Kimata K. Inhibitory effects of PG-H/aggrecan and PG-M/versican on avian neural crest cell migration. *FASEB J* 1996;10:293–301. [PubMed: 8641562]
13. Macé K, Saxod R, Feuerstein C, Sadoul R, Hemming FJ. Chondroitin and keratan sulfates have opposing effects on attachment and outgrowth of ventral mesencephalic explants in culture. *J Neurosci Res* 2002;70:46–56. [PubMed: 12237863]
14. Aquino JB, Hjerling-Leffler J, Koltzenburg M, Edlund T, Villar MJ, Ernfors P. In vitro and in vivo differentiation of boundary cap neural crest stem cells into mature Schwann cells. *Exp Neurol* 2006;198:438–449. [PubMed: 16442526]
15. Mirsky R, Woodhoo A, Parkinson DB, Arthur-Farraj P, Bhaskaran A, Jessen K. Novel signals controlling embryonic Schwann cell development, myelination and dedifferentiation. *J Periph Nerv Syst* 2008;13:122–135.
16. Schreiner S, Cossais F, Fischer K, et al. Hypomorphic Sox10 alleles reveal novel protein functions and unravel developmental differences in glial lineages. *Development* 2007;134:3271–3281. [PubMed: 17699610]

17. Kioussi C, Gross MK, Gruss P. Pax3: a paired domain gene as a regulator in PNS myelination. *Neuron* 1995;15:553–562. [PubMed: 7546735]
18. Blanchard AD, Sinanan A, Parmantier E, et al. Oct-6(SCIP/Tst-1) is expressed in Schwann cell precursors, embryonic Schwann cells, and postnatal myelinating Schwann cells: comparison with Oct-1, Krox-20 and Pax-3. *J Neurosci Res* 1996;45:630 – 640. [PubMed: 8951674]
19. Peirano RI, Goerich DE, Riethmacher D, Wegner M. Protein zero gene expression is regulated by glial transcription factor Sox10. *Mol Cell Biol* 2000;20:3198 –3209. [PubMed: 10757804]
20. Parkinson DB, Bhaskaran A, Arthur-Farraj P, et al. c-Jun is a negative regulator of myelination. *J Cell Biol* 2008;181:625– 637. [PubMed: 18490512]
21. Ho W-H, Armanini MP, Nuijens A, Phillips HS, Osheroff PL. Sensory and motor neuron-derived Factor: a novel heregulin variant highly expressed in sensory and motor neurons. *J Biol Chem* 1995;270:14523–14532. [PubMed: 7782315]
22. Adlkofer K, Lai C. Role of neuregulins in glial cell development. *Glia* 2000;29:104 –111. [PubMed: 10625327]
23. Carroll SL, Miller ML, Frohnert PW, Kim SS, Corbett JA. Expression of neuregulins and their putative receptors, ErbB2 and ErbB3, is induced during Wallerian degeneration. *J Neurosci* 1997;17:1642– 1659. [PubMed: 9030624]
24. Taveggia C, Zanazzi G, Petrylak A, et al. Neuregulin-a type III determines the ensheathment fate of axons. *Neuron* 2005;47:681– 694. [PubMed: 16129398]
25. Michailov GV, Sereda MW, Brinkmann BG, et al. Axonal neuregulin-1 regulates myelin sheath thickness. *Science* 2004;304:700 –703. [PubMed: 15044753]
26. Willem M, Garratt AN, Novak B, et al. Control of peripheral nerve myelination by the β -Secretase BACE1. *Science* 2006;314:664 –666. [PubMed: 16990514]
27. Spiegel I, Adamsky K, Eshed Y, et al. A central role for Necl4 (SynCAM4) in Schwann cell-axon interaction and myelination. *Nat Neurol* 2007;10:861– 869.
28. Wang H, Copeland NG, Gilbert DJ, Jenkins NA, Tessier-Lavigne M. Netrin-3, mouse homolog of human NTN2L, is highly expressed in sensory ganglia and shows differential binding to netrin receptors. *J Neurosci* 1999;19:4938 – 4947. [PubMed: 10366627]
29. Deiner MS, Kennedy TE, Fazeli A, Serafini T, Tessier-Lavigne M, Sretavan DW. Netrin and DCC mediate axon guidance locally at the optic disc: loss of function leads to optic nerve hypoplasia. *Neuron* 1997;19:575–589. [PubMed: 9331350]
30. Vielmetter J, Kayyem JF, Roman JM, Dreyer WJ. Neogenin, an avian cell surface protein expressed during terminal neuronal differentiation, is closely related to the human tumor suppressor molecule deleted in colorectal cancer. *J Cell Biol* 1994;127:2009 –2020. [PubMed: 7806578]
31. Leonardo DE, Hinck L, Masu M, Keino-Masu K, Ackerman SL, Tessier-Lavigne M. Vertebrate homologues of *C. elegans* UNC-5 are candidate netrin receptors. *Nature* 1997;386:833– 838. [PubMed: 9126742]
32. Hong K, Hinck L, Nishiyama M, Poo M, Tessier-Lavigne M, Stein E. A ligand-gated association between cytoplasmic domains of UNC5 and DCC family receptors converts netrin-induced growth cone attraction to repulsion. *Cell* 1999;97:927–941. [PubMed: 10399920]
33. Tang X, Jang S-W, Okada M, et al. Netrin mediates neuronal survival through PIKE-L interaction with the independence receptor UNC5B. *Nat Cell Biol* 2008;10:698 –706. [PubMed: 18469807]
34. Yebra M, Montgomery AMP, Diaferia GR, et al. Recognition of the neural chemoattractant netrin-1 by integrins $\alpha 6 \beta 4$ and $\alpha 3 \beta 1$ regulates epithelial cell adhesion and migration. *Dev Cell* 2003;5:695– 707. [PubMed: 14602071]
35. Armstrong SJ, Wiberg M, Terenghi G, Kingham PJ. Laminin activates NF-kB in Schwann cells to enhance neurite outgrowth. *Neurosci Lett* 2008;439:42– 46. [PubMed: 18502047]
36. Chan KY, Jarvelainen M, Chang JH, Edenfield MJ. A cryodamage model for studying corneal nerve regeneration. *Invest Ophthalmol Vis Sci* 1990;31:2008 –2021. [PubMed: 2210997]
37. Tervo T, Vannas A, Tervo K, Holden BA. Histochemical evidence of limited reinnervation of human corneal grafts. *Acta Ophthalmol (Copenh)* 1985;63:207–214. [PubMed: 3890460]
38. Erie JC, McLaren JW, Hodge DO, Bourne WM. Recovery of corneal subbasal nerve density after PRK and LASIK. *Am J Ophthalmol* 2005;140:1059 –1064. [PubMed: 16376651]

39. Shimeld C, Efsthathiou S, Hill T. Tracking the spread of a lacZ-tagged Herpes Simplex virus Type 1 between the eye and the nervous system of the mouse: comparison of primary and recurrent infection. *J Virol* 2001;75:5252–5262. [PubMed: 11333907]
40. Müller LJ, Pels L, Vrensen GF. Ultrastructural organization of human corneal nerves. *Invest Ophthalmol Vis Sci* 1996;37:476 – 488. [PubMed: 8595948]
41. Peavy D, Bobos S, Gebhardt B, Beuerman R. Immunohistochemical differentiation of Schwann and epithelial basal lamina in the rabbit cornea. *Neurosci Lett* 1987;79:40 – 46. [PubMed: 3313118]
42. Conrad AH, Zhang Y, Walker AR, et al. Thyroxine affects expression of KSPG-related genes, the carbonic anhydrase II gene, and KS sulfation in the embryonic chicken cornea. *Invest Ophthalmol Vis Sci* 2006;47:120 –132. [PubMed: 16384953]
43. Koo SJ, Clark-Alderfer JD, Tanaka H, et al. Species-specific immunostaining of embryonic corneal nerves: techniques for inactivating endogenous peroxidases and demonstration of lateral diffusion of antibodies in the plane of the corneal stroma. *J Neurosci Methods* 1998;85:63–71. [PubMed: 9874142]
44. Dunlevy JR, Beales MP, Berryhill BL, Cornuet PK, Hassell JR. Expression of the keratocan sulfate proteoglycans lumican, keratocan, osteoglycin/mimecan during chick corneal development. *Exp Eye Res* 2000;70:349 –362. [PubMed: 10712821]
45. Cheng Y-C, Cheung M, Abu-Elmad MM, Orme A, Scotting PJ. Chick Sox10, a transcription factor expressed in both early neural crest cells and central nervous system. *Dev Brain Res* 2000;121:233–241. [PubMed: 10876038]
46. Etchevers HC, Vincent C, Le Douarin NM, Couly GF. The cephalic neural crest provides pericytes and smooth muscle cells to all blood vessels of the face and forebrain. *Development* 2001;128:1059 –1068. [PubMed: 11245571]
47. Conrad AH, Conrad GW. The keratocan gene is expressed in both ocular and non-ocular tissues during early chick development. *Matrix Biol* 2003;22:323–337. [PubMed: 12935817]
48. Moorman AFM, Houweling AC, de Boer PAJ, Christoffels VM. Sensitive nonradioactive determination of mRNA in tissue sections: novel application of the whole-mount in situ hybridization protocol. *J Histochem Cytochem* 2001;49:1– 8. [PubMed: 11118473]
49. Zannettino ACW, Roubelakis M, Weldon KJ, et al. Novel mesenchymal and hematopoietic cell isoforms of the SHP-2 docking receptor, PZR: identification, molecular cloning and effects on cell migration. *Biochem J* 2003;370:537–549. [PubMed: 12410637]
50. Cao T, Racz P, Szaulter KM, et al. Mutation in MpzL3, a gene encoding a predicted adhesion protein, in the rough coat (rc) mice with severe skin and hair abnormalities. *J Invest Dermatol* 2007;127:1375–1386. [PubMed: 17273165]
51. Mirsky R, Jessen KR. The biology of non-myelin-forming Schwann cells. *Ann N Y Acad Sci* 1986;486:132–146. [PubMed: 3551735]
52. Britsch S, Goerich DE, Riethmacher D, et al. The transcriptional factor Sox10 is a key regulator of peripheral glial development. *Genes Dev* 2001;15:66 –78. [PubMed: 11156606]
53. Relaix F, Rocancourt D, Mansouri A, Buckingham M. A pax3/pax7-dependant population of skeletal muscle progenitor cells. *Nature* 2005;435:948 –953. [PubMed: 15843801]
54. Stark MR, Sechrist J, Bronner-Fraser M, Marcelle C. Neural tube-ectodermal interactions are required for trigeminal placode formation. *Development* 1997;124:4287– 4295. [PubMed: 9334277]
55. Hornyak TJ, Hayes DJ, Chiu L-Y, Ziff EB. Transcription factors in melanocyte development: distinct roles for Pax-3 and Mitf. *Mech Dev* 2001;101:47–59. [PubMed: 11231058]
56. Arcuri C, Giambanco I, Bianchi R, Donato R. Annexin V, annexin VI, S100A1 and S100B in developing and adult avian skeletal muscles. *Neuroscience* 2002;109:371–388.
57. Saito T, Ikeda T, Nakamura K, Chung U-I, Kawaguchi H. S100A1 and S100B, transcriptional targets of Sox trio, inhibit terminal differentiation of chondrocytes. *EMBO Rep* 2007;8:504 –509. [PubMed: 17396138]
58. Atanassova P. Immunohistochemical expression of S-100 protein in human embryonal fat cells. *Cells Tissues Organs* 2001;169:355–360. [PubMed: 11490114]
59. Steiner J, Bernstein HG, Bielau H, et al. Evidence for a wide extra-astrocytic distribution of S100B in human brain. *BMC Neurosci* 2007;8:2–11. [PubMed: 17199889]

60. Park HR, Min SK. Expression of S100A2 and S100B proteins in epithelial tumors of the skin. *J Cutan Pathol* 2003;30:373–378. [PubMed: 12834486]
61. Feng J-M, Givogri IM, Bongarzone ER, et al. Thymocytes express the Golli products of the myelin basic protein gene and levels of expression are stage specific. *J Immunol* 2000;165:5443–5450. [PubMed: 11067896]
62. Marty MC, Aqlli F, Rutin J, Fritz R, Trisler D, Pessac B. The myelin basic protein gene is expressed in differentiated blood lineages and in hemopoietic progenitors. *Proc Natl Acad Sci USA* 2002;99:8856 – 8861. [PubMed: 12084930]
63. Campagnoni AT, Pribyl TM, Campagnoni CW, et al. Structure and developmental regulation of the Gollimbp, a 105-kilobase gene that encompasses the myelin basic protein gene and is expressed in cells in the oligodendrocyte lineage in the brain. *J Biol Chem* 1993;268:4930 – 4938. [PubMed: 7680345]
64. Hamrah P, Dana MR. Corneal antigen-presenting cells. *Chem Immunol Allergy* 2007;92:58 –70. [PubMed: 17264483]
65. Verhagen C, Mor F, Coihlen IR. T cell immunity to myelin basic protein induces anterior uveitis in Lewis rats. *J Neuroimmunol* 1994;53:65–71. [PubMed: 7519633]
66. Grimpe B, Pressman Y, Lupa MD, Horn KP, Bunge MB, Silver J. The role of proteoglycans in Schwann cell/astrocyte interactions and the regeneration failure at PNS/CNS interfaces. *Mol Cell Neurosci* 2005;28:18 –29. [PubMed: 15607938]
67. Chernousov MA, Rothblum K, Stahl RC, Evans A, Prentiss L, Carey DJ. Glypican-1 and $\alpha 4(V)$ collagen are required for Schwann cell myelination. *J Neurosci* 2006;26:508 –517. [PubMed: 16407548]
68. Sakisaka T, Takai Y. Biology and pathology of nectins and nectin-like molecules. *Curr Opin Cell Biol* 2004;16:513–521. [PubMed: 15363801]
69. Maurel P, Einheber S, Galinska J, et al. Nectin-like proteins mediate axon-Schwann cell interactions along the internode and are essential for myelination. *J Cell Biol* 2007;178:861– 874. [PubMed: 17724124]
70. Thompson H, Camand O, Barker D, Erskine L. Slit proteins regulate distinct aspects of retinal ganglion cell axon guidance within dorsal and ventral retina. *J Neurosci* 2006;26:8082– 8091. [PubMed: 16885222]
71. Ma L, Tessier-Lavigne M. Dual branch-promoting and branch-repelling actions of Slit/Robo signaling on peripheral and central branches of developing sensory axons. *J Neurosci* 2007;27:6843–6851. [PubMed: 17581972]
72. Nguyen-Ba-Charvet KT, Brose K, Marillat V, Sotelo C, Tessier-Lavigne M, Chedotal A. Sensory axon response to substrate-bound Slit2 is modulated by laminin and cyclic GMP. *Mol Cell Neurosci* 2001;17:1048 –1058. [PubMed: 11414793]
73. Liang Y, Annan RS, Carr SA, et al. Mammalian homologues of the *Drosophila* slit protein are ligands of the heparan sulfate proteoglycan glypican-1 in brain. *J Biol Chem* 1999;274:17885–17892. [PubMed: 10364234]
74. Gad JM, Keeling SL, Wilks AF, Tan S-S, Cooper HM. The expression patterns of guidance receptors, DCC and Neogenin, are spatially and temporally distinct throughout mouse embryogenesis. *Dev Biol* 1997;192:258 –273. [PubMed: 9441666]
75. Müller LJ, Marfurt CF, Kruse F, Tervo TMT. Corneal nerves: structure, contents and function. *Exp Eye Res* 2003;76:521–542. [PubMed: 12697417]

Figure 1.

Corneal QPCR expression of some nerve growth-regulating protein families and SC attractant modifiers did not vary markedly over the period of corneal innervation. The SLIT/ROBO family was more highly expressed than was the NTN/UNC family. Neuroattractant NEO receptor was the most highly expressed NTN receptor, suggesting that NTNs may function as neuroattractants in the cornea. BACE1 expression did not increase during development.

Figure 2.

QPCR expressions of genes normally upregulated in myelinating SCs were not significantly upregulated in corneal cells during corneal innervation, SC-specific *CADM4* expression was not significantly upregulated, and *PAX3* expression remained higher than *EGR2* and *POU3F1* expressions throughout.

Figure 3.

QPCR-detected expressions of genes for alternative NTN-receptor integrin pairs *ITGNA6/ITGNB4* and *ITGNA3/ITGNB1* changed in parallel with pair mates during corneal development. Expressions of *ITGNA6/ITGNB4* were similar and rose 10-fold, suggesting that these integrins function together. Expression of *ITGNB1* was approximately 100-fold higher than *ITGNA3* expression throughout corneal development, suggesting that some *ITGNB1* may function with *ITGNA3*, whereas other *ITGNB1* could be involved in yet to be defined corneal functions.

Figure 4.

QPCR revealed that mRNAs for some SC markers were expressed moderately and rose slightly during corneal development, whereas other SC marker mRNAs were weakly expressed. *S100* and *MBP* expressions began moderately and rose approximately twofold during corneal innervation, whereas *SCMP*, *MAG*, and *GFAP* were weakly expressed and did not increase in expression, suggesting that corneal SCs remain in the immature stage throughout corneal development. *MPZL* isoforms were more highly expressed than SC marker mRNAs in the cornea, suggesting that they play significant roles in corneal development.

Figure 5.

Localizations of nerve growth- or cell adhesion-related gene expressions in the E14 chick cornea. In situ hybridization was performed on E14 corneal sections using labeled probes for *KERA*, *SLIT2*, and *MPZL1*. (A–C) *KERA*, the gene for the KSPG core protein keratocan, was expressed in corneal endothelium and stroma but not in corneal epithelium. (D–F) *SLIT2*, the gene for a soluble neuro-regulatory ligand, was expressed in the anterior corneal stroma and the corneal epithelium, but not in the posterior stroma or endothelium. (G–I) *MPZL1*, the gene for a MPZ-like transmembrane protein thought to be involved in cell–cell adhesion and migration, was expressed primarily in the peripheral and mid–corneal epithelium (G, arrowheads), especially in the basal epithelial layer (I), but not in the corneal stroma or endothelium. ep, epithelium; st, stroma. Bar, 100 μm.

Figure 6.

SC marker proteins were detected in the limbus and pericorneal tissue, but not in the mid and central regions of the cornea. Corneas were dissected, fixed, incubated with selected SC marker protein antibodies, and developed with secondary antibody. (A) E14 cornea incubated with anti- β -tubulin specific for nerves. (B) E19 cornea incubated with anti-MPZ antibody. (C) E19 cornea incubated with anti-S100 antibody. (D) E19 cornea incubated with anti-MAG antibody. (E) E17 cornea incubated with anti-SCMP antibody. *White arrows*: pericorneal nerve ring in the limbus outside the cornea. *White arrowheads*: pericorneal nerves defasciculating from the pericorneal ring and penetrating to the periphery of the cornea. *Black arrowheads*: mid and central corneal nerves (A) or area of the cornea (B–E). Bar, 500 μ m.

Figure 7.

SC marker mRNAs were expressed within pericorneal nerve bundles. E14 pericorneal tissue was developed by in situ hybridization techniques to reveal expression of KSPG core protein mRNA or SC marker-mRNA, or by immunocytochemical techniques to identify neuronal protein expression. (A) *LUM* KSPG core protein mRNA expression. (B) Neuronal class III β -tubulin protein expression. (C) SC marker *SOX10* mRNA expression. (D) SC marker *PAX3* mRNA expression. (E) SC marker *MBP* mRNA expression. (F) SC marker *S100* mRNA expression. Circle: pericorneal nerve bundle. Bar, 30 μ m.

Figure 8.

SC marker *SOX10* mRNA was expressed in embryonic corneal stroma and epithelium, whereas SC marker *PAX3*, *S100*, and *MBP* mRNAs were expressed only in corneal epithelium. E14 corneal tissue was developed by in situ hybridization techniques to reveal KSPG core protein mRNA or SC marker-mRNA expressions, or by immunocytochemical techniques to identify neuronal protein expression. (A–C) *LUM* KSPG core protein mRNA was expressed by cells in corneal stroma and anterior iris, but not in a pericorneal nerve bundle (B, circle with arrow) or corneal epithelium. (D–F) Neuronal class III β -tubulin protein was expressed in nerve segments in the corneal stroma (D, F, black arrowheads), iris, limbus, a pericorneal nerve bundle (E, circle with arrow), passing from stroma to epithelium (F, white arrowhead), and the basal and anterior epithelium. (G–I) SC marker *SOX10* mRNA was expressed by cells in the anterior corneal stroma (G, I, black arrowheads), iris, limbus, a pericorneal nerve bundle (H, circle with arrow), passing from stroma to epithelium (I, white arrowheads), and basal and anterior epithelium. (J–L) SC marker *PAX3* mRNA was expressed

by cells in corneal basal and anterior epithelium (**J, L**), iris, limbus, and a pericorneal nerve bundle (**K**, *circle with arrow*), but not by cells in the corneal stroma (**L**). (**M–O**) SC marker *S100* mRNA was expressed by cells in the corneal basal and anterior epithelium (**M, O**), the iris, the limbus, and a pericorneal nerve bundle (**N**, *circle with arrow*), but not by cells in the corneal stroma (**O**). (**P–R**) SC marker *MBP* mRNA was expressed by cells in the corneal basal and anterior epithelium (**P, R**), the iris, the limbus, and a pericorneal nerve bundle (**Q**, *circle with arrow*), but not by cells in the corneal stroma (**R**). Bar: (**A, D, G, J, M, P**) 300 μm ; (**B, E, H, K, N, Q**) 100 μm ; (**C, F, I, L, O, R**) 30 μm .

Figure 9.

In embryonic iris and cornea, nerves and SC marker– expressing cells were sometimes in contact with one another and sometimes not in apparent contact with one another. E14 iris and corneal tissue was developed by in situ hybridization to reveal expression of KSPG core protein mRNA or SC marker mRNA, and then also was double stained by immunocytochemistry to simultaneously identify neuronal protein expression. (A, B) *LUM* KSPG core protein mRNA (blue) and neuronal class III β -tubulin protein (red) expressions in iris (A) and corneal stroma and epithelium (B). (C, D) SC marker *SOX10* mRNA (blue) and neuronal class III β -tubulin protein (red) expressions in iris (C) and corneal stroma and epithelium (D). (E, F) SC marker *PAX3* mRNA (blue) and neuronal class III β -tubulin protein (red) expressions in iris (E) and corneal stroma and epithelium (F). (G, H) SC marker *S100* mRNA (blue) and neuronal class III β -tubulin protein (red) expressions in iris (G) and corneal stroma and epithelium (H). (I, J) SC marker *MBP* mRNA (blue) and neuronal class III β -tubulin protein (red) expressions in iris (I) and corneal stroma and epithelium (J). Large white arrows: nerve segments in apparent contact with SC marker-expressing cells. Small white arrows: SC marker-expressing cells in apparent contact with nerve segments. Small black arrows: SC marker-expressing cells not in apparent contact with nerve segments. ep, epithelium; st, stroma. Bar: (A, C, E, G, I) 50 μ m; (B, D, F, H, J) 20 μ m.

Table 1

Genes Expressed by SCs at Different Stages of Differentiation

SC Differentiation Stage	Genes Expressed
SC precursors	<i>SOX10</i> <i>PAX3</i> <i>MPZ</i>
Immature SCs	<i>SOX10</i> <i>PAX3</i> <i>POU3F1</i> <i>S100</i> <i>GFAP</i> <i>MPZ</i>
Mature myelinating SCs	<i>SOX10</i> <i>EGR2</i> <i>S100</i> <i>MBP</i> <i>MPZ</i> <i>MAG</i>
Mature nonmyelinating SCs	<i>SOX10</i> <i>PAX3</i> <i>S100</i> <i>GFAP</i> β 1-Integrin

Real-Time PCR Primers for SC-Related Genes

Table 2

Symbol	Gene Name/gi number	Forward Primer	Reverse Primer
<i>GAPDH</i>	Glyceraldehyde-3-phosphate-dehydrogenase/gi: 4105595	5'-GCTGAGAACGGGAAACTTGTGA	5'-GCACCTGCATCTGCCCATTT
<i>BACE1</i>	Beta-site amyloid precursor protein-cleaving enzyme 1/gi: 45444680	5'-AGAGTATCGTGGACAGTG	5'-GTAGAGGGACAGAACAGG
<i>CADM1</i>	Cell adhesion molecule 1 (formerly Nect2)/gi: 118101954	5'-GTTCAACGGCTTTACTCAGTTG	5'-GGTCCACGCTCTCTATTTCG
<i>CADM2</i>	Cell adhesion molecule 2 (formerly Nect3)/gi: 118083731	5'-CCTTAAACAGCCAGCACAAAC-3'	5'-GACAACTACAGCCACTATTCC-3'
<i>CADM3</i>	Cell adhesion molecule 3 (formerly Nect1)/gi: 47022790	5'-CGGAGAGGGTGAGAAATGG	5'-CGAGAGTAAATCTGTGGGTGG
<i>CADM4</i>	Cell adhesion molecule 4 (formerly Nect4)/gi: 25827035	5'-ACTCCATCAGCAGAGCATTTG	5'-AGAGGAACCGCAGCATGAGG
<i>EGR2</i>	Early growth response 2 (Krox-20 homologue)/gi: 13430084	5'-AACCCAGACCTTCACCTAC	5'-CACGATGTTGATGATGCC
<i>GFAP</i>	Glial fibrillary acidic protein/gi: 50760636	5'-ACGGCGCAACCAACGAGAAG	5'-TCCAGCACCGACACCTTATTG
<i>ITGNA3</i>	Integrin alpha 3/gi: 45383819	5'-GAGCGACGAATGGCAGAC	5'-AATAGATGATGTTGGCAGTGAAAG
<i>ITGNA6</i>	Integrin alpha 6/gi: 46048713	5'-GGGTGTAACCTGTCCAAAG	5'-ACTCCATTATCCATATCATCC
<i>ITGNB1</i>	Integrin beta 1/gi: 86129417	5'-TTCCTCTGTTGCTTGATATGG	5'-GTTGGTTCTCTCTCTTTCG
<i>ITGNB4</i>	Integrin beta 4/gi: 118099661	5'-GAACATCAGCATCAACAC	5'-GAGATTATCCAAGTCATCAG
<i>MAG</i>	Myelin associated glycoprotein/gi: 245729	5'-GGCTTCTATCGGTGGTTTATG	5'-GTGAAACGGCGGATGATGG
<i>MBP</i>	Myelin basic protein/gi: 45384133	5'-CATTACTTAGTGCCCAATTAGGTTTC	5'-GTCAGTCGTCTGTTACTCTAC
<i>MPZL1</i>	Myelin protein zero-like 1/gi: 118083691	5'-TTGATGTCACCAAGTTAAGC	5'-AGCATAGACACACAGACTC
<i>MPZL3</i>	Myelin protein zero-like 3/gi: 118101994	5'-GTCCGGCTTTCCTCTTCCTC	5'-CGTAACCTCGCACCTCAG
<i>NEO1</i>	Neogenin 1/gi: 118095507	5'-GATACAGATGGAGGATTC	5'-TTAGTAAAGTGAAGATGGC
<i>NTN1</i>	Netrin 1/gi: 46048658	5'-AGGAGCGGTGATGGACTG	5'-CCCTCAATACGAGATGCC
<i>NTN2L</i>	Netrin 2-like/gi: 529420	5'-AAACCCCTTCGTGGCTCAG	5'-ATCGCAGTCCCGTGTGG
<i>NTN4</i>	Netrin 4/gi: 118082520	5'-CTGAACCCCAAGTCCTTAG	5'-GCTCCCAAATTCGCATTTC
<i>PAX3</i>	Paired box 3/gi: 45383597	5'-CCGTGCTAGATGGAGGAAAG	5'-TGGAGGTGGGCTGATAGG
<i>POU3F1</i>	POU class 3 homeobox 1: (formerly OCT6)/gi: 71896128	5'-CTTCTCGCAGACCAACCATC	5'-CTCCTCCAGCCACTTGTTC
<i>ROBO1</i>	Roundabout, axon guidance receptor homolog 1/gi: 15281514	5'-AATGTAGGCAAGAGGTG	5'-GATCATCAGAAATTCATATCTC
<i>ROBO2</i>	Roundabout, axon guidance receptor homolog 2/gi: 15281516	5'-AGAGAGAGAAAAGAAAGGGGTAG	5'-TCGGAAGAACAAAGGAGAGG
<i>SCMP</i>	Schwann cell myelin protein/gi: 245729	5'-GGCTTCTATCGGTGGTTTATG	5'-GTGAACCGCGGATGATGG
<i>S100</i>	S100 calcium binding protein/gi: 118093315	5'-CCTTGGCTACTTGAGACATTTC	5'-AAGCAGGAGGATTCATAGGC
<i>SLIT2</i>	Slit homolog 2/gi: 15281510	5'-GTTATCACACCTGGCTATTG	5'-TTAGATGGAGTTGTAAGTAGAAAG
<i>UNC5A</i>	Unc-5 homolog A/gi: 118097387	5'-CCTACGGCACCTTCAACTTC	5'-CGGGAGGGATCAGCAAAGC
<i>UNC5B</i>	Unc-5 homolog B/gi: 71895020	5'-GTCAATCAACAAGGCAGAG	5'-AATGGTGAGGACAAACAGG
<i>UNC5C</i>	Unc-5 homolog C/gi: 45383262	5'-ACTCGGATGATGTTGCTCTC	5'-AGTCACGGTGGTTCTTGC

Table 3

PCR Primers for SC-Related Gene In Situ Fragments

Symbol	Gene Name/gi Number	Forward Primer	Reverse Primer
<i>KERA</i>	Keratocan/gi: 45383763	5'-AGTCCATTCTGTGTGGACTCG	5'-TGTGATCAAGGTGAAGGTGC
<i>LUM</i>	Lumican/gi: 118082483	5'-TTGCCAGTATGACTATGGTCC	5'-TGTACATCTCCTGTGGCAGG
<i>SLIT2</i>	Slit homolog 2/gi: 15281510	5'-CTTCAGATTGCCACAGACGA	5'-ATGTCCTGCAGGCATTAC
<i>MPZL1</i>	Myelin protein zero-like 1/gi: 118083691	5'-AATGGAAAGCCGTACACTGG	5'-TATGCACATGCATTGGTCCT
<i>SOX10</i>	SRY (sex determining region Y)-related HMG-box gi: 6760468	5'-TGATGACCAAGATCTGTCCG	5'-TCGGTGTAATGACCTCCAGG
<i>PAX3</i>	Paired box 3/gi: 45383597	5'-TGCGTCTCCAAGATCCTCTG	5'-GCTGCTCTCTGGGTTTGAAG
<i>S100</i>	S100 calcium binding protein/gi: 118093315	5'-GCAGCGTGGCGTAGAAAC	5'-CCACAGAGGTCACAGGAGTC
<i>MBP</i>	Myelin basic protein/gi: 45384133	5'-ACAACCCTGTAGTCCACTTCTTC	5'-TCACATACCTTGAACCTCGCAATG