Published in final edited form as:

N Engl J Med. 2009 July 16; 361(3): 245–254. doi:10.1056/NEJMoa0809578.

Disclosure of *APOE* Genotype for Risk of Alzheimer's Disease

Robert C. Green, M.D., M.P.H., J. Scott Roberts, Ph.D., L. Adrienne Cupples, Ph.D., Norman R. Relkin, M.D., Ph.D., Peter J. Whitehouse, M.D., Ph.D., Tamsen Brown, M.S., Susan LaRusse Eckert, M.S., Melissa Butson, Sc.M., A. Dessa Sadovnick, Ph.D., Kimberly A. Quaid, Ph.D., Clara Chen, M.H.S., Robert Cook-Deegan, M.D., and Lindsay A. Farrer, Ph.D. for the REVEAL Study Group*

G. Annas,

Boston University School of Medicine, Boston

D. Bhatt,

Cleveland Clinic Foundation, Cleveland

B. Biesecker,

Johns Hopkins School of Public Health, Baltimore

D. Blacker,

Harvard School of Public Health, Boston

E. Cox,

Weill Cornell Medical College, New York

K. Christensen,

University of Michigan School of Public Health, Ann Arbor

J.G. Davis,

Weill Cornell Medical College, New York

G.-A. Fasaye,

Howard University, Washington, DC

P. Griffith,

Morehouse School of Medicine, Atlanta

S. Hiraki,

Boston University School of Medicine, Boston

E. Juengst,

Case Western Reserve University School of Medicine, Cleveland

J. Karlawish,

University of Pennsylvania School of Bioethics, Philadelphia

E. Linnenbringer,

University of Michigan, Ann Arbor

T. Marteau,

Kings College, London

Copyright © 2009 Massachusetts Medical Society.

Address reprint requests to Dr. Green at Boston University School of Medicine, 715 Albany St., L-320, Boston, MA 02118, or at rcgreen@bu.edu.

*Members of the Risk Evaluation and Education for Alzheimer's Disease (REVEAL) Study Group are listed in the Appendix.

No other potential conflict of interest relevant to this article was reported.

E. Neimeister,
Weill Cornell Medical College, New York

T. Obisesan,
Howard University, Washington, DC

R. Petersen,
Mayo Alzheimer's Disease Research Center, Rochester, MN

S. Post,
Case Western Reserve University School of Medicine, Cleveland

L. Ravdin,
Weill Cornell Medical College, New York

R.P. Reilly,
Shriver Center, Waltham, MA

D. Roter,
Johns Hopkins Bloomberg School of Public Health, Baltimore

C. Royal,
Duke University, Durham, NC

R. Stern,
Boston University School of Medicine, Boston

E. Topol,
Cleveland Clinic Lerner College of Medicine, Cleveland

W. Uhlmann,
University of Michigan, Ann Arbor

M. Williams, and
Morehouse School of Medicine, Atlanta

L. Wright
Medical College of Georgia, Athens

Boston University School of Medicine (R.C.G., T.B., L.A.F.), Boston University School of Public Health (R.C.G., L.A.C., C.C., L.A.F.), and Harvard Medical School Genetics Training Program (R.C.G.) — all in Boston; the University of Michigan School of Public Health, Ann Arbor (J.S.R.); Weill Cornell Medical College, New York (N.R.R.); Case Western Reserve University School of Medicine, Cleveland (P.J.W., M.B.); Columbia University School of Medicine, New York (S.L.E.); the University of British Columbia, Vancouver Hospital and Health Sciences Centre, Vancouver, BC, Canada (A.D.S.); Indiana University School of Medicine, Indianapolis (K.A.Q.); and Duke University, Durham, NC (R.C.-D.).

Abstract

Background—The apolipoprotein E (*APOE*) genotype provides information on the risk of Alzheimer's disease, but the genotyping of patients and their family members has been discouraged. We examined the effect of genotype disclosure in a prospective, randomized, controlled trial.

Methods—We randomly assigned 162 asymptomatic adults who had a parent with Alzheimer's disease to receive the results of their own *APOE* genotyping (disclosure group) or not to receive such results (nondisclosure group). We measured symptoms of anxiety, depression, and test-related distress 6 weeks, 6 months, and 1 year after disclosure or nondisclosure.

Results—There were no significant differences between the two groups in changes in time-averaged measures of anxiety (4.5 in the disclosure group and 4.4 in the nondisclosure group, $P = 0.84$), depression (8.8 and 8.7, respectively; $P = 0.98$), or test-related distress (6.9 and 7.5, respectively; $P = 0.61$). Secondary comparisons between the non-disclosure group and a disclosure subgroup of subjects carrying the *APOE* $\epsilon 4$ allele (which is associated with increased risk) also revealed no significant differences. However, the $\epsilon 4$ -negative subgroup had a significantly lower level of test-related distress than did the $\epsilon 4$ -positive subgroup ($P = 0.01$). Subjects with clinically meaningful changes in psychological outcomes were distributed evenly among the nondisclosure group and the $\epsilon 4$ -positive and $\epsilon 4$ -negative subgroups. Baseline scores for anxiety and depression were strongly associated with post-disclosure scores of these measures ($P < 0.001$ for both comparisons).

Conclusions—The disclosure of *APOE* genotyping results to adult children of patients with Alzheimer's disease did not result in significant short-term psychological risks. Test-related distress was reduced among those who learned that they were *APOE* $\epsilon 4$ -negative. Persons with high levels of emotional distress before undergoing genetic testing were more likely to have emotional difficulties after disclosure. (ClinicalTrials.gov number, NCT00571025.)

Knowledge of the results of genetic-susceptibility testing may cause anxiety, depression, and other types of distress. Nevertheless, gene variants that are associated with risks of common diseases are being rapidly discovered, and genetic testing is now marketed to consumers.¹⁻³ A variant of the gene *APOE*, which encodes apolipoprotein E, is associated with an increased susceptibility to Alzheimer's disease,⁴ and we hypothesized that testing for the presence of this gene would be useful for evaluating the effect of genetic-risk assessment.⁴ By consensus, *APOE* testing is not currently recommended for asymptomatic persons; a major concern is the emotional effect of risk disclosure.⁵⁻⁸

In the Risk Evaluation and Education for Alzheimer's Disease (REVEAL) study, we hypothesized that persons who learned about their *APOE* genotype through an education-and-disclosure protocol would not have greater symptoms of anxiety, depression, or test-related distress than those not receiving such information.

Methods

Study Population and Instruments

We recruited adult children of a living or deceased parent with Alzheimer's disease through self-referral or telephone calls to families in research registries.⁹ As part of the screening process, we interviewed the subjects and administered standardized tests to evaluate their cognitive ability, academic achievement, and levels of anxiety and depression. We excluded subjects who scored 1.3 SD below norms on the Repeatable Battery for the Assessment of Neuropsychological Status or the Wide Range Achievement Test 3; higher than 20 on the Beck Anxiety Inventory (BAI), which ranges from 0 to 63, with higher scores indicating greater anxiety; or higher than 26 on the Center for Epidemiological Studies Depression Scale (CES-D), which ranges from 0 to 60, with higher scores indicating greater depression. Ancestry was determined by self-report obtained through a multiple-choice question, with the following options: white, American Indian, Asian, black, and Hispanic; we prespecified the designation of "other" in the event that a subject selected more than one option.

The BAI^{10,11} is a 21-item scale designed to distinguish symptoms of anxiety from those of depression and to be sensitive to change. This test is based on self-reported severity of a given anxiety symptom during the previous week, with a clinical cutoff score of 16 (moderate anxiety). The CES-D^{12,13} measures depressive symptoms in studies of nonclinical populations, with a clinical cutoff score of 16 to 20.^{14,15} We estimated that 5-point differences on either the

BAI or the CES-D would be a sensitive indicator of clinically meaningful change and that with a sample of 46 persons per group, the study would have a statistical power of 80% to detect this difference on either scale. The Impact of Event Scale (IES) is a 15-item self-report instrument assessing test-related distress^{16,17} that is commonly used in genetics research. The scores range from 0 to 75, with higher scores indicating greater distress.¹⁸⁻²⁵ A total score of 20 to 40 may indicate significant distress, and a 5-point difference is a conservative measure of clinically meaningful change. We also developed a series of original questions to gauge changes in risk perception, positive and negative effects of the genetic disclosure, and whether the subject would make the same decision again to undergo genetic testing.

Study Design

The REVEAL study group — an interdisciplinary team of experts in clinical trials, neurology, genetics, genetic counseling, health psychology, biostatistics, and bioethics — designed the study, drawing on surveys conducted with relatives of patients with Alzheimer's disease.²⁶⁻²⁸ We created risk curves for the disclosure process that were specific for age and sex, showing the lifetime cumulative incidence of Alzheimer's disease and the remaining risk of Alzheimer's disease for each subject (cumulative incidence from current age to the age of 85 years).²⁹

The study was conducted from 2000 through 2003 at sites in Boston, Cleveland, and New York. It was approved by the institutional review board at each center and was coordinated by a board-certified or board-eligible genetic counselor at each site. Informed consent was obtained first by telephone, then in writing (Fig. 1, and the Supplementary Appendix, available with the full text of this article at NEJM.org). A 90-minute, semiscripted group session that was led by the genetic counselor described the limitations of *APOE* testing, the absence of a medical benefit of such testing, and the format for communication of the risk. All subjects later met individually with the genetic counselor for the drawing of blood samples, which were sent to Athena Diagnostics for *APOE* genotyping. Subjects were then randomly assigned to receive the genotyping results (the disclosure group) or not to receive the results (the nondisclosure group). Subjects in the nondisclosure group were individually shown two charts: one showing the incidence of Alzheimer's disease in the general population according to age and another showing the sex- and age-specific incidence of the disease among first-degree relatives of patients with Alzheimer's disease.²⁹ Subjects in the disclosure group were shown the same curves with an additional line for their genotype-specific risk (Fig. 1 in the Supplementary Appendix). They also received their lifetime cumulative incidence risk by the age of 85 years. Subjects were told their *APOE* genotype and were given written reports of their lifetime cumulative incidence risk and remaining incident risk.²⁹

Outcome Measures

The prespecified primary outcomes were changes in subjects' anxiety and depression symptoms, as measured by the BAI and the CES-D, respectively. The prespecified secondary outcome was test-related distress, as measured by the IES.

We administered the BAI and the CES-D before randomization and 6 weeks, 6 months, and 1 year after disclosure or nondisclosure of risk. We administered the IES at 6 weeks, 6 months, and 1 year. The primary analysis compared the two randomized groups. A secondary analysis compared the subgroup of subjects in the disclosure group who were informed that they carried at least one $\epsilon 4$ allele, which is associated with an increased risk of Alzheimer's disease (the $\epsilon 4$ -positive subgroup), with either a subgroup of subjects who were informed that they did not carry an $\epsilon 4$ allele (the $\epsilon 4$ -negative subgroup) or with the nondisclosure group.

Safety Monitoring

Throughout the study, genetics counselors monitored all subjects for adverse psychological effects. We created an independent external ethics and safety board to review the protocol, monitor study progress, and establish criteria for the reporting of adverse events to the institutional review board at each site. For example, subjects whose BAI or CES-D scores were more than 16 or increased by more than 15 points were immediately interviewed, with referral as appropriate. The chair of the ethics and safety board reviewed the results annually to screen for adverse or unanticipated events.

Statistical Analysis

We used two-sided t-tests or chi-square tests to compare baseline variables between the two initial randomization groups (the disclosure group and the nondisclosure group) and between the nondisclosure group and the $\epsilon 4$ -positive and $\epsilon 4$ -negative subgroups. We compared the rate of withdrawal from the study among the groups. Prespecified primary analyses compared scores on the BAI and the CES-D in the disclosure group with those in the nondisclosure group and included data from all time points, with the use of longitudinal analysis of mixed-effects models that were adjusted for age, sex, years of education, time, and baseline outcome score (if available). In the intention-to-treat analysis, missing values were calculated with the Markov chain Monte Carlo method of multiple imputation with the use of PROC MI statistical software, version 9.1 (SAS Institute).

To assess trends over time, we added interaction terms between group and time as covariates to longitudinal models. The model for the primary analysis was also run separately for each time point. This process was repeated for the IES scores. All three outcomes were then examined in the same manner for the two disclosure subgroups and the nondisclosure group. Although the study was not originally powered for equivalence testing, equivalence was demonstrated post hoc when a confidence interval for a group difference did not include 5 points in either direction.³⁰ We examined raw changes in scores on each outcome measure to calculate the percentage of subjects whose changes in scores exceeded clinically significant thresholds (Fig. 3 in the Supplementary Appendix).

Results

Subjects

Of the subjects who participated in the informational interview and educational session, 61% and 84%, respectively, underwent phlebotomy (Fig. 1). Twelve subjects withdrew from the study after phlebotomy and before randomization, and seven subjects with low neurocognitive scores and two subjects with high depression scores were excluded. The remaining 162 subjects were randomly assigned in a 2:1 ratio to either the disclosure group (111 subjects) or the nondisclosure group (51 subjects) (Table 1).

After randomization, 14 subjects withdrew from the study, citing study-related reasons. Of the variables listed in Table 1, only the baseline BAI score showed a trend toward an association with withdrawal from the study (i.e., those who were less anxious were more likely to withdraw; $P=0.07$). Some subjects in the nondisclosure group were dissatisfied at not receiving their genotyping results; of these subjects, 8 (16%) withdrew, as did 6 subjects (5%) in the disclosure group ($P=0.04$). Of the 53 subjects in the $\epsilon 4$ -positive subgroup, 3 (6%) withdrew from the study, as did 3 of 58 subjects (5%) in the $\epsilon 4$ -negative subgroup. Of the 111 subjects in the disclosure group, 3 (3%) had the $\epsilon 4/\epsilon 4$ genotype, 46 (41%) had the $\epsilon 3/\epsilon 4$ genotype, 53 (48%) had the $\epsilon 3/\epsilon 3$ genotype, 5 (4%) had the $\epsilon 2/\epsilon 3$ genotype, and 4 (4%) had the $\epsilon 2/\epsilon 4$ genotype; none of the subjects had the $\epsilon 2/\epsilon 2$ genotype. Subjects with the $\epsilon 4/\epsilon 4$ genotype were given a

higher risk estimate¹³ but were included in the $\epsilon 4$ -positive group in analyses. Data that were collected after randomization but before withdrawal were included in the analyses.

For the end points of scores on the BAI and the CES-D, adjusted group means for the disclosure and nondisclosure groups did not differ significantly either with the use of the time-averaged longitudinal model or at any individual time point (Table 2). Changes from baseline in time-averaged scores for anxiety and depression did not differ significantly between the two groups: for the anxiety score, 4.5 in the disclosure group and 4.4 in the nondisclosure group ($P=0.84$); and for the depression score, 8.8 and 8.7, respectively ($P=0.98$).

Interaction analyses indicated that between-group differences in scores were stable over time. We designed the prespecified analyses to allow detection of significant differences, and observing none, we then carried out a post hoc analysis for equivalence by examining 95% confidence intervals. All the confidence intervals excluded a difference of 5 points or more for scores on both the BAI (within 3 points) and the CES-D (within 2 points). Post-disclosure scores on the BAI and the CES-D were strongly associated with respective baseline scores on these measures ($P<0.001$ for both comparisons).

There were no significant differences between the nondisclosure group and the two disclosure subgroups ($\epsilon 4$ -positive and $\epsilon 4$ -negative) in the overall model on the BAI or the CES-D at any time point (Tables 3 and 4), with all 95% confidence intervals excluding a difference of 5 points or more. We observed no significant differences between the nondisclosure group and the $\epsilon 4$ -positive group over time using the longitudinal model or at any time point. Adjusted means were very similar to unadjusted means. (Tables show only adjusted means; raw means are shown in Figure 2 in the Supplementary Appendix.)

Adjusted mean IES scores for the disclosure and nondisclosure groups did not differ significantly at any time point or over time (6.9 in the disclosure group and 7.5 in the nondisclosure group, $P=0.61$). All 95% confidence intervals excluded clinically meaningful differences of 5 points or more, except for the scores at 6 months, which showed a trend toward less distress in the disclosure group. Adjusted IES scores in the $\epsilon 4$ -negative subgroup were lower than those in the nondisclosure group at 6 months ($P=0.01$), with a similar trend over time ($P=0.09$) (Table 4). A comparison of IES scores in the $\epsilon 4$ -negative group with those in the $\epsilon 4$ -positive group showed significant differences over time (4.8 in the $\epsilon 4$ -negative group vs. 8.5 in the $\epsilon 4$ -positive group, $P=0.01$) and at 6 weeks ($P=0.02$) and 6 months ($P=0.01$), with a marginally significant difference at 12 months ($P=0.06$).

On the IES, we could not demonstrate equivalence to within 5 points with 95% confidence at any time point for comparisons between the non-disclosure group and the $\epsilon 4$ -negative subgroup or between the $\epsilon 4$ -positive subgroup and the $\epsilon 4$ -negative subgroup. A comparison of the nondisclosure group with the $\epsilon 4$ -positive subgroup showed no significant differences over time or at any time point. A difference of 5 points could be excluded with 95% confidence at 6 months, 12 months, and over time but not at 6 weeks, when scores for distress were higher in the $\epsilon 4$ -positive subgroup. The results of all intention-to-treat analyses were similar to the results of analyses that included only subjects for whom scores at all time points were available (data not shown).

There was no significant change in our findings after adjustment for race, years of education, marital status, study site, referral source, family history, caregiver status, or mean age of the affected parent at the onset of disease. Lower overall BAI scores were associated with lower baseline BAI scores ($P<0.001$) and also with self-referral to the study ($P=0.002$). Lower overall CES-D scores were associated with lower baseline CES-D scores ($P<0.001$), with a younger age at the onset of the parent's symptoms of Alzheimer's disease ($P=0.003$), and with self-

referral ($P=0.008$). Lower overall IES scores were associated with male sex ($P=0.01$). No other covariates were significantly associated with outcome.

Subjects in the $\epsilon 4$ -positive group were more likely than those in either the nondisclosure group or the $\epsilon 4$ -negative subgroup to report a high perceived risk and an overall negative effect of learning their genotype (Table 1 in the Supplementary Appendix). This finding suggests that the subjects in the $\epsilon 4$ -positive group understood the information they received with respect to risk and that they had some negative feelings about receiving their results. Nevertheless, subjects in the $\epsilon 4$ -positive subgroup were no less likely than those in the $\epsilon 4$ -negative group to say that they would undergo testing again.

The distributions of changes in scores were similar among the two disclosure subgroups and the overall disclosure group at all time points (Fig. 3 in the Supplementary Appendix). Outcome scores, combined with prespecified safety criteria, triggered an assessment of adverse psychological effects in 13 subjects: 3 in the nondisclosure group, 4 in the $\epsilon 4$ -negative subgroup, and 6 in the $\epsilon 4$ -positive group (4 with the $\epsilon 3/\epsilon 4$ genotype and 2 with the $\epsilon 4/\epsilon 4$ genotype). None of the subjects attributed their psychological state to concern about disclosure but instead cited personal events or experiences, such as family illness and job-related stress, that were not related to the study.

Discussion

Subjects who were randomly assigned to undergo risk assessment with *APOE* disclosure did not have greater anxiety, depression, or test-related distress than those who were assigned to undergo risk assessment without *APOE* disclosure. Post hoc equivalence within 5 points was demonstrated at all visits over time for all outcomes, with the exception of the 6-month time point for the secondary outcome of the IES score, which showed a trend toward less distress in the disclosure group.

Subjects who learned they were $\epsilon 4$ -positive and were therefore at increased risk for Alzheimer's disease showed no more anxiety, depression, or test-related distress than those who did not learn their genotype, although strict equivalence could not be demonstrated for test-related distress at 6 weeks, which reflected, in part, transient test-related distress among $\epsilon 4$ -positive subjects at this time point. The comparison of subjects in the $\epsilon 4$ -positive subgroup with those in the $\epsilon 4$ -negative subgroup revealed significant but not clinically meaningful differences on test-related distress, driven by reduced distress among $\epsilon 4$ -negative subjects at 6 months and 12 months. The change in the IES score from baseline to 6 weeks also differed significantly between the $\epsilon 4$ -positive subgroup and the $\epsilon 4$ -negative subgroup, which we partly attributed to an increase in distress in the $\epsilon 4$ -positive group. On all outcome measures, mean scores in the two disclosure subgroups and the overall disclosure group were well below clinical thresholds for concern. Subjects with outcome scores above prespecified safety thresholds were evenly distributed among the nondisclosure group and the $\epsilon 4$ -positive and $\epsilon 4$ -negative subgroups, and these subjects did not cite genotype disclosure as contributing to their psychological distress.

Additional questions about risk perception and the effect of testing 6 weeks after disclosure suggested that subjects understood that their risk was higher or lower according to their genotype, and they had the expected negative or positive feelings about this news. Thus, subjects were not immune to the negative implications of learning that they had an increased risk, but these feelings were not associated with clinically significant psychological distress.

These data support the psychological safety of disclosing data regarding genetic-counseling protocols to screened adult children of patients with Alzheimer's disease who request such information, despite the frightening nature of the disease and the fact that disclosure has no clear medical benefit. Larger studies that follow subjects for more than 1 year will be required

to detect uncommon and long-term effects, such as delayed emotional repercussions and injudicious life decisions.

APOE is the most robust risk marker available for Alzheimer's disease⁴; it is also associated with impaired memory among persons without dementia³¹ and with progression to Alzheimer's disease among persons with mild cognitive impairment.³² Surveys³³⁻³⁵ indicate that the public is interested in *APOE* genotyping and that 15% of primary care physicians who treat patients with Alzheimer's disease have already received requests for genotyping. If *APOE* genotyping is discovered to predict treatment efficacy or a risk of side effects, the level of interest is likely to increase.

Although visions of personalized medicine suggest that genetic risk markers will empower patients to improve their health through preventive practices and early interventions,³⁶ there is concern that the understanding of risk among both the lay public and medical professionals is exceedingly poor,³⁷ that genetic tests offering probabilistic estimates for risks of common diseases in the absence of a family history or environmental risks may be misunderstood, and that the psychological harm of such misunderstanding may outweigh the benefits, particularly with disorders such as Alzheimer's disease, for which no medical interventions are available.³⁸⁻⁴⁰ These concerns are amplified by the recent emergence of companies offering direct-to-consumer genetic testing, with most of them evaluating single-nucleotide polymorphisms that indicate the *APOE* genotype,¹⁻³ in the absence of guidelines for deciding which associations between genes and disease have sufficient clinical validity and usefulness to justify disclosure and with no gauge of the effect of such disclosure. Caution is thus warranted, and empirical data are valuable.

Our study is limited in that it concerns a single polymorphism that is robustly associated with Alzheimer's disease. In addition, most of the subjects were of fairly homogeneous and similar ancestry. If *APOE* genotyping had been provided without genetic counseling or to subjects who had no family history of Alzheimer's disease, the results might have been different. In addition, the exclusion of subjects with low neurocognitive scores and high depression scores may have influenced the results. Within these constraints, our results suggest that disclosure of genotyping information provides a benefit to those who are negative for a susceptibility variant and causes transient, modest distress to those with a positive result.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Supported by grants (R01-HG/AG02213, to Dr. Green; and R01-AG09029, to Dr. Farrer) from the Ethical, Legal and Social Implications (ELSI) Branch of the National Human Genome Research Institute and the National Institute on Aging (NIA); by an NIA Mentoring Award (K24 AG027841, to Dr. Green); by a grant (P30-AG13846) from the Boston University Alzheimer's Disease Center; by a grant (M01-RR00533) from the Boston University General Clinical Research Center; and by a grant (P50-HG003391, to Duke University) from the Centers of Excellence in ELSI Research.

Drs. Relkin and Farrer report receiving consulting fees from Smart Genetics, which provided direct-to-consumer *APOE* genotyping from March 2008 through October 2008, when it ceased operations.

We thank Joe Massaro, Ph.D., Suzanne Hendrix, Ph.D., and Steve Piantadosi, Ph.D, for their helpful comments on the manuscript.

Appendix

Members of the REVEAL study group are as follows: G. Annas, Boston University School of Medicine, Boston; D. Bhatt, Cleveland Clinic Foundation, Cleveland; B. Biesecker, Johns Hopkins School of Public Health, Baltimore; D. Blacker, Harvard School of Public Health, Boston; E. Cox, Weill Cornell Medical College, New York; K. Christensen, University of Michigan School of Public Health, Ann Arbor; J.G. Davis, Weill Cornell Medical College, New York; G.-A. Fasaye, Howard University, Washington, DC; P. Griffith, Morehouse School of Medicine, Atlanta; S. Hiraki, Boston University School of Medicine, Boston; E. Juengst, Case Western Reserve University School of Medicine, Cleveland; J. Karlawish, University of Pennsylvania School of Bioethics, Philadelphia; E. Linnenbringer, University of Michigan, Ann Arbor; T. Marteau, Kings College, London; E. Neimeister, Weill Cornell Medical College, New York; T. Obisesan, Howard University, Washington, DC; R. Petersen, Mayo Alzheimer's Disease Research Center, Rochester, MN; S. Post, Case Western Reserve University School of Medicine, Cleveland; L. Ravdin, Weill Cornell Medical College, New York; R.P. Reilly, Shriver Center, Waltham, MA; D. Roter, Johns Hopkins Bloomberg School of Public Health, Baltimore; C. Royal, Duke University, Durham, NC; R. Stern, Boston University School of Medicine, Boston; E. Topol, Cleveland Clinic Lerner College of Medicine, Cleveland; W. Uhlmann, University of Michigan, Ann Arbor; M. Williams, Morehouse School of Medicine, Atlanta; L. Wright, Medical College of Georgia, Athens.

References

1. Wolfberg AJ. Genes on the Web — direct-to-consumer marketing of genetic testing. *N Engl J Med* 2006;355:543–5. [PubMed: 16899772]
2. Hunter DJ, Khoury MJ, Drazen JM. Letting the genome out of the bottle — will we get our wish? *N Engl J Med* 2008;358:105–7. [PubMed: 18184955]
3. Offit K. Genomic profiles for disease risk: predictive or premature? *JAMA* 2008;299:1353–5. [PubMed: 18349097]
4. Farrer LA, Cupples LA, Haines JL, et al. Effects of age, sex and ethnicity on the association between apolipoprotein E genotype and Alzheimer disease: a meta-analysis. *JAMA* 1997;278:1349–56. [PubMed: 9343467]
5. Farrer LA, Brin MF, Elsas L, et al. Statement on use of apolipoprotein E testing for Alzheimer disease. *JAMA* 1995;274:1627–9. [PubMed: 7474250]
6. Post SG, Whitehouse PJ, Binstock RH, et al. The clinical introduction of genetic testing for Alzheimer disease: an ethical perspective. *JAMA* 1997;277:832–6. [PubMed: 9052715]
7. McConnell LM, Koenig BA, Greely HT, Raffin TA. Genetic testing and Alzheimer disease: has the time come? *Nat Med* 1998;4:757–9. [PubMed: 9662356]
8. Relkin NR, Tanzi R, Breitner J, et al. Apolipoprotein E genotyping in Alzheimer's disease. *Lancet* 1996;347:1091–5. [PubMed: 8602063]
9. Roberts JS, Barber M, Brown TM, et al. Who seeks genetic susceptibility testing for Alzheimer's disease: findings from a multisite, randomized clinical trial. *Genet Med* 2004;6:197–203. [PubMed: 15266207]
10. Beck, AT.; Steer, RA. Beck Depression Inventory. Psychological Corporation; New York: 1987.
11. Beck AT, Epstein N, Brown G, Steer RA. An inventory for measuring clinical anxiety: psychometric properties. *J Consult Clin Psychol* 1988;56:893–7. [PubMed: 3204199]
12. Radloff LS. The CES-D Scale: A self report depression scale for research in the general population. *Appl Psychol Meas* 1977;1:385–401.
13. Santor DA, Zuroff DC, Ramsay JO, Cervantes P, Palacios J. Examining scale discriminability in the BDI and CES-D as a function of depressive severity. *Psychol Assess* 1995;7:131–9.
14. Vernon SW, Gritz ER, Peterson SK, et al. Correlates of psychologic distress in colorectal cancer patients undergoing genetic testing for hereditary colon cancer. *Health Psychol* 1997;16:73–86. [PubMed: 9028817]

15. Lerman D, Narod S, Schulman K, et al. BRCA1 testing in families with hereditary breast-ovarian cancer: a prospective study of patient decision making and outcomes. *JAMA* 1996;275:1885–92. [PubMed: 8648868]
16. Horowitz M, Wilner N, Alvarez MA. Impact of Event Scale: a measure of subjective stress. *Psychosom Med* 1979;41:209–18. [PubMed: 472086]
17. Zilberg NJ, Weiss DS, Horowitz MJ. Impact of Event Scale: a cross-validation study and some empirical evidence supporting a conceptual model of stress response syndromes. *J Consult Clin Psychol* 1982;50:407–14. [PubMed: 7096742]
18. Payne K, Nicholls S, McAllister M, Macleod R, Donnai D, Davies LM. Outcome measurement in clinical genetics services: a systematic review of validated measures. *Value Health* 2008;11:497–508. [PubMed: 18489673]
19. DudokdeWit AC, Tibben A, Duivenvoorden HJ, Niermeijer MF, Passchier J. Predicting adaptation to presymptomatic DNA testing for late onset disorders: who will experience distress? *J Med Genet* 1998;35:745–54. [PubMed: 9733033]
20. Michie S, Bobrow M, Marteau TM. Predictive genetic testing in children and adults: a study of emotional impact. *J Med Genet* 2001;38:519–26. [PubMed: 11483640]
21. Thewes B, Meiser B, Hickie IB. Psycho-metric properties of the Impact of Event Scale amongst women at increased risk for hereditary breast cancer. *Psychooncology* 2001;10:459–68. [PubMed: 11747058]
22. Meiser B, Butow P, Friedlander M, et al. Psychological impact of genetic testing in women from high-risk breast cancer families. *Eur J Cancer* 2002;38:2025–31. [PubMed: 12376208]
23. Schwartz MD, Peshkin BN, Hughes C, Main D, Isaacs C, Lerman C. Impact of BRCA1/BRCA2 mutation testing on psychologic distress in a clinic-based sample. *J Clin Oncol* 2002;20:514–20. [PubMed: 11786581]
24. Smith CO, Lipe HP, Bird TD. Impact of presymptomatic genetic testing for hereditary ataxia and neuromuscular disorders. *Arch Neurol* 2004;61:875–80. [PubMed: 15210524]
25. Meiser B, Collins V, Warren R, et al. Psychological impact of genetic testing for hereditary non-polyposis colorectal cancer. *Clin Genet* 2004;66:502–11. [PubMed: 15521977]
26. Roberts JS. Anticipating response to predictive genetic testing for Alzheimer's disease: a survey of first-degree relatives. *Gerontologist* 2000;40:43–52. [PubMed: 10750312]
27. Roberts JS, Connell CM, Cisewski D, Hips YG, Demissie S, Green RC. Differences between African Americans and whites in their perceptions of Alzheimer disease. *Alzheimer Dis Assoc Disord* 2003;17:19–26. [PubMed: 12621316]
28. Hips YG, Roberts JS, Farrer LA, Green RC. Differences between African Americans and whites in their attitudes toward genetic testing for Alzheimer's disease. *Genet Test* 2003;7:39–44. [PubMed: 12820701]
29. Cupples LA, Farrer L, Sadovnick D, Relkin N, Whitehouse P, Green RC. Estimating risk curves for first-degree relatives of patients with Alzheimer's disease: the REVEAL study. *Genet Med* 2004;6:192–6. [PubMed: 15266206]
30. Jones B, Jarvis P, Lewis JA, Ebbutt AF. Trials to assess equivalence: the importance of rigorous methods. *BMJ* 1996;313:36–9. [PubMed: 8664772][Erratum, *BMJ* 1996;313:550.]
31. Small BJ, Rosnick CB, Fratiglioni L, Bäckman L. Apolipoprotein E and cognitive performance: a meta-analysis. *Psychol Aging* 2004;19:592–600. [PubMed: 15584785]
32. Petersen RC. Mild cognitive impairment: current research and clinical implications. *Semin Neurol* 2007;27:22–31. [PubMed: 17226738]
33. Roberts JS, LaRusse SA, Katzen H, et al. Reasons for seeking genetic susceptibility testing among first-degree relatives of people with Alzheimer disease. *Alzheimer Dis Assoc Disord* 2003;17:86–93. [PubMed: 12794385]
34. Neumann PJ, Hammitt JK, Mueller C, et al. Public attitudes about genetic testing for Alzheimer's disease. *Health Aff (Millwood)* 2001;20(5):252–64. [PubMed: 11558711]
35. Chase GA, Geller G, Havstad SL, Holtzman NA, Bassett SS. Physicians' propensity to offer genetic testing for Alzheimer's disease: results from a survey. *Genet Med* 2002;4:297–303. [PubMed: 12172396]

36. Collins FS, Green ED, Guttmacher AE, Guyer MS. A vision for the future of genomics research. *Nature* 2003;422:835–47. [PubMed: 12695777]
37. Julian-Reynier C, Welkenhuysen M, Hagoel L, Decruyenaere M, Hopwood P. Risk communication strategies: state of the art and effectiveness in the context of cancer genetic services. *Eur J Hum Genet* 2003;11:725–36. [PubMed: 14512961]
38. Khoury MJ, McCabe LL, McCabe ERB. Population screening in the age of genomic medicine. *N Engl J Med* 2003;348:50–8. [PubMed: 12510043]
39. Haga SB, Khoury MJ, Burke W. Genomic profiling to promote a healthy lifestyle: not ready for prime time. *Nat Genet* 2003;34:347–50. [PubMed: 12923535]
40. Burke W, Psaty BM. Personalized medicine in the era of genomics. *JAMA* 2007;298:1682–4. [PubMed: 17925520]

Figure 1. Enrollment and Outcomes

Of the 21 subjects who were excluded before randomization, 7 had low neurocognitive scores, and 2 had high depression scores. The other 12 subjects withdrew from the study for other reasons.

Table 1

Characteristics of the Subjects at Baseline.*

Characteristic	Randomization Groups (N = 162)		Disclosure Subgroups (N = 111)	
	Nondisclosure Group (N = 51)	Disclosure Group (N = 111)	$\epsilon 4$ -Positive (N = 53)	$\epsilon 4$ -Negative (N = 58)
Age — yr				
Mean	55.3±9.0	52.0±10.0	50.4±8.6	53.5±11.0
Range	37–78	30–76	34–72	30–76
Female sex — no. (%)	40 (78)	77 (69)	41 (77)	36 (62)
White race — no. (%) [†]	46 (90)	106 (96)	50 (94)	56 (97)
Education — yr				
Mean	16.8±2.5	16.7±2.2	16.8±2.1	16.6±2.3
Range	10–22	12–22	12–21	12–22
Currently married — no. (%)	31 (61)	74 (67)	39 (74)	35 (60)
BAI score [‡]	4.6±4.5	4.2±5.0	3.5±4.3	4.8±5.5
CES-D score [§]	5.2±4.9	6.7±5.6	5.5±4.3	7.8±6.3
Site — no. (%)				
Boston	17 (33)	39 (35)	17 (32)	22 (38)
New York	18 (35)	37 (33)	16 (30)	21 (36)
Cleveland	16 (31)	35 (32)	20 (38)	15 (26)
Self-referred to study — no. (%)	35 (69)	80 (72)	37 (70)	43 (74)
More than 1 relative with Alzheimer's disease — no. (%)	28 (55)	70 (63)	37 (70)	33 (57)
Caregiver for relative with Alzheimer's disease — no. (%) [¶]	40 (78)	82 (75)	35 (67)	47 (81)
Age at onset of Alzheimer's disease in affected parent — yr				
Mean	74.6±9.0	70.8±8.6	68.8±7.7	72.8±9.0
Range	55–90	55–91	55–83	55–91

* Plus-minus values are means ±SD.

[†] Race was self-reported.[‡] Scores on the Beck Anxiety Inventory (BAI) range from 0 to 63, with higher scores indicating greater anxiety.[§] Scores on the Center for Epidemiological Studies Depression Scale (CES-D) range from 0 to 60, with higher scores indicating greater depression.[¶] In this category, the denominators were 110 in the disclosure group and 52 in the $\epsilon 4$ -positive subgroup.^{||} The age at onset was known for 44 (86%) of subjects in the nondisclosure group, 97 (87%) in the disclosure group, 48 (91%) in the $\epsilon 4$ -positive subgroup, and 49 (84%) in the $\epsilon 4$ -negative subgroup.

Table 2

Measures of Anxiety, Depression, and Test-Related Distress in the Randomized Groups.*

Measure	Nondisclosure Group (N = 51)	Disclosure Group (N = 111)	Difference between Disclosure Group and Nondisclosure Group (95% CI)	Adjusted P Value
BAI score [†]				
Longitudinal model (time-averaged)	4.4±0.5	4.5±0.3	0.1 (−1.0 to 1.2)	0.84
At 6 wk	4.4±0.7	4.8±0.4	0.4 (−1.2 to 2.0)	0.60
At 6 mo	4.6±0.6	4.2±0.4	−0.4 (−1.8 to 1.0)	0.58
At 12 mo	4.2±0.6	4.3±0.4	0.1 (−1.3 to 1.6)	0.88
CES-D score [‡]				
Longitudinal model (time-averaged)	8.7±0.8	8.8±0.5	0.1 (−1.7 to 1.8)	0.98
At 6 wk	9.3±1.0	8.7±0.7	−0.6 (−3.0 to 1.8)	0.62
At 6 mo	8.7±1.0	9.2±0.7	0.5 (−1.8 to 2.9)	0.66
At 12 mo	8.0±0.9	8.4±0.6	0.4 (−1.8 to 2.6)	0.74
IES score [§]				
Longitudinal model (time-averaged)	7.5±1.1	6.9±0.7	−0.7 (−3.3 to 2.0)	0.61
At 6 wk	6.8±1.4	7.2±0.9	0.4 (−2.9 to 3.8)	0.80
At 6 mo	8.9±1.3	6.3±0.9	−2.6 (−5.7 to 0.5)	0.10
At 12 mo	7.7±1.5	6.7±0.9	−1.0 (−4.2 to 2.2)	0.54

* Plus-minus values are means ±SE. Scores were adjusted for age, sex, years of education, and baseline score if it was available (there was no baseline score for the measure of test-related distress). Missing values were calculated with the Markov chain Monte Carlo method of multiple imputation.

[†] Scores on the Beck Anxiety Inventory (BAI) range from 0 to 63, with higher scores indicating greater anxiety.

[‡] Scores on the Center for Epidemiological Studies Depression Scale (CES-D) range from 0 to 60, with higher scores indicating greater depression.

[§] Scores on the Impact of Event Scale (IES) range from 0 to 75, with higher scores indicating greater distress.

Table 3

Measures of Anxiety, Depression, and Test-Related Distress in the Nondisclosure Group and in the Two Disclosure Subgroups with Known *APOE* $\epsilon 4$ Status.*

Measure	Nondisclosure Group (N = 51)	$\epsilon 4$ -Positive Subgroup (N = 53)	$\epsilon 4$ -Negative Subgroup (N = 58)
BAI score [†]			
At 6 wk	4.4±0.7	5.2±0.7	4.5±0.6
At 6 mo	4.6±0.6	4.6±0.6	3.9±0.6
At 12 mo	4.2±0.6	4.4±0.6	4.2±0.6
CES-D score [‡]			
At 6 wk	9.3±1.0	9.0±1.0	8.5±0.9
At 6 mo	8.7±1.0	9.6±1.0	8.9±1.0
At 12 mo	8.0±0.9	8.3±0.9	8.5±0.9
IES score [§]			
At 6 wk	6.7±1.4	9.4±1.3	5.2±1.3
At 6 mo	8.9±1.3	8.6±1.2	4.2±1.2
At 12 mo	7.7±1.5	8.5±1.3	5.1±1.2

* Plus-minus values are means ±SE. Scores were adjusted for age, sex, years of education, and baseline score if it was available (there was no baseline score for the measure of test-related distress). Missing values were calculated with the Markov chain Monte Carlo method of multiple imputation.

[†] Scores on the Beck Anxiety Inventory (BAI) range from 0 to 63, with higher scores indicating greater anxiety.

[‡] Scores on the Center for Epidemiological Studies Depression Scale (CES-D) range from 0 to 60, with higher scores indicating greater depression.

[§] Scores on the Impact of Event Scale (IES) range from 0 to 75, with higher scores indicating greater distress.

Table 4

Differences in Measures of Anxiety, Depression, and Test-Related Distress among the Nondisclosure Group and the Two Disclosure Subgroups with Known *APOE* $\epsilon 4$ Status.*

Measure [†]	Difference between $\epsilon 4$ -Negative Group and Nondisclosure Group (95% CI) P Value	Difference between $\epsilon 4$ -Positive Group and Nondisclosure Group (95% CI) P Value	Difference between $\epsilon 4$ -Positive Group and $\epsilon 4$ -Negative Group (95% CI) P Value
BAI score [‡]			
Longitudinal model			
(time-averaged)	-0.1 (-1.4 to 1.1)	0.4 (-0.9 to 1.7)	0.5 (-0.7 to 1.7)
At 6 wk	0.1 (-1.8 to 2.0)	0.8 (-1.1 to 2.7)	0.7 (-1.1 to 2.5)
At 6 mo	-0.7 (-2.4 to 0.9)	-0.02 (-1.7 to 1.7)	0.7 (-0.9 to 2.3)
At 12 mo	0.02 (-1.6 to 1.7)	0.2 (-1.5 to 1.9)	0.2 (-1.4 to 1.8)
CES-D score [‡]			
Longitudinal model			
(time-averaged)	-0.1 (-2.1 to 1.9)	0.1 (-1.9 to 2.2)	0.2 (-1.7 to 2.2)
At 6 wk	-0.8 (-3.6 to 1.9)	-0.3 (-3.2 to 2.5)	0.5 (-2.2 to 3.2)
At 6 mo	0.2 (-2.6 to 3.0)	0.9 (-2.0 to 3.7)	0.7 (-2.1 to 3.4)
At 12 mo	0.5 (-2.0 to 3.0)	0.2 (-2.3 to 2.8)	-0.2 (-2.7 to 2.3)
IES score [§]			
Longitudinal model			
(time-averaged)	-2.5 (-5.5 to 0.4)	1.4 (-1.7 to 4.4)	3.7 (1.0 to 6.5)
At 6 wk	-1.6 (-5.3 to 2.2)	2.7 (-1.2 to 6.5)	4.2 (0.6 to 7.8)
At 6 mo	-4.7 (-8.2 to -1.1)	-0.3 (-3.9 to 3.2)	4.3 (0.9 to 7.8)
At 12 mo	-2.6 (-6.4 to 1.2)	0.8 (-3.1 to 4.7)	3.4 (-0.2 to 7.0)

* Mean scores were adjusted for age, sex, years of education, and baseline score if it was available (there was no baseline score for the measure of test-related distress). Missing values were calculated with the Markov chain Monte Carlo method of multiple imputation.

[†] Scores on the Beck Anxiety Inventory (BAI) range from 0 to 63, with higher scores indicating greater anxiety.

[‡] Scores on the Center for Epidemiological Studies Depression Scale (CES-D) range from 0 to 60, with higher scores indicating greater depression.

[§] Scores on the Impact of Event Scale (IES) range from 0 to 75, with higher scores indicating greater distress.