Published in final edited form as:

Microb Pathog. 2007 ; 43(5-6): 198–207. doi:10.1016/j.micpath.2007.05.007.

γ -Glutamyl transpeptidase has a role in the persistent colonization of the avian gut by *Campylobacter jejuni*

If H.A. Barnes^{a,b}, Mary C. Bagnall^a, Darren D. Browning^b, Stuart A. Thompson^b, Georgina Manning^a, and Diane G. Newell^{a,*}

^aVeterinary Laboratories Agency (Weybridge), Woodham Lane, New Haw, Addlestone, Surrey, KT15 3NB, UK

^bDepartment of Biochemistry & Molecular Biology, Medical College of Georgia, 1120 15th Street, Augusta, GA 30912-2100, USA

Abstract

The contribution of γ -glutamyl transpeptidase (GGT) to *Campylobacter jejuni* virulence and colonization of the avian gut has been investigated. The presence of the *ggt* gene in *C. jejuni* strains directly correlated with the expression of GGT activity as measured by cleavage and transfer of the γ -glutamyl moiety. Inactivation of the monocistronic *ggt* gene in *C. jejuni* strain 81116 resulted in isogenic mutants with undetectable GGT activity; nevertheless, these mutants grew normally *in vitro*. However, the mutants had increased motility, a 5.4-fold higher invasion efficiency into INT407 cells *in vitro* and increased resistance to hydrogen peroxide stress. Moreover, the apoptosis-inducing activity of the *ggt* mutant was significantly lower than that of the parental strain. *In vivo* studies showed that, although GGT activity was not required for initial colonization of 1-day-old chicks, the enzyme was required for persistent colonization of the avian gut.

Keywords

Campylobacter jejuni; γ -glutamyl transpeptidase (*ggt*); Avian colonization

1. Introduction

Campylobacter jejuni is found in the intestinal tracts of a large number of food-producing animals, but appears to be adapted to the avian gut [1–3]. Colonization with *C. jejuni* in poultry is asymptomatic. However, infection with this organism in humans is a common cause of acute bacterial enteritis particularly in industrialized countries [4,5]. It is commonly assumed that the handling of contaminated poultry carcasses, and the consumption of undercooked poultry meat, are major sources of campylobacteriosis.

Quantitative risk assessment has demonstrated that the control and prevention of human campylobacteriosis will be best achieved by intervention to reduce or eradicate colonization

© 2007 Elsevier Ltd. All rights reserved.

*Corresponding author. Tel.: +44 1932357547; fax: +44 1932357268. d.newell@vla.defra.gsi.gov.uk (D.G. Newell).

Publisher's Disclaimer: This article was published in an Elsevier journal. The attached copy is furnished to the author for non-commercial research and education use, including for instruction at the author's institution, sharing with colleagues and providing to institution administration. Other uses, including reproduction and distribution, or selling or licensing copies, or posting to personal, institutional or third party websites are prohibited. In most cases authors are permitted to post their version of the article (e.g. in Word or Tex form) to their personal website or institutional repository. Authors requiring further information regarding Elsevier's archiving and manuscript policies are encouraged to visit: <http://www.elsevier.com/copyright>

at the poultry flock level [6]. Although improved biosecurity may contribute to this, it seems likely that complementary methods, such as vaccination or decontamination, may also be required [2,7]. Such targeted intervention strategies require fundamental understanding of those bacterial factors associated with the campylobacter properties enabling colonization and survival. To date approaches to the identification of such factors have largely been pragmatic with genes selected for characterization on the basis of previously published observations in other enteric bacteria. However, with the availability of the genome sequence of the *C. jejuni* reference strain NCTC11168 [8], post-genomic approaches are now being adopted [9–11]. In one such approach genomic subtractive hybridization between two strains, 81116 and NCTC11168 was undertaken to identify genes present in 81116 but absent in NCTC11168 [11], which is poorly able to colonise chickens. More recent studies have shown that the ability of the NCTC11168 isolate to colonize birds is highly dependent on the variant used [9,10]. Thus colonization is not only due to the presence or absence of genes but also whether those genes are transcribed. Nevertheless, given the difference in colonization potential between strains 81116 and NCTC11168, it has been speculated that such novel genes could contribute to aspects of colonization in chicks [12–16]. One such novel gene was a 369 bp DNA fragment (insert 236), unique to strain 81116, and which was identified and predicted to be part of a *C. jejuni* γ -glutamyl transpeptidase (*ggt*) gene [11].

The enzyme γ -glutamyl transpeptidase (GGT) is present in both prokaryotes and eukaryotes. Most of our knowledge is based on studies in mammalian tissues where GGT is a membrane-bound enzyme that plays a major role in glutathione (γ -glutamyl-L-cysteinylglycine) degradation in the γ -glutamyl cycle [17]. GGT cleaves and transfers the γ -glutamyl moiety from glutathione to amino acids or peptides. Glutathione is an antioxidant molecule, which plays an important part in providing vital cellular protection against the reactive oxygen species (ROS), such as hydrogen peroxide, generated by aerobic respiration [18]. Thus in eukaryotes, GGT-dependent breakdown of glutathione aids maintenance of cellular glutathione levels and increased cellular resistance to hydrogen peroxide-induced injury [19]. Not surprisingly, eukaryotic cells, once depleted of glutathione, have an increased susceptibility to oxidant-mediated killing [20,21]. Therefore, GGT appears to have an important role in combating oxidative stress.

A number of GGT-encoding genes from both mammalian and bacterial species have been sequenced and share extensive amino acid homology. However, there are two major taxon-associated differences. Firstly, the N-termini of the bacterial GGTs are signal peptides, and consequently the enzyme is thought to be either periplasmic or associated with the inner membrane [22]. In contrast, the N-termini of mammalian GGTs are anchor domains, which interact with plasma membranes [17,23]. Secondly, mammalian GGT is glycosylated [17]. In contrast bacterial GGTs are not known to be glycosylated. However, *C. jejuni* is unusual in that protein glycosylation pathways are active [24] and therefore the *C. jejuni* GGT could be glycosylated. Both bacterial and mammalian GGTs are first translated as precursor proteins, which in prokaryotes then undergo two proteolytic cleavages, the first of which results in cleavage of the signal peptide and the second processes the GGT to form the two subunits of the enzyme.

In *Helicobacter pylori*, GGT is involved in colonization of the gastric mucosa of mice [25–27], potentially by participating in the de novo synthesis of essential amino acids and thus enabling survival *in vivo*. In addition, GGT also possesses an apoptosis-inducing activity [28] and upregulates COX-2 and EGF-related peptide expression in human gastric cells [29]. Interestingly, in *Neisseria meningitidis*, GGT appears to provide an advantage for bacterial multiplication during environmental cysteine shortage by supplying cysteine from environmental peptides [30]. GGT is also involved in osmoadaptation in *Escherichia coli* [31].

In this study, the *ggt* gene from *C. jejuni* strain 81116 has been identified, fully sequenced and an isogenic mutant generated. The effect of this mutation on GGT activity and bacterial growth rate, motility, resistance to hydrogen peroxide stress, rate of internalization into INT407 cells, apoptosis and avian gut colonization, have been investigated.

2. Materials and methods

2.1. Bacterial strains

C. jejuni strain 81116 was isolated from a patient with diarrhea during a water-borne outbreak [32]. Strains NCTC11168 and 81-176 were both derived from human cases of campylobacteriosis [8,33]. Other isolates included in this investigation were randomly selected from the collection of *C. jejuni* strains held at the Veterinary Laboratories Agency (Weybridge), including isolates from the faeces of humans ($n = 32$), broilers ($n = 34$), cattle ($n = 32$), pigs ($n = 15$), sheep ($n = 18$), ostrich ($n = 1$), and dog ($n = 6$), as well as isolates from in and around poultry broiler houses ($n = 4$).

2.2. Media and growth conditions

All *C. jejuni* strains were cultured on 10% (vol/vol) sheep blood agar plates containing 250 μg actidione ml^{-1} and Skirrow's supplement (10 μg of vancomycin ml^{-1} , 2.5 IU of polymyxin B ml^{-1} , and 5 μg of trimethoprim) (BASA) [34] at 42°C in a microaerobic environment (7.5% [vol/vol]-CO₂, 7.5% [vol/vol] O₂, and 85% [vol/vol] N₂) for 20 or 40 h. *E. coli* strain TOPO10F' (Invitrogen) was grown on Luria-Bertani (LB) (Difco) agar or LB broth at 37°C under atmospheric conditions. When required, the media was supplemented with 100 μg ampicillin ml^{-1} or 50 μg kanamycin ml^{-1} . *C. jejuni* and *E. coli* were stored as frozen cultures at -80°C in 1% (w/v) proteose peptone water (Difco) containing 10% (v/v) glycerol or LB broth containing 50% (v/v) glycerol [35], respectively.

For assessing growth rates, strains were grown in biphasic broths (100 ml of brain heart infusion agar with 1% (w/v) yeast extract (BHI/YE) overlaid with 200 ml of BHI/YE broth). Viable counts of bacteria were taken after 0, 4, 7, 22, 30 and 50 h of growth.

2.3. PCR and DNA sequencing

PCR amplifications were performed using HotStarTaq master mix (Qiagen) as recommended by the manufacturer. For colony PCR, a loopful of freshly grown bacteria was suspended in 200 μl of water and 1 μl of the suspension was directly used (without prior boiling) as template with HotStarTaq (Qiagen). Degenerate PCR and chromosomal walking were performed as previously described [36,37]. Sequencing was carried out on duplicate samples with BigDye Terminator mix (Applied Biosystems) according to the manufacturer's instructions, the sequences analyzed using an ABI 3700 DNA sequencer (Applied Biosystems). Contigs were assembled using the SeqMan program and protein molecular mass calculated by the Protean Program (Lasergene version 5; DNASTAR Inc., Madison USA). The SignalP program at www.cbs.dtu.dk/services/signalP/ was used to identify the signal sequence.

2.4. Construction of *ggt* mutant

The standard and recombinant vectors and oligos used in this study are listed in Table 1. All standard methods of DNA manipulation were performed according to the protocols of Ausubel et al. [35]. The *ggt* gene in *C. jejuni* strain 81116 was disrupted by insertional mutagenesis with a kanamycin cassette. Briefly, *ggt* was amplified from genomic DNA isolated from *C. jejuni* strain 81116 using oligo primers If50 and If66 (Table 1b). The 1.5 kbp DNA fragment generated was cloned into pCR-ScriptTMSK(+). The *ggt* gene was mutated using inverse PCR oligos If67 and If68 (Table 1b), which introduced a unique *Bgl*III restriction site [38] 532 bp

downstream from the *ggt* start codon. The PCR fragment was digested with *Bgl*II and self ligated. The resulting plasmid was digested with *Bgl*II and a kanamycin resistance gene with *Bam*HI ends from pJMK30 (kindly provided by J. Ketley, University of Leicester) was inserted and the kanamycin cassette was in the opposite orientation to *ggt*. The suicide vector pHA-*ggt* was introduced into *C. jejuni* strain 81116 by electroporation, as previously described [38].

2.5. In vitro phenotypic studies

All the phenotypic assays were carried out in triplicate. Experiments were performed three times for verification. Statistical analysis of the data was carried out using GraphPad Prism™ software version 2.01 (San Diego, CA, USA).

Thermotrace method—The ‘GGT reagent’ (Alpha Laboratories Ltd) was used, as recommended by the manufacturer, to spectrophotometrically determine the specific activity for GGT. The protein concentration of whole *C. jejuni* cells ($\sim 10^7$ – 10^9 cfu), disrupted by freeze-thawing in 1 M Tris pH 8.0, was predetermined using the ‘Protein Assay Dye Reagent’ (Bio-Rad Laboratories Ltd). The sample was then diluted 1 in 10 with water and a 50 μ l aliquot was mixed with 950 μ l of prewarmed GGT reagent. The spectrophotometer was maintained at the appropriate temperature (32, 37, 42 or 47°C). Cleavage of the substrate (L- γ -glutamyl-3-carboxy-4-nitroanilide) due to GGT activity induced the appearance of a yellow color due to the formation of 5-amino-2-nitrobenzoate, which was quantified by absorbance at 405 nm. The final activity value was determined as U/ml/mg of whole bacterial cell protein. One unit was defined as the quantity of enzyme that released 1 μ mol of 5-amino-2-nitrobenzoate per min per mg of protein at the relevant temperature. For the colorimetric GGT assay *C. jejuni* isolates were incubated at 37°C for 30 min and visually monitored. In the presence of GGT activity the solution changed color from clear to yellow.

In vitro invasion assay—Bacterial invasion into INT407 cells was studied using the gentamicin protection assay [38] using a range of MOIs (50–1200) as previous studies have indicated that invasion efficiency varies with the inoculum [39]. Briefly, the INT407 cells were cultured in a 24-well tissue culture plate at 37°C with 5% (v/v) CO₂ until confluent monolayers of approx. 5×10^5 cells per well were established. The cells were rinsed with Hank's Balanced Salt Solution (HBSS) and inoculated with *C. jejuni*. Tissue culture plates were centrifuged at 600g, 22°C for 15 min to eliminate variations in motility between strains. Infected monolayers were incubated at 37°C in 5% (v/v) CO₂ for 3 h to allow the bacteria to invade the cells. The monolayers were washed three times with HBSS and incubated for a further 2 h in maintenance media containing gentamicin at 250 μ gml⁻¹. Finally, the monolayers were washed in HBSS and lysed with 1 ml of 1% (v/v) Triton X-100 in PBS. The suspensions were serially diluted and the number of viable, internalized bacteria was determined by counting the resultant colonies on blood plates.

Gentamicin susceptibility assay—Bacterial susceptibility to gentamicin was determined on 10% (vol/vol) sheep blood agar plates supplemented with serial two-fold dilutions of gentamicin (0–64 μ gml⁻¹).

Motility Assay—The motility of *C. jejuni* strains and mutants was determined as a measure of swarming on semisolid motility (SSM) media [40]. Briefly, a blunt hypodermic needle was dipped into the strain to be tested and then stabbed into the center of a SSM plate. The plates were incubated in microaerobic conditions at 37°C and the diameter of the colony in mm was determined after 24 and 48 h. Statistical significance was assessed using the two-tailed-*T*-test.

Hydrogen peroxide susceptibility—Hydrogen peroxide (final concentration of 0.5 mM) was added directly to bacterial cells grown in Mueller Hinton broth (10^9 cfu ml⁻¹) and incubated under microaerobic conditions at 42°C. Viable counts of bacteria were taken after 0, 20, 40, 60, 80, 100 and 120 min.

Apoptosis assay—The normal colonic epithelial cell line, CCD 841 CoN was obtained from the American Type Culture Collection (ATCC, Manassas, VA; no. CRL-1790™). Cells were grown in RPMI 1640 supplemented with 10% fetal bovine serum and 1% penicillin/streptomycin and cultured in 6-well plates. Once the cells were 40–60% confluent, *C. jejuni* grown to an OD₆₀₀ of 1.0 were added and incubated at 37°C in 5% (v/v) CO₂ for 4 days. The cell nuclei were stained with 4', 6-diamidino-2-phenylindole (DAPI) and used to assess both apoptosis and relative cell density by fluorescence quantitation. Nuclei with highly condensed and fragmented chromatin were considered apoptotic. Apoptotic cells were enumerated by counting 400 cells in multiple randomly selected fields.

2.6. Chick colonization

The 1-day-old chick model of colonization [41] was used to determine colonization potential. Briefly, eggs from specific-pathogen-free chickens (Lohmann's) were hatched in isolators. Groups of eight chicks were maintained in separate isolators with unlimited food and water. At 1-day old, chicks were each dosed orally, by gavage, with approximately 10^4 cfu of *C. jejuni* in 0.1 ml of 0.1 M PBS, pH 7.2. Bacteria were grown overnight on blood agar plates under microaerobic conditions at 42°C. Chicks were sacrificed 5 days or 3 weeks after dosing, and bacterial colonization levels were determined by plating serial dilutions of caecal contents. The non-parametric Mann–Whitney test was used to assess the statistical significance of differences in colonization levels.

3. Results

3.1. Identification of the genetic location of the ggt gene in *C. jejuni*

Insert 236, previously identified by subtractive hybridization [11], encoded a putative polypeptide with 72% amino acid identity to the *H. pylori* γ -glutamyl transpeptidase (GGT) polypeptide (GenBank accession no. AE000511). There was also a significant degree of similarity with GGT from *E. coli* (GenBank accession no. P18956) and *Salmonella enterica* serovar Typhimurium (GenBank accession no. NP_462452) and therefore a degenerate primer, If66 (Table 1) was designed to a conserved amino acid region, as near as possible to the N-terminus of these aligned proteins. The degenerate primer was used with a primer (If50) designed to the known *C. jejuni* *ggt* DNA sequence [11], which aligned to the amino acids at the C-terminus of the GGT sequence. Degenerate PCR was performed using genomic DNA from strain 81116 with the aforementioned primers and sequencing of the resulting 1.5 kbp PCR product identified a DNA sequence, which encoded a putative polypeptide with significant amino acid similarity (76%) to GGT from *H. pylori*.

Chromosomal walks on genomic DNA from strain 81116, using primer If49 identified the remaining *ggt* sequence and additional flanking DNA, which had no similarity to any sequence in strain NCTC11168, but had 54% amino acid identity and 67% similarity (Blast *P* value $2e^{-62}$) to a cytochrome c-type biogenesis protein (YCF5) from *Wolinella succinogenes* (GenBank accession no. NP_908253.1). The DNA sequence of *ggt* and this flanking DNA, obtained from strain 81116, was compared to the *ggt* genetic region (30 kbp) identified and kindly provided by Emily Kay, Sanger Center, Cambridge, from strain 81-176. Comparison of the *ggt* sequences from strains 81-176 and 81116 revealed 99.5% similarity at the DNA level (data not shown). In addition, the cytochrome c biogenesis DNA sequence obtained from strain 81116 was 99% identical to that in 81-176 (data not shown). This indicated that the

genomic location of *ggt* in strain 81116 was similar to that in strain 81-176 (Fig. 1). Following the identification of the full *ggt* sequence from strain 81116 (GenBank accession no. AY623656), this gene was also identified in another *C. jejuni* strain (GenBank accession no. AJ786772) which was isolated from chicken caecal contents. Comparison of the GGT amino acid sequences from three strains (GenBank accession no. AY623656, AJ786772 and from strain 81-176), confirmed the high degree of conservation of *ggt* within the *C. jejuni* species at both the DNA and amino acid levels. Bioinformatical analysis of the *C. jejuni ggt* gene from strain 81116 revealed a 1671 base-pair open reading frame encoding a poly-peptide with a calculated mass of 60300 Da. The first 17 amino acids at the N-terminus of GGT exhibit a typical signal sequence. A second post-translational cleavage site is predicted between positions Asn-370 and Thr-371, which would subsequently result in the formation of the large and small subunits of the mature enzyme, consistent with other bacterial and mammalian GGT orthologues [17,22].

3.2. The GGT-specific activity for strain 81116

Strain 81116 was assayed for GGT activity by the rate of formation of 5-amino-2-nitrobenzoate, using the Thermotrace method (Alpha Laboratories) at exponential and stationary phases of growth (20 and 40 h from the predetermined growth curve, respectively). The GGT activity at 32, 37, 42 and 47°C, was also determined. The results are summarized in Table 2. GGT activity was observed at both growth phases but higher (approx. 1.7 to 1.9-fold) in the exponential phase, regardless of temperature. GGT activity was also observed at all the temperatures tested but appeared to be maximal at 42°C, though this was not statistically significant.

3.3. Prevalence of *ggt* and GGT activity among *C. jejuni* isolates

One hundred and thirty four *C. jejuni* strains, from various sources, were tested for the prevalence of the *ggt* gene by colony-PCR using the primers If50 and If100 (Table 1b). These primers were designed to detect the regions of highest conservation observed within the sequenced *ggt* genes. Overall 19.4% of the strains generated the 1.5 kbp *ggt* PCR product (Table 3). This prevalence was considerably lower than the 54.5% indicated by a similar previously published study [11] but represents a substantially larger strain sample size. However, because the level of conservation of the primer regions is only known over a few strains, false negative PCRs are possible. Therefore, 19.4% must be considered as the minimal prevalence of positivity for *ggt*. Nevertheless, it is interesting to note that *ggt*-positive strains predominated in human isolates (37.93%).

The *C. jejuni* isolates previously screened for the presence of *ggt* [11] were tested in the colorimetric GGT assay. The yellow color change was only detected in strains positive for *ggt* (data not shown). Thus the presence of the gene, as determined by PCR, correlated directly with the expression of the GGT activity.

3.4. Construction of a *C. jejuni* mutant deficient in GGT activity

To determine the function of GGT in *C. jejuni*, a mutant was constructed in strain 81116. The *ggt* gene was disrupted by insertion of a kanamycin resistance cassette at 532 bp downstream from the *ggt* start codon, using the suicide vector pIHA-*ggt*. PCR and Southern blot analysis (data not shown) confirmed a double crossover and successful insertion of the kanamycin resistance cassette. Southern blot analysis also showed that *C. jejuni* strain 81116 carried only one copy of the *ggt* gene. In strain 81116, *ggt* gene is monocistronic and cannot be transcriptionally linked with its downstream gene, which is transcribed in the opposite direction. It is therefore extremely unlikely that gene replacement would have any effect on the expression of flanking genes (Fig. 1). However, in order to prevent any effect on the flanking genes, the PCR primers were designed within the *ggt* gene so that flanking genes and intergenic

regions including potential promoters would remain undisrupted in the mutant. Nevertheless, as complementation is difficult with *Campylobacter*, another independently isolated *ggt* mutant was used as a control for secondary mutations. Neither of the kanamycin-resistant mutants had detectable GGT-specific activity. Comparison of the growth curves in biphasic broth or morphologies as observed by electron microscopy indicated no differences between the wild-type and mutant strains (data not shown).

3.5. Invasion assay using INT407 cells

The potential role of the GGT of *C. jejuni* 81116 in virulence was investigated by comparing the mutant and wild-type strains in an *in vitro* assay of invasion (Fig. 2). At all MOIs tested, higher numbers of the *ggt* mutant had internalized INT407 cells. At the optimal MOI of 200 the apparent invasion efficiency was 5.4-fold higher for the *ggt* mutant (0.0092%) compared to the wild-type (0.0017%).

3.6. Gentamicin resistance

An alteration in the gentamicin resistance may have influenced the invasion assay results and therefore was tested. Both the *ggt* mutant and wild-type strains were susceptible to 0.125 μgml^{-1} gentamicin and were unable to grow on 0.25 μgml^{-1} gentamicin (data not shown). This showed that the increased invasion efficiency of the *ggt* mutant was not due to increased gentamicin resistance.

3.7. Motility

Motility may also have been a factor influencing the INT407 invasion phenotypes. The average diameter of the zone of motility of the wild-type was 25.42 mm $\pm$ 3.95 at 24 h and 58.85 mm $\pm$ 6.89 at 48 h. The *ggt* mutant was significantly more motile than the parent strain; at 24 h the diameter was 32.85 mm $\pm$ 2.61 (P 0.0018) and 48 h it was 69.86 mm $\pm$ 2.91 (P = 0.0045). This difference may have caused the increased invasiveness but any such effect should have been minimized by centrifugation of the bacteria onto the monolayer.

3.8. Hydrogen peroxide susceptibility

GGT activity has been shown to be associated with hydrogen peroxide resistance [19]. The sensitivity of the *ggt* mutant to hydrogen peroxide, at a final concentration of 0.5 mM, was considerably less than the wild-type strain (Fig. 3). After 20 min, levels of viable mutant cells were consistently around 3 log₁₀ higher than those of the wild-type strain, indicating that the inactivation of GGT increases the bacterium's resistance to hydrogen peroxide stress. Such differences could also influence the invasion efficiencies.

3.9. Apoptosis-inducing activity of GGT

Previous studies in *H. pylori* have shown that GGT is involved in the induction of apoptosis [28]. Therefore, the ability of the *C. jejuni* wild-type strain to induce apoptosis of CCD 841 CoN epithelial cells was compared with that of the *ggt* mutant. The percentage of CCD 841 CoN cells with either highly condensed or fragmented nuclei was dose-dependently higher following incubation with the wild-type strain. (Fig. 4(a)). However, the relative CCD 841 CoN cell number decreased following incubation with either *C. jejuni* strain but to a much higher degree with the wild-type strain (Fig. 4(b)). Furthermore, phase contrast microscopy analysis showed that incubation with both the *ggt* mutant and wild-type strain altered the normal cell morphology and significantly reduced the density of adherent cells (Fig. 4(c–h)) compared with the control monolayer. Only the control cells become 95–100% confluent following the incubation period. Interestingly, nuclear DAPI staining revealed an increased population of cells with highly condensed and fragmented nuclei following infection with the wild-type strain. In contrast, a greater proportion of enlarged cell nuclei were observed

following incubation with the *ggt* mutant. Overall, these results indicate that GGT plays a significant role in *C. jejuni*-mediated apoptosis.

3.10. Chick colonization model

The colonization potential of the *ggt* mutant was determined using a well-established 1-day-old chick model of colonization. Five days post-oral-challenge, with a dose of approximately 10^4 cfu, both the wild-type strain and *ggt* mutant had similar colonization potentials with 88% of chicks colonized (Fig. 5(a)). The geometric mean level of colonization by the *ggt* mutant was $4.74 \log_{10}$ cfu g^{-1} of caecal contents, which was lower than the wild-type strain at $6.11 \log_{10}$ cfu g^{-1} . However, this was not statistically significant ($P = 0.1049$). Nevertheless, after three weeks post-oral challenge, the wild-type strain colonized 100% of chicks to maximal levels of $8-9 \log_{10}$ cfu g^{-1} (Fig. 5(b)) while the colonization level in all birds by the *ggt* mutant was undetectable ($P = 0.0002$). These results indicate a lack of persistence of the GGT mutant in the avian gut.

4. Discussion

Although the importance of γ -glutamyl transpeptidase in glutathione metabolism and amino acid transport in mammalian tissues has been recognized for many years [17], there is little information about the role of this enzyme in bacteria. To date, several prokaryotes, including *E. coli*, *Pseudomonas aeruginosa*, *Bacillus subtilis*, *H. pylori*, and *T. denticola*, have been shown to exhibit GGT activity [22,25,42–44] but the physiological role of GGT in these bacteria remains unclear.

In this study, the γ -glutamyl transpeptidase (*ggt*) gene has been identified and characterized in *C. jejuni* strain 81116. The role of GGT in *C. jejuni* was investigated using two defined *ggt* mutants generated in *C. jejuni* strain 81116. Both mutants had identical properties. Neither *ggt* mutant expressed any detectable GGT activity. Two-dimensional protein analysis also confirmed that GGT was no longer present (data not shown). Interestingly, in the wild-type strain, two adjacent GGT protein spots of the same molecular mass, but different isoelectric points, were observed by the two-dimensional electrophoresis method (data not shown), suggesting that GGT may be post-translationally modified, and possibly glycosylated, which would be consistent with the known post-translational glycosylation system in *C. jejuni*.

Results with the mutants demonstrated that GGT is not essential for *in vitro* growth, which also agrees with a recently published study using *C. jejuni* strain 81-176 [45]. Although characterization of *ggt* mutants in *C. jejuni* and in *N. meningitidis* [46] indicates that GGT is not required for bacterial growth under normal *in vitro* conditions, the expression of active GGT in *E. coli* is essential for the utilization of exogenous γ -glutamyl peptides [47]. It therefore seems likely that, although not essential, GGT may contribute to the organism's ability to cope with growth limiting factors during environmental survival.

Only 19.4% of 134 randomly selected *C. jejuni* strains contained the *ggt* gene and prevalence was highest in human isolates (37.93%). This suggests that there may be some advantage for those *C. jejuni* possessing GGT to colonize the human host. In contrast, in the closely related *Helicobacter* species all strains synthesize a catalytically active GGT, regardless of host sources, even though this was also inessential for growth [25].

In order to assess the role of GGT, a representative *ggt* mutant was investigated in a number of assay systems, reflecting properties related to virulence, survival and colonization potential. To date there are no acceptable models of virulence for *C. jejuni* [48]. However, surrogate models, such as invasion, have been utilized as indicators of potential of this organism to cause enteric disease.

The apparent higher recovery of *ggt* mutant cells, as compared to the wild-type, during the INT407 cell invasion assay, was initially surprising and contradicts the study on a *ggt* mutant in strain 81-176 in which the ability to invade T84 intestinal epithelial cells was indistinguishable [45]. This discrepancy is most likely due to the differences in cell lines used and genetic background of strains. Nevertheless, the combined effects of increased motility and resistance to hydrogen peroxide of the *ggt* mutant in strain 81116 most likely contributed to the enhanced invasion phenotype. Previous reports have shown motility as a major factor influencing invasion [38,49]. In addition, prolonged intracellular survival within host cells has been linked with increased hydrogen peroxide resistance [50]. Previous studies [51] have indicated that loss of GGT activity in bacteria is initially accompanied by a rise in intracellular glutathione levels, potentially enhancing the organism's resistance to host cell-mediated oxidative stress. Although the association of GGT with that of the antioxidant glutathione in prokaryotes has not been clearly established, in some bacteria protection against peroxides is related to the ability to acquire glutathione [52]. Conversely, an essential role for GGT in glutathione metabolism has been reported in the periodontal bacterial pathogen, *Treponema denticola* [42,53], in which GGT-mediated hydrolysis of glutathione is a source for the H₂S necessary to reduce injury or death as a consequence of some environmental stresses.

Additional results indicated that the *C. jejuni* GGT enzyme has an apoptosis-inducing activity, which may be involved in pathogenesis. Clearly, the induction by *C. jejuni* of apoptotic cell death and changes to cell morphology is multifactorial [54,55]. The *C. jejuni* cytolethal distending toxin (CDT) causes epithelial cells to become blocked in the G₂/M phase, where the cells appear enlarged and obvious mitotic cells are absent [54,56]. Coincidentally, a greater proportion of enlarged cells were observed following infection with the *ggt* mutant and whether this is due to an indirect effect on CDT has yet to be investigated. The role of such effects on host cells is unclear but apoptotic cells may be a source of essential nutrients to *C. jejuni*, thus contributing to pathogen survival in the gut.

The colonization potentials of the wild type and mutant strains in the avian intestine were also investigated. Interestingly, although initial colonization was not markedly different between the variants, the *ggt* mutant was unable to sustain colonization of the chicken gut. The dose response curve for colonization with *C. jejuni* strain 81116 wild type is highly reproducible, and with the dose level of 10⁴ cfu, colonization once established consistently persists for at least 40 days after which it may decline [2]. A similar effect was also observed for the *ggt* mutant in strain 81-176 in mixed-infection mouse colonization experiments [45]. Why GGT should be required for only persistent, but not initial, intestinal mucosal colonization is unclear and confusing when considering the *in vitro* phenotypes. Nevertheless, the presence or absence of GGT may, at least in part, explain differences in colonization phenotype reported by others [57].

In contrast to the avian studies, GGT-negative mutants of *H. pylori* tested in adult mammalian *in vivo* models were either non-colonizing [25] or initiated poor but sustained gastric mucosal colonization [26]. Such differences may reflect the separate ecological niches of these closely related organisms. It, therefore, seems possible that the presence of GGT in some *C. jejuni* strains confers an advantage in the *in vivo* environment by possibly sustaining glutathione levels (via recycling) to reduce the sensitivity to host inflammatory responses or as a means of providing a more effective supply of the essential amino acids, glycine and cysteine [58].

In conclusion, this study has identified the *ggt* gene in *C. jejuni* strain 81116 and partly characterized the function of GGT in this strain using an isogenic mutant. Although genetic tools for *C. jejuni* remain somewhat limited, future analysis involving attempts to complement *ggt* need to be undertaken to confirm these observations. Nevertheless, a number of interesting indicators of the role of GGT in *C. jejuni* infections have been provided. Of particular interest

was the role in sustaining avian gut colonization. Although, colonization by *C. jejuni* is clearly multifactorial in nature, the identification of bacterial colonization factors, which enable persistent avian gut colonization, such as *ggt*, may in the future enable the development of targeted strategies for intervention to control and prevent this major public health problem.

Acknowledgments

We thank Kenneth Stapleton for assistance with the 1-day-chick model and Emily Kay (Sanger Center, Cambridge) for kindly providing the *ggt* sequence from strain 81-176. This project (No. OZ0607) was mainly funded by the Department of Environmental Food and Rural Affairs (DEFRA), United Kingdom (to D.G. Newell) and also partly by National Institutes of Health grant AI055715 (to S.A. Thompson).

References

1. Stanley KN, Jones K. High frequency of metronidazole resistance among strains of *Campylobacter jejuni* isolated from birds. *Lett Appl Microbiol* 1998;27:247–50. [PubMed: 9830138]
2. Newell, DG.; Wagenaar, JA. Poultry infections and their control at the farm level.. In: Nachamkin, I.; Blaser, MJ., editors. *Campylobacter*. Vol. 2nd ed.. ASM Press; Washington, DC: 2000. p. 497-509.
3. Petersen L, Nielsen EM, Engberg J, On SL, Dietz HH. Comparison of genotypes and serotypes of *Campylobacter jejuni* isolated from Danish wild mammals and birds and from broiler flocks and humans. *Appl Environ Microbiol* 2001;67:3115–21. [PubMed: 11425730]
4. Tompkins DS, Hudson MJ, Smith HR, Eglin RP, Wheeler JG, Brett MM, et al. A study of infectious intestinal disease in England: microbiological findings in cases and controls. *Commun Dis Public Health* 1999;2:108–13. [PubMed: 10402744]
5. Friedman, CR.; Neimann, J.; Wegenaar, HC.; Tauxe, R. Epidemiology of *Campylobacter jejuni* infection in the United States and other industrialized nations.. In: Naschamkin, I.; Blaser, MJ., editors. *Campylobacter*. ASM Press; Washington DC: 2001. p. 121-38.
6. Rosenquist H, Nielsen NL, Sommer HM, Norrung B, Christensen BB. Quantitative risk assessment of human campylobacteriosis associated with thermophilic *Campylobacter* species in chickens. *Int J Food Microbiol* 2003;83:87–103. [PubMed: 12672595]
7. Newell DG, Fearnley C. Sources of *Campylobacter* colonization in broiler chickens. *Appl Environ Microbiol* 2003;A69:4343–51. [PubMed: 12902214]
8. Parkhill J, Wren BW, Mungall K, Ketley JM, Churcher C, Basham D, et al. The genome sequence of the food-borne pathogen *Campylobacter jejuni* reveals hypervariable sequences. *Nature* 2000;403:665–8. [PubMed: 10688204]
9. Carrillo CD, Taboada E, Nash JHE, Lanthier P, Kelly J, Lau PC, et al. Genome-wide expression analyses of *Campylobacter jejuni* NCTC11168 reveals coordinate regulation of motility and virulence by *flhA*. *J Biol Chem* 2004;279:20327–38. [PubMed: 14985343]
10. Gaynor EC, Cawthraw S, Manning G, MacKichan JK, Falkow S, Newell DG. The genome-sequenced variant of *Campylobacter jejuni* NCTC 11168 and the original clonal clinical isolate differ markedly in colonization, gene expression, and virulence-associated phenotypes. *J Bacteriol* 2004;186:503–17. [PubMed: 14702320]
11. Ahmed IH, Manning G, Wassenaar TM, Cawthraw S, Newell DG. Identification of genetic differences between two *Campylobacter jejuni* strains with different colonization potentials. *Microbiology* 2002;148:1203–12. [PubMed: 11932464]
12. Dingle KE, Colles FM, Wareing DRA, Ure R, Fox AJ, Bolton FE, et al. Multilocus Sequence Typing System for *Campylobacter jejuni*. *J Clin Microbiol* 2001;39:14–23. [PubMed: 11136741]
13. Wassenaar TM, Blaser MJ. Pathophysiology of *Campylobacter jejuni* infections of humans. *Microbes Infect* 1999;1:1023–33. [PubMed: 10617934]
14. Poly F, Threadgill D, Stintzi A. Identification of *Campylobacter jejuni* ATCC 43431-specific genes by whole microbial genome comparisons. *J Bacteriol* 2004;186:4781–95. [PubMed: 15231810]
15. Poly F, Threadgill D, Stintzi A. Genomic diversity in *Campylobacter jejuni*: identification of *C. jejuni* 81-176-specific genes. *J Clin Microbiol* 2005;43:2330–8. [PubMed: 15872262]

16. Stintzi A, Marlow D, Palyada K, Naikare H, Panciera R, Whitworth L, et al. Use of genome-wide expression profiling and mutagenesis to study the intestinal lifestyle of *Campylobacter jejuni*. *Infect Immun* 2005;73:1797–810. [PubMed: 15731081]
17. Tate SS, Meister A. Gamma-glutamyl transpeptidase: catalytic, structural and functional aspects. *Mol Cell Biochem* 1981;39:357–68. [PubMed: 6118826]
18. O'Donovan DJ, Fernandes CJ. Mitochondrial glutathione and oxidative stress: implications for pulmonary oxygen toxicity in premature infants. *Mol Genet Metab* 2000;71:352–8. [PubMed: 11001827]
19. Shi M, Gozal E, Choy HA, Forman HJ. Extracellular glutathione and gamma-glutamyl transpeptidase prevent H₂O₂-induced injury by 2,3-dimethoxy-1,4-naphthoquinone. *Free Radical Biol Med* 1993;15:57–67. [PubMed: 8103030]
20. Izawa S, Inoue Y, Kimura A. Oxidative stress response in yeast: effect of glutathione on adaptation to hydrogen peroxide stress in *Saccharomyces cerevisiae*. *FEBS Lett* 1995;368:73–6. [PubMed: 7615092]
21. Meister A. On the antioxidant effects of ascorbic acid and glutathione. *Biochem Pharmacol* 1992;44:1905–15. [PubMed: 1449510]
22. Xu K, Strauch MA. Identification, sequence, and expression of the gene encoding gamma-glutamyltranspeptidase in *Bacillus subtilis*. *J Bacteriol* 1996;178:4319–22. [PubMed: 8763966]
23. Ikeda Y, Fujii J, Anderson ME, Taniguchi N, Meister A. Involvement of Ser-451 and Ser-452 in the catalysis of human gamma-glutamyl transpeptidase. *J Biol Chem* 1995;270:22223–8. [PubMed: 7673200]
24. Szymanski CM, Yao R, Ewing CP, Trust T, Guerry JP. Evidence for a system of general protein glycosylation in *Campylobacter jejuni*. *Mol Microbiol* 1999;32:1022–30. [PubMed: 10361304]
25. Chevalier C, Thiberge JM, Ferrero RL, Labigne A. Essential role of *Helicobacter pylori* gamma-glutamyltranspeptidase for the colonization of the gastric mucosa of mice. *Mol Microbiol* 1999;31:1359–72. [PubMed: 10200957]
26. McGovern KJ, Blanchard TG, Gutierrez JA, Czinn SJ, Krakowka S, Youngman P. Gamma-glutamyltransferase is a *Helicobacter pylori* virulence factor but is not essential for colonization. *Infect Immun* 2001;69:4168–73. [PubMed: 11349094]
27. Gong M, Ho B. Prominent role of gamma-glutamyl-transpeptidase on the growth of *Helicobacter pylori*. *World J Gastroenterol* 2004;20:2994–6. [PubMed: 15378780]
28. Shibayama K, Kamachi K, Nagata N, Yagi T, Nada T, Doi Y, et al. A novel apoptosis-inducing protein from *Helicobacter pylori*. *Mol Microbiol* 2003;47:443–51. [PubMed: 12519194]
29. Busiello I, Acquaviva R, Di Popolo A, Blanchard TG, Ricci V, Romano M, et al. *Helicobacter pylori* gamma-glutamyltranspeptidase upregulates COX-2 and EGF-related peptide expression in human gastric cells. *Cell Microbiol* 2004;6:255–67. [PubMed: 14764109]
30. Takahashi H, Hirose K, Watanabe H. Necessity of meningococcal gamma-glutamyl aminopeptidase for *Neisseria meningitidis* growth in rat cerebrospinal fluid (CSF) and CSF-like medium. *J Bacteriol* 2004;186:244–7. [PubMed: 14679245]
31. Smirnova GV, Krasnykh TA, Oktyabrsky ON. Role of glutathione in the response of *Escherichia coli* to osmotic stress. *Biochemistry (Moscow)* 2001;66:973–8. [PubMed: 11703177]
32. Palmer SR, Gully PR, White JM, Pearson AD, Suckling WG, Jones DM, et al. Water-borne outbreak of *Campylobacter* gastroenteritis. *Lancet* 1983;1:287–90. [PubMed: 6130305]
33. Korlath JA, Osterholm MT, Judy LA, Forfang JC, Robinson RA. A point-source outbreak of campylobacteriosis associated with consumption of raw milk. *J Infect Dis* 1985;152:592–6. [PubMed: 4031557]
34. Skirrow MB. *Campylobacter* enteritis: a 'new' disease. *Br Med J* 1977;2:9–11. [PubMed: 871765]
35. Ausubel, SF.; Brent, R.; Kingston, RE.; Moore, DD.; Seidman, JG.; Smith, JA., et al., editors. *Current Protocols in Molecular Biology*. Wiley; New York, NY: 1994.
36. Sachadyn P, Sobiewska G, Kur J. Thermal profile with alternately raised and lowered annealing temperature improves the PCR amplification using highly degenerate primers. *Acta Biochim Pol* 1998;45:691–4. [PubMed: 9918495]
37. Screaton GR, Bangham CR, Bell JI. Direct sequencing of single primer PCR products: a rapid method to achieve short chromosomal walks. *Nucleic Acids Res* 1993;21:2263–4. [PubMed: 8502578]

38. Wassenaar TM, Bleumink-Pluym NM, van der Zeijst BA. Inactivation of *Campylobacter jejuni* flagellin genes by homologous recombination demonstrates that *flaA* but not *flaB* is required for invasion. *EMBO J* 1991;10:2055–61. [PubMed: 2065653]
39. Hu L, Kopecko DJ. *Campylobacter jejuni* 81-176 associates with microtubules and dynein during invasion of human intestinal cells. *Infect Immun* 1999;67:4171–82. [PubMed: 10417189]
40. Goossens H, Vlaes L, Galand I, Van den Borre C, Butzler JP. Semisolid blood-free selective-motility medium for the isolation of *Campylobacters* from stool specimens. *J Clin Microbiol* 1989;27:1077–80. [PubMed: 2745681]
41. Wassenaar TM, van der Zeijst BA, Ayling R, Newell DG. Colonization of chicks by motility mutants of *Campylobacter jejuni* demonstrates the importance of flagellin A expression. *J Gen Microbiol* 1993;139:1171–5. [PubMed: 8360610]
42. Chu L, Dong Z, Xu X, Cochran DL, Ebersole JL. Role of glutathione metabolism of *Treponema denticola* in bacterial growth and virulence expression. *Infect Immun* 2002;70:1113–20. [PubMed: 11854190]
43. Ishiye M, Yamashita M, Niwa M. Molecular cloning of the gamma-glutamyltranspeptidase gene from a *Pseudomonas* strain. *Biotechnol Prog* 1993;9:323–31. [PubMed: 7765305]
44. Suzuki H, Kumagai H, Echigo T, Tochikura T. DNA sequence of the *Escherichia coli* K-12 gamma-glutamyltranspeptidase gene, *ggt*. *J Bacteriol* 1989;171:5169–72. [PubMed: 2570061]
45. Hofreuter D, Tsai J, Watson RO, Novik V, Altman B, Benitez M, et al. Unique features of a highly pathogenic *Campylobacter jejuni* strain. *Infect Immun* 2006;74:4694–707. [PubMed: 16861657]
46. Riou JY, Buissiere J, Richard C, Guibourdenche M. Gamma-glutamyl-transferase activity in the family “*Neisseriaceae*”. *Ann Microbiol (Paris)* 1982;133:387–92. [PubMed: 6125120]
47. Suzuki H, Hashimoto W, Kumagai H. *Escherichia coli* K-12 can utilize an exogenous gamma-glutamyl peptide as an amino acid source, for which gamma-glutamyltranspeptidase is essential. *J Bacteriol* 1993;175:6038–40. [PubMed: 8104180]
48. Newell DG. Animal models of *Campylobacter jejuni* colonization and disease and the lessons to be learned from similar *Helicobacter pylori* models. *Symp Ser Soc Appl Microbiol* 2001:57S–67S.
49. Golden NJ, Acheson DWK. Identification of motility and autoagglutination *Campylobacter jejuni*. *Infect Immun* 2002;70:1761–71. [PubMed: 11895937]
50. Wu H, Seib KL, Edwards JL, Apicella MA, McEwan AG, Jennings MP. Azurin of pathogenic *Neisseria* spp. is involved in defense against hydrogen peroxide and survival within cervical epithelial cells. *Infect Immun* 2005;73:8444–8. [PubMed: 16299348]
51. Nakayama R, Kumagai H, Tochikura T. Leakage of glutathione from bacterial cells caused by inhibition of gamma-glutamyltranspeptidase. *Appl Environ Microbiol* 1984;47:653–7. [PubMed: 6144289]
52. Vergauwen B, Pauwels F, Van Beeumen JJ. Glutathione and catalase provide overlapping defenses for protection against respiration-generated hydrogen peroxide in *Haemophilus influenzae*. *J Bacteriol* 2003;185:5555–62. [PubMed: 12949108]
53. Chu L, Xu X, Dong Z, Cappelli D, Ebersole JL. Role for recombinant gamma-glutamyltransferase from *Treponema denticola* in glutathione metabolism. *Infect Immun* 2003;71:335–42. [PubMed: 12496183]
54. Hickey TE, Majam G, Guerry P. Intracellular survival of *Campylobacter jejuni* in human monocytic cells and induction of apoptotic death by cytolethal distending toxin. *Infect Immun* 2005;73:5194–7. [PubMed: 16041038]
55. Zhu J, Meinersmann RJ, Hiatt KL, Evans DL. Apoptotic effect of outer-membrane proteins from *Campylobacter jejuni* on chicken lymphocytes. *Curr Microbiol* 1999;38:244–9. [PubMed: 10069862]
56. Whitehouse CA, Balbo PB, Pesci EC, Cottle DL, Mirabito PM, Pickett CL. *Campylobacter jejuni* cytolethal distending toxin causes a G2-phase cell cycle block. *Infect Immun* 1998;66:1934–40. [PubMed: 9573072]
57. Ringoir DD, Korolik V. Colonisation phenotype and colonization potential differences in *Campylobacter jejuni* strains in chickens before and after passage *in vivo*. *Vet Microbiol* 2003;92:225–35. [PubMed: 12523984]

58. Fang FC. Perspectives series: host/pathogen interactions. Mechanisms of nitric oxide-related antimicrobial activity. *J Clin Invest* 1997;99:2818–25. [PubMed: 9185502]
59. van Vliet AH, Wooldridge KG, Ketley JM. Iron acquisition and regulation in *Campylobacter jejuni*. *J Bacterio* 1998;180:5291–8.

Fig. 1.
The arrangement of *ggt* and adjacent genes in strains 81116 and 81-176. Arrows denote predicted direction of transcription.

Fig. 2.

Comparison of the capacity of the *ggt* mutant to invade INT407 cells compared to the wild-type strain 81116. Invasion is represented as the total number of bacteria surviving after the gentamicin treatment. The MOI at 200 was compared for the mutant and wild-type strain. Symbols: ▲, *ggt* mutant; ■, wild-type.

Fig. 3. Susceptibility of the *ggt* mutant to hydrogen peroxide compared to the wild-type strain 81116. This was determined as the numbers of bacteria ($\log_{10}$ cfu ml^{-1}) that survived exposure to 0.5 mM H_2O_2 over a period of 120 min. The assay was performed three times and the results from one assay using triplicate samples is shown. Symbols: $\blacktriangle$, *ggt* mutant; $\blacksquare$, wild-type.

Fig. 4. (a) DAPI assay for *C. jejuni* infection induced apoptosis and (b) the relative number of adherent colonic epithelial (CCD 841 CoN) cells. The assay was performed three times. Symbols: ▲, *ggt* mutant; ■, wild-type. Phase-contrast microscopy analysis of colonic epithelial (CCD 841 CoN) cells: (c) control uninfected cells, (d) infection with wild-type strain and (e) *ggt* mutant. DAPI stained epithelial nuclei: (f) control uninfected cells, (g) infection with wild-type strain and (h) *ggt* mutant.

Fig. 5.

Chick caecal colonization at (a) 5 days post-challenge and (b) three weeks post-challenge by the *ggt* mutant compared to the wild-type strain 81116. Colonization was determined as the number of bacteria recovered from the caecum, expressed as cfu per gram of caecal contents. The geometric mean level of colonization for each group is shown by a+. The dotted line represents the limit of detection of 100 cfu. Both wild-type and mutant were given to the chicks at a dose of $\sim 1 \times 10^4$ cfu per 100 μ l 0.1 M PBS pH 7.2. Symbols: $\blacktriangle$, *ggt* mutant; $\blacksquare$, wild-type.

Table 1

(a) Standard and recombinant vectors used in this study. (b) Primers used in this study (T_m, temperature in °C; CW, chromosomal walking; Kan, kanamycin resistance; Amp, ampicillin resistance)

(a) Vectors	Comment	Reference
pCR-Script™ SK(+)	Cloning vector, Amp	Stratagene Europe
pJMK30	<i>C. coli</i> Kan cassette in pUC19	van Vliet et al. [59]
pCR-Script- <i>ggt</i>	<i>ggt</i> in cloning vector, Amp	This study
pIHA- <i>ggt</i>	<i>ggt</i> suicide vector, Amp, Kan	This study

(b) Oligo	T _m	Sequence 5' to 3'	Comment
If49	54.9	gtc tct atg cca act atc a	Standard/CW primer
If50	53.7	ggg taa ata aga agt tag aat tc	Standard primer
If66	62.7	ggn ggn aay gcn rtn gay gcn	Degenerate PCR primer
If67	64.2	gaa gat ctc ata tct tta gct tct agc atg	Inverse PCR primer 1
If68	65.6	gaa gat cta aga tgg cag cac tta taa aag	Inverse PCR primer 2
If69	63.4	ggc agc act tat aaa agc gg	Sequencing primer 1
If70	63.1	cct ggt ttg att gaa aaa tca tc	Sequencing primer 2
If81	56.9	gtg atg taa atg cta tca tgg	Sequencing primer for CW product
If100	66.3	ctt gat aaa ggc gga aat gcc	Standard primer

Table 2

The GGT specific activity of strain 81116 at different temperatures and at exponential phase or stationary phase

	Exponential phase (20 h) (U/ml/mg)	Stationary phase (40 h) (U/ml/mg)
32 °C	21.40	12.49
37 °C	22.95	12.12
42 °C	25.63	13.76
47 °C	22.25	12.01

The sensitivity of the assay is 0.001 Abs/min per U/L.

Quantitative determination of γ -glutamy transpeptidase activity expressed as units per ml per mg of total protein

Table 3Presence of γ -glutamyl transpeptidase gene in *C. jejuni* from various sources

<i>C. jejuni</i> source	Numbers (percentage) positive
Poultry	7/31 (22.58%)
Bovine	2/31 (6.45%)
Human	11/29 (37.93%)
Ovine	3/18 (16.67%)
Porcine	1/15 (6.67%)
Other	2/10 (20.00%)
Total	26/134 (19.40%)

The number (percentage) of isolates from a particular source carrying *ggt* gene is given.