CORRESPONDENCE

Cardiac Surgery in the Elderly Patient

by PD Dr. med. Ivar Friedrich, PD Dr. med. Andreas Simm, Joachim Kötting, Dr. med. Frank Thölen, Burkhard Fischer, Prof. Dr. med. Rolf-Edgar Silber in volume 25/2009

Cognitive Impairments After Bypass Surgery

With regard to the overall results of cardiosurgical interventions and the quality of life of affected patients, postoperative cognitive impairments are enormously important. Substantial impairments to patients' cognitive functioning are found in the immediate postoperative period as well as in the longer term after the intervention. Newman et al reported early postoperative incidence rates of cognitive impairments of more than 50%, and of 24% even after 6 months after bypass surgery. Impairments to cognitive functioning occurred in elderly people who had no such impairments before the procedure, and deteriorations of cognitive status were noted in patients with pre-existing impairments (1, 2).

While it has been proved that elderly patients benefit from surgical myocardial revascularization with regard to improving cardiac function, Philipps-Bute showed that impairments to neurocognitive functioning as a consequence of cardiac surgery have negative effects on patients' quality of life in the medium and long term: in 36% of patients, any benefits derived from the improvement in cardiac function are negated by the cognitive deterioration. Interestingly, the use of the heart-lung machine (pump-oxygenator; cardiopulmonary bypass pump) itself does not seem to negatively affect a patient's neurocognitive status after cardiac surgery. "Off-pump" surgical techniques have not shown measurable risk reductions (2, 3).

The older the patient is in whom surgery is planned, the more important for the overall result it is to consider his or her preoperative comorbidities, especially impairments to cognitive function and to mobility. In the context of the preoperative stratification of risk factors in older patients about to have cardiac surgery, clinically manifest or latent cognitive deficits should be detected, and timely intervention should be planned for the eventuality of postoperative delirium, so as to reduce postoperative cognitive deterioration in as far as is possible. Older patients with heart disease benefit from an interdisciplinary approach, and the importance of individual comorbidities needs to be balanced, in an interdisciplinary fashion, against quality of life in the long term.

DOI: 10.3238/arztebl.2009.0693a

REFERENCES

1. Newman MF, Kirchner JL, Phillips-Bute B, et al.: Neurological Outcome Research Group and the Cardiothoracic Anesthesiology Research Endeavors Investigators. Longitudinal assessment of neurocognitive function after coronary-artery bypass surgery. *N Engl J Med* 2001; 344: 395–402.

2. Phillips-Bute B, Mathew JP, Blumenthal JA, Grocott HP, Laskowitz DT, Jones RH, Mark DB, Newman MF: Association of neurocognitive function and quality of life 1 year after coronary artery bypass graft (CABG) surgery. *Psychosom Med* 2006; 68: 369–75.
3. Jensen BØ, Rasmussen LS, Steinbrüchel DA: Cognitive outcomes in elderly high-risk patients 1 year after off-pump versus on-pump coronary artery bypass grafting. A randomized trial. *Eur J Cardiothorac Surg* 2008; 34: 1016–21.
4. Friedrich I, Simm A, Kötting J, Thölen F, Fischer B, Silber RE: Cardiac surgery in the elderly patient. *Dtsch Arztebl Int* 2009; 106(25): 416–22.

Dr. med. Gernot Heusinger von Waldegg
Dr. med. Andreas Klement

Dr. med. Gernot Heusinger von Waldegg
Klinik für Geriatrie
Pfeiffersche Stiftungen
Pfeifferstr 10
39114 Magdeburg, Germany
gernot.heusingervonwaldegg@pfeiffersche-stiftungen.org

Geriatric Perspective

We congratulate Friedrichs et al on their article. On the basis of data from the BQS (Bundesgeschäftsstelle Qualitätssicherung, the Federal Office for Quality Assurance) they convincingly showed the notable drop in mortality of elderly patients after bypass surgery. They also point out, rightly, that current, commonly used risk scores—such as the EuroSCORE—notably overestimate postoperative mortality and are thus of limited usefulness for assessing risk. The authors therefore suggest using a new risk score based on the BQS data.

We think that such a score may be equally inadequate in trying to assess the operative risk in elderly patients. Several studies have shown that functional variables that are collected in the context of geriatric assessment have a substantial influence on the hospital mortality and morbidity of elderly patients. Such factors include dependence on others in activities of everyday life, cognition, mobility, sarcopenia, and symptoms of depression (1, 2). As Silber and colleagues mentioned, considering these factors makes much sense in assessing the operative risk of elderly patients. In June 2009, our group concluded a cohort study evaluating the prospective validity of these parameters, and we have presented initial results at the congress of the American College of Cardiology (3). Sarcopenia (using hand-grip strength as a marker), cognitive impairments, and impaired activities of everyday life were associated with a two- to threefold mortality compared with patients without such functional impairments. We wish to encourage all colleagues to include these factors in their work so as to appropriately take into account the particular risk constellation of elderly and very old patients.

DOI: 10.3238/arztebl.2009.0693b

REFERENCES

1. Fried LP, Tangen CM, Walston, et al.: Frailty in older adults: evidence for a phenotype. *J Gerontol A Biol Sci Med Sci* 2001; 56A: M146–56.
2. Inouye S, Peduzzi P, Robison, et al.: Importance of functional measures in predicting mortality among older hospitalized patients. *JAMA* 1998; 279: 1187–93.
3. Frilling B, von Renteln-Kruse W, Riess FC: Prognostic value of geriatric assessment prior to cardiac surgery of the elderly. *JACC* 2009; 53: A403–18.
4. Friedrich I, Simm A, Kötting J, Thölen F, Fischer B, Silber RE: Cardiac surgery in the elderly patient. *Dtsch Arztebl Int* 2009; 106(25): 416–22.

PD Dr. med. Friedrich-Christian Rieß

Herzzentrum, Albertinen-Krankenhaus, Hamburg

Prof Dr. med. Wolfgang von-Renteln Kruse

Medizinisches Zentrum, Albertinen-Krankenhaus, Hamburg

Dr. med. Birgit Frilling

Zentrum für Geriatrie und Gerontologie

an der Universität Hamburg

Albertinen-Haus

Selhopsweg 8–22

22459 Hamburg, Germany

In Reply:

Thus far, the classic scoring systems have used postoperative mortality as their target parameter for a successful intervention. Thus far, scoring systems for assessing postoperative mortality have not focused primarily on the risk of morbidity and reduction in quality of life after cardiac surgery in very old patients. The postoperative result depends crucially on a patient's biological age. The extent of frailty comprises various areas, such as, among others, muscular weakness, cognition or depression, which may be present in individually varying degrees (1). For this reason, the postoperative risk assessment needs to be extended because specifically geriatric risk factors seem to have a substantial influence on postoperative morbidity after cardiac surgery. Frilling et al showed this very convincingly, reporting the uncomplicated assessment of risk constellations by measuring muscular reserve (manual strength) and by testing patients' mental and cognitive status. A scoring system especially for elderly patients would be highly desirable, to quantify the known, specifically geriatric, factors that influence operative risk and thus enable a more precise assessment of the surgical risk. The parameters should be able to be captured by simple methods in the context of routine clinical examinations. Further to a thorough risk

assessment, this would enable individually tailored advice on suitable therapeutic options. Additionally, targeted preoperative conditioning may potentially reduce the operative risk (2).

In addition to postoperative mortality, surgical procedures in elderly patients should be approached with a view to the patients' quality of life. Even if a patient's physical integrity is fully restored and cardiac output is notably improved by the procedure, his or her quality of life may be impaired for the long term. This is due to cognitive impairments, as justifiably introduced into the discussion by Heusinger von Waldegg and Klement. The statement "Since his cardiac surgery, Father has not been who he used to be" is well known to many general practitioners/family physicians. Pre-existing impairments that were mostly asymptomatic before the procedure may deteriorate as a result of the surgery (3). If the risk constellation is high then less invasive procedures should be considered in order to achieve a good functional outcome, such as off-pump coronary artery bypass grafting (OPCAB surgery), minimally invasive aortic valve implantation, or a hybrid intervention. In defining an indication for cardiac surgery in very elderly patients, it is crucial to give greater weight to the patient's postoperative quality of life. This can be achieved only by conducting research in collaboration with geriatric specialists.

DOI: 10.3238/arztebl.2009.0694

REFERENCES

1. Afilalo J, Karunanathan S, Eisenberg MJ, Alexander KP, Bergmann H: Role of frailty in patients with cardiovascular disease. *Am J Cardiol* 2009; 103: 1616–21.
2. Daniels R, van Rossum E, de Witte L, Kempen GJ, van den Heuvel W: Interventions to prevent disability in frail community-dwelling elderly: a systematic review. *BMC Health Serv Res* 2008; 8: 278.
3. Stroobant N, Vingerhoets G: Pre-existing cognitive impairment in candidates for cardiac surgery: an overview. *Heart* 2009; (Epub ahead of print).
4. Friedrich I, Simm A, Kötting J, Thölen F, Fischer B, Silber RE: Cardiac surgery in the elderly patient. *Dtsch Arztebl Int* 2009; 106(25): 416–22.

PD Dr. med. Ivar Friedrich

Klinik für Herz- und Thoraxchirurgie

Universitätsklinikum Halle

Ernst-Grube-Str. 40

06108 Halle, Germany

ivar.friedrich@medizin.uni-halle.de

Conflict of interest statement

The authors of all contributions declare that no conflict of interest exists according to the guidelines of the International Committee of Medical Journal Editors.