Re: M Lovell, K Harris, T Forbes, et al; on behalf of the Peripheral Arterial Disease Coalition. Peripheral arterial disease: Lack of awareness in Canada. *Can J Cardiol* 2009;25(1):39-45.

The level of knowledge of risk factors for peripheral arterial disease also depends on subjects' individual clinical situations

To the Editor:

We congratulate Lovell et al (1) for their interesting survey conducted to assess the level of knowledge of peripheral arterial disease (PAD) in Canada. Along with the United States survey (2), they highlighted the poor level of awareness of the disease and the risk factors associated with this condition. We would like to confront these issues with our findings in France.

We submitted a questionnaire to 150 consecutive patients referred to our vascular laboratory for ultrasound assessment of the lower limb arteries. They either had symptoms suggestive of PAD or had several cardiovascular disease (CVD) risk factors requiring PAD screening. The questionnaire collected the reported CVD risk factors. They were also asked in an open question to quote the risk factors they consider to be important to avoid PAD and/or to limit its worsening. Although these patients were at high risk for PAD, their response rates were even more worrisome than those obtained in the Canadian population survey. Factors quoted the most were smoking (39%), high cholesterol/lipid-rich diet (27%) and alcohol abuse (23%). The response rates regarding diabetes and hypertension were appallingly low (8% and 2%, respectively). Interestingly, 13% considered standing for long periods of time to be a risk factor for PAD, and 6% also considered cold baths to be a way to limit PAD deterioration, suggesting some confusion between PAD and symptoms related to chronic venous disease. To our knowledge, there are no other reports on this possible confusion between these two vascular diseases affecting the lower limbs. We believe that this possible confusion should be systematically screened for during the patient's education.

Another important issue not raised in either North American survey (1,2) was the subjects' level of awareness about a risk factor according to whether they had the risk factor themselves (eg, for a smoker to identify smoking as a risk factor). Table 1 displays the response rates regarding the four main modifiable CVD risk factors. For at least the two main risk factors for PAD – smoking and diabetes – it appears that the level of awareness about the hazards related to these two factors is higher among those who are directly concerned with them. A similar analysis in both the Canadian (1) and United States (2) surveys would be interesting to focus our health education efforts on those who would benefit the most.

Victor Aboyans MD PhD
Philippe Lacroix MD
Marc Laskar MD
Dupuytren University Hospital
Limoges, France

TABLE 1
Risk factors for peripheral arterial disease (PAD) quoted by patients suspected of having PAD, according to their own risk factors

Factors quoted	Patients with the risk factor, %	Patients without the risk factor, %
Smoking	45*	34
Diabetes	15*	2
High cholesterol	29	25
Hypertension	3	0

* $P < 0.05$ versus those without that risk factor

REFERENCES

1. Lovell M, Harris K, Forbes T, et al; on behalf of the Peripheral Arterial Disease Coalition. Peripheral arterial disease: Lack of awareness in Canada. *Can J Cardiol* 2009;25:39-45.
2. Hirsch AT, Murphy TP, Lovell MB, et al; Peripheral Arterial Disease Coalition. Gaps in public knowledge of peripheral arterial disease: The first national PAD public awareness survey. *Circulation* 2007;116:2086-94.

From the Authors:

Thank you for your comments. PAD awareness is dangerously low compared with other CVDs, and needs to be addressed. Your results are lower than ours, notably in people who are suspected of having PAD, which is very concerning. You raise two interesting questions – confusion about vascular disease and subjects' awareness of risk factors depending on whether they had the risk factor themselves, which would be interesting to investigate.

Our survey assessed descriptors of PAD knowledge among the 178 survey respondents who reported to be 'somewhat or very familiar with PAD'. We do not know whether these patients have PAD or have a family member who has PAD. Among this group, knowledge of known PAD risk factors was poor. Nearly one-half of respondents did not know that diabetes could lead to PAD, and 41.6% did not link smoking with PAD. Many did not know that high blood pressure (48.9%) and high cholesterol (51.4%) can cause PAD. Survey respondents cited both obesity or lack of exercise as potential causes of PAD.

Your comments reinforce the need for providing educational programs on PAD online, locally and in national public awareness programs worldwide.

Marge Lovell RN
Division of Vascular Surgery
London Health Sciences Centre
University of Western Ontario
London, Ontario