Published in final edited form as:

Psychophysiology. 2009 May ; 46(3): 655–667.

The instructed context of a motor task modulates covert response preparation and shifts of spatial attention

Elena Gherri[°], Josè Van Velzen[^], and Martin Eimer[°]

[°]School of Psychology, Birkbeck College, University of London

[^]Goldsmiths College, University of London

Abstract

We investigated how covert response preparation is modulated by the instructed cognitive context of a motor task. Participants prepared left-hand or right-hand movements towards or away from the body midline, as indicated by a response cue (S1) presented prior to a Go-Nogo stimulus (S2). Different participants were instructed that response cues specified the response hand or movement direction, respectively. This emphasis on effector versus movement direction selection modulated lateralised ERP components triggered during the S1-S2 interval. Attention shifts during movement preparation were assessed by measuring ERPs to irrelevant visual probes. Enhanced N1 components were found for probes near the effector when effector selection was emphasized, but for probes near the movement target location when movement direction selection was emphasized. Results demonstrate strong top-down contextual biases on motor control and on the locus of spatial attention during response preparation.

Keywords

Attention; Motor control; response preparation; Cognitive control; Event-related potentials

Introduction

During the preparation of a motor response, an abstract action goal is translated into the appropriate motor program that specifies the dimensions on which a movement varies (Keele, 1968; 1981; Rosenbaum, 1983). These movement dimensions are computed through a complex series of sensorimotor transformations where sensory spatial information about the current effector and target locations is eventually translated into an output pattern of motor commands and peripheral muscle activation. Studies of the neural mechanisms that are activated during response preparation have frequently used the “instructed delay period” paradigm where a sensory cue specifies parameters of the movement to be prepared, and is followed after a delay period by an imperative stimulus that signals whether to execute or withhold the prepared movement. Electrophysiological studies in macaques have found activity in several movement-related regions, such as motor and premotor cortices and posterior parietal cortex, during the instructed delay period (Alexander & Crutcher, 1990; Andersen, 1995; Kalaska & Crammond, 1995; Kurata, 1993; Prut & Fetz, 1999; Riehle & Requin, 1989; Wise, 1985). These premotor and parietal brain areas are characterized by highly specific neural connections (e.g. Matelli & Luppino, 2001) which form a series of largely segregated anatomic circuits (Luppino & Rizzolatti, 2000; Fadiga, Fogassi, Gallese

& Rizzolatti, 2000; Rizzolatti, Luppino & Matelli, 1998), each dedicated to a specific sensorimotor transformation. For instance, a parieto-frontal circuit dedicated to the implementation of reaching movements connects the dorsal premotor cortex (PMd) to a specialised parietal area (parietal reach region, PRR) located within the medial intra-parietal sulcus of macaque superior parietal lobe (Andersen & Buneo, 2002). Evidence from functional neuroimaging studies suggests a similar functional organization in humans (see Culham & Kanwisher, 2001, for a review). For example, the fronto-parietal activation found during the instructed delay interval of a reaching movement may be homologous the activity of neural circuits that subserve reaching in monkeys (Filimon, Nelson, Hagler & Sereno, 2007).

It is important to note that the sensorimotor transformations involved in the selection and activation of motor responses are not simply stimulus-response reflexes that are elicited in a rigid and invariant fashion whenever a specific type of movement is being prepared in response to a certain kind of stimulus (Jeannerod, 1997). In order to be flexible, the control of goal-directed behaviour needs to take into account the contextual setting of motor tasks. Accordingly, sensorimotor processes involved in response selection and response preparation have been shown to be modulated in a top-down fashion by cognitive and motivational factors, such as the selection and evaluation of specific types of response-relevant sensory information, the presence of predictive relationships between different spatial and temporal parameters of a motor task, or the anticipated reward value of responses based on past experience. Recent neurophysiological studies have uncovered initial evidence for such top-down effects of cognitive content on motor control processes in prefrontal and primary motor areas (see Johnson, Mason, & Ebner, 2001, for a review) as well as in parietal cortex (area LIP, see Gottlieb, 2007, for a review): When executed in different sensory, decisional, or motivational contexts, the selection of the same movement parameters can be accompanied by a quantitatively or qualitatively different activation pattern of anterior and posterior motor control regions.

In the present study, we used event-related brain potential (ERP) markers of cortical processes involved in the covert preparation of manual reaching movements to investigate whether and how such processes might be affected by a subtle manipulation of the contextual setting for a motor task. Because of their excellent temporal resolution, ERPs provide a useful tool to investigate covert response preparation processes. A number of recent studies have recorded ERPs during the instructed delay period of simple motor tasks that required the preparation of finger lift or key press responses (Eimer, Forster, Van Velzen & Prabhu, 2005; Eimer & Van Velzen, 2006; Eimer, Van Velzen, Gherri & Press, 2006; Praamstra, Boutsen & Humphreys, 2005; Mathews, Dean & Sterr, 2006; Wauschkuhn, Wascher & Verleger, 1997; Van der Lubbe, Wauschkuhn, Wascher, Niehoff, Kompf & Verleger, 2000). These studies have uncovered two lateralised ERP components that elicited during covert response preparation, and are likely to reflect preparatory activity in dorsal premotor and occipitotemporal areas involved in response programming. About 350 ms after the presentation of a response cue, ERPs were found to be more negative at anterior electrodes over the hemisphere contralateral to the cued response side relative to ipsilateral ERPs. Almost simultaneously, an ERP component of opposite polarity (an enhanced positivity contralateral to the cued response side) emerged at lateral posterior recording sites. Interestingly, these lateralised ERP components found during covert response preparation were remarkably similar to ERP components observed in other studies during cued shifts of spatial attention in tasks where the side of a forthcoming target was indicated by a central symbolic cue (c.f., Harter, Miller & Price, 1989; Yamaguchi, Tachiya & Kobayashi, 1994; Nobre, Sebestyen & Miniussi, 2000; Hopf & Mangun, 2000; Eimer, Van Velzen & Driver, 2002). In these tasks, the enhanced anterior negativity contralateral to the side of a cued attentional shift was termed Anterior Directing Attention Negativity

(ADAN), and the posterior contralateral positivity was termed Late Directing Attention Positivity (LDAP). Because ERP components triggered during the instructed delay period of a motor task are very similar to the ADAN and LDAP observed during cued attention shifts, the same labels are now also being used to refer to the lateral anterior and posterior components that are elicited during covert response preparation.

The fact that similar lateralized ERP components are triggered during instructed shifts of spatial attention and during covert movement preparation provides support for the claim of the premotor theory of attention (Rizzolatti, Riggio & Sheliga, 1994) that shared sensorimotor control mechanisms are involved in attentional orienting and in the programming of motor responses. According to this theory, spatial attention is tightly coupled to functionally specialized brain circuits that direct action in space. The main difference between attention and action is that in the former case the pragmatic map is activated but the motor program for that specific action is not executed (Rizzolatti et al., 1994; Sheliga, Riggio & Rizzolatti, 1994; Sheliga, Craighero, Riggio & Rizzolatti, 1997). If the same sensorimotor circuits are responsible for directing both attention and action towards the same location in space, attentional orienting and spatially directed response preparation should give rise to a very similar pattern of ERP activity, and this is exactly what has been observed in several recent studies (Eimer et al., 2006; Praamstra et al., 2005; Praamstra, 2006). Additional evidence for the assumption that covert response preparation and shifts of spatial attention are closely linked comes from previous studies where task-irrelevant visual probes were presented during the response preparation interval, either adjacent to the hand involved in an anticipated response or adjacent to the opposite uncued hand (Eimer et al., 2006; Eimer & Van Velzen, 2006). Here, enhanced visual N1 components were elicited to visual probes presented next to the cued response hand, relative to probes presented near to the opposite uncued hand. Such N1 modulations strongly suggest that attention is shifted towards the location of the hand that is involved in an upcoming response, resulting in a spatially selective modulation of visual processing for probe stimuli that are presented near to this hand.

While there is now converging evidence that covert manual response preparation is associated with a systematic and replicable pattern of lateralised ERP components (ADAN, LDAP), it is not yet known whether these components are associated with the preparation of specific and dissociable spatial parameters of a prepared response. Response preparation usually involves the activation of an effector, the specification of a movement path, and the selection of the target location for a movement. Components such as the ADAN and LDAP that are triggered during response planning might be linked to one or more of these parameters. This issue could not be addressed in most previous ERP studies of response preparation where simple actions such as finger lift or button press responses were used, effector and movement target locations were spatially coincident, and movement direction was constant. To begin to dissociate the effects of different movement parameters on ERP components elicited during response preparation, we have recently measured these components under conditions where response hand and movement direction were cued independently (Gherri et al., 2007). Participants had to prepare and then execute one of four possible reaching movements with their right or left hand towards a target located on the right or left hemifield, and response precues provided either partial or full information about response hand or movement direction. The anterior ADAN component was found to be equally sensitive to effector and movement direction information. In contrast, the posterior LDAP component was substantially larger when response precues specified the effector for an upcoming movement than when they signaled movement direction instead, although this component was also reliably present when only movement direction information was provided. This suggests that the LDAP predominantly reflects mechanisms underlying effector selection, with only a limited contribution from processes involved in the

specification of movement direction. The aim of the present study was to gain further insights into the processes involved in effector and movement direction selection during covert manual response preparation. Most importantly, we investigated for the first time with ERP measures whether and how these processes are modulated by top-down contextual factors. In different blocks, participants had to perform two different types of movement along the horizontal plane. In blocks with outward movements, they had to move one of their hands from a starting position located 5 cm to the left or right of the body midline towards an outer target location on the same side (20 cm from the midline). During blocks with inward movements, start and target locations were reversed, that is, movements were now directed from an outer starting position towards the body midline (see Figure 1). At the start of each trial, a centrally presented visual cue (S1) signaled which of the two alternative movements had to be prepared. This response cue was followed after a 1000 ms interval by an imperative Go-Nogo stimulus (S2) that instructed participants to either execute or withhold the prepared movement. In addition, a task-irrelevant visual probe stimulus was presented on each trial. This probe appeared randomly and with equal probability at one of the four start or end positions for a movement on the left or right side, either during the covert response preparation interval (900 ms after cue onset), or during movement execution (200 ms after the onset of a manual response).¹

The critical manipulation of the present experiment was designed to investigate the role of top-down contextual factors during covert manual response preparation. Participants were randomly assigned to two groups that were given different instructions with respect to the nature of the response information provided by the cue. One group of participants was told that the response cue signaled whether the left or right hand had to be prepared for an upcoming response (effector cue instruction). The other group of participants was informed that the cue specified the direction (leftward or rightward) of an upcoming movement (direction cue instruction). Thus, even though cues always provided full response information for both groups of participants (i.e., they specified unequivocally which of the two alternative movements was to be prepared in a given trial), these two instructions either emphasized the selection of an effector or the selection of movement direction. If cortical processes involved in response selection can be biased in a top-down fashion by this subtle contextual manipulation, the pattern of ERP effects obtained during covert response preparation should differ systematically as a function of which instruction was provided.

Two sets of analyses were conducted. To obtain components sensitive to covert response preparation (ADAN, LDAP), ERPs triggered in the S1-S2 interval were measured for cues signaling a response on the left or right side, separately for inward and outward movements.² To investigate spatially specific modulations of visual processing during the covert response preparation interval that indicate shifts of spatial attention towards the cued response side, ERPs were also obtained in response to the visual probe stimuli on the cued and uncued side, separately for probes presented at the start and end positions for manual movements.

¹Visual probes were presented with equal probability during response preparation or during response execution in order to reduce their potential to be used as a temporal marker (i.e., as a warning signal that S2 presentation is imminent). However, ERPs were only measured for probes presented during the response preparation interval, but not for probes presented during response execution, as the latter are contaminated by motor, somatosensory, and visual components associated with response execution and vision of the moving hand.

²In addition to the two lateralized components on which our analyses were focused (ADAN and LDAP), another component that is typically elicited during the S1-S2 interval of cued response tasks is the contingent negative variation (CNV; Walter, Cooper, Aldridge, McCallum & Winter, 1964). The CNV was not analysed in our study because it is essentially non-lateralised (for a review see Brunia & Van Boxtel, 2001), whereas our critical manipulations (movement types, response cue instructions) had an inherent left-right component, and were designed to affect lateralised components such as ADAN and LDAP. CNV amplitudes are typically larger for more difficult tasks (e. g., Brunia, 1993), whereas task difficulty did not differ between different movement types or response cue instructions in our study (see behavioural performance).

ADAN and LDAP components were predicted to be differentially affected by movement type (inward versus outward movements). The ADAN component was expected to be more pronounced during blocks where outward movements were being prepared, because for these movements, the spatial codes involved in effector and movement direction selection were always congruent (i.e., the left hand had to be moved towards the left, or the right hand towards the right). In contrast, during blocks with inward movements, these two codes were incongruent (i.e., the left hand had to be moved towards the right, or the right hand towards the left). If the ADAN component reflects processes involved in the selection of the left versus right hand, as well as processes underlying the selection of a leftward versus rightward movement direction, as suggested by our previous study (Gherri et al., 2007), it should be strongly affected by the congruency versus incongruency of their respective spatial codes. In contrast to the ADAN, the posterior LDAP component primarily reflects effector selection (Gherri et al., 2007). Furthermore, results from an ERP study where ADAN and LDAP components were measured during cued shifts of tactile attention towards the left or right hand (Eimer, Forster, Fieger, & Harbich, 2004) have suggested that the selection processes that give rise to the LDAP are mediated by representations of *external* space. In this study, the LDAP (but not the ADAN) was larger in amplitude when hands were positioned far apart than when they were located closely together, in line with the assumption that external spatial coordinates play an important role for the LDAP component. If hand posture affected LDAP amplitudes in a similar fashion also during covert movement preparation, this component should be larger during the preparation of inward movements, where the external location of hands is more peripheral, than during outward movement preparation, where the hands are located close to the body midline (see Figure 1). The critical new question addressed in the present study was whether any such effects of movement type on ADAN and LDAP components would be modulated in a top-down fashion by the relative emphasis on effector selection or movement direction selection that differed between the two instruction groups.

If situational context of a motor task, as specified via response cue instructions, can affect response programming, such a top-down bias might also have an impact on shifts of spatial attention that are triggered as a result of manual response preparation processes. To investigate this possibility, we compared the effects of covert response preparation on visual ERPs to irrelevant probe stimuli for the two instruction groups. Based on earlier findings (Eimer et al., 2006; Eimer & Van Velzen, 2006), we expected to find enhanced N1 components for visual probes presented on the side cued for an upcoming movement relative to probes on the opposite uncued side. Such a result would further confirm the presence of spatially selective response preparation effects on visual processing, as postulated by the premotor theory of attention. The critical new question was whether the locus of these attentional modulations of visual ERPs would shift from the initial starting position of a hand to the target position for an intended movement, or vice versa, depending on which instruction was provided. When effector selection is emphasized (effector cue instruction), effects of response preparation on visual processing, as reflected by enhanced N1 amplitudes, should be found for probes at the initial effector position, but not for movement target locations. In contrast, when movement direction is emphasized (direction cue instruction), the opposite pattern might emerge, with visual N1 enhancements now focused at the target location for a movement.

Methods

Participants

Twenty-five healthy volunteers participated in this experiment. Three were excluded due to poor eye gaze control in the cue-target interval (see below), and two others were excluded due to an insufficient number of trials after artifact rejection. Thus twenty participants

remained in the sample. These were randomly assigned to two experimental groups of 10 participants that differed with respect to the response instructions given (see below). The group receiving effector cue instructions contained three males and seven females (19 – 34 years old; average age: 25.3 years; one left-handed participant). The group receiving direction cue instructions contained five males and females (21 – 35 years old; average age: 28.6 years; all right-handed). All participants had normal or corrected-to-normal vision by self-report and were naïve as to the purpose of the experiment.

Apparatus, Stimuli and Procedure—Participants sat in a dimly lit sound attenuated cabin in front of a computer screen at a viewing distance of approximately 55 cm. A custom-made response device was placed on the table just below the computer screen (see Figure 1). The device contained four response keys (two start keys and two target keys) that were horizontally aligned on a wooden panel (60 × 48 cm). This panel was tilted at an angle of 30° and its midline was aligned with the screen centre. Two infra-red keys were used as start keys while the two target keys were normal response buttons. Infra-red keys consisted of a transmitter and a receiver LED located on either side of the middle segment of each index finger in the resting position. A response was registered when the index finger was lifted from the resting position allowing the light beam of the transmitter LED to reach the receiver LED. Start and target keys were located in different positions in blocks that required inward or outward movements. During outward movements, start keys were located 5.5 cm to the left and right of the centre of the panel, while the two target keys were placed further outwards (25.5 cm from the centre). During inward movements, these positions were swapped, with start keys located at the two outer positions, and target keys at the two inner positions. Thus, the distance between each start and corresponding target key (movement length) was kept constant at 20 cm.

Each trial started with a 100 ms presentation of a visual response precue (S1) at fixation. This cue consisted of two adjacent triangles (visual angle: $3.5^\circ \times 2.5^\circ$). One triangle was red, the other blue, and they always pointed in opposite directions ('<>' or '><'). A central fixation cross, located in the space between the two triangles, was present throughout the experimental blocks. The required movement for each trial was signalled by the direction of one of the triangles. For half of the participants, blue triangles were relevant, while for the other half red triangles were relevant. Relevant left-pointing or right-pointing triangles were presented with equal probability to the left or right of fixation. On each trial, the response cue was followed after an empty interval of 1000 ms by the letter 'G' or 'S' (S2), indicating to execute ('go') or withheld ('stop') the movement. This letter was presented at fixation for 100 ms, replacing the fixation cross during this period. To investigate whether covert manual response preparation results in spatially selective modulation of visual processing, an irrelevant visual probe stimulus was presented on each trial, either 900 ms after cue onset (i.e., 200 ms prior to the onset of the imperative Go-Nogo stimulus), or 200 ms after movement onset, as recorded by the infra-red keys. Probes consisted of 100 ms illumination of one of four ensembles of green LEDs that were positioned 2.5 cm above each start and target key on the left and right side (see Figure 1). Participants were instructed to completely ignore these probe stimuli throughout.

The experiment consisted of 16 experimental blocks of 96 trials each. A Go S2 was presented on 80 trials per block, and a Nogo S2 on the remaining 16 trials. On each trial an irrelevant visual probe was presented with equal probability and in random order next to one of the four start or target keys, either 900 ms cue onset or 200 ms after movement onset. Thus every block contained five Go trials and one Nogo trials for each combination of cued response side (left versus right), probe location (next to hands / start keys versus next to movement target keys), probe side (left versus right), and probe onset (early versus late). Trials were presented in a pseudo-random sequence. In eight successive blocks, participants

performed one of two different types of movements. In 'Inward movement' blocks, they had to prepare a movement from an outer start key towards an inner target key (i.e., towards the body midline). In 'Outward movement' blocks, they had to prepare a movement from an inner start key towards an outer target key (i.e., away from the body midline). The order in which these two blocked movement type conditions were delivered was counterbalanced across participants. One training block of 48 trials was run before each movement type condition.

Participants were randomly assigned to two different instruction groups. Ten participants were told that the response cue specified which effector (right versus left hand) to prepare in anticipation of an upcoming Go S2 ('Effector cue instruction'). The other ten participants were told instead that the response cue indicated the direction of the movement (leftward versus rightward) to be prepared ('Direction cue instruction'). These task instructions were shown on the computer screen prior the start of each block. All participants placed their forearms on the response panel with their index fingers on the start keys (resting position), and maintained a body posture such that their midline was aligned with the centre of the screen and response panel. They were explicitly encouraged to maintain central eye fixation, to use the information provided by S1 to covertly prepare the cued movement, while maintaining the resting position with their hands on the start keys until the imperative stimulus (S2) was displayed on the screen.

On Go trials, participants had to perform a movement from one start to one target key, on the side that was indicated by the response cue, press the target key, and then place their hand back on the infra-red start key in the resting position infra-red start keys in order to start the next trial. Therefore, the time interval between the onset of the imperative stimulus on one trial and the onset of the response cue on the next trial was a variable function of total movement time. The next response cue was presented 600 ms after the hand had returned to the start key. On Nogo trials, participants had to leave their hands on the start keys and to wait for the next response cue presentation. Here, the interval between the onset of the imperative stimulus on one trial and the onset of the response cue on the next trial was randomly selected between 1850 ms and 2050 ms.

EEG Recording and Data analysis

EEG was recorded with Ag-AgCl electrodes and linked-earlobe reference from Fpz, F7, F3, Fz, F4, F8, FC5, FC6, T7, C3, Cz, C4, T8, CP5, CP6, P7, P3, Pz, P4, P8, and Oz (according to the 10-20 system), and from OL and OR (located halfway between O1 and P7, and O2 and P8, respectively). Horizontal EOG was recorded bipolarly from the outer canthi of both eyes. Electrode impedance was kept below 5 k Ω , and the impedances of the earlobe electrodes were kept as equal as possible. Data were recorded with a bandpass filter from 0 to 40 Hz, and a digitisation rate of 200 Hz. Trials with eyeblinks (Fpz exceeding ± 80 μ V), horizontal eye movements (HEOG exceeding ± 30 μ V), or other artefacts (a voltage exceeding ± 80 μ V at any electrodes) in the S1-S2 interval were excluded prior to data analysis. To detect systematic deviations of eye position indicating residual tendencies to move the eyes towards the side of the cued response, averaged HEOG waveforms in the S1-S2 interval in response to left versus right cues were examined for each participant. HEOG deviations exceeding ± 3 μ V led to the disqualification of three participants. For the 20 participants kept in the sample, overall residual HEOG deviation (calculated as a difference between grand averaged ERP elicited by left versus right cues) was below 0.45 μ V throughout the S1-S2 interval. ERPs were averaged relative to a 100 ms pre-S1 baseline prior for the 1000 ms time interval following S1 onset.

One set of analyses focused on *ADAN and LDAP components elicited during the covert response preparation interval*. ERP mean amplitude values were obtained for a time window

between 400 and 700 ms after S1 onset (where the ADAN and the early phase of the LDAP was observed in earlier response precueing experiments), and for a 700-900 ms post-cue time window (where the later phase of the LDAP was previously observed). An initial analysis assessed the overall presence and distribution of ADAN and LDAP components on the basis of ERP mean amplitudes to response cues instructing participants to prepare a response on the left versus right side, collapsed across both movement types (inward versus outward) and instruction (effector versus direction cue), with a 3-way repeated measures ANOVAs conducted separately for lateral anterior, central, and posterior sites. In these analyses the factors cued response side (left versus right), laterality (electrode ipsilateral versus contralateral to cued response side) and electrode site (F7/8 versus F3/4 versus FC5/6, for the anterior analysis, C3/4 versus T7/8 versus CP5/6, for the central analysis, and OL/R versus P3/4 versus P7/8, for the posterior analysis) were used as within-subjects variables. The presence of ADAN and LDAP components will be reflected by significant main effects of the factor laterality. Next, we analyzed the effects of movement type (inward versus outward), and, critically, possible top-down effects of response instruction on these lateralised components. These analyses were restricted to the electrode pairs where ADAN or LDAP components were elicited most reliably. ERP mean amplitude values obtained for the 400 and 700 ms and for the 700-900 ms post-cue time windows were entered into a 4-way mixed design ANOVA with the factor instruction (effector versus direction cue instruction) as between-subject variable and cued response side (left versus right), laterality (electrode ipsilateral versus contralateral to cued response side), and movement type (inward vs. outward) as within-subject variables.

The second set of analyses was conducted for *visual ERPs in response to probe stimuli* presented 900 ms after S1 onset. These were computed for the 300 ms interval after probe onset (relative to a 100 ms baseline prior to probe onset), separately for each combination of probe location (next to hands / start keys versus next to target keys) and movement preparation (probe located in hemispace of cued movement versus opposite hemispace), but were collapsed across both movement types (inward and outward).³ Because inspection of these visual ERPs suggested that they might be affected by overlapping slow wave activity, a high-pass filter (2 Hz, 48 dB/oct) was applied to the raw EEG data (prior to EEG epoching, baseline correction, artefact rejection, and averaging) to remove this activity. All analyses of visual ERPs reported here are based on these filtered data. Mean amplitudes were computed within latency window centered on the peak amplitude of P1 and N1 components (P1: 90 – 120 ms post-stimulus; N1: 140 – 180 ms post-stimulus), and were analyzed with a 5-way mixed design ANOVAs, separately for midline electrodes (Fz, Cz, Pz) and for lateral anterior, central and posterior sites. Analyses included the factor instruction (effector versus direction cue instruction) as a between-subjects variable and the factors probe location (next to hands / start keys versus next to target keys), cued response side (probe located on the side of the cued movement versus on the opposite side), hemisphere (left versus right, for lateral electrodes only) and electrode site (F7/8, F3/4, FC5/6, for anterior electrodes; C3/4, T7/8, CP5/6 for central electrodes; OL/R, P3/4, P7/8 for posterior electrodes; Fz, Cz, Pz for midline electrodes) as within subjects variables.

Behavioural performance was assessed to confirm that there were no systematic differences in task difficulty between inward and outward movements and between effector and direction cue instructions. Both reaction times (RTs, i.e., latencies of start key responses) and movement times (MTs, i.e., intervals between start key and target key responses) were

³Note that as a result of collapsing across inward and outward movements, these visual ERPs show the combined response to probe stimuli at near and far locations. For example, for blocks where inward movements were required, probes next to the start key / effector were located at the two outer locations, whereas these probes were located at the two inner locations in blocks with outward movements (see Figure 1).

measured. All trials with missing and premature responses (i.e., start key responses prior to S2 onset), with RTs shorter than 100 ms (anticipations) or longer than 1000 ms, and with MTs exceeding 1000 ms were excluded from analysis. RTs and MTs recorded in the remaining correct Go trials were separately entered into a 3-way mixed design ANOVAs with instruction (effector versus direction cue) as between-subjects variable and movement type (inward versus outward) and response hand (left versus right) as within-subjects variables. In these analyses, main effects of the factors movement type or instruction would indicate that there were task difficulty differences between conditions.

All statistical analyses were conducted with a sample of 20 participants. Follow-up analyses conducted separately for each instruction group were carried out over a sample of 10 participants each. For all analyses, Greenhouse-Geisser adjustments to the degrees of freedom were applied where appropriate.

Results

Behavioural performance

RTs did not differ between inward and outward movements ($M=379$ ms, $SD=60$ ms and $M=371$ ms, $SD=53$ ms, respectively, $F(1,18)=.6$, $MSE=1848$, $p=.4$), or between groups with effector ($M=370$ ms, $SD=58$ ms) and direction cue instructions ($M=380$ ms, $SD=50$ ms; $F(1,18)=.17$, $MSE=11703$, $p=.7$). A main effect of response hand ($F(1,18)=15.0$, $MSE=194$, $p<.001$) was due to the fact that RTs were faster for left-hand than for right-hand movements ($M=370$ ms, $SD=50$ ms versus $M=380$ ms, $SD=56$ ms). MTs did not differ between effector and direction cue instruction groups ($M=266$ ms, $SD=57$ and $M=275$ ms, $SD=146$ ms, respectively, $F(1,18)=.03$, $MSE=49156$, $p=.9$), but were faster for outward relative to inward movements ($M=257$ ms, $SD=102$ ms versus $M=283$ ms, $SD=119$ ms; $F(1,18)=7.9$, $MSE=1732$, $p<.011$). This difference was only present with effector cue instructions ($M=242$ ms, $SD=52$ ms versus $M=290$ ms, $SD=67$ ms) but not with direction cue instructions ($M=277$ ms, $SD=159$ ms versus $M=273$ ms, $SD=136$ ms), as reflected by a movement type $\times$ instruction interaction ($F(1,18)=5.5$, $MSE=466$, $p<.030$). In contrast to RTs, MTs were shorter for right-hand as compared to left-hand movements ($M=262$ ms, $SD=106$ ms versus $M=279$ ms, $SD=111$ ms; main effect of response hand: $F(1,18)=12$, $MSE=466$, $p<.003$). Premature responses (before S2 onset) occurred on the 1% of all trials. Anticipations were measured in 0.1%, RTs beyond 1000 ms in 0.5%, and MTs beyond 1000 ms in 0.16% of all Go trials. On 17.7% of all Nogo trials, an index finger movement was registered by one of the infra-red response keys. On 16.4% of all Go trials, response onset latencies were longer than 1000ms, or no manual responses were recorded.⁴

Lateralised ERPs components triggered during reaching movement preparation

Figure 2 shows ERPs in the 900 ms interval after response cue onset at lateral anterior, central and posterior electrodes ipsilateral (dashed line) and contralateral (solid line) to the side of cued movement. Waveforms are collapsed across different movement types (inward versus outward) and instruction groups (effector versus direction cue instructions). As expected, both ADAN and LDAP components were elicited during the covert movement preparation interval (see Figure 2, right panels for scalp distribution). The presence of the ADAN was confirmed by a main effect of laterality at anterior electrode pairs during the 400-700 ms post-cue interval ($F(1,18)=8.1$, $MSE=.16$, $p<.011$). There was also a laterality $\times$ electrode site interaction ($F(2,36)=15.7$, $MSE=.08$, $p<.001$), and follow-up analyses conducted separately for each anterior electrode pair confirmed the presence of a reliable

⁴The relatively high percentage of False Alarms and missed responses is in part due to the fact that the infrared system used to measure manual response onset did not always operate with absolute reliability.

ADAN at F3/4 ($F(1,18)=30.5$, $MSE=.07$, $p<.001$) and at F7/8 ($F(1,18)=5.1$, $MSE=.08$, $p<.036$) but not at FC5/6 ($F(1,18)=.18$, $MSE=.09$, $p=.68$).

As can be seen in Figure 2, the LDAP component was present at central and posterior electrode pairs. In the 400-700 post cue interval, main effects of laterality (both $F(1,18)>15.7$, $MSE<1.1$, both $p<.001$) were obtained at lateral central and posterior electrodes, reflecting the early phase of the LDAP. These were accompanied by laterality $\times$ electrode site interactions (both $F(2,36)>9.7$; both $p<.002$), and follow-up analyses revealed the presence of a significant early LDAP at all posterior electrode site (all $F(1,18)>27.7$, $MSE<.59$, all $p<.001$), as well as at T7/8 and CP5/6 (both $F(1,18)>9.7$, $MSE<.24$, both $p<.006$), approaching significance at C3/4 ($F(1,18)=3.8$, $MSE=.21$, $p=.067$). During the later analysis window (700-900 ms post-stimulus), a significant laterality effect was present at posterior electrode sites ($F(1,18)=25.7$, $MSE=1.03$, $p<.001$), reflecting the later phase of the LDAP. This effect only approached significance at central electrode sites ($F(1,18)=3.9$, $MSE=.7$, $p=.065$).

To investigate the effects of movement type (inward versus outward) and instruction (effector versus direction cue) on ADAN and LDAP components, additional analyses were carried out for the two electrode pairs where these components were most reliably elicited (F3-F4 for the ADAN, P7-P8 for the LDAP, see Figure 2). Figure 3 shows ERPs elicited at these electrodes during inward and outward movement preparation (panels A-D), together with difference waveforms obtained by subtracting ERPs at electrodes contralateral to the side of cued movement from ipsilateral electrodes (panels E-F). In these difference waveforms, a contralateral negativity is indicated by positive values (downward-going deflections), and a contralateral positivity by negative values (upward-going deflections).

As can be seen in Figure 3 (top panels), the ADAN component was clearly present at F3-F4 (main effect of laterality in the 400-700 ms post-cue interval: $F(1,18)=30.5$, $MSE=.07$, $p<.001$), but was more pronounced during outward as compared to inward movement preparation. This was substantiated by a significant laterality $\times$ movement type interaction ($F(1,18)=6.7$, $MSE=.04$, $p<.02$). However, follow-up analyses conducted separately for each movement type revealed that the ADAN was significantly present not only during outward movement preparation ($F(1,18)=29.9$, $MSE=.065$, $p<.001$), but also when inward movements were being prepared ($F(1,18)=9.7$, $MSE=.04$, $p<.006$). In contrast, the difference between effector and direction cue instructions had no effect on the ADAN (laterality $\times$ instruction, $F(1,18)=2.4$, $MSE=.07$, $p=.14$), and there were also no laterality $\times$ movement type $\times$ instruction interaction, $F(1,18)=.16$, $MSE=.04$, $p=.7$.⁵

Figure 3 (bottom panel) shows the LDAP component at lateral posterior electrode pair P7-P8. During the 400-700 ms measurement interval, a main effect of laterality ($F(1,18)=28.9$, $MSE=.6$, $p<.001$) confirmed the presence of the early phase of the LDAP, which was not modulated by the difference between inward and outward movements (laterality $\times$ movement type: $F(1,18)=.15$, $MSE=.11$, $p=.7$). In the subsequent 700-900 ms time window, the late phase of the LDAP was reliably present (main effect of laterality: $F(1,18)=23.4$, $MSE=.59$, $p<.001$). As can be seen in Figure 3 (panel F), this late LDAP was more pronounced during inward as compared to outward movement preparation, as reflected by a significant laterality $\times$ movement type interaction ($F(1,18)=5.2$, $MSE=.22$, $p<.035$). Follow-up analyses conducted separately for each movement type revealed that the late phase of the LDAP was significantly present not only during inward movement preparation

⁵Follow-up analyses conducted separately for the two different movement types confirmed the absence of significant laterality $\times$ instruction interactions during inward movements ($F(1,18)=2.7$, $MSE=.04$, $p=.12$), as well as during outward movements ($F(1,18)=.84$, $MSE=.065$, $p=.37$).

($F(1,18)=19.8$, $MSE=.59$, $p<.001$), but also when outward movements were being prepared ($F(1,18)=15.6$, $MSE=.24$, $p<.001$).

Importantly, and in contrast to the ADAN, the LDAP was systematically affected by the type of instruction that participants received with respect to the response cue. This differential instruction effect is illustrated in Figure 4, which shows ERPs elicited at P7-P8 during the preparation of inward and outward movements, and the resulting difference waveforms, separately for the group of participants that had received direction cue instructions (top panels), and for the effector cue instruction group (bottom panels). When participants were told that the response cue specified movement direction (see Figure 4, top panels), the early phase the LDAP component was more pronounced during outward movement preparation (i.e., movements directed away from the body midline) than during inward movement preparation (i.e., movements towards the body midline). In contrast, with effector cue instructions (see Figure 4, bottom panels), the early LDAP was larger during inward movement preparation (when hands were located further apart) than during outward movement preparation (when hands were located close to the midline). This is further illustrated in panels F and H of Figure 4, which shows the absolute size of the LDAP amplitudes for the 400-700 ms and the 700-900 ms time intervals, obtained by subtracting mean ERP amplitudes at electrode ipsilateral to the cued response side from mean amplitudes obtained at the corresponding contralateral electrode, separately for inward (black bars) and outward movements (grey bars). This pattern of results was substantiated by a significant laterality $\times$ instruction $\times$ movement type interaction ($F(1,18)=9.5$, $MSE=.114$, $p<.006$) in the 400-700 ms post-cue time window. Follow-up analyses were conducted separately for the two different instruction groups. For participants receiving direction cue instructions, a reliable movement $\times$ laterality interaction ($F(1,9)=7.4$, $MSE=.056$, $p<.024$) confirmed that the early LDAP was larger during outward movement preparation. For participants with effector cue instructions, this interaction was almost significant ($F(1,9)=4.0$, $MSE=.17$, $p=.07$), due to the strong trend for a larger early LDAP during the preparation of inward movements. During the late phase of the LDAP (700-900 ms post-cue), no laterality $\times$ instruction, or laterality $\times$ instruction $\times$ movement type interactions were present ($F(1,18)=.02$, $MSE=.62$, $p=.9$ and $F(1,18)=.62$, $MSE=.22$, $p=.44$, respectively).

Visual ERPs to task irrelevant probe stimuli

Figure 5 shows visual ERPs elicited by irrelevant visual probe stimuli presented 900 ms after response cue onset, displayed separately for participants with direction cue instructions (top) and effector cue instructions (bottom), and for trials where probes were presented next to the hands / start keys (left), or next to the target keys (right). Solid lines represent ERPs to probes presented on the side of a cued movement, while dashed lines represent ERPs to probes on the opposite uncued side. As can be seen from Figure 5, advance information about the side of a to-be-prepared response had a systematic effect on the processing of visual probes, as N1 components tended to be generally larger for probes that were presented on the cued response side. However, and importantly, the spatial locus of this response preparation effect on visual N1 amplitudes was strongly modulated by response cue instructions. For participants who were told that the response cue specified the direction of an upcoming movement (direction cue instruction), a modulatory effect of response preparation on visual N1 components was present for probes presented next to the cued versus uncued movement target location, whereas no such effect emerged for probes that were presented next to the hands / start keys (Figure 5, top panel). In contrast, participants who were informed that the cue specified the hand to be prepared (effector cue instruction) showed larger effects of response preparation on visual N1 components when probes were presented close to the hands than when they were located next to the response target keys (Figure 5, bottom panel).

These informal observations were substantiated by statistical analyses. While there were no significant response cueing effects on visual P1 amplitude (90 – 120 ms post stimulus), the N1 component to visual probes was strongly modulated by movement preparation. In the N1 time range (140 – 180 ms post stimulus), main effects of cued response side on ERP mean amplitudes were present at lateral posterior, lateral central, lateral frontal and midline sites (all $F(1,18) > 5$, all $MSE < 1.95$, all $p < .04$), demonstrating that N1 amplitudes were generally enhanced for visual probes that were presented on the side of a cued manual reaching movement. However, this effect was jointly modulated by response cue instructions (emphasizing effector or movement direction) and the location of the visual probes (next to the hands or next to the target keys), as reflected by cued response side $\times$ probe location $\times$ instruction interactions that were significant at all lateral posterior, lateral central and midline sites (all $F(1,18) > 6.6$, all $MSE < .7$, all $p < .019$) while this interaction approached significance at lateral frontal electrodes ($F(1,18) = 4.2$, $MSE = .7$, all $p = .055$).

This three-way interaction was further investigated in follow-up analyses carried out separately for each of the four combinations of response cue instruction (effector versus direction cue) and probe location (next to the hands versus next to the target keys). For participants who were informed that the cue specified the direction of an upcoming response, a significant N1 modulation for probes presented close to the cued versus uncued target key emerged at all electrode sites (all $F(1,9) > 9.6$, all $MSE < 1.1$, all $p > .013$), while no such modulation emerged when probes were presented close to the cued versus uncued hand (all $F(1,9) < 1$, $MSE < 2$, all $p > .65$). For participants who were told that the response cue indicated the hand to be prepared for an upcoming movement, significant N1 modulations were found for probes presented close to the cued versus uncued hand at lateral frontal, lateral central, lateral posterior and midline electrode sites (all $F(1,9) > 5.7$, $MSE < .8$, all $p < .041$). In contrast, no reliable effects of cued response side emerged for probes that were presented close to the cued versus uncued target key (all $F(1,9) < 3.3$, $MSE < 1.5$, all $p > .103$).

Discussion

To investigate cortical processes activated during the programming of manual reaching movements, and to find out whether and how such processes might be susceptible to a top-down bias by contextual factors, we measured ERP markers of covert manual response preparation (ADAN and LDAP components) and spatially selective visual processing (visual N1 component) in an instructed delay period paradigm, where participants prepared manual reaching movements, as instructed by response cues presented at the start of each trial. Analyses of behavioural data confirmed that there were no systematic differences in task difficulty between different movement types or response cue instructions, thus making it unlikely that ERP components elicited during the S1-S2 interval would be affected by factors such as anticipated cognitive effort or task complexity.

Several results obtained in this study confirmed findings from other recent ERP investigations of covert manual response preparation (e.g., Eimer et al., 2005, 2006; Eimer & Van Velzen, 2006; Gherri et al., 2007; Praamstra et al., 2005; Praamstra, 2006; Mathews et al., 2006; Wauschkühn et al., 1997; Van der Lubbe et al., 2000). As expected, ADAN and LDAP components were triggered in the interval between the response cue and the subsequent imperative Go-Nogo stimulus (see Figure 2), thus confirming that these components can be reliably observed during covert response preparation. The anterior ADAN component was larger in blocks where outward movements had to be prepared than in blocks where inward movements were required (Figure 3, top panel). With outward movements, the spatial codes involved in effector and movement direction selection were always compatible (e.g., when preparing a leftward movement with the left hand), whereas these codes were always incompatible for inward movements (e.g., when preparing a

rightward movement with the left hand). The observation that the ADAN component was attenuated when spatial codes were incompatible provides further evidence that the ADAN reflects the joint contribution of processes involved in the selection of the left versus right hand, and of processes that underlie the selection of a leftward versus rightward direction for a movement (see also Gherri et al., 2007).

In contrast to the ADAN, the late phase of the LDAP component was more pronounced for inward as compared to outward movements (Figure 3, bottom panel). This difference is most likely due to the fact that initial hand positions were different for these two types of movements. During the preparation of inward movements, hands were placed much further apart than during the preparation of outward movements, where they were positioned close together (see Figure 1). The distance of effectors from the body midline has previously been found to modulate the LDAP during covert shifts of tactile attention to the left or right hand (Eimer et al., 2004). The current results demonstrate that effectors position affects LDAP amplitudes not just during spatial orienting, but also during covert response preparation, and thus suggest that the LDAP primarily reflects the selection of response hands, rather than movement direction (as already suggested by our previous findings, see Gherri et al., 2007). Furthermore, they also imply that this selection is mediated by representations that code the location of the hands in terms of external space (i.e., their distance from the body midline). This conclusion is further supported by the results of a recent unpublished study from our lab. This study was similar to the experiment reported here, except that start and end positions for all reaching movements were now equidistant from the body midline, so that movement paths were always parallel to the median plane. In different blocks, these movements were either executed near to the body midline (5 cm to the left or right), or at a distance of 25 cm on the left or right side. While the ADAN was unaffected by this distance manipulation, the LDAP was significantly larger during the preparation of far as compared to near movements, again suggesting that this component is sensitive to the representation of effector locations in an external spatial coordinate system.

The critical new question addressed in the present study was whether cortical covert response preparation processes are modulated by top-down contextual effects. For different groups of participants, instructions either stressed the link between response cues and effector selection (effector cue instruction), or the link between response cues and the direction of an upcoming movement (direction cue instruction). Even though all response cues were in fact informationally equivalent regardless of which instruction was provided (i.e., they always specified which of two alternative reaching movements would have to be executed in response to an upcoming Go stimulus), this instruction manipulation still had systematic effects on the posterior LDAP component, as well as on visual N1 components triggered in response to task-irrelevant probe stimuli. The differential effect of instructions on the LDAP is shown in Figure 4. When instructions emphasized effector selection, the initial phase of the LDAP component in the 400-700 ms time window after response cue onset was larger for inward than for outward movements (Figure 4, bottom panel), that is, for the movement type where hands were wide apart during response preparation. This result is in line with the general assumption that the LDAP primarily reflects the selection of effector locations that are mediated by representations of external space (see above). However, and critically, when instructions emphasized movement direction selection instead, this pattern was reversed, as the early phase of the LDAP was now more pronounced during the preparation of outward movements. When coded in terms of movement direction, outward movements are associated with a stronger spatial left/right bias than inward movements, as outward movements were directed towards the left or right periphery, while inward movements were directed towards the body midline (Figure 4, top panel). Overall, the results as shown in Figure 4 provide clear evidence for a top-down contextual bias on cortical response selection processes: Depending on which instruction

was provided, priority was given either to the selection of an effector, or to the selection of a specific response direction. LDAP amplitudes were enhanced whenever the prioritised response parameter was associated with a more lateral spatial location, and attenuated whenever this parameter specified a location closer to the body midline.

While these differential effects of response cue instructions on the posterior LDAP component are indicative of a contextual modulation of cortical response preparation processes, even more direct evidence for such a top-down bias comes from the pattern of ERP modulations observed in response to task-irrelevant probe stimuli. As can be seen in Figure 5, visual N1 components were generally enhanced when probes were presented on the side that was cued for an upcoming response relative to probes presented on the opposite uncued side. This finding confirms previous observations that covert manual response preparation results in spatially specific modulations of visual processing (Eimer et al., 2006; Eimer & Van Velzen, 2006), and thus provides further evidence that response preparation processes are closely linked to attentional shifts towards response-relevant locations, as postulated by the premotor theory of attention (Rizzolatti et al., 1994). The critical new finding was that such attentional modulations of visual processing are not always confined to locations next to the effector involved in a cued response, but that their locus in external space can instead be shifted by contextual factors. For the group of participants that were given effector cue instructions (Figure 5, bottom panel), spatially selective effects of response preparation on visual N1 components were reliably elicited when probes were located next to the hands, but not when they were presented next to the target keys. This difference might simply be due to the fact that these visual probes were always presented during the covert response preparation interval when hands were still in their resting position. However, the N1 results observed for the group that received direction cue instructions instead (Figure 5, top panel) shows that this explanation cannot be correct. For this group, response preparation effects on visual N1 components were only present when probes were presented next to the target location for a cued movement, but not when probes were presented next to their starting position, even though hands were still located there. As these spatially selective modulations of visual N1 amplitudes are likely to reflect shifts of attention that are triggered during covert manual response preparation, this pattern of results suggests that such attention shifts are subject to a strong top-down contextual bias. When the importance of effector selection is stressed, the focus of spatial attention remains close to the effector throughout the response preparation interval. In contrast, when the importance of movement direction is emphasized instead, attention is shifted towards the goal location of an anticipated reaching movement well before this movement is initiated.

In summary, the present experiment has found new and compelling evidence for strong top-down contextual biases that are active during response programming, and systematically affect the covert preparation of manual reaching movements. These results show that even when full response information is available, sensorimotor processes involved in preparation of motor responses are by no means rigid and automatic, but are instead modulated in a top-down fashion by cognitive factors related to expectations, strategies, and the instructed context for a motor task.

Acknowledgments

This research was supported by a grant from the Medical Research Council (MRC). M.E. holds a Royal Society-Wellcome Research Merit Award.

References

1. Alexander GE, Crutcher MD. Preparation for movement: Neural representations of intended direction in three motor areas of the monkey. *Journal of Neurophysiology*. 1990; 64:133–150. [PubMed: 2388061]
2. Andersen RA. Encoding of intention and spatial location in the posterior parietal cortex. *Cerebral Cortex*. 1995; 5:457–469. [PubMed: 8547792]
3. Andersen RA, Buneo CA. Intentional maps in the posterior parietal cortex. *Annual Review Neuroscience*. 2002; 25:189–220.
4. Brunia CHM. Waiting in readiness: gating in attention and motor preparation. *Psychophysiology*. 1993; 30:327–339. [PubMed: 8327618]
5. Brunia CHM, Van Boxtel GJM. Wait and see. *International Journal of Psychophysiology*. 2001; 43:59–75. [PubMed: 11742685]
6. Culham JC, Kanwisher NG. Neuroimaging of cognitive functions in human parietal cortex. *Current Opinion in Neurobiology*. 2001; 11:157–63. [PubMed: 11301234]
7. Eimer M, Van Velzen J. Covert manual response preparation triggers attentional modulations of visual, but not auditory processing. *Clinical Neurophysiology*. 2006; 117:1063–1074. [PubMed: 16516545]
8. Eimer M, Forster B, Fieger A, Harbich S. Effects of hand posture on preparatory control processes and sensory modulations in tactile-spatial attention. *Clinical Neurophysiology*. 2004; 115:596–608. [PubMed: 15036056]
9. Eimer M, Forster B, Van Velzen J, Prabhu G. Covert manual response preparation triggers attentional shifts: ERP evidence for the premotor theory of attention. *Neuropsychologia*. 2005; 43:957–966. [PubMed: 15716166]
10. Eimer M, Van Velzen J, Driver J. Crossmodal interactions between audition, touch and vision in endogenous spatial attention: ERP evidence on preparatory states and sensory modulations. *Journal of Cognitive Neuroscience*. 2002; 14:254–271. [PubMed: 11970790]
11. Eimer M, Van Velzen J, Gherri E, Press C. Manual response preparation and saccade programming are linked to attention shifts: ERP evidence for covert attentional orienting and spatially specific modulations of visual processing. *Brain Research*. 2006; 1105:7–19. [PubMed: 16448629]
12. Fadiga L, Fogassi L, Gallese V, Rizzolatti G. Visuomotor neurons: ambiguity of the discharge or 'motor' perception? *International Journal of Psychophysiology*. 2000; 35:165–77. [PubMed: 10677645]
13. Filimon F, Nelson JD, Hagler DJ, Sereno MI. Human cortical representations for reaching: mirror neurons for execution, observation, and imagery. *Neuroimage*. 2007; 37:1315–28. [PubMed: 17689268]
14. Gherri E, Van Velzen J, Eimer M. Dissociating effector and movement direction selection during the preparation of manual reaching movements: evidence from lateralized ERP components. *Clinical Neurophysiology*. 2007; 118:2031–49. [PubMed: 17646131]
15. Gottlieb J. From thought to action: the parietal cortex as a bridge between perception, action and cognition. *Neuron*. 2007; 53:9–16. [PubMed: 17196526]
16. Harter MR, Miller SL, Price NJ, LaLonde ME, Keyes AL. Neural processes involved in directing attention. *Journal of Cognitive Neuroscience*. 1989; 1:223–237.
17. Hopf JM, Mangun GR. Shifting visual attention in space: an electrophysiological analysis using high spatial resolution mapping. *Clinical Neurophysiology*. 2000; 111:1241–1257. [PubMed: 10880800]
18. Jeannerod, M. *The cognitive neuroscience of action*. Blackwell; Oxford: 1997.
19. Johnson MT, Mason CR, Ebner TJ. Central processes for the multiparametric control of arm movements in primates. *Current Opinion in Neurobiology*. 2001; 11:684–8. [PubMed: 11741018]
20. Kalaska JF, Crammond DJ. Deciding not to GO: neuronal correlates of response selection in a GO/NOGO task in primate premotor and parietal cortex. *Cerebral Cortex*. 1995; 5:410–28. [PubMed: 8547788]
21. Keele SW. Movement control in skilled motor performance. *Psychological Bulletin*. 1968; 70:387–403.

22. Keele, SW. Behavioral analysis of movement. In: Brooks, VB., editor. *Handbook of Physiology*. American Physiological Society; Bethesda, MD: 1981. p. 1391-1414.
23. Kurata K. Premotor cortex of monkeys: set- and movement-related activity reflecting amplitude and direction of wrist movements. *Journal of Neurophysiology*. 1993; 69:187–200. [PubMed: 8433130]
24. Luppino G, Rizzolatti G. The organization of the frontal motor cortex. *News in Physiological Science*. 2000; 15:219–224.
25. Matelli M, Luppino G. Parietofrontal circuits for action and space perception in the macaque monkey. *Neuroimage*. 2001; 14:27–32.
26. Mathews S, Dean PJ, Sterr A. EEG dipole analysis of motor-priming foreperiod activity reveals separate sources for motor and spatial attention components. *Clinical Neurophysiology*. 2006; 117:2675–83. [PubMed: 17011820]
27. Nobre AC, Sebestyen GN, Miniussi C. The dynamics of shifting visuospatial attention revealed by event-related potentials. *Neuropsychologia*. 2000; 38:964–974. [PubMed: 10775707]
28. Perrin F, Pernier J, Bertrand O, Echallier JF. Spherical Splines for Scalp Potential and Current Density Mapping. *Electroencephalography and Clinical Neurophysiology*. 1989; 72:184–187. [PubMed: 2464490]
29. Praamstra P. Prior information of stimulus location: effects on ERP measures of visual selection and response selection. *Brain Research*. 2006; 1072:153–160. [PubMed: 16406014]
30. Praamstra P, Boutsen L, Humphreys GW. Frontoparietal control of spatial attention and motor intention in human EEG. *Journal of Neurophysiology*. 2005; 94:764–774. [PubMed: 15744008]
31. Prut Y, Fetz EE. Primate spinal interneurons show pre-movement instructed delay activity. *Nature*. 1999; 401:590–4. [PubMed: 10524626]
32. Riehle A, Requin J. Monkey primary motor and premotor cortex: single-cell activity related to prior information about direction and extent of an intended movement. *Journal of Neurophysiology*. 1989; 61:534–49. [PubMed: 2709098]
33. Rizzolatti G, Luppino G, Matelli M. The organization of the cortical motor system: new concepts. *Electroencephalography and Clinical Neurophysiology*. 1998; 106:283–96. [PubMed: 9741757]
34. Rizzolatti, G.; Riggio, L.; Sheliga, BM. Space and selective attention. In: Umiltà, C.; Moscovitch, M., editors. *Attention and performance XV*. MIT Press; Cambridge, MA: 1994. p. 231-265.
35. Rosenbaum, DA. The movement precuing technique: Assumptions, applications, and extensions. In: Magill, RA., editor. *Memory and control of action*. North-Holland; Amsterdam: 1983. p. 231-274.
36. Sheliga BM, Craighero L, Riggio L, Rizzolatti G. Effects of spatial attention on directional manual and ocular responses. *Experimental Brain Research*. 1997; 112:339–51.
37. Sheliga BM, Riggio L, Rizzolatti G. Orienting of attention and eye movements. *Experimental Brain Research*. 1994; 98:507–22.
38. Van der Lubbe RH, Wauschkuhn B, Wascher E, Niehoff T, Kompf D, Verleger R. Lateralized EEG components with direction information for the preparation of saccades versus finger movements. *Experimental Brain Research*. 2000; 132:163–178.
39. Walter WG, Cooper R, Aldridge VJ, McCallum WC, Winter AL. Contingent negative variation: an electric sign of sensori-motor association and expectancy in the human brain. *Nature*. 1964; 203:380–384. [PubMed: 14197376]
40. Wauschkuhn B, Wascher E, Verleger R. Lateralised cortical activity due to preparation of saccades and finger movements: a comparative study. *Electroencephalography and Clinical Neurophysiology*. 1997; 102:114–124. [PubMed: 9060862]
41. Wise SP. The primate premotor cortex: Past, present, and preparatory. *Annual Review of Neuroscience*. 1985; 8:1–19.
42. Yamaguchi S, Tsuchiya H, Kobayashi S. Electroencephalographic activity associated with shifts of spatial attention. *Brain*. 1994; 117:553–562. [PubMed: 8032865]

Figure 1.

Schematic representation of the different movement types and instruction conditions. During inward movement (left side), start keys where hands were positioned during response preparation were located far apart, while target keys (i.e., end locations of movements) were positioned close to the body midline. During outward movements (right side), these positions were reversed as start keys and hands were now located close to the body midline, and movement targets were in the two outer positions. Top panels show the direction cue instruction condition where participants were informed that response cues specified the direction of a reaching movement towards the left or right side (represented by grey thick arrows). Bottom panels show the effector cue instruction condition where participants were told that response cues indicated whether the left or right hand had to be prepared for an upcoming movement (represented by start keys filled in grey). The location of the irrelevant visual probes that were presented either during response preparation or response execution is represented by the small circles above each start and target key.

ERPs during covert movement preparation

Figure 2.

Grand-averaged ERPs elicited in the 900 ms interval following response cue onset at lateral anterior, central and posterior electrode pairs ipsilateral (dashed line) and contralateral (solid line) to the side of the cued response. ERPs are collapsed across trials with inward and outward movements, and both response instruction groups. The measurement windows used to compute ADAN amplitudes (400–700 ms after cue onset) and LDAP amplitudes (400–700 and 700–900 ms post-cue) are marked with boxes. Right panels show scalp distributions of ADAN and LDAP components in the 400–700 ms and 700–900 ms intervals. Maps represent differences between brain activity observed over hemisphere ipsi- and contralateral to the cued response side. They were constructed by spherical splines interpolation (see Perrin, Pernier, Bertrand & Echallier, 1989) after mirroring the difference waveforms to obtain symmetrical but inverse amplitude values for both hemispheres. Amplitudes range between -0.6 to $0.6 \mu V$, contour lines represent changes of $0.1 \mu V$.

ERPs during covert movement preparation

Figure 3.

Grand-averaged ERPs elicited in the 900 ms interval after response cue onset at anterior F3/4 (top panels) and posterior P7/8 (bottom panels) electrode pairs. Panels A, B, C and D show ERPs ipsilateral (dashed line) and contralateral (solid line) to the side of the cued response, separately represented for inward and outward movement preparation. Panels E and F (right side) show difference waveforms generated by subtracting ERPs elicited contralateral from ERPs elicited ipsilateral to the side of the cued response, separately for inward (solid line) and outward movement preparation (dashed line). Enhanced negativities contralateral to the side of the cued response are reflected by positive values (downward deflections) and enhanced contralateral positivities are reflected by negative values (upward deflections). Box markers indicate the LDAP measurement windows.

ERPs during covert movement preparation

Figure 4.

Grand-averaged ERPs elicited in the 900 ms interval after response cue onset at posterior electrode pair P7/8, shown separately for the participants with direction cue instructions (top panels), and for participants with effector cue instructions (bottom panels). Panels A to D show ERPs ipsilateral (dashed line) and contralateral (solid line) to the side of the cued response, separately for inward and outward movement preparation. Panels E and G (right side) show difference waveforms generated by subtracting ERPs elicited contralateral from ERPs elicited ipsilateral to the side of the cued response, separately for inward (solid line) and outward movement preparation (dashed line). Box markers indicate the LDAP measurement windows. Panels F and H (right side) show LDAP amplitudes obtained at P7/8 by subtracting ERP mean amplitudes at electrode ipsilateral to the cued response side from mean amplitudes obtained at the corresponding contralateral electrode in the 400-700 ms and 700-900 ms time intervals, separately for inward movement (black bars) and outward movements (grey bars).

ERPs triggered by visual probe stimuli

Figure 5.

Grand-averaged visual ERPs elicited by visual probe stimuli delivered 900 ms after response cue onset, obtained on the basis of high-pass filtered raw EEG data, and shown separately for probes located next to the initial position of the hands on the start keys (left side), and for probes located next to the target keys (right side). Solid lines show ERPs triggered by visual probes presented on the side of a cued response, while dashed lines show ERPs elicited on the opposite uncued side. Visual ERPs are shown separately for the group of participants that received direction cue instructions (top panels), and for the group that received effector cue instructions (bottom panels).