Sexually antagonistic chromosomal cuckoos

William R. Rice^{1,*}, Sergey Gavrilets^{2,3}
and Urban Friberg¹

¹Department of Ecology, Evolution and Marine Biology, University of California, Santa Barbara, CA 93106, USA

²Department of Ecology and Evolutionary Biology, and ³Department of Mathematics, University of Tennessee, Knoxville, TN 37996, USA

*Author for correspondence (rice@lifesci.ucsb.edu).

The two kinds of sex chromosomes in the heterogametic parent are transmitted to offspring with different sexes, causing opposite-sex siblings to be completely unrelated for genes located on these chromosomes. Just as the nest-parasitic cuckoo chick is selected to harm its unrelated nest-mates in order to garner more shared resources, sibling competition causes the sex chromosomes to be selected to harm siblings that do not carry them. Here we quantify and contrast this selection on the X and Y, or Z and W, sex chromosomes. We also develop a hypothesis for how this selection can contribute to the decay of the non-recombining sex chromosome.

Keywords: sexual conflict; sibling competition; sex chromosomes; sexually antagonistic zygotic drive; degenerate Y chromosome; Hamilton's rule

1. INTRODUCTION

The asymmetrical transmission of a parent's X and Y, or Z and W, sex chromosomes to opposite-sex siblings, when coupled with sibling competition, selects for a meiotic-drive-like process called sexually antagonistic zygotic drive (hereafter SA-zygotic drive; Miller *et al.* 2006; Rice *et al.* 2008). Meiotic drive operates via the differential success of haploid gametes from the same parent (reviewed in Hurst *et al.* 1996; Burt & Trivers 2006). SA-zygotic drive operates in an analogous way, but in the next generation, via differential success of diploid siblings. Just as meiotic drive elements are selected to harm the gametes that do not carry them, so too are the sex chromosomes selected to harm the competing siblings of the sex in which they are not represented. SA-zygotic drive can be mediated through (i) the heterogametic parent via sex-specific parental investment and/or epigenetic parental effects that harm only one sex of siblings and (ii) the competing siblings themselves, via sex-specific antagonistic interactions (Rice *et al.* 2008). Here we focus on interactions between siblings. Throughout, the term 'sex chromosomes' refers to their regions that do not recombine in the heterogametic sex. For simplicity, but without loss of generality, we will assume that males are the heterogametic sex.

Electronic supplementary material is available at <http://dx.doi.org/10.1098/rsbl.2009.0061> or via <http://rsbl.royalsocietypublishing.org>.

One contribution of 16 to a Special Feature on 'Sexual conflict and sex allocation: evolutionary principles and mechanisms'.

Many cuckoo species of birds are brood parasites that lay their eggs in the nests of other bird species (hosts) and these surrogate parents rear their young. Cuckoo's chicks have evolved to be highly harmful to their unrelated nest-mates, e.g. ejecting them from the nest, pecking them and out-competing them for resources via faster development and super-stimulating begging displays (Soler & Soler 2000). Paternal X and Y sex chromosomes, in opposite-sex siblings, have the same absence of relatedness as that between cuckoo and host chicks. Correspondingly, paternally inherited sex chromosomes are selected, as cuckoos, to harm brood-mates they are not transmitted through. Here we quantify the selective constraints on SA mutations occurring on sex chromosomes that code for phenotypes that harm sisters (Y-linked) or brothers (X-linked). We use our results to motivate a new hypothesis by which SA-zygotic drive can contribute to the evolution of degenerate Y and W sex chromosomes.

2. MATERIAL AND METHODS

(a) Model: application of Hamilton's rule

To quantify the constraints on the accumulation of Y-linked mutations that harm sisters, and X-linked mutations that harm brothers, we begin with Hamilton's rule (Hamilton 1964) for the evolution of altruism: $C/B < r$, where C is the fitness cost of an altruistic behaviour; B is the fitness benefit to the individual receiving the altruism; and r is the level of relatedness (proportion shared genes) between the two individuals. To convert Hamilton's rule to the case of a focal individual harming one sex of sibling, we multiply C and B (numerator and denominator) by -1 , so the 'altruism' is converted to harm and the 'cost' to the focal individual becomes a benefit. Let B_H be fitness benefit to the focal individual of harming another individual and C_H be the fitness cost to the individual receiving the harm. Hamilton's rule when applied to harmful interactions between individuals becomes: $B_H/C_H < r$, or when expressed as a cost-benefit ratio $C_H/B_H < 1/r$.

(b) Model: population genetic approach

We consider a randomly mating population in which each family has two siblings. With an even sex ratio, the probabilities that these are sisters, brothers or one of each sex are $1/4$, $1/4$ and $1/2$, respectively. We focus on a single diallelic locus with a mutant allele (which is rare initially) decreasing the fitness (viability) of the brother from 1 to $1 - C_H$, while simultaneously increasing the fitness of its carrier (whether male or female) from 1 to $1 + B_H$. We assume that in families with two brothers, both carrying mutant alleles, their effects on each other combine multiplicatively resulting in fitness $(1 + B_H)(1 - C_H)$.

(i) Autosomal locus

Allele a produces no antagonistic effects and allele A produces a brother-harming phenotype in both sexes. Let h be the frequency of genotype Aa , which we assume is small. The frequency of Aa by aa matings is then $2h$ and one can show that in the next generation (see the electronic supplementary material),

$$h' = h \frac{8 + 4B_H - 2C_H - B_H C_H}{8}.$$

(ii) X-linked locus

We assume male heterogamety, that allele x has no antagonistic effects and a rare allele X produces a brother-harming phenotype. In this case, there are only two female genotypes, Xx and xx , at non-trivial frequencies at the adult stage, which we define to be u and $1 - u$, respectively. There are also two male genotypes at the adult stage, X and x , at frequencies v and $1 - v$, respectively. Only three types of mating occur at non-trivial frequency: $Xx \times x$ at frequency u ; $xx \times X$ at frequency v ; and $xx \times x$ at frequency $1 - u - v$. Under these conditions, one can show that in the next generation (see the electronic supplementary material),

$$u' = (u + 2v) \frac{2 + B_H}{4},$$

$$v' = u \frac{(2 + B_H)(2 - C_H)}{8}.$$

3. RESULTS

Consider a new mutation in a large outbred population. When Y-linked, $r=0$ between brothers and sisters, so sister-harming mutations will be favoured by selection so long as they help brothers. For the X chromosome, the same logic applies to an imprinted brother-harming mutation that is expressed exclusively when inherited from the father (although it would be selected only one-third of the time). But in the more general case when the X-linked mutation is not imprinted, Hamilton's rule must be met when averaged across the sexes, i.e. the constraints are $C_H/B_H < 1/0$ when the X is paternally inherited and $C_H/B_H < 2$ when inherited from the mother. Since two-thirds of X chromosomes reside in females, the weighted average result is $(1/3) \times B_H + (2/3) \times B_H > 0 + (1/3) \times C_H$, which implies that $C_H/B_H < 3$. These X and Y constraints are unchanged with multiple paternity. For an autosomal gene, the constraint is $C_H/B_H < 2$ (see [Mock & Parker \(1997\)](#) for a more general treatment of the autosomes). In summary, selection favours any sister-harming mutation that is Y-linked so long as some benefit accrues to brothers (cost/benefit $< \infty$). Brother-harming X-linked mutations, lacking a special form of imprinting, are more strongly constrained than the Y, but less constrained than the autosomes ([figure 1](#)).

Using our population genetic model for an autosomal locus, one can show that the brother-harming allele *A* invades when rare if (see the electronic supplementary material)

$$C_H < \frac{4B_H}{2 + B_H}. \quad (3.1)$$

With X-linkage, a brother-harming *X* allele invades when rare if (see the electronic supplementary material)

$$C_H < \frac{\left(9 - \left[\frac{6 - B_H}{2 + B_H}\right]^2\right)}{4}. \quad (3.2)$$

These results converge on those based on Hamilton's rule when B_H is small: on the autosomes $C_H/B_H < 2$ and on the X $C_H/B_H < 3$ (left side of [figure 1](#)). With stronger selection, however, there is stronger constraint on both the X and autosomes (right side of [figure 1](#)).

Searching the literature, we found specific examples of neither Y-linked mutations that cause brothers to harm their sisters nor X-linked mutations causing both sexes to harm brothers. There are, however, reports in which siblings of the opposite sex are more aggressive to one another than same-sex siblings (e.g. [Dunn & Kendrick 1981](#); [Bortolotti 1986](#); [Drummond *et al.* 1991](#); [Legge 2000](#); [Moura 2003](#)). However, we also found several examples where aggression was more pronounced between siblings of the same sex (e.g. [Minnett *et al.* 1983](#); [Frank *et al.* 1991](#); [Snowdon & Pickhard 1999](#)).

4. DISCUSSION

Because X chromosomes in sisters are more closely related than those in brothers, and than the autosomes in both sexes, the homogametic sex is expected to be more cooperative ([Kawecki 1991](#))—a prediction

Figure 1. The constraints on the accumulation of new mutations that are Y-linked and harm sisters, X-linked and harm brothers or autosomal and harm one sex of sibling. Y-linked mutations accumulate over the entire parameter space, X and autosomal mutations only when they map below the curves. Black curves denote X-linkage, grey curves autosomal linkage; solid curves are based on Hamilton's rule and dashed curves are based on a population genetic model.

with at least some empirical support (see [Haig \(2000\)](#) for a review of this literature). Here we have shown that X chromosomes are also expected to more readily accumulate selfish mutations that harm brothers ([figure 1](#)). Although we were unable to find any studies that documented X-coded, brother-harming phenotypes, we found no studies that specifically screened for this phenotype.

In contrast to the X, Y chromosomes have no restrictions on accumulating sister-harming mutations (full parameter space in [figure 1](#)). From the perspective of X and autosomes, Y chromosomes are selected to accumulate ultra-selfish mutations that harm sisters that do not carry them, irrespective of the cost-benefit ratio (C_H/B_H ; [figure 1](#)). Despite this unconstrained selection, we were unable to document established cases of sister-harming Y chromosomes in the literature, but, as for the X, we found no genetic screens that directly assayed for this Y-coded phenotype. We next propose a model, as a hypothesis, in which antagonistic coevolution keeps Y-coded, sister-harming phenotypes rare, and thereby leads to the decay of the Y.

[Hamilton \(1967\)](#) proposed that Y chromosomes degenerated because the X and autosomes were selected to shut down Y-linked meiotic drive. [Charlesworth \(1978\)](#) dismissed this scenario as a general explanation for the degeneration of Y chromosomes because of the apparent paucity of genes that can evolve to cause meiotic drive in males, and because of the cost of shutting down a large region of the nascent Y chromosome (due to hemizygous gene expression without dosage compensation). However, Hamilton's logic, when applied to SA-zygotic drive, may be a more potent factor leading to the degeneration of Y chromosomes. In mice, more than 80 per cent of the genome (structural genes) is expressed in the brain ([Sunkin & Hohmann 2007](#)). As a consequence, many genes can potentially contribute to behaviour, at least some of which could influence

Figure 2. Coevolution between selfish Y chromosomes that code for harmful sibling–sibling interactions that harm sisters, and the X and autosomes that are selected to silence them.

sister-harming behavioural phenotypes. Suppose that a sister-harming mutation accumulated on a nascent Y chromosome, leading to SA-zygotic drive. When it harmed sisters more than when it helped brothers, the entire genome would be selected to silence the mutation (figure 2). One mechanism by which genomes are known to shut down harmful genes, such as transposable elements, is via epigenetic silencing (Slotkin & Martienssen 2007). Recent evidence indicates that transacting, ncRNA-based silencing of only one allele at a diploid locus is feasible (Rinn *et al.* 2007). Recurrent silencing of sister-harming, Y-linked genes could, in principle, contribute to the degeneration of the Y sex chromosome (figure 2). The effect of such silencing would be magnified when high sequence similarity between alleles on the nascent X and Y required the ncRNA to target sequences flanking a sister-harming, Y-linked allele (that had diverged sufficiently to be uniquely targeted), leading to collateral silencing of any intervening genes. In this case many, and perhaps most, silenced genes on the Y would not code for sister-harming phenotypes.

Our study illustrates why the evolution of nascent sex chromosomes is ‘dangerous’ to the autosomes in all species with sibling competition: paternally inherited sex chromosomes are selected to harm the sex of siblings that do not carry them. This selection is more constrained on the X than the Y. Its ramifications may include elevated hostility among siblings, especially those of opposite sex, and antagonistic, intragenomic coevolution that, similar to Hill Robertson effects (Bachtrog & Charlesworth 2000; Nicolas *et al.* 2005), contributes to the decay of the Y.

Funding for this project came from the National Science Foundation (W.R.R. and S.G.), the National Institutes of Health (W.R.R. and S.G.) and the Wenner-Gren Foundations (U.F.).

- Bachtrog, D. & Charlesworth, B. 2000 Reduced levels of microsatellite variability on the neo-Y chromosome of *Drosophila miranda*. *Curr. Biol.* **10**, 1025–1031. (doi:10.1016/S0960-9822(00)00656-4)
- Bortolotti, G. R. 1986 Influence of sibling competition on nestling sex-ratios of sexually dimorphic birds. *Am. Nat.* **127**, 495–507. (doi:10.1086/284498)
- Burt, A. & Trivers, R. 2006 *Genes in conflict: the biology of selfish genetic elements*. Cambridge, MA: Belknap Press of Harvard University Press.
- Charlesworth, B. 1978 Model for evolution of Y chromosomes and dosage compensation. *Proc. Natl Acad. Sci. USA* **75**, 5618–5622. (doi:10.1073/pnas.75.11.5618)
- Drummond, H., Osorno, J. L., Torres, R., Chavelas, C. G. & Larios, H. M. 1991 Sexual size dimorphism and sibling competition: implications for avian sex-ratios. *Am. Nat.* **138**, 623–641. (doi:10.1086/285238)
- Dunn, J. & Kendrick, C. 1981 Social behavior of young siblings in the family context: differences between same-sex and different-sex dyads. *Child Dev.* **52**, 1265–1273. (doi:10.2307/1129515)
- Frank, L. G., Glickman, S. E. & Licht, P. 1991 Fatal sibling aggression, precocial development, and androgens in neonatal spotted hyenas. *Science* **252**, 702–704. (doi:10.1126/science.2024122)
- Haig, D. 2000 Genomic imprinting, sex-biased dispersal, and social behavior. *Ann. NY Acad. Sci.* **907**, 149–163.
- Hamilton, W. D. 1964 The genetical evolution of social behaviour. I. *J. Theor. Biol.* **7**, 1–16. (doi:10.1016/0022-5193(64)90038-4)
- Hamilton, W. D. 1967 Extraordinary sex ratios. *Science* **156**, 477–488. (doi:10.1126/science.156.3774.477)
- Hurst, L. D., Atlan, A. & Bengtsson, B. O. 1996 Genetic conflicts. *Q. Rev. Biol.* **71**, 317–364. (doi:10.1086/419442)
- Kaweck, T. J. 1991 Sex-linked altruism: a stepping-stone in the evolution of social behavior. *J. Evol. Biol.* **4**, 487–500. (doi:10.1046/j.1420-9101.1991.4030487.x)
- Legge, S. 2000 Siblicide in the cooperatively breeding laughing kookaburra (*Dacelo novaeguineae*). *Behav. Ecol. Sociobiol.* **48**, 293–302. (doi:10.1007/s002650000229)
- Miller, P. M., Gavrillets, S. & Rice, W. R. 2006 Sexual conflict via maternal-effect genes in ZW species. *Science* **312**, 73. (doi:10.1126/science.1123727)
- Minnett, A. M., Vandell, D. L. & Santrock, J. W. 1983 The effects of sibling status on sibling interaction: influence of birth order, age spacing, sex of child, and sex of sibling. *Child Dev.* **54**, 1064–1072. (doi:10.2307/1129910)
- Mock, D. W. & Parker, G. A. 1997 *The evolution of sibling rivalry*. Oxford Series in Ecology and Evolution. Oxford, UK; New York, NY: Oxford University Press.
- Moura, A. C. D. 2003 Sibling age and intragroup aggression in captive *Saguinus midas midas*. *Int. J. Primatol.* **24**, 639–652. (doi:10.1023/A:1023748616043)
- Nicolas, M. *et al.* 2005 A gradual process of recombination restriction in the evolutionary history of the sex chromosomes in dioecious plants. *PLoS Biol.* **3**, 47–56. (doi:10.1371/journal.pbio.0030004)
- Rice, W. R., Gavrillets, S. & Friberg, U. 2008 Sexually antagonistic ‘zygotic drive’ of the sex chromosomes. *PLoS Genet.* **4**, e1000313. (doi:10.1371/journal.pgen.1000313)
- Rinn, J. L., Wang, J. K., Allen, N., Mikels, A. J., Nusse, R., Helms, J. A. & Chang, H. Y. 2007 Site-specific induction of epidermal fates by dermal HOX transcriptional program. *J. Invest. Dermatol.* **127**, S109.
- Slotkin, R. K. & Martienssen, R. 2007 Transposable elements and the epigenetic regulation of the genome. *Nat. Rev. Genet.* **8**, 272–285. (doi:10.1038/nrg2072)
- Snowdon, C. T. & Pickhard, J. J. 1999 Family feuds: severe aggression among cooperatively breeding cotton-top tamarins. *Int. J. Primatol.* **20**, 651–663. (doi:10.1023/A:1020796517550)
- Soler, J. J. & Soler, M. 2000 Brood-parasite interactions between great spotted cuckoos and magpies: a model system for studying coevolutionary relationships. *Oecologia* **125**, 309–320. (doi:10.1007/s004420000487)
- Sunkin, S. M. & Hohmann, J. G. 2007 Insights from spatially mapped gene expression in the mouse brain. *Hum. Mol. Genet.* **16**, R209–R219. (doi:10.1093/hmg/ddm183)