Published in final edited form as:

Org Lett. 2009 November 5; 11(21): 4774–4776. doi:10.1021/ol901963v.

A Synthesis of the Carbon Skeleton of Maoecrystal V

Paul J. Krawczuk, Niklas Schöne, and Phil S. Baran*

Department of Chemistry, The Scripps Research Institute, 10550 North Torrey Pines Rd., La Jolla, CA 92037

Abstract

An enantioselective synthesis of the maoecrystal V (1) carbon skeleton is described. The key transformations include arylation of a 1,3-dicarbonyl compound with a triarylbismuth(V)dichloride species, oxidative dearomatization of a phenol, and a subsequent intramolecular Diels-Alder reaction.

Maoecrystal V (**1**, Figure 1) was isolated and characterized in 2004 by Sun and coworkers from the Chinese medicinal herb *Isodon Eriocalyx*.¹ It possesses an unusual architecture rendering it the most modified naturally occurring *ent*-kauranoid isolated to date.² The structure was confirmed by X-ray crystallographic analysis to contain five highly congested rings and seven stereocenters, two of which are adjacent all carbon quaternary centers located in its interior. One of these all carbon quaternary centers is also the bridgehead carbon of the [2,2,2]-bicyclic moiety. This letter reports an early approach to **1** pursued in our laboratory from November 2007 to May 2008.³

Our initial retrosynthesis, shown in Scheme 1A, began by disconnecting the tetrahydrofuran ring and simplifying the enone of the A ring to an olefin leading to **2**. Next, an intramolecular Diels-Alder⁴ disconnection of the bicyclic ring system would generate a cyclohexadiene/acrylate conjugate **3**. It was initially envisioned that silyl enol ether **3** could be used as the diene component in the Diels-Alder reaction. In order to construct this substrate via the corresponding enone **4**, an oxidative coupling of enolates or related substrates was proposed (i.e. **5** + **6**).⁵

Initial studies (Scheme 1B) with a model system using β -ketoaldehyde **7** and enol ether **8** led to the generation of the C-10 quaternary center by treatment with cerium ammonium nitrate.⁶ Following reduction of the formyl group and esterification, silylation of the ketone under a variety of conditions led only to the undesired regioisomeric dienol ether **9** (presumably due to the sterically hindered α -proton at C-9). To circumvent this problem, the diene was further dehydrogenated (Pd/C, O₂) and the resulting aromatic moiety was oxidized using a Wessely

pbaran@scripps.edu.

Supporting Information Available: Detailed experimental procedures, copies of all spectral data, and full characterization. This material is available free of charge via the Internet at <http://pubs.acs.org>.

oxidation⁷ (Pb(OAc)₄) to furnish cyclohexadienone **10** in 59% yield. A Diels–Alder reaction delivered polycycle **11** in *ca.* 60% yield (3 mg scale) upon heating to 165°C.⁸ X-ray crystallography confirmed the structure as that corresponding to the undesired regioisomeric outcome (based on the position of the geminal dimethyl group relative to the bicyclic portion of the molecule). Encouraged by the outcome of the intramolecular Diels–Alder, a revised route was devised to access the correct carbon framework found in **1** (Scheme 1C).

The sequence begins with β -ketoaldehyde **12**, derived in three steps from a readily available symmetrical 1,3-diketone.⁹ Barton arylation¹⁰ of **12** with **13**⁹ led to **14** in 67% yield as a 7:3 readily separable mixture of diastereomers. In a similar fashion to the model study, reduction (to afford **15**) and acylation led to **16**. Removal of the MOM group with TFA and Wessely oxidation of the intermediate hemiketal **17** yielded an inconsequential mixture of diastereomeric acetates **18/19** in 53% yield over two steps.

Dienes **18** and **19** were heated in a microwave reactor in *o*-DCB to 165°C for 1.5 hours in the presence of BHT, to give the expected *endo*-cycloadducts in 79% and 69% yields. The stereochemistry of the cycloadducts was verified by X-ray crystallographic analysis.¹¹ Once the structures had been determined, both adducts were elaborated further towards **1** by hydrogenation of the bicyclic olefin (H₂, Pd/C). Next, the acetate group was excised using SmI₂ to give an intermediate samarium enolate that was protonated by methanol to give the α -methyl ketone as a mixture of diastereomers in 76% yield (*dr* = 17:3 with **25** as major).¹² An X-ray crystal structure of **26** (C-16-*epi*-**25**) was obtained from the minor diastereomer as shown in Scheme 1C, establishing that the major isomer possesses the proper orientation of the C-16 methyl group in **1**.

Certain elements of the sequential Barton arylation/Wessely oxidation/Diels–Alder Strategy (i.e. **12**→**14**, **17**→**18/19**, **18/19**→**25**) reported herein may well find use in an eventual synthesis of **1**.¹³

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Dr. D.-H. Huang (TSRI) and Dr. L. Pasternack (TSRI) for NMR spectroscopic assistance, and Dr. G. Siuzdak (TSRI) for assistance with mass spectroscopy. Dr. Arnold Rheingold (UCSD) and Dr. Antonio DiPasquale (UCSD) are acknowledged for X-ray crystallographic assistance. Financial support for this work was provided by Bristol-Myers Squibb, the NIH (CA134785) and the German Research Foundation (DFG) (postdoctoral fellowship to N.S.)

References

1. Li S-H, Wang J, Niu X-M, Shen Y-H, Zhang H-J, Sun H-D, Li M-L, Tian Q-E, Lu Y, Cao P, Zheng Q-T. *Org. Lett* 2004;6:4327–4330. [PubMed: 15524475]
2. Sun H-D, Huang S-X, Han Q-B. *Nat. Prod. Rep* 2006;23:673–698. [PubMed: 17003905]
3. During this period one of the corresponding authors of the preceding letter was employed as a postdoctoral associate (C.-C. Li) in this laboratory: Gong J, Lin G, Li C-C, Yang Z. *Org. Lett*. 2009, DOI: 10.1021/ol9014392. This work was reported in an NIH proposal (received 8/11/2008 and reviewed on 10/9/2008 by the BCMB-B study section) and a final report to the DFG (submitted: 7/17/2009).
4. (a) Nicolaou KC, Snyder SA, Montagnon T, Vassilikogiannakis G. *Angew. Chem., Int. Ed* 2002;41:1668–1698. (b) Roush, WR. *Comprehensive Organic Synthesis*. Trost, BM., editor. Vol. 5. Oxford: Pergamon; 1991. p. 513–550.
5. (a) Baran PS, DeMartino MP. *Angew. Chem., Int. Ed* 2006;45:7083–7086. (b) DeMartino MP, Chen K, Baran PS. *J. Am. Chem. Soc* 2008;130:11546–11560. [PubMed: 18680297]

6. Nair V, Mathew J. J. Chem. Soc., Perkin Trans 1995;1:187–188.
7. (a) Wessely F, Sinwel F. Monatsh. Chem 1950;81:1055–1070. (b) Barnes-Seeman D, Corey EJ. Org. Lett 1999;1:1503–1504. [PubMed: 10825998] (c) Nicolaou KC, Simonsen KB, Vassilikogiannakis G, Baran PS, Vidali VP, Pitsinos EN, Couladouros EA. Angew. Chem., Int. Ed 1999;38:3555–3559. (d) Magdziak D, Meek SJ, Pettus TRR. Chem. Rev 2004;104:1383–1429. [PubMed: 15008626]
8. For recent examples using *ortho*-quinols as dienophiles see: (a) Nicolaou KC, Toh Q-Y, Chen DY-K. J. Am. Chem. Soc 2008;130:11292–11293. [PubMed: 18665593] (b) Gao S-Y, Chittimalla SK, Chuang GJ, Liao C-C. J. Org. Chem 2009;74:1632–1639. [PubMed: 19199660] (c) Dory YL, Roy A-L, Soucy P, Deslongchamps P. Org. Lett 2009;11:1197–1200. [PubMed: 19228005] (d) Snyder SA, Kontes F. J. Am. Chem. Soc 2009;131:1745–1752. [PubMed: 19191699].
9. See Supporting Information for details.
10. (a) Barton DHR, Lester DJ, Motherwell WB, Papoula MTB. J. Chem. Soc., Chem. Comm 1980:246–247. (b) Barton DHR, Blazejewski J-C, Charpiot B, Finet J-P, Motherwell WB, Papoula MTB, Stanforth SP. J. Chem. Soc., Perkin Trans 1985;1:2667–2675. (c) Barton DHR, Finet J-P, Giannotti C, Halley F. J. Chem. Soc., Perkin Trans 1987;1:241–249. (d) Combes S, Finet J-P. Tetrahedron 1999;55:3377–3386. (e) Fedorov AY, Finet J-P. J. Chem. Soc., Perkin Trans 2000;1:3775–3778. (f) Finet J-P, Fedorov AY. J. Organomet. Chem 2006;691:2386–2393.
11. For adduct **21** a crystal was grown directly, whereas adduct **20** was crystallized after removal of the TBS group as alcohol **22**.
12. Molander GA, Hahn G. J. Org. Chem 1986;51:1135–1138.
13. While this paper was in review, two other groups reported progress towards **1**: (a) Peng F, Yu M, Danishefsky SJ. Tetrahedron Lett. 2009 (b) Nicolaou KC, Dong L, Deng L, Talbot AC, Chen DY-K. Chem. Commun. 2009accepted.

Figure 1.
Representations of maoecrystal V.

A: Retrosynthetic Analysis **B: Model Study** **C: Synthesis of the Carbon Skeleton** **Scheme 1.**

Maoecrystal V: (A) Retrosynthetic Analysis, (B) Model Study, and (C) Synthesis of the Complete Carbon Skeleton

^aDetails: reagents and conditions (a) CAN, NaHCO₃, MeOH, 0°C, 46%, (b) Li(*t*-BuO)₃AlH, THF, -78°C, 53%, (c) acryloyl chloride, Et₃N, DMAP, DCM, 0°C, 36%, (d) TBSOTf, Et₃N, DCM, 63%, (e) Pd/C, O₂, PhMe, 110°C, 99%, (f) Pb(OAc)₄, AcOH, 59%, (g) 165°C, *o*-DCB, *ca.* 60%, (h) Ar₃BiCl₂(**13**), DBU, PhMe, 67%, (i) Li(*t*-BuO)₃AlH, THF, -78°C, 72%, (j) acryloyl chloride, DIEPA, DMAP, DCM, -78°C, 69%, (k) TFA, DCM, 0°C, 65%, (l) Pb(OAc)₄, AcOH, 81%, dr = 3:7, (m) 165°C, *o*-DCB, BHT 79% for **20**, 69% for **21** (n) BF₃·OEt₂, DCM, 82%, (o) H₂, Pd/C, EtOAc, 97% for **23**, 99% for **24** (p) SmI₂, MeOH, THF, 0°C, 76%, dr = 17:3. CAN = cerium ammonium nitrate, DMAP = 4-dimethylaminopyridine, DCM = dichloromethane, TBSOTf = *tert*-butyldimethylsilyl trifluoromethylsulfonate, *o*-DCB = *ortho*-dichlorobenzene, DIEPA = diisopropylethylamine, DBU = 1,8-diazobicyclo[5.4.0]undec-7-ene, TFA = trifluoroacetic acid, BHT = 2,6-*di-tert*-butyl-*p*-cresol.