Published in final edited form as:

Biochim Biophys Acta. 2009 ; 1789(11-12): 709–718. doi:10.1016/j.bbagr.2009.09.005.

Interaction of intestinal and pancreatic transcription factors in the regulation of CFTR gene expression

Victoria A McCarthy¹, Christopher J Ott², Marios Phylactides^{1,†}, and Ann Harris^{1,2}

¹Paediatric Molecular Genetics, Weatherall Institute of Molecular Medicine, University of Oxford, John Radcliffe Hospital, Oxford, UK

²Children's Memorial Research Center, Northwestern University Feinberg School of Medicine, Chicago, IL. USA

Summary

The tissue-specific regulation of the cystic fibrosis transmembrane conductance regulator gene (*CFTR*) is coordinated by intronic and extragenic cis-acting elements that influence its transcriptional activity. The promoter apparently lacks sequences to drive cell-type specific expression. We previously identified a number of intronic elements that were associated with DNase I hypersensitive sites (DHS) and bound the hepatocyte nuclear factor 1 (HNF1) transcription factor. Moreover, we demonstrated the likely involvement of HNF1 in the regulation of *CFTR* expression *in vivo*. Here we investigate DHS in introns 16 and 17a of the *CFTR* gene, which are evident in intestinal and pancreatic cell lines, and determine the transcription factors that interact with these sites. Of particular interest were factors known to interact with HNF1 in co-ordinated expression of genes in the gastrointestinal tract. We demonstrate that though sequences within these DHS bind HNF1, CDX2 and PBX1 *in vitro*, only PBX1 show a robust *in vivo* interaction. These data contribute to our understanding of the complexity of cell-type-specific *CFTR* regulatory mechanisms.

Introduction

The molecular mechanisms that control expression of the *CFTR* gene are complex and poorly defined though the gene shows tight tissue specific and temporal regulation. No genetic elements conferring tissue-specificity on *CFTR* expression have been found in the promoter region. However, we previously identified DNase I hypersensitive sites (DHS) across the whole locus that may be associated with regulatory elements (reviewed in [1]). Several of these DHS have been evaluated further and we identified hepatocyte nuclear factor 1 alpha (HNF1 α) as an important transcription factor involved in *CFTR* expression [2]. We predicted that though playing an important role, HNF1 α was probably part of a complex of transcription factors and co-factors that regulated *CFTR* expression and that the precise components of the complex would be cell type-dependent.

© 2009 Elsevier B.V. All rights reserved.

Correspondence to: Ann Harris.

²Current address: Human Molecular Genetics Program, Children's Memorial Research Center, Northwestern University, 2300 Children's Plaza, Box 211, Chicago, IL 60614-3394 US, Phone: 773 755 6582; Fax: 773-755-6593, ann-harris@northwestern.edu

[†]Current address: Cyprus Institute of Neurology and Genetics, Nicosia, Cyprus.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

One of the DHS regions that was shown to bind HNF1 α in vitro was in intron 17a and corresponded to a potential regulatory element that was evident in some CFTR - expressing intestinal (Caco2), pancreatic duct (Capan 1 and NP31) and airway (Calu3) cell lines but not in lymphoblastoid cell lines that lack CFTR expression [2], [4]. The presence of this DHS in several pancreatic duct cell lines in addition to the Caco2 line led us to re-evaluate the region to identify other intestinal/pancreatic duct enriched transcription factors that might be functioning in complex with HNF1 α . Moreover, our recent data demonstrating the importance of HNF1 in the mechanism of action of cell-type-specific enhancer in the first intron of CFTR further confirmed the importance of this transcription factor in tissue-specific regulation of the gene [3].

First, the accuracy of the original map location of the intron 17a DHS was reevaluated, together with that of a neighbouring DHS in intron 16 that was evident in the same cell types [4]. Since the initial data were generated on a large genomic fragment there was potential for the precise DHS to be somewhat inaccurate. Comparative genomics was utilized to align regions of cross species homology between the intron 16 and intron 17a DHS regions in cow, pig, sheep and human CFTR. Next, based on the cross-species homologies DNase I footprinting was carried out over 1kb spanning each conserved region that was associated with a DHS. Regions that exhibited footprints were then evaluated for consensus binding sites for known transcription factors. We identified predicted binding sites for several factors that are known to be involved in the regulation of expression of genes involved in intestinal function. These factors were shown to bind specifically to the CFTR intronic regions involved in the DHS. They include CDX1/2 and PBX1 that are known to interact with HNF1 α in co-ordinating tissue-specific gene expression [5]. CDX1 and CDX2 are both members of the caudal-related homeobox gene family that contributes to intestinal differentiation, proliferation and intestine-specific gene transcription [6]. They show sequence homology to the *Drosophila melanogaster* caudal gene that is involved in segmentation. CDX1 and CDX2 recognize the same DNA-binding motif and the ratio of the two factors is important in the differentiation of the intestinal epithelium. CDX1 and CDX2 may act as positive and/or negative regulators of the same gene. CDX2 is involved in regulation of expression of genes (like CFTR) that are implicated in the modulation of water/ion transport across the intestinal epithelium and it interacts directly with HNF1 α [5]. Many genes that are expressed in the intestine show regulation by CDX1 and CDX2 and mutations in these genes are associated with some tumors of intestinal origin. PBX1 is a member of the TALE (three-amino acid loop extension) class of homeodomain transcription factors, which are components of hetero-oligomeric complexes that regulate developmental gene expression and maintain differentiated states. PBX1 is essential for normal pancreatic development in humans and mice [7]. The likely involvement of CDX2 and PBX1 in CFTR regulation provides further support for our working model of tissue-specific control of the CFTR gene: multiple cis-acting elements interact with each other and the CFTR promoter in vivo, and the subtleties of cell type-specificity are achieved by individual elements that are associated with unique DHS within the gene.

Materials and methods

Cell culture

The 37566 lymphoblastoid and Capan 1 cell lines were cultured in RPMI 1640 supplemented with 10% fetal calf serum (FCS). The Caco2 [8] and Calu3 [9] cell lines were cultured in DMEM supplemented with 10% FCS. The T84 cell line [10] was cultured in a 1:1 mixture of DMEM and Ham's F12, 10% FCS.

Protein Resources

Caco2 and T84 nuclear extracts were prepared as described previously [2]. In vitro translated (IVT) proteins were generated by using the TNT-Kit (Promega) with relevant template plasmids: pCDX2 (kindly donated by D. Silberg) and pHNF1 (kindly donated by G. Crabtree) and pPBX1 (kindly donated by D. Zappavigna).

DNase I footprinting

To localize putative regulatory elements by in vitro DNA footprinting, primers were designed to clone overlapping DNA fragments of around 300 bp covering the area of each DHS (Supplementary data, Table 1A – C). Fragments were amplified using *Pfu* DNA polymerase (Stratagene) and cloned into the *SrfI* site of pPCR-script Amp SK with the PCR-Script™ Amp cloning kit (Stratagene). Inserts were sequence verified. Footprint probes were excised from the vector, with *ClaI* and *SacI* or with *NotI* and *EcoRI* to label the sense or antisense strand, respectively and gel purified. DNase I footprinting reactions were carried out as described previously [2].

Electrophoretic mobility shift assays (EMSA)

EMSA probes were designed on the basis of the DNase I foot-printing results and are summarized in Supplementary data, Tables 2A–C. Complementary oligonucleotides with or without a short 5' overhang, as described, were annealed to generate double stranded probes. EMSA reactions and supershifts were performed as described previously [2].

Western blots

Western blots were performed as described previously [2]. Antibodies were anti HNF1 α / β (Santa Cruz, sc-8986), PBX1(sc-889) and CDX2 (BioGenex-88). Enhanced chemiluminescence reagents (ECL) and film (Amersham Pharmacia Biotech Ltd., UK) were used for the final detection of antibodies.

Chromatin immunoprecipitation (ChIP)

ChIP was performed as described previously [3], [11]. Chromatin was sonicated to an average size of 500 bp. Immunoprecipitations were performed with antibodies specific for HNF1 (Santa Cruz, sc-8986), PBX1 (sc-889) and CDX2 (BioGenex-88) or IgG (Santa Cruz, sc-2027). Enrichment was analyzed using specific primers, PowerSYBR Green (Applied Biosystems), and a 7500 Real-time PCR machine (Applied Biosystems). Primers for ChIP experiments are shown in Supplementary data, Table 3.

Results

We previously identified DNase I hypersensitive sites across the *CFTR* locus in a number of cell types including Caco2 colon carcinoma cells [4]. These included sites in intron 16 (at 3120 + 3 kb, where 3120 is the last base in exon 16) and 17a (at 3271 + 0.7 kb, where 3271 is the last base in exon 17a) that were also evident in the pancreatic adenocarcinoma cell line Capan 1 though not in non-epithelial cells. Subsequent comparative genomics analyses revealed an additional region of cross-species homology in intron 16 (intron 16hom, located at 3120 + 1.1 kb) that was much more highly conserved than the 3120 + 3 kb DHS region. We therefore evaluated the DNA-protein interactions of all three regions in order to identify transcription factors (TF) that might coordinate the function of these cis-acting elements.

The core regions of the DHS in introns 16 and 17a contain multiple predicted binding sites for CDX2, PBX1 and HNF1 α

Overlapping 250–300 bp fragments spanning each DHS (Figures 1A and B) were evaluated by *in vitro* DNase I footprinting using nuclear extracts from Caco2 colon carcinoma cells that express abundant CFTR, Capan 1 pancreatic adenocarcinoma cells that express low levels of CFTR and MCF7 breast carcinoma cells or 37566 lymphoblastoid cells that do not express CFTR. Protected regions, indicating protein-DNA interactions, were found within each of the DHS. The locations of these DNase I footprints are shown in Figure 2 and Figure 3 and the predicted transcription factor binding sites that coincide with them, derived from the Transfac 4.0 database [12] are shown in Supplementary data, Table 4. We next specifically pursued predicted binding sites for the intestinal transcription factors HNF1 and CDX2, also PBX1, which is known to interact with HNF1.

Figure 2A shows a CDX2 binding site (aaataatTTTAgtatgaag) predicted in PR2 that was detected in the intron 16 DHS probes 3 (shown) and 4 using Caco2 nuclear extracts. Though no other protected regions in the 3120 + 3kb DHS element corresponded to predicted TF binding sites of interest, multiple relevant footprints were seen in the region of homology in intron 16 at 3120 + 1.1 kb (Figure 3A–E). PR1 is predicted to encompass a binding site for the PBX1 homeodomain transcription factor (acacCTATcactc) on the forward strand and a binding site for a PBX1/Meis heterodimer (tcccTGATgatagatg) on the reverse strand (Figure 3A). PR2 also corresponds to a predicted site for PBX1 on the reverse strand (ccacCAATcagtc) (Figure 3B) as does PR6 in probe 16homC on the forward strand (tggtCAATcaaat) (Figure 3C). Eight footprints were identified in probe 16 hom D (Figure 3D): PR7 corresponded to potential binding sites for PBX1 (gtgtCAATcattt) on the negative strand and for a PBX1/Meis heterodimer (gaaaTGATgacacaaa) on the positive strand. Probe 16hom E showed two footprints with nuclear extracts from Caco2 cells (Figure 3E): PR14 corresponds to predicted binding sites for HNF1 (tGATAattattgaggc) on the forward strand and PBX1 (ttcaCAATcaagt) on the negative strand; PR15 encompasses binding sites for HNF1 on both strands ((gGTTAatttttaaatag) and (tGTTAatttataaaatg)).

The DHS 17a (3271 + 0.7 kb) DHS showed multiple relevant footprinted regions (Figure 2B, 2C). Data on two predicted HNF1 sites in probes 17a-C and 17a-D (17a FP2 and FP3) were previously published in the context of our analysis of the role of HNF1 in regulation of *CFTR* gene expression [2] and so will not be discussed further here. Probe 17a-D shows an additional relevant footprint with Caco2 nuclear extracts (Figure 2B): PR3 encompasses two predicted HNF1-binding sites (tGTTAatttattaat and gGTAAactcataata on the negative strand and positive strands respectively). Probe 17a-E shows an additional 4 footprints in this DHS region, using nuclear extracts from Caco2 cells: PR6 corresponds to a predicted CDX2 site on the forward strand (ttcatttaTAATAaac) which was also seen on the overlapping portion of probe 17a-A (Figure 1B, data not shown); PR7 and PR9 also encompass predicted CDX2 binding sites on the reverse strand (tacctatTTTAAaaccaaa) and forward strand respectively (ttaaactTTTAcacatat); PR8 corresponds to an HNF1 site (gGTCAattaataagaaa) on the reverse strand.

CDX2, PBX1 and HNF1 α bind to the DHS in introns 16 and 17a *in vitro*

The predicted binding sites for CDX2, PBX1 and HNF1 that were identified in the DNase I footprinting experiments were evaluated further.

In vitro evaluation of CDX2 binding—PR2 in probe 16-4 was predicted to be a binding site for CDX2. An oligonucleotide probe, 16FP2CDX2 (Supplementary data, Table 2A) encompassing PR2 was used in electrophoretic mobility shift assays (EMSA) (Figure 4A and B). In Figure 4A, nuclear extracts from Caco2 cells that express CDX2 (Supplementary Figure

1) generated several complexes with 16FP2CDX2. Complex A was thought to involve CDX2 as it was competed more efficiently by a CDX2 consensus oligonucleotide (lane 3) than by self (lane 4). Further, another oligonucleotide 17aFP6CDX2 (see Supplementary table 2A) encompassing a predicted CDX2 binding site in intron 17a (17aPR6) competed complex A more efficiently (lane 9) than the same oligonucleotide containing a mutation in the CDX2 consensus site (17aFP6CDX2M, lane 10). A supershift reaction with an antibody to CDX2 (Fig. 4A, lane 11) abolished complex A, further supporting involvement of CDX2 at this site (the β tubulin isotype control antibody had no effect on complex A). Though PR2 in probe 16-4 does not contain a predicted binding site for HNF1, complex B may include HNF1, since it is partially competed by the HNF1 consensus oligonucleotide (Fig. 4A, lane 6) but not the CDX2 consensus (Fig. 4A, lane 3). Further, an oligonucleotide (16homFP15FAC1, supplementary table 2C) encompassing the two predicted HNF1 sites at PR15 in the 16 hom E probe also partially competed complex B (Fig. 4A, lane 7). However a supershift reaction with an antibody specific for HNF1 failed to alter the mobility of complex B (Fig. 4A, lane 13).

In vitro translated CDX2 also bound to the 16FP2CDX2 oligonucleotide (Figure 4B, lane 3) generating complex A at an equivalent mobility to that generated with Caco2 nuclear extracts. The complex was competed by self (Fig 4B, lane 2), the CDX2 consensus (lane 3) and partially by the 17aFP6CDX2 oligonucleotide (lane 9), but not mutant self (16FP2CDX2M, lane 5), mutant 17aFP6CDX2 (17aFP6CDX2M, lane 10), HNF1 (lane 6) or 16homFP15FAC1 (lane 7) encompassing the two HNF1 sites. An antibody to CDX2 generated a definitive supershift to complex C (lane 11), which was not seen with the isotype control antibody. These data confirm that CDX2 is one of the transcription factors that can bind to the intron 16 (3120 + 3kb) DHS in vitro.

The CDX2 transcription factor is also predicted to bind at three footprints in the DHS17a region (3271 + 0.7kb): PR6, PR7 and PR9 probe 17a-E (Figure 2C). The PR6 region was evaluated further while PR7 and PR9 were not as both were very weak footprints and the latter is also very close to the end of the region of interest. Probe 17aFP6CDX2 was generated to evaluate the potential interaction of CDX2 with this site in vitro. Figure 5 shows that the most abundant protein complexes (B) that bind to this probe in nuclear extracts from Caco2 cells are not CDX2 alone, consistent with the data presented above on 16FP2CDX2. However, a minor complex shown by arrow A likely corresponds to CDX2 since it is competed by the CDX2 consensus (Figure 5, lanes 3) but not by the consensus with a mutation in the predicted CDX2 binding site (CDX2M, lane 4). Furthermore addition of an antibody specific for CDX2 to the reaction abolished complex A (lane 12). In vitro translated CDX2 also generated a complex with mobility of complex A (data not shown). Further analyses of the major complexes (B) shows that the major component again appears to be HNF1, despite the lack of a predicted HNF1 binding site in 17aFP6CDX2: the B complexes are competed efficiently by the HNF1 consensus (lane 9) and by another predicted HNF1 binding site in the 16hom region (16homHNF1, lane 7) but not by the same sequence with a mutation in the HNF1 binding site (16homHNF1M, lane 8). Conclusively, an antibody specific for HNF1 supershifted part of the complex to C (lane 14).

In vitro evaluation of PBX1 binding—Several footprints in the 16 hom region were predicted to bind PBX1 or a PBX1/Meis heterodimer. These include PR1, PR2, PR6, PR7 and PR14. The site in PR1, which was a strong footprint that did not coincide with predicted binding sites for any other transcription factors, was evaluated further by EMSA (Figure 6) using probe 16homFP1PBX1. Nuclear extracts from Caco2 cells generated a complex (A, Figure 6) that was competed by self (lanes 3, 4), and a consensus PBX1 binding site oligonucleotide (lanes 7,8) but not by self with a mutation in the consensus PBX1 site (16HomFP1 mut, lanes 5,6) or mutant PBX1 (PBX1 mut, lanes 9,10). An antibody specific for PBX1 did not supershift the

complex efficiently in this experiment, only demonstrating a very weak complex at B, though this complex was more evident in some other experiments. Further, a consensus PBX1 oligonucleotide generated a specific complex of similar mobility to that generated by 16homFP1PBX1 and this complex was effectively competed by the latter (data not shown).

In vitro evaluation of HNF1 binding—As we observed previously at other DHS in the *CFTR* gene there is an abundance of HNF1 binding sites among the footprinted regions. Several of these were investigated further. PR14 and 15 in probe 16hom E both corresponded to predicted HNF1 binding sites. PR15 contained 2 sites, one on each strand and was of mechanistic interest. Figure 7A shows a complex (A) formed with nuclear extracts from Caco2 cells by the 16homFP15FAC1 probe, that is competed by self and the 17aFP3FAC1 probe that we showed previously to bind HNF1 in vitro [2]. The reciprocal experiment using 17aFP3FAC1 as a probe and competing with 16homFP15FAC1 showed consistent results and in both cases the complex was supershifted (B) by an antibody specific for HNF1 (lanes 5, 10). The two predicted HNF1 sites in FP15 were next analysed further independently of each other (Figure 7B,C): the complex A, generated with the 16homFP15FAC1 probe, though effectively competed by self (Figure 7B, lane 3) and the HNF1 consensus (lane 5) was competed less efficiently by the 16homFP15FAC2 probe (lane 4). 16homFP15FAC1M, with a mutation in the HNF1 binding site did not compete the complex (lane 6), which was supershifted by an antibody reacting with both HNF1 α and β (lane 7). In contrast, though the 16homFP15FAC2 probe (Figure 7C) generated a complex of similar mobility to that seen with the FAC1 probe, which was partially competed by self (lane 3) but not with self containing a mutation in the predicted HNF1 binding site (16homFP15FAC2M, lane 6), this complex was competed ineffectively by 16homFP15FAC1 (lane 2) and the HNF1 consensus (lane 5) and none of the HNF1 antibodies generated a supershift (lanes 7–9). Since HNF1 functions as a dimer we decided to further investigate the potential interaction of these two HNF1 sites in a single oligonucleotide probe, 16hom FP15HNF1 (see Figure 8). In panel A, two complexes (A) were formed between this probe and Caco2 nuclear extracts which were competed by self (lane 3), HNF1 consensus (lane 8) and 16homFP15FAC1 (lane 4) but not 16homFP15FAC1M (lane 5). The latter reproducibly enhanced the stability of the slower migrating complex A1. However, 16homFP15FAC2 was only slightly more effective at competing the complexes at A (lane 6) than its mutant counterpart 16homFP15FAC2M (lane 8). The slower migrating complexes at A1 supershifted (B) with an antibody reacting with HNF1 α and β (lane 9). In Figure 8B, the complex generated with probe 16homFP15HNF1 was competed with mutant versions that disrupted the FAC1 HNF1 site (16homFP15HNF1T), the FAC2 HNF1 site (16homFP15HNF1B) or both sites (16homFP15HNF1M). Probe 16homFP15HNF1T (lane 4) was a less effective competitor than the 16homFP15HNF1 wildtype probe (lane 3), though 16homFP15HNF1B competed more effectively (lane 5), suggesting that the second HNF1 site was less crucial in complex formation. Of interest in this experiment was the observation that the HNF1 consensus (lane 7) competed the slower mobility complex at A1 more effectively than the faster migrating one (A2), and this was confirmed by the HNF1 consensus mutant (lane 6). Again the slower migrating complexes at A1 supershifted with an antibody reacting with HNF1 α and β (lane 8), while the faster migrating complex did not, suggesting it may involve another factor.

In addition to the HNF1 sites that we investigated previously [2] in the intron 17a DHS, two further sites were evaluated here (Supplementary Figure 2 and Supplementary Figure 3). The data in Supplementary Figure 2 demonstrate that in the 17aFP3FAC1 probe, which contains two HNF1 sites (on opposite strands) both sites bind HNF1, though the 5' site is apparently dominant (as it is a less effective competitor when mutated). Binding of HNF1 in vitro was also seen to probe 17AFP8HNF1 which encompasses a predicted HNF1 binding site in probe 17a-E PR8 (Supplementary Figure 3). Interestingly, the main complex generated with this probe was also effectively competed by the 17aFP6CDX2 probe that contains binds CDX2 at

a predicted consensus site (Figure 5) but not by a mutant version of this probe 17aFP6CDX2M, suggesting that though there is no predicted CDX2 binding site in the 17aFP8 probe, CDX2 is in complex with HNF1 at this site.

In vivo interaction of CDX2, HNF1 and PBX1 with potential cis-regulatory elements

Chromatin immunoprecipitation experiments were performed to investigate the binding of CDX2, HNF1 and PBX1 with the elements that were characterized using *in vitro* techniques. Since ChIP experiments can often yield misleading results in the absence of relevant controls for non-specific interactions we not only performed no antibody and rIgG controls for each experiment but each antibody was evaluated for interactions with well characterized binding site in other genes. The Valosin-containing peptide (VCP) promoter for PBX1 [13], the sucrose isomaltase (SI) promoter for CDX2 [5] and the intron 1 enhancer in CFTR for HNF1 [3]. Though the PBX1 antibody effectively enriched sequences from the VCP promoter (~ 30 fold in Capan 1 cells, Figure 9) and the HNF1 antibody enriched the CFTR intron 1 enhancer (~4.5 fold in Caco2 cells) only modest enrichment (~ 2 fold over rIgG) of the SI promoter was seen in Caco2 cells with the CDX2 antibody (Biogenex) (Supplementary Figure 4) and none in Capan 1 cells, despite the fact that they express CDX2 (see supplementary figure 1). Though this antibody was used in other ChIP experiments [14], its relative inefficiency in ChIP for the SI promoter in a cell line expressing abundant CDX2 prevents a conclusive result when used to evaluate the *in vivo* binding of CDX2 to the other elements. Another antibody used in different ChIP experiments (BL 3195 Bethyl labs [15]) was even less efficient in our hands. Though we have previously shown *in vivo* binding of HNF1 to the enhancer element at 185 + 10kb in the first intron of *CFTR* [3] and this element provides a positive control for the HNF1 ChIP experiments (Figure 9, DHS1), binding of this transcription factor to the 17aDHS region could not be demonstrated in Caco2 or Capan 1 cells. However, data shown in Figure 9 demonstrate that PBX1 binds *in vivo* in Capan 1 cells to 16Hom PBX1 site characterized *in vitro* in Figure 6 and possibly to other PBX1 binding sites predicted *in silico* in this region. Moreover, we determine enrichment of the DHS17a region with the anti-PBX1 antibody, though we have not identified a predicted binding site for this factor within this sequence. This may be due to the presence of PBX1 binding sites that were not revealed by DNase I footprinting, the existence of PBX1 binding sites flanking the DHS region, or the direct interaction of PBX1 with other proteins involved in the regulatory complex that binds at DHS17a.

Discussion

Recent progress in research aimed at elucidating the mechanisms of regulation of tissue specific expression of *CFTR* [3], [16], (Ott et al. in submission) has provided a working model of how the gene is controlled, however, many of the details remain to be elaborated. We have shown that multiple intronic, cell-type-specific enhancers that lie between 10 and ~ 200 kb from the promoter are in direct physical association with this element when the *CFTR* gene is active, but not in the inactive locus. The full range of cis-acting elements and many of the trans-acting factors interacting with these elements still need to be defined.

To further elucidate these elements and the transcription factors that are recruited to activate *CFTR* expression in intestinal and pancreatic cells, we re-examined DHS in introns 16 and 17a that we defined in earlier work [4]. The intron 17a element does not have enhancer activity as measured in luciferase reporter gene assays [17] though the DHS in intron 16 and the region of homology in intron 16 have not been evaluated for this function. DNase I footprinting of regions spanning the predicted core of each DHS revealed *in vitro* binding of some novel and some familiar transcription factors.

Of particular interest were binding sites for hepatocyte nuclear factor 1 and proteins that are known to directly interact with HNF1, since there is now ample evidence for the involvement of this transcription factor in intestinal regulation of *CFTR* expression [2], [3], [15]. In this context, the observation of multiple predicted binding sites for HNF4 in the footprinted regions of introns 16 and 17a may also be of relevance (Supplementary data, Table 4), though it was not pursued in the current experiments. However, HNF1 and HNF4 are known to interact directly and play an important role in the transcriptional network involved in the maintenance of the hepatocyte phenotype [18], [19]. Among other well-characterized interacting partners of HNF1 are chromatin-modifying enzymes such as histone acetyl transferases and transcriptional co-activators including CBP/p300 [20]. Moreover, for a number of genes that show cell-type specific regulation in the intestine, other intestinal transcription factors such as the caudal-related homeobox proteins CDX1 and CDX2 act co-operatively with HNF1 [5], [21], [22] suggesting that these proteins may interact directly. PBX1 is essential for the normal development of the pancreas and is involved in maintenance of differentiated function in a number of cell types in which *CFTR* is expressed. Hence it was logical to pursue the potential function of the in silico predicted binding sites for these proteins in the mechanism of *CFTR* regulation.

Our data demonstrate the in vitro binding of HNF1, CDX2 and PBX1 to intronic cis-acting regulatory elements of the *CFTR* gene that were initially identified through DHS mapping. Though our ChIP data provide evidence for the in vivo binding of PBX1 to elements in intron 16, we were not able to show binding of CDX2 and HNF1 to the DHS cores in introns 16 and 17. It is of interest that though we have not identified predicted binding sites for PBX1 in DHS17a, this region is also enriched during ChIP for PBX1, suggesting that the protein may be involved in the complex interacting at that DHS. This phenomenon has been described for other factors and a number of models exist to explain the recruitment of transcription factors to sites that lack a consensus binding motif [23].

It seems probable, since we have clearly demonstrated the in vivo interaction of HNF1 with multiple intronic regulatory elements in the *CFTR* gene [3], (Ott et al. in submission), that the lack of binding of HNF1 to DHS17a in the current experiments may be accounted for by technical limitations: firstly, the antibodies that are available for these intestinal transcription factors, though relatively specific, are not ChIP-grade reagents and therefore have significant potential to fail to demonstrate enrichment at individual TF binding sites; second it is now clear that a large complex of proteins are interacting in the *CFTR* active chromatin hub [3], [16], (Ott et al. in submission). In this context, the ChIP technique may not be sufficiently sensitive to detect individual interactions of transcription factors with unique binding sites. We envisage that in the active *CFTR* locus, once the chromatin domain has been established, likely by a few dominant cis-acting elements [16], (Ott et al. in submission), a dynamic series of interactions bring different, distal cis-acting regulatory elements into close proximity with the *CFTR* promoter, thus co-ordinating the subtleties of cell-type-specific control.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported in part by the Cystic Fibrosis Trust, UK, the Cystic Fibrosis Foundation USA, NIH R01 HL094585, Children's Memorial Research Center and a Medical Research Council Scholarship (V.A.M). We are grateful to Drs D. Silberg, V. Zappavigna, and G. Crabtree for cloned transcription factors.

References

1. McCarthy VA, Harris A. The CFTR gene and regulation of its expression. *Pediatr Pulmonol* 2005;40:1–8. [PubMed: 15806593]
2. Mouchel N, Henstra SA, McCarthy VA, Williams SH, Phylactides M, Harris A. HNF1a is involved in regulation of expression of the CFTR gene. *Biochemical Journal* 2004;378:909–918. [PubMed: 14656222]
3. Ott CJ, Suszko M, Blackledge NP, Wright JE, Crawford GE, Harris A. A complex intronic enhancer regulates expression of the CFTR gene by direct interaction with the promoter. *J Cell Mol Med* 2009;13:680–692. [PubMed: 19449463]
4. Smith DJ, Nuthall HN, Majetti ME, Harris A. Multiple potential intragenic regulatory elements in the CFTR gene. *Genomics* 2000;64:90–96. [PubMed: 10708521]
5. Boudreau F, Rings EH, van Wering HM, Kim RK, Swain GP, Krasinski SD, Moffett J, Grand RJ, Suh ER, Traber PG. Hepatocyte nuclear factor-1 alpha, GATA-4, and caudal related homeodomain protein Cdx2 interact functionally to modulate intestinal gene transcription. Implication for the developmental regulation of the sucrase-isomaltase gene. *J Biol Chem* 2002;277:31909–31917. [PubMed: 12060663]
6. Silberg DG, Swain GP, Suh ER, Traber PG. Cdx1 and cdx2 expression during intestinal development. *Gastroenterology* 2000;119:961–971. [PubMed: 11040183]
7. Kim SK, MacDonald RJ. Signaling and transcriptional control of pancreatic organogenesis. *Curr Opin Genet Dev* 2002;12:540–547. [PubMed: 12200159]
8. Fogh J, Wright WC, Loveless JD. Absence of HeLa cell contamination in 169 cell lines derived from human tumors. *J Natl Cancer Inst* 1977;58:209–214. [PubMed: 833871]
9. Fogh, J.; Trempe, G. *Human Tumor Cells In Vitro*. New York: Plenum Press; 1975.
10. Dharmasathaphorn K, McRoberts JA, Mandel KG, Tisdale LD, Masui H. A human colonic tumor cell line that maintains vectorial electrolyte transport. *Am J Physiol* 1984;246:G204–G208. [PubMed: 6141741]
11. Blackledge NP, Carter EJ, Evans JR, Lawson V, Rowntree RK, Harris A. CTCF mediates insulator function at the CFTR locus. *Biochem J* 2007;408:267–275. [PubMed: 17696881]
12. Wingender E, Chen X, Hehl R, Karas H, Liebich I, Matys V, Meinhardt T, Pruss M, Reuter I, Schacherer F. TRANSFAC: an integrated system for gene expression regulation. *Nucleic Acids Res* 2000;28:316–319. [PubMed: 10592259]
13. Qiu Y, Tomita Y, Zhang B, Nakamichi I, Morii E, Aozasa K. Pre-B-cell leukemia transcription factor 1 regulates expression of valosin-containing protein, a gene involved in cancer growth. *Am J Pathol* 2007;170:152–159. [PubMed: 17200190]
14. Jonckheere N, Vincent A, Perrais M, Ducourouble MP, Male AK, Aubert JP, Pigny P, Carraway KL, Freund JN, Renes IB, Van Seuning I. The human mucin MUC4 is transcriptionally regulated by caudal-related homeobox, hepatocyte nuclear factors, forkhead box A, and GATA endodermal transcription factors in epithelial cancer cells. *J Biol Chem* 2007;282:22638–22650. [PubMed: 17553805]
15. Paul T, Li S, Khurana S, Leleiko NS, Walsh MJ. The epigenetic signature of CFTR expression is coordinated via chromatin acetylation through a complex intronic element. *Biochem J* 2007;408:317–326. [PubMed: 17848139]
16. Blackledge NP, Ott CJ, Gillen AE, Harris A. An insulator element 3' to the CFTR gene binds CTCF and reveals an active chromatin hub in primary cells. *Nucleic Acids Res* 2009;37:1086–1094. [PubMed: 19129223]
17. Phylactides M, Rowntree R, Nuthall H, Ussery D, Wheeler AHA. Evaluation of potential regulatory elements identified as DNase I hypersensitive sites in the CFTR gene. *Eur J Biochem* 2002;269:553–559.
18. Ktistaki E, Talianidis I. Modulation of hepatic gene expression by hepatocyte nuclear factor 1. *Science* 1997;277:109–112. [PubMed: 9204893]
19. Duncan SA, Navas MA, Dufort D, Rossant J, Stoffel M. Regulation of a transcription factor network required for differentiation and metabolism. *Science* 1998;281:692–695. [PubMed: 9685261]

20. Soutoglou E, Papafotiou G, Katrakili N, Talianidis I. Transcriptional activation by hepatocyte nuclear factor-1 requires synergism between multiple coactivator proteins. *J Biol Chem* 2000;275:12515–12520. [PubMed: 10777539]
21. Isshiki S, Kudo T, Nishihara S, Ikehara Y, Togayachi A, Furuya A, Shitara K, Kubota T, Watanabe M, Kitajima M, Narimatsu H. Lewis type 1 antigen synthase (beta3Gal-T5) is transcriptionally regulated by homeoproteins. *J Biol Chem* 2003;278:36611–36620. [PubMed: 12855703]
22. Gregory PA, Lewinsky RH, Gardner-Stephen DA, Mackenzie PI. Coordinate regulation of the human UDP-glucuronosyltransferase 1A8, 1A9, and 1A10 genes by hepatocyte nuclear factor 1alpha and the caudal-related homeodomain protein 2. *Mol Pharmacol* 2004;65:953–963. [PubMed: 15044625]
23. Farnham PJ. Insights from genomic profiling of transcription factors. *Nat Rev Genet.* 2009

Figure 1A

Figure 1B

Figure 1. Linear maps of the DHS regions analyzed in intron 16 and 17a

A. Intron 16, B. Intron 17a. The diagrams are to scale as marked and the numbers above the gene show the locations on AC000111 (A) and AC000061 (B) respectively. A, Overlapping fragments A – E (ROH in intron16) and 1–5 (DHS16) were evaluated by DNase 1 footprinting. B, Fragments A – E were analyzed across DHS 17a. ROH=region of cross species homology.

Figure 2. Identification of DNase 1 footprints in DHS 16 fragment 4 and DHS17a fragments D and E

The probes and DNA strand shown are as marked above each panel. Footprints are highlighted with a bar to the side of the panel and the corresponding predicted transcription factor binding site(s) are shown only for PRs to be analysed further. For each gel, lane 1, AG ladder, lane 2, probe alone, lane 3, probe with DNase I, lane 4 and 5, probe with DNase I and 30µg or 60µg respectively of nuclear extract from Caco2 cells. A. 16-4 PR2: predicted CDX2 (- strand). B. 17a-D PR3: HNF1 on both strands. C. 17a-E PR6: CDX2 (+ strand); PR7 and PR9: CDX2 (- and + strands respectively); PR8: HNF1 (- strand).

Figure 3. Identification of DNase 1 footprints in 16hom fragments B, C, D and E

The probes and DNA strand shown are as marked above each panel. Footprints are highlighted with a bar to the side of the panel and the corresponding predicted transcription factor binding site(s) are shown only for PRs to be analysed further. For each gel, lane 1, AG ladder, lane 2, probe alone, lane 3, probe with DNase I, lane 4 and 5, probe with DNase I and 30µg or 60µg respectively of nuclear extract from Caco2 cells. In 16 hom B: A. PR1: predicted PBX1 binding site (+ strand) and PBX1/Meis site (on - strand); B. PR2: predicted PBX1 site (- strand). C. In 16homC: PR6: PBX1. D. In 16hom D, PR7: PBX1 and PBX1/Meis and COUP (+ strand). E. In 16hom E, PR14: HNF1 (+ strand); PR15 HNF1 on both strands.

Figure 4. Electrophoretic mobility shift assays demonstrate binding of CDX2 to DNase I footprint FP2 in DHS16
The sequences of the probes and competitors used are shown in Supplementary Table 2A. For all probes, M denotes a mutation in the predicted transcription factor binding site. For each panel the lanes are marked and + denotes with, and – without each competitor or antibody as shown. A, Caco2 nuclear extract, B, *in vitro* translated (IVT) CDX2. In supershift reactions 2μg of antibody was used and βtubulin (βtub) is an isotype control antibody. Arrows A and B denote the major specific complexes formed by interaction of the probe with proteins in the Caco2 nuclear extracts, with A likely including CDX2; complex C denotes a supershift with the CDX2 antibody and IVT CDX2.

Figure 5. Electrophoretic mobility shift assays demonstrate binding of CDX2 to DNase I footprint FP6 in DHS17a

The sequences of the probes and competitors used are shown in Supplementary Table 2A. For all probes, M denotes a mutation in the predicted transcription factor binding site. All lanes except the first include Caco2 nuclear extract and each competitor or antibody as shown. In supershift reactions 2 μ g of antibody was used and β tubulin (β tub) is an isotype control antibody. Arrows A and B denote the major specific complexes formed by interaction of the probe with proteins in the Caco2 nuclear extracts, with A likely including CDX2; complex C denotes a supershift with the HNF1 antibody.

Figure 6. Electrophoretic mobility shift assays demonstrate binding of PBX1 to DNase I footprint FP6 in 16hom

The sequences of the probes and competitors used are shown in Supplementary Table 2C. For all probes, M denotes a mutation in the predicted transcription factor binding site. All lanes except the first include Caco2 nuclear extract and each competitor or antibody as shown. In supershift reactions anti-RAR is an isotype control antibody. Arrowed complex A is the major specific complexes formed by interaction of the probe with proteins in the Caco2 nuclear extracts, and likely includes PBX1; complex B denotes a faint supershift with the PBX1 antibody.

Figure 7. Electrophoretic mobility shift assays demonstrate binding of HNF1 to DNase I footprint FP15 in 16hom

The sequences of the probes and competitors used are shown in Supplementary Table 2B. For all probes, M denotes a mutation in the predicted transcription factor binding site. For each panel the lanes are marked and + denotes with, and - without each competitor or antibody as shown. All lanes except probe alone contain Caco2 nuclear extract. Arrowed complexes A are the major specific complexes formed by interaction of the probe with proteins in the Caco2 nuclear extracts; complex B denotes a supershift with the HNF1 antibody.

Figure 8. Electrophoretic mobility shift assays demonstrate binding of HNF1 to both cognate sites in DNase I footprint FP15 in 16hom, though with different efficiencies

The sequences of the probes are shown and competitors used are shown in Supplementary Table 2B. For the 16homFP15HNF1 probe, B and T denote single mutations in each predicted transcription factor binding site and M denotes the double mutant. For each panel the lanes are marked and + denotes with, and - without each competitor or antibody as shown. All lanes except probe alone contain Caco2 nuclear extract. Arrowed complexes A1 and A2 are the major specific complexes formed by interaction of the probe with proteins in the Caco2 nuclear extracts; complex B denotes a supershift with the HNF1 antibody. Supershifts with anti-GR (glucocorticoid receptor) in A and anti-PBX1 in B are isotype controls.

Figure 9. ChIP shows in vivo binding of PBX1 to 16hom and DHS 17a

ChIP analysis using antibodies specific for HNF1, CDX2 and PBX1 in Capan 1 cells; no antibody and rabbit IgG were control and values are relative to the former with a value of 1. ChIP was followed by real time PCR using PowerSYBR Green and primer sets at the indicated positions of the *CFTR* locus. Primer sets at the VCP promoter and *CFTR* intron 1 (185 +10 kb DHS) were used as positive controls for PBX1 and HNF1 binding, respectively. Primers are shown in Supplementary Table 3. Error bars represent standard error of the mean.