Published in final edited form as:

J Allergy Clin Immunol. 2009 November ; 124(5): 1108–1110. doi:10.1016/j.jaci.2009.06.021.

Bovine and Porcine Gelatin Sensitivity in Milk and Meat-Sensitized Children

Jelena Bogdanovic, Ph.D.^a, Neal A. Halsey, M.D.^b, Robert A. Wood, M.D.^a, and Robert G. Hamilton, Ph.D.^a

^aJohns Hopkins University School of Medicine, Departments of Medicine, Pediatrics and Pathology, Baltimore, MD

^bJohns Hopkins University Bloomberg School of Public Health, Baltimore, MD

Capsule Summary

Cross-reactive bovine and porcine gelatin-specific IgE antibody is more common in cow's milk and beef or pork meat-sensitized individuals than previously known and a potential risk factor for allergic reactions to gelatin-containing products (foods, vaccines).

Keywords

gelatin; porcine; bovine; meat; cow's milk; IgE antibody; prevalence; cross-reactivity

To the Editor

Gelatin is a protein derived from collagen and it is obtained principally from cow and pig bones and hides and fish skin. It is a common ingredient in foods such as jellies, sweets, yogurt and frozen desserts. It is also found in lunch meats, and is used extensively as clarifying agents in wine, juices and other beverages. Bovine and porcine gelatins, in particular, also have numerous applications throughout the pharmaceutical industry as integral components in drug capsules, plasma expanders, and stabilizers in vaccines, including MMR, varicella, yellow fever, rabies, and some influenza vaccines. Severe allergic reactions, including anaphylaxis, have been reported following intravenous administration of modified fluid gelatins as plasma substitutes.¹ Post-vaccination allergic reactions to MMR and varicella vaccines have been linked to the gelatin excipient.^{2–3} Systemic allergic reactions have also been observed with the ingestion of gelatin-containing foods and administration of gelatin-containing medical products (e.g., suppositories). These gelatin exposures have been associated with sensitization as evidenced by the induction of gelatin-specific IgE antibodies.^{2–5}

American⁶ and Finnish⁷ groups have reported that 27% and 14–28%, respectively, of children who experienced systemic reactions after measles, mumps and rubella vaccination, had gelatin-specific IgE antibodies. In contrast, a Japanese study reported that 86% of children, who

© 2009 American Academy of Allergy, Asthma and Immunology. Published by Mosby, Inc. All rights reserved.

Corresponding author Robert G. Hamilton, Ph.D., Johns Hopkins University School of Medicine, Johns Hopkins Asthma & Allergy Center, Room 1A20, 5501 Hopkins Bayview Circle, Baltimore, MD 21224, Telephone: 410-550-2031, Fax: 410-550-2030, E-mail: rhamilt2@jhmi.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

manifested an immediate-type hypersensitivity reaction following receipt of a gelatin-containing vaccine (measles, rubella, mumps, or varicella), had detectable gelatin-specific IgE in their blood.⁸ Type I hypersensitivity reactions to gelatin have been even reported with specific IgE levels as low as 0.8 kUa/L.⁸

In the current study, we chose to not study fish gelatin sensitivity, since the gelatins used in medical applications are almost exclusively bovine and porcine. We hypothesized that subjects who are sensitized to beef and pork meat and/or cow's milk are at greater risk for sensitization to bovine and porcine gelatin. Moreover, we hypothesized that there is cross-reactivity between bovine and porcine gelatin. These hypotheses were investigated using serological techniques to determine the prevalence and extent of cross-reactivity of bovine and porcine gelatin-specific IgE antibodies among children with confirmed sensitivity (IgE antibodies > 0.35 kUa/L) to pork or beef meat and cow's milk and a clinical history of cow's milk allergy.

Serum from children (n=141; 3 months to 17 years, median age: 4 years, 74% male, total IgE range 19–49,457 kU/L, median: 909 kU/L) were selected for evaluation in the current study on the basis of a positive IgE antibody serology to cow's milk, beef and/or pork meat. The exception was one subject who had a weak bovine and porcine gelatin-specific IgE < 0.5 kUa/L in the absence of detectable IgE anti-milk or beef and pork meat. Each subject had a positive history sufficient to warrant evaluation for sensitization to cows' milk, beef and/or pork meat. Bovine and porcine gelatin, beef and pork meat, and cow's milk-specific IgE levels were quantified by ImmunoCAP250 (Phadia, Kalamazoo, Michigan, USA, analytical sensitivity = 0.1 kUa/L). We chose the conservative 0.35 kUa/L threshold to identify positive IgE antibody responses. Cross-reactivity of bovine and porcine gelatin specific IgE antibody was studied by competitive cross-inhibition using soluble homologous and heterologous gelatins at 2 mg/ml (Sigma-Aldrich, St. Louis, MO) or Phadia diluent as the sham negative control.

In this selected population of beef and/or pork meat-sensitized children, 93% of children had beef meat- (range: 0.35–99 kUa/L, median: 2.4 kUa/L), 84% pork meat- (range: 0.39–266 kUa/L; median: 2.3 kUa/L), and 79% both beef and pork-meat specific IgE antibodies. Ninety-seven percent of the subjects were also sensitized to cow's milk (milk-specific IgE: range: 0.39–464 kUa/L). There was a significant correlation ($r=0.66$, $p<0.001$) between the level of IgE anti-beef and IgE anti-pork (meat) in the 111 subjects who had both antibody specificities. The IgE anti-milk levels also correlated weakly, but significantly with the levels of beef meat-specific-IgE ($r=0.41$, $p<0.001$, $n=111$) and pork meat-specific IgE ($r=0.30$, $p=0.002$, $n=100$) in the same serum.

Bovine gelatin-specific IgE was detected in 21 (16%, range 0.35 to 4.12 kUa/L; median: kUa/L) of the 130 beef meat-specific IgE positive children. Only four sera that contained low levels of bovine gelatin-specific IgE (0.42–1.06 kUa/L) had no detectable beef meat-specific IgE. No significant correlation was detected between the quantitative level of bovine gelatin-specific IgE and beef-specific IgE ($r=-0.11$). The majority of IgE anti-bovine gelatin positive sera (88%) also contained detectable porcine gelatin-specific IgE. Within each serum that was positive for both, the quantitative levels of IgE anti-bovine and IgE anti-porcine gelatin were significantly correlated ($r=0.73$, $p<0.001$, $n=21$), suggesting antibody cross-reactivity among gelatins. Competitive inhibition experiments (Table I) verified partial IgE antibody cross-reactivity among the gelatins. While homologous bovine gelatin inhibited bovine gelatin-specific IgE binding to solid phase bovine gelatin by 72–85%, heterologous porcine gelatin only partially inhibited (13–47%) equivalent levels of bovine gelatin-specific IgE binding. Our data confirm partial cross-reactivity seen by others between bovine and mammalian gelatins⁹. Our observations, however, are the first empirical demonstration of cross-reactivity between bovine and porcine gelatin.

Porcine gelatin-specific IgE was detected in 44 (38%, range 0.35–17.98 kUa/L; median: kUa/L) of the 116 pork meat-specific IgE positive children. Only 4 sera negative for pork meat-specific IgE were positive for porcine gelatin-specific IgE (0.36–6.60 kUa/L). No significant correlation was detected between the quantitative levels of porcine gelatin- and pork meat-specific IgE ($r=-0.16$). Homologous solubilized porcine gelatin inhibited porcine gelatin-specific IgE antibody binding to the solid phase porcine gelatin by 35–85% (Table I). Bovine gelatin was a more effective cross-inhibitor of IgE anti-porcine binding to porcine gelatin allergosorbent (81–94% cross-inhibition). The less effective self-inhibition of porcine gelatin-specific IgE binding to the porcine ImmunoCAP by soluble porcine gelatin may have resulted from one of two factors. First, higher concentrations of porcine gelatin could not be used because it gelled at concentrations above 2 mg/ml. Second, the source of porcine gelatin used for insolubilization onto the ImmunoCAP allergosorbent was different than that which was available for soluble antigen competitive inhibition.

In conclusion, 16% and 38% of beef and pork meat sensitized children, respectively, have IgE antibodies to gelatins that are cross-reactive. The presence of IgE anti-gelatin may place them at risk for potential allergic reactions following exposure to gelatin containing foods, vaccines or other medical products. However, the true clinical significance of these will require further, prospective study.

Acknowledgments

Declaration of all sources of funding: This project was supported by internal funds within the Johns Hopkins University School of Medicine and by contract no. 1406-04-08-CT-20000 from the CDC.

References

1. Vervloet D, Senft M, Dugue P, Arnaud A, Charpin J. Anaphylactic reactions to modified fluid gelatins. *J Allergy Clin Immunol* 1983;71:535–540. [PubMed: 6189875]
2. Kelso JM, Jones RT, Yunginger JW. Anaphylaxis to measles, mumps, and rubella vaccine mediated by IgE to gelatin. *J Allergy Clin Immunol* 1993;91:867–872. [PubMed: 8473675]
3. Sakaguchi M, Yamanaka T, Ikeda K, Sano Y, Fujita H, Miura T, et al. IgE-mediated systemic reactions to gelatin included in the varicella vaccine. *J Allergy Clin Immunol* 1997;9:263–264. [PubMed: 9042057]
4. Wang J, Sicherer SH. Anaphylaxis following ingestion of candy fruit chews. *Ann Allergy Asthma Immunol* 2005;94:530–533. [PubMed: 15945554]
5. Yamada A, Ohshima Y, Tsukahara H, Hiraoka M, Kimura I, Kawamitsu T, et al. Two cases of anaphylactic reaction to gelatin induced by a chloral hydrate suppository. *Pediatr Int* 2002;44:87–89. [PubMed: 11982879]
6. Pool V, Miles Braun M, Kelso JM, Mootrey G, Chen RT, Yunginger JW, et al. Prevalence of anti-gelatin IgE antibodies in people with anaphylaxis after measles-mumps-rubella vaccine in the United States. *Pediatrics* 2002;110:e71. [PubMed: 12456938]
7. Patja A, Mäkinen-Kiljunen S, Davidkin I, Paunio M, Peltola H. Allergic reactions to measles-mumps-rubella vaccination. *Pediatrics* 2001;107:e27. [PubMed: 11158501]
8. Sakaguchi M, Nakayama T, Fujita H, Toda M, Inouye S. Minimum estimated incidence in Japan of anaphylaxis to live virus vaccines including gelatin. *Vaccine* 2001;19:431–436. [PubMed: 11027805]
9. Sakaguchi M, Hori H, Ebihara T, Irie S, Yanagida M, Inouye S. Reactivity of immunoglobulin E in bovine gelatin-sensitized children to gelatins from various animals. *Immunology* 1999;96:286–290. [PubMed: 10233707]

IgE antibody cross-reactivity between bovine and porcine gelatins, as measured by the ImmunoCAP-250 autoanalyzer.

Sample No. *	IgE anti-bovine gelatin		IgE anti-porcine gelatin	
	Uninhibited kUa/L	Inhibited with:	Uninhibited kUa/L	Inhibited with:
		Bovine gelatin kUa/L (%inhibition)		Bovine gelatin kUa/L (%inhibition)
1	4.1	0.9 (79%)	16.6	2.0 (88%)
2	3.0	0.5 (85%)	7.7	1.5 (81%)
3	4.1	1.2 (72%)	18.0	2.5 (86%)
4	1.0	0.2 (80%)*	3.7	0.2 (94%)
5	0.5	<LOD ()	6.6	0.8 (88%)
3.5 (79%)				
3.4 (56%)				
2.7 (85%)				
1.1 (69%)				
4.3 (35%)				

LOD = Limit of Detection

* The percentage of inhibition could not be accurately determined

* Note : sera from only the 5 subjects noted in the table were in sufficient quantities to permit repetitive inhibition.