Published in final edited form as:

J Neuroimmunol. 2009 August 18; 213(1-2): 123–130. doi:10.1016/j.jneuroim.2009.05.014.

Potential of a unique antibody gene signature to predict conversion to clinically definite Multiple Sclerosis

Elizabeth M. Cameron^a, Sade Spencer^a, Jonathan Lazarini^a, Christopher T. Harp^a, E. Sally Ward^b, Mark Burgoon^c, Gregory P. Owens^c, Michael K. Racke^d, Jeffrey L. Bennett^c, Elliot M. Frohman^a, and Nancy L. Monson^{a,b}

^a Department of Neurology, University of Texas Southwestern Medical Center, Dallas TX 75154

^b Department of Immunology, University of Texas Southwestern Medical Center, Dallas TX 75154

^c Department of Neurology, University of Colorado Denver, Denver, CO, 80262

^d Department of Neurology, Ohio State University Medical Center, Columbus, OH, 43210

Abstract

We identified a unique antibody gene mutation pattern (i.e. “signature”) in CSF B cells from MS patients not present in control populations. Prevalence of the signature in CSF B cells of patients at risk to develop MS predicted conversion to MS with 91% accuracy in a small cohort of clinically isolated syndrome patients. If confirmed, signature prevalence would be a novel genetic diagnostic tool candidate for patients with early demyelinating disease of the central nervous system.

Keywords

Multiple sclerosis; B lymphocytes; Antibodies; Gene rearrangement; Mutational signature

1. Introduction

B cells have historically been implicated in the pathogenesis of multiple sclerosis (MS) since elevated central nervous system (CNS) gamma immunoglobulins were first described in MS patients in the 1940s (Kabat et al., 1948). Additional evidence of B cell involvement in MS pathogenesis is extensive (reviewed in (Antel and Bar-Or, 2006; Duddy and Bar-Or, 2006; Franciotta et al., 2008; Frohman et al., 2006; Martin Mdel and Monson, 2007; McLaughlin and Wucherpfennig, 2008; Owens et al., 2006)) and has been recently substantiated by the efficacy of rituximab (Rituxan), a B cell depleting antibody, in a cohort of patients with relapsing remitting MS (RRMS) (Hauser et al., 2008). Furthermore, Rituxan and intravenous immunoglobulin, drugs that solely affect B cells or their antibody products, have been reported to decrease severity of disease in MS patients refractory to benefit with corticosteroids, interferon-beta, and mitoxantrone (Achiron, 2008; Leussink et al., 2008; Stuve et al., 2005; Tselis et al., 2008).

To whom correspondence should be addressed: Nancy.Monson@UTSouthwestern.edu, Phone Number: (214) 648-9413, Fax Number: (214) 648-9129.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Several groups investigating the role of B cells in MS have hypothesized that the distribution of genes used to generate antibodies in B cells from the cerebrospinal fluid (CSF) and brain lesions of MS patients are different from expected distributions. Indeed, the distributions are different in some cases, particularly with regard to a family of variable heavy chains (VH4), which are significantly increased in frequency compared to expected distributions (Baranzini et al., 1999; Colombo et al., 2000; Harp et al., 2007; Monson et al., 2005; Owens et al., 1998; Owens et al., 2003; Owens et al., 2007; Qin et al., 1998; Ritchie et al., 2004). Additionally, MS CSF B cells show extensive clonal expansion and high mutational frequencies in the CSF B cell pool from this population of patients (Baranzini et al., 1999; Colombo et al., 2000; Monson et al., 2005; Owens et al., 2003; Qin et al., 1998; Ritchie et al., 2004), and the antibodies these cells produce bind to neuroantigens (Kolln et al., 2006; Lambracht-Washington et al., 2007). In contrast, VH4 expressing B cells in the periphery of healthy donors (Brezinschek et al., 1995; Brezinschek et al., 1997), MS patients (Owens et al., 2007), and VH4 expressing B cells in the CSF of patients with other neurological diseases (OND) are present at expected frequencies (Table 1 and (Harp et al., 2007)).

Since antibody gene mutation patterns are influenced by antigen driven selection, we hypothesized that VH4 expressing CSF-derived B cells of MS patients would harbor antibody gene mutation patterns that would be distinct from VH4 expressing peripheral B cells derived from healthy controls. To address this contention, we characterized antibody gene mutations in a VH4 subdatabase extracted from the parent heavy chain antibody database consisting of 373 CSF-derived B cells from 11 patients with definite MS. Our analysis revealed a unique pattern of antibody gene replacement mutations in CSF B cells from MS patients that was not prevalent in antibody gene repertoires from CSF B cells of OND patients. Furthermore, prevalence of this conspicuous signature in B cell antibody repertoires from patients with a first inflammatory demyelinating episode (a clinically isolated syndrome; CIS) can predict conversion to clinically definite MS (CDMS) within 3–18 months after initial sampling.

2. Materials and methods

2.1 Patient description

CSF was collected from ten RRMS patients, one PPMS patient (M484), three patients with other neurological diseases (OND341, ataxia; OND758, headache, and OND116, chronic inflammatory demyelinating polyneuropathy), and two patients with one demyelinating event suggestive of MS (i.e. Clinically Isolated Syndrome (CIS)) at UT Southwestern Medical Center (UTSWMC) (Harp et al., 2007; Monson et al., 2005) in accordance with the UTSWMC Institutional Review Board (IRB). CSF was collected from nine patients with CIS at University of Colorado Denver (UCD) as previously described (Bennett et al., 2008) in accordance with the UCD IRB. The CIS patients had a single episode of demyelination (optic neuritis, brainstem or spinal cord syndrome), and the majority had multiple lesions on MRI satisfying the dissemination in space criterion of the McDonald criteria. None of the patients had received immunomodulatory agents for at least one month prior to lumbar puncture. A second relapse confirming a multiple sclerosis diagnosis had not occurred at the time of sample acquisition, thus not fulfilling the dissemination in time criterion (McDonald et al., 2001; Polman et al., 2005). Subsequent diagnosis of definite MS was made using the revised McDonald criteria (Polman et al., 2005). Conversion to definite MS was not revealed to the antibody sequence analysis team until after signature score predictions had been calculated.

2.2 MS and CIS B Cell Antibody Database Generation

At UTSWMC, antibody repertoires were generated from CD19⁺ CSF B cells using single cell PCR as previously described (Harp et al., 2007; Monson et al., 2005). The MSCSF database consists of antibody rearrangements from 373 CD19⁺ CSF B cells from 10 RRMS and 1 PPMS

patient recruited at UTSWMC. The CISCSF database consists of antibody rearrangements from 304 CD19⁺ CSF B cells from 10 CIS patients (ON4–8 did not have a CD19⁺ CSF B cell antibody repertoire) and 228 CD138⁺ CSF plasma cells from 7 CIS patients (CIS132, CIS429, ON4–10 and ON3–4 did not have CD138⁺ CSF plasma cell antibody repertoires). To clarify, antibody repertoires from CIS patients at UCD were generated from both single CD19⁺ CSF B cells and single CD138⁺ CSF plasma cells (Bennett et al., 2008), while antibody repertoires from CIS patients at UTSWMC were generated from single CD19⁺ CSF B cells only. Since the resultant databases (CIS CD19⁺ CSF from UTSWMC, CIS CD19⁺ and CIS CD138⁺ CSF from UCD) were similar in mutational frequency, variable heavy chain (VH) gene family usage, and heavy chain Joining segment (JH) usage, the two databases were combined for analysis (Supplemental Table 1).

2.3 Control B Cell Antibody Database Generation

The healthy control peripheral blood (HCPB) antibody database has been used in multiple studies (Brezinschek et al., 1997; Brezinschek et al., 1998; Dorner et al., 1997; Dorner et al., 1998a; Dorner et al., 1998b; Dorner et al., 1998c; Farner et al., 1999; Hansen et al., 2000; Harp et al., 2007; Monson et al., 2005; Monson et al., 2000) and consists of 348 CD19⁺ or CD19⁺IgM⁺ peripheral B cells from two healthy control donors. The memory HCPB antibody database (mHCPB) consists of 205 sequences from the HCPB antibody database that contain 4 or more mutations (less than 98% homology to the germline sequence, n=123) combined with sequences from a HCPB antibody database generated from class-switched IgD⁺CD27⁺ memory B cells (n=82) (Tian et al., 2007) (Genbank 535266–535274, 535324–535368, 535381–535408, and 535416–535418). As expected, the class-switched IgD⁺CD27⁺ memory B cell database had a higher percentage of mutated codons that resulted in a replacement than the mHCPB database (compare 64.7% vs 70.3%, $p=0.002$ by χ^2 test). The OND CD19⁺ CSF antibody database consists of 65 sequences. UCD and UTSWMC cell isolation and IgH amplification was performed similarly. All sequences were re-confirmed by our laboratory using IgBlast (those obtained from UCD and from GenBank) (<http://www.ncbi.nlm.nih.gov/igblast/>), and only codons 24–93 were considered in the analysis.

2.4 Mutation analyses

Frequency of replacement mutations (RF) was calculated as the number of replacement mutations at each codon position divided by the total number of replacement mutations in each VH4 subdatabase and displayed as a percentage. The MSCSF database contains 373 sequences with 2,475 replacement mutations, and the CISCSF database contains 302 CD19⁺ and 226 CD138⁺ sequences with 4081 replacement mutations (2,052 in CD19⁺ and 2,029 in CD138⁺). The HCPB database contains 348 sequences with 1,086 replacement mutations, and the mHCPB database contains 205 sequences with 1,857 replacement mutations. The ONDCSF database contains 65 sequences with 482 replacement mutations, and the MSPB database contains 156 sequences with 392 replacement mutations. In total, 1,675 sequences and 10,373 replacement mutations were analyzed in this manner. Table 1 contains VH4 sequence numbers and Table 2 legend contains number of VH4 replacement mutations. Codon domains and numbers were defined by Kabat (Kabat et al., 1983), and Tomlinson in V-base (vbase.mrc-cpe.cam.ac.uk/), respectively.

2.5 Statistical strategy for signature identification

Codons included in the signature were identified using three criteria. First, we identified codons that had statistically different RF values in the MSCSF_{VH4} database compared to HCPB_{VH4} by Goodness of Fit test where the expected frequency is the RF calculated in HCPB_{VH4}. Twenty-four codons passed this criterion. Next, codon positions that had an RF in both the

MSCSF_{VH4} and HCPB_{VH4} databases that was less than the average + 2 S.D. of the memory HCPB_{VH4} subdatabase were excluded. Thus, since the average $\pm$ S.D. RF of the memory HCPB_{VH4} database was 0.68 ± 0.59 , any individual codon RF less than 1.86 in both databases was excluded. Fourteen codons passed this additional criterion. Eight of these 14 codons (31B, 32, 40, 56, 57, 60, 81, and 89) were defined as “hot” since the RF at that codon position within the MSCSF_{VH4} database was statistically higher compared to the HCPB_{VH4} database. Six of these 14 codons (30, 43, 52, 77, 82 and 82a) were defined as “cold” since the RF at that codon position within the MSCSF_{VH4} database was statistically less compared to the HCPB_{VH4} database. Two of the 6 “cold” codons (52 and 82a) were excluded because the RF value in the MSCSF_{VH4} database at that codon position was significantly higher than 1.86 (the average + 2 S.D. of the memory HCPB_{VH4} subdatabase). The overall signature consequently consisted of codons 30, 31B, 32, 40, 43, 56, 57, 60, 77, 81, 82, and 89. This analysis was not biased by differences in the prevalence of particular codons (31B in particular), as individual VH4 gene frequencies in MSCSF were similar to HCPB by χ^2 test using a Bonferroni corrected p-value of 0.004 (data not shown).

2.5 Statistical computation of the signature score

Signature scores were generated by calculating Z-scores for the RF values at the 6 codons within the signature (31B, 40, 56, 57, 81 and 89) that had the most significant difference in RF compared to HCPB_{VH4} at each codon position. The Z-score formula is: (RF at codon X minus the average RF in HCPB_{VH4}) divided by the standard deviation of the average RF in HCPB_{VH4}. For example, the average RF in HCPB_{VH4} within the 6 signature codons was 1.6 ± 0.9 and so an RF of 4.4 at codon 31B would be assigned a score of 3.1 ((Z score=(4.4-1.6)/0.9)). Individual z-scores at each of the 6 codon positions were then added to generate the composite signature z-score. The average composite signature score in the MSCSF_{VH4} database was 10.9 ± 2.0 and so any signature score of an individual CIS patient above 6.8 (average – 2 S.D.) was predicted to convert to CDMS. Of note, both the ONDCSF_{VH4} signature score (at 4.5), and the MSPB_{VH4} score (at 2.0) were below the threshold for MS conversion. CD19⁺ CSF B cell and CD138⁺ CSF plasma cell mutation positions both contributed to each CIS patient's signature score, while the MSCSF_{VH4} signature scores were only composed of CD19⁺ CSF B cells.

2.6 VH4 structure

A human VH4–30.4 antibody structure described in (Guddat et al., 1993) was obtained from the Protein Data Bank (www.rcsb.org) under the identification moniker 1MCO, and adapted using the RasMol program (mc2.cchem.berkeley.edu/Rasmol/) to highlight codons within the designated signature of the heavy chain variable region.

3. Results

The 51 antibody heavy chain variable genes are subdivided into 7 different families ((Cook and Tomlinson, 1995) (<http://www.ncbi.nlm.nih.gov/igblast/>)), and it has been well-established that peripheral blood B cells from healthy donors utilize VH antibody genes most often from the VH3 family (“HCPB” in Table 1 and (Brezinschek et al., 1995; Brezinschek et al., 1997; Huang et al., 1992; Kraj et al., 1997; Wardemann et al., 2003; Yurasov et al., 2005)). In contrast, it has been reported by us and others that B cells in the CSF of MS patients often utilize VH4 antibody genes more frequently than those in the VH3 family (“MSCSF” in Table 1 and (Baranzini et al., 1999; Colombo et al., 2000; Harp et al., 2007; Monson et al., 2005; Owens et al., 1998; Owens et al., 2003; Owens et al., 2007; Qin et al., 1998; Ritchie et al., 2004)). The CISCSF antibody database consisting of CD19⁺ B cells only had a similar frequency of B cells that utilize VH4 family genes in comparison to HCPB (26.2% vs. 21.8%, $p=0.20$ by χ^2 test) (Table 1); in contrast, when CD138⁺ plasma cells were included, the CISCSF

had a higher frequency of B cells that utilize VH4 family genes in comparison to HCPB (35.0% vs. 21.8%, $p=0.00001$ by χ^2 test) (Table 1). Some individual CIS patient CSF B cell antibody repertoires were enriched for B cells utilizing VH4 family genes in comparison to the random expected frequency, as reported previously (Bennett et al., 2008). CSF-derived B cell antibody repertoires from patients with Other Neurological Diseases (OND) were not enriched for VH4-expressing CSF B cells in comparison to HCPB (23.1% vs. 21.8%, $p=0.83$ by χ^2 test) or mHCPB (23.1% vs. 16.6%, $p=0.24$ by χ^2 test), indicating that VH4 over-expression in the CSF of MS patients was not due to bias in the ability of VH4 expressing B cells to enter the CNS.

3.1 Identification of codons within MSCSF that are enriched for replacement mutations

Since VH4 expressing B cells are enriched in the CSF of MS patients, we hypothesized that mutational analysis would reveal a pattern (i.e. “signature”) of antibody gene replacement mutations that is unique to VH4 expressing B cells from the CNS of MS patients in comparison to HCPB. In order to test this hypothesis, the percentage of replacement mutations (RF) at each codon within the VH4 subdatabase extracted from the parent database (MSCSF_{VH4}) was determined and compared to the RF at each codon position within the VH4 subdatabase extracted from the parent HCPB database (HCPB_{VH4}). Replacement frequencies were used so that only those mutations resulting in an amino acid change would be considered. In addition, codon amino acid replacement can result from 1, 2, or 3 nucleotide changes within the codon, and so replacement frequencies limit bias based on the number of nucleotides in a codon that are mutated to generate a replacement. Hot spots were defined as those codon positions within MSCSF_{VH4} with a statistically higher RF at a particular codon position in comparison to HCPB_{VH4} (Table 2). Using this approach, 8 codon positions (31B, 32, 40, 56, 57, 60, 81, and 89) were identified that have a total RF value in MSCSF_{VH4} (25.0%) that was statistically higher than in HCPB_{VH4} (12.6%) ($p=0.001$ by χ^2 test). Cold spots were defined as those codon positions within MSCSF_{VH4} with a statistically lower RF at a particular codon position in comparison to HCPB_{VH4} (Table 2). Four codons (30, 43, 77 and 82) were identified as cold spots that have a total RF value in MSCSF_{VH4} (5.1%) that was statistically less than in HCPB_{VH4} (8.5%) ($p=0.001$ by χ^2 test).

Individual MS patient RFs within the 8 hot spot codons of the signature ranged from 22.5 to 34.1 percent (data not shown), indicating that some individual patient MSCSF repertoires had a greater enrichment of replacements at these 8 codon positions than others. Also, the variability of RF values within the 8 hot spot codons of the signature in individual VH4 genes in MSCSF_{VH4} ranged from 14.5 to 36 percent (data not shown), indicating that some individual VH4 genes had a greater enrichment of replacements at these 8 hot spot codon positions than others. Previous analysis had identified codon 56 as a replacement mutation hotspot in HCPB (Dorner et al., 1997; Dorner et al., 1998a), which intensified as a hot spot in MSCSF_{VH4} since a significantly greater percentage of replacement mutations were found in MSCSF_{VH4} at codon 56 compared to HCPB_{VH4}. Of note, there was a 7.0 fold increase in replacement accumulation at codon 31B in the MSCSF_{VH4} database in comparison to HCPB_{VH4} that is likely due to the use of this codon by only a subset of VH4 genes (4–30, 4–31, 4–39 and 4–61). When the analysis was restricted to those B cells expressing VH4 genes that contain codon 31B, there was a 3.1-fold increase in RF of the MSCSF_{VH4} database compared to HCPB_{VH4} ($p<0.001$).

An example of a signature-enriched VH4 antibody gene rearrangement from a CSF-derived B cell of an MS patient is provided in Figure 1. Of note, 5 of the 8 hot spot codons of the signature retained higher RF values in MSCSF compared to the memory HCPB database (31B, 40, 56, 57, and 60), emphasizing that the signature does not simply reflect enrichment of memory B cells in the CSF.

3.2 Potency of signature score to predict development of clinically definite MS

We reasoned that prevalence of the signature would allow us to identify patients at risk to develop MS who subsequently convert to CDMS. Current criteria for diagnosis of MS requires dissemination of lesions both in time and space (Barkhof et al., 1997; Polman et al., 2005; Tintore et al., 2000). When MRI lesions alone are not sufficient to confirm diagnosis, CSF abnormalities can be used to meet the criteria of dissemination in space (Polman et al., 2005; Siritho and Freedman, 2009). Risk of conversion to clinically definite MS in patients who have had a single demyelinating event is 50 to 90% if the patient has an abnormal MRI (Beck et al., 2003; Brex et al., 2002; Cole et al., 1998; O’Riordan et al., 1998; Soderstrom et al., 1998), but of those patients with a normal MRI, up to 29% had oligoclonal bands and converted to CDMS (Cole et al., 1998).

In order to test whether signature prevalence could predict conversion to MS, we generated CSF B cell repertoires from patients who had one demyelinating event that placed them “at risk” to develop MS. Such patients are typically diagnosed with CIS. CD19⁺ B cell and CD138⁺ plasma cell repertoires from the CSF of two CIS patients at UTSWMC and nine patients at UCHSC were generated and analyzed for RF values within the 6 codons of the signature defined in the MSCSF_{VH4} database that had the most significant difference in RF compared to HCPB_{VH4} at each codon position (codons 31B, 40, 56, 57, 81 and 89). RF values were combined using a signature score that accounts for RF variance as described in Materials and Methods. The average signature score in the MSCSF_{VH4} database was 10.9 ± 2.0 (range 7.6 to 11.9), and so any individual CIS patient score that was 6.8 (average signature score – 2 S.D.) or higher was predicted to develop MS (Figure 2 and Table 3). Notably, the signature score from a pool of VH4 expressing CSF B cells of 3 OND patients was 4.5, and the signature score from a pool of VH4 expressing peripheral blood B cells of 3 CDMS patients was 2.0, and thus did not reach the 6.8 signature score threshold. Also, signature scores based on CD19⁺ B cell sequences only (in the patients where this was possible) did not change predictions based on signature score. This was expected since there is significant overlap in the antibody gene repertoires of CD19⁺ B cells and CD138⁺ plasma cells from the CSF of the same patients (Martin Mdel and Monson, 2007; Ritchie et al., 2004), suggesting that the memory B cell pool present in the CSF is the reservoir for differentiation of plasma cells in the CSF.

As indicated in Table 3, prediction of conversion to CDMS using the antibody gene signature score was accurate in 8 of 8 CIS patients that converted to CDMS. Lack of signature prevalence also accurately predicted that 2 of 2 patients who had recently experienced a first demyelinating event (ON3-1 and ON4-10) would not develop CDMS, and indeed, have not developed CDMS up to 2 years after initial sampling. One additional patient who had recently experienced a first demyelinating event (ON3-4) had a high signature score (11.3), but had not converted to CDMS at the two-year follow-up. The antibody gene signature yielded a sensitivity of 100%, specificity of 67%, positive predictive value of 89%, negative predictive value of 100%, and accuracy of 91%, as defined by others applying the McDonald Criteria to identify CIS patients that would convert to MS (Dalton et al., 2002). Most patients in this cohort converted to CDMS within 3–6 months of repertoire sampling, although in the case of CIS132, conversion to CDMS was not confirmed until 17 months after antibody repertoire sampling (Table 3). MRI, OCB and VH4/VH2 bias are also useful in assessing probability of MS conversion (Bennett et al., 2008; Freedman et al., 2005; Frohman et al., 2003; Korteweg et al., 2006; Paolino et al., 1996; Soderstrom et al., 1998), but were not considered in calculating the signature score.

4. Discussion

The intense somatic hypermutation accumulation in MSCSF_{VH4} enabled us to identify a unique antibody gene signature enriched for replacement accumulation at codons 31B, 32, 40, 56, 57,

60, 81 and 89 that was not observed in HCPB_{VH4} or ONDCSF_{VH4}. Of note, any residual effect of naïve B cells on the RF calculation was minimized by tabulating only those sequences with mutations resulting in amino acid replacements. This approach minimized bias in the signature that may have reflected enrichment of mutation accumulation in CNS derived B cells (which are mostly memory and thus have high mutation rates) compared to peripheral B cells (which are mostly naïve and thus have low mutation rates). In addition, 5 of the 8 hot codons of the signature retain higher RF values in the MSCSF database compared to the memory HCPB database. Finally, if signature score reflected enrichment of mutation accumulation due to the repertoire's high memory representation, then all signature scores from the CISCSF antibody repertoires should have been high since they were all heavily enriched for memory B cells. This was not the case, since CISCSF repertoires ON3-1 and ON4-10, despite being heavily enriched for memory B cells (with mutation frequencies of 5.2% and 6.7%, respectively), had signature scores below the threshold of 6.8 (ON3-1 score=6.4, ON4-10 score=2.2).

It was compelling to investigate whether the antibody gene signature may be of value to identify CIS patients who would subsequently develop MS, since early and accurate diagnosis of MS is of tantamount importance in clinical care (Stuve et al., 2008). Signature prevalence could be used to identify patients who would be diagnosed with CDMS within 3–18 months of experiencing their first demyelinating event. Of note, patient ON3-4 had a signature score that indicated this patient would convert to CDMS (score=11.3), but did not demonstrate a lesion load by MRI, banding by OCB, or VH4/VH2 bias, and had not developed CDMS up to 2 years after sampling was performed (Table 3). It will be interesting to determine whether this patient is diagnosed with CDMS over time. It is also important to note that the majority of patients in this cohort already had evidence of MS risk as indicated by positive MRI and OCB. Patient ON4-7, however, did not present with brain lesions by MRI, but had a signature score that indicated this patient would be diagnosed with CDMS (score=10.2). Indeed, this patient did convert to CDMS within 5 months of CSF B cell antibody repertoire sampling, and provides a reasonable example of how signature prevalence may predict CDMS diagnosis in patients that either do not present with brain lesions by MRI, or who are not evaluated by MRI at this stage of their disease. It will be interesting to determine whether the combination of MRI and signature prevalence would be useful in predicting MS conversion. Signature prevalence may also provide an evaluation mechanism to identify the most appropriate patient candidates to receive B cell depletion therapies, for example. Certainly this is a priority since a recent investigation demonstrated significant efficacy of Rituxan in RRMS patients (Hauser et al., 2008).

Given the urgency for the early identification of MS and the rapid initiation of disease modifying therapy, presentation of a molecular signature in the CSF B cells of CIS patients who develop MS may provide a unique tool for identifying at-risk individuals. However, wide implementation of the current form of this approach would be problematic since the AGS scores presented here were generated using a specialized technique that is labor intensive (single cell PCR) and requires fresh CSF for sampling. Developing other approaches to generate AGS data that maintains accuracy, but does not require a specialized laboratory to perform, is attainable and of paramount importance.

In addition, many early MS patients have atypical clinical presentations or unremarkable MRI scans, and patients with alternative inflammatory conditions may mimic idiopathic demyelinating disease. In these circumstances, the advent of a molecular diagnostic signature would increase diagnostic sensitivity and specificity. Investigating the utility of the antibody gene signature in such patients is ongoing in our laboratory.

The presence of a mutational signature among clonally expanded VH4 germline antibodies in MSCSF may be helpful in understanding disease pathogenesis. For example, the VH4 germline

mutational signature may be the direct result of antigen targeting in the humoral immune response. Therefore, determining the antigen specificity of signature-enriched antibodies from CSF B cells of patients with definite MS and CIS is one of the first steps towards dissecting whether signature-enriched B cells have the potential to participate in MS pathogenesis. Of note, 5 of the 8 signature codons (31B, 32, 56, 57 and 60) we identified as having a unique accumulation of amino acid replacements in MSCSF_{VH4} are predicted to have direct antigen contact since they reside in CDRs (Supplemental Figure 1). Dissecting the relative contribution of replacement mutations at each of these signature codons as well as those outside of the antigen-binding region will address the impact of both codon classifications (direct and indirect antigen binding capacity) on antigen binding affinity.

In summary, a unique signature of antibody gene replacement mutations was identified in the MSCSF_{VH4} database that is not observed in healthy control peripheral blood or CSF-derived VH4-expressing B cells from patients with other neurological diseases. Prevalence of the signature was accurate in identifying CIS patients that would convert to CDMS, but needs to be tested on a larger cohort of patients at both high and low risk to develop MS. Identifying the antigen specificity of signature-enriched CSF B cells from these patients may also reveal a unique group of antigens that are central to initiation of humoral autoimmunity in the CNS. It is likely that the MS-specific VH4 antibody gene signature provides both a new focus of investigation to further elucidate the role of B cells and their antibody products in MS and a new candidate molecular diagnostic tool for MS.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors thank the patients who consented to sampling for this study. Special thanks to Dr. Olaf Stüve for critical review of this manuscript. Bonnie Darnell, Angie Mobley and Elizabeth Curry are thanked for their technical expertise in cell sorting, and Tanya Hendricks and Timothy Ahearne for repertoire analysis support.

This study was supported by grants from the National Institutes of Health (NIH) to NLM (RO1 NS 40993) and MKR (RO1 NS 37513, RO1 AI 47133, and K24 NS 44250), the National Multiple Sclerosis Society (NMSS) to NLM (RG3267) and JLB (RG3908), the Yellow Rose Foundation (NLM and MKR), the Wadsworth Foundation (to NLM) and Howson funds (to NLM). EMC and CH were supported by NIH NRSA5 T32 AI 005284-28 from NIAID. The authors have no conflicting financial interests.

References

- Achiron A. Winning combination: the additive/synergistic benefits of IVIg in corticosteroid refractory optic neuritis. *Eur J Neurol* 2008;15:1145. [PubMed: 18973610]
- Antel J, Bar-Or A. Roles of immunoglobulins and B cells in multiple sclerosis: from pathogenesis to treatment. *J Neuroimmunol* 2006;180:3–8. [PubMed: 16934338]
- Baranzini SE, Jeong MC, Butunoi C, Murray RS, Bernard CC, Oksenberg JR. B cell repertoire diversity and clonal expansion in multiple sclerosis brain lesions. *J Immunol* 1999;163:5133–5144. [PubMed: 10528220]
- Barkhof F, Filippi M, Miller DH, Scheltens P, Campi A, Polman CH, Comi G, Ader HJ, Losseff N, Valk J. Comparison of MRI criteria at first presentation to predict conversion to clinically definite multiple sclerosis. *Brain* 1997;120 (Pt 11):2059–2069. [PubMed: 9397021]
- Beck RW, Trobe JD, Moke PS, Gal RL, Xing D, Bhatti MT, Brodsky MC, Buckley EG, Chrousos GA, Corbett J, Eggenberger E, Goodwin JA, Katz B, Kaufman DI, Keltner JL, Kupersmith MJ, Miller NR, Nazarian S, Orengo-Nania S, Savino PJ, Shults WT, Smith CH, Wall M. High- and low-risk profiles for the development of multiple sclerosis within 10 years after optic neuritis: experience of the optic neuritis treatment trial. *Arch Ophthalmol* 2003;121:944–949. [PubMed: 12860795]

- Bennett JL, Haubold K, Ritchie AM, Edwards SJ, Burgoon M, Shearer AJ, Gilden DH, Owens GP. CSF IgG heavy-chain bias in patients at the time of a clinically isolated syndrome. *J Neuroimmunol* 2008;199:126–132. [PubMed: 18547652]
- Brex PA, Ciccarelli O, O’Riordan JI, Sailer M, Thompson AJ, Miller DH. A longitudinal study of abnormalities on MRI and disability from multiple sclerosis. *N Engl J Med* 2002;346:158–164. [PubMed: 11796849]
- Brezinschek HP, Brezinschek RI, Lipsky PE. Analysis of the heavy chain repertoire of human peripheral B cells using single-cell polymerase chain reaction. *J Immunol* 1995;155:190–202. [PubMed: 7602095]
- Brezinschek HP, Foster SJ, Brezinschek RI, Dorner T, Domiati-Saad R, Lipsky PE. Analysis of the human VH gene repertoire. Differential effects of selection and somatic hypermutation on human peripheral CD5(+)/IgM+ and CD5(-)/IgM+ B cells. *J Clin Invest* 1997;99:2488–2501. [PubMed: 9153293]
- Brezinschek HP, Foster SJ, Dorner T, Brezinschek RI, Lipsky PE. Pairing of variable heavy and variable kappa chains in individual naive and memory B cells. *J Immunol* 1998;160:4762–4767. [PubMed: 9590222]
- Cole SR, Beck RW, Moke PS, Kaufman DI, Tourtellotte WW. The predictive value of CSF oligoclonal banding for MS 5 years after optic neuritis. Optic Neuritis Study Group. *Neurology* 1998;51:885–887. [PubMed: 9748050]
- Colombo M, Dono M, Gazzola P, Roncella S, Valetto A, Chiorazzi N, Mancardi GL, Ferrarini M. Accumulation of clonally related B lymphocytes in the cerebrospinal fluid of multiple sclerosis patients. *J Immunol* 2000;164:2782–2789. [PubMed: 10679121]
- Cook GP, Tomlinson IM. The human immunoglobulin VH repertoire. *Immunol Today* 1995;16:237–242. [PubMed: 7779254]
- Dalton CM, Brex PA, Miszkiel KA, Hickman SJ, MacManus DG, Plant GT, Thompson AJ, Miller DH. Application of the new McDonald criteria to patients with clinically isolated syndromes suggestive of multiple sclerosis. *Ann Neurol* 2002;52:47–53. [PubMed: 12112046]
- Dorner T, Brezinschek HP, Brezinschek RI, Foster SJ, Domiati-Saad R, Lipsky PE. Analysis of the frequency and pattern of somatic mutations within nonproductively rearranged human variable heavy chain genes. *J Immunol* 1997;158:2779–2789. [PubMed: 9058813]
- Dorner T, Brezinschek HP, Foster SJ, Brezinschek RI, Farner NL, Lipsky PE. Comparable impact of mutational and selective influences in shaping the expressed repertoire of peripheral IgM+/CD5- and IgM+/CD5+ B cells. *Eur J Immunol* 1998a;28:657–668. [PubMed: 9521076]
- Dorner T, Brezinschek HP, Foster SJ, Brezinschek RI, Farner NL, Lipsky PE. Delineation of selective influences shaping the mutated expressed human Ig heavy chain repertoire. *J Immunol* 1998b;160:2831–2841. [PubMed: 9510186]
- Dorner T, Foster SJ, Farner NL, Lipsky PE. Somatic hypermutation of human immunoglobulin heavy chain genes: targeting of RGYW motifs on both DNA strands. *Eur J Immunol* 1998c;28:3384–3396. [PubMed: 9808208]
- Duddy M, Bar-Or A. B-cells in multiple sclerosis. *Int MS J* 2006;13:84–90. [PubMed: 17101076]
- Farner NL, Dorner T, Lipsky PE. Molecular mechanisms and selection influence the generation of the human V lambda J lambda repertoire. *J Immunol* 1999;162:2137–2145. [PubMed: 9973488]
- Franciotta D, Salvetti M, Lolli F, Serafini B, Aloisi F. B cells and multiple sclerosis. *Lancet Neurol* 2008;7:852–858. [PubMed: 18703007]
- Freedman MS, Thompson EJ, Deisenhammer F, Giovannoni G, Grimsley G, Keir G, Ohman S, Racke MK, Sharief M, Sindic CJ, Sellebjerg F, Tourtellotte WW. Recommended standard of cerebrospinal fluid analysis in the diagnosis of multiple sclerosis: a consensus statement. *Arch Neurol* 2005;62:865–870. [PubMed: 15956157]
- Frohman EM, Goodin DS, Calabresi PA, Corboy JR, Coyle PK, Filippi M, Frank JA, Galetta SL, Grossman RI, Hawker K, Kachuck NJ, Levin MC, Phillips JT, Racke MK, Rivera VM, Stuart WH. The utility of MRI in suspected MS: report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. *Neurology* 2003;61:602–611. [PubMed: 12963748]
- Frohman EM, Racke MK, Raine CS. Multiple sclerosis--the plaque and its pathogenesis. *N Engl J Med* 2006;354:942–955. [PubMed: 16510748]

- Guddat LW, Herron JN, Edmundson AB. Three-dimensional structure of a human immunoglobulin with a hinge deletion. *Proc Natl Acad Sci U S A* 1993;90:4271–4275. [PubMed: 8483943]
- Hansen A, Dorner T, Lipsky PE. Use of immunoglobulin variable-region genes by normal subjects and patients with systemic lupus erythematosus. *Int Arch Allergy Immunol* 2000;123:36–45. [PubMed: 11014970]
- Harp C, Lee J, Lambracht-Washington D, Cameron E, Olsen G, Frohman E, Racke M, Monson N. Cerebrospinal fluid B cells from multiple sclerosis patients are subject to normal germinal center selection. *J Neuroimmunol* 2007;183:189–199. [PubMed: 17169437]
- Hauser SL, Waubant E, Arnold DL, Vollmer T, Antel J, Fox RJ, Bar-Or A, Panzara M, Sarkar N, Agarwal S, Langer-Gould A, Smith CH. B-cell depletion with rituximab in relapsing-remitting multiple sclerosis. *N Engl J Med* 2008;358:676–688. [PubMed: 18272891]
- Huang C, Stewart AK, Schwartz RS, Stollar BD. Immunoglobulin heavy chain gene expression in peripheral blood B lymphocytes. *J Clin Invest* 1992;89:1331–1343. [PubMed: 1556192]
- Kabat EA, Glusman M, Knaub V. Quantitative Estimation of the Albumin and Gamma Globulin in Normal and Pathologic Cerebrospinal Fluid by Immunochemical Methods. *Am J Med* 1948;4:653–662. [PubMed: 18856761]
- Kabat, EA.; Wu, TT.; Bilofsky, H.; Reid-Miller, M.; Perry, H. Sequences of Proteins of Immunological Interest. United States Department of Health and Human Services; Washington, D.C: 1983.
- Kolln J, Ren HM, Da RR, Zhang Y, Spillner E, Olek M, Hermanowicz N, Hilgenberg LG, Smith MA, van den Noort S, Qin Y. Triosephosphate isomerase- and glyceraldehyde-3-phosphate dehydrogenase-reactive autoantibodies in the cerebrospinal fluid of patients with multiple sclerosis. *J Immunol* 2006;177:5652–5658. [PubMed: 17015754]
- Korteweg T, Tintore M, Uitdehaag B, Rovira A, Frederiksen J, Miller D, Fernando K, Filippi M, Agosta F, Rocca M, Fazekas F, Enzinger C, Matthews P, Parry A, Polman C, Montalban X, Barkhof F. MRI criteria for dissemination in space in patients with clinically isolated syndromes: a multicentre follow-up study. *Lancet Neurol* 2006;5:221–227. [PubMed: 16488377]
- Kraj P, Rao SP, Glas AM, Hardy RR, Milner EC, Silberstein LE. The human heavy chain Ig V region gene repertoire is biased at all stages of B cell ontogeny, including early pre-B cells. *J Immunol* 1997;158:5824–5832. [PubMed: 9190934]
- Lambracht-Washington D, O'Connor KC, Cameron EM, Jowdry A, Ward ES, Frohman E, Racke MK, Monson NL. Antigen specificity of clonally expanded and receptor edited cerebrospinal fluid B cells from patients with relapsing remitting MS. *J Neuroimmunol* 2007;186:164–176. [PubMed: 17451814]
- Leussink VI, Lehmann HC, Meyer zu Horste G, Hartung HP, Stuve O, Kieseier BC. Rituximab induces clinical stabilization in a patient with fulminant multiple sclerosis not responding to natalizumab. Evidence for disease heterogeneity. *J Neurol* 2008;255:1436–1438. [PubMed: 18685916]
- Martin Mdel P, Monson NL. Potential role of humoral immunity in the pathogenesis of multiple sclerosis (MS) and experimental autoimmune encephalomyelitis (EAE). *Front Biosci* 2007;12:2735–2749. [PubMed: 17127276]
- McDonald WI, Compston A, Edan G, Goodkin D, Hartung HP, Lublin FD, McFarland HF, Paty DW, Polman CH, Reingold SC, Sandberg-Wollheim M, Sibley W, Thompson A, van den Noort S, Weinshenker BY, Wolinsky JS. Recommended diagnostic criteria for multiple sclerosis: guidelines from the International Panel on the diagnosis of multiple sclerosis. *Ann Neurol* 2001;50:121–127. [PubMed: 11456302]
- McLaughlin KA, Wucherpfennig KW. B cells and autoantibodies in the pathogenesis of multiple sclerosis and related inflammatory demyelinating diseases. *Adv Immunol* 2008;98:121–149. [PubMed: 18772005]
- Monson NL, Brezinschek HP, Brezinschek RI, Mobley A, Vaughan GK, Frohman EM, Racke MK, Lipsky PE. Receptor revision and atypical mutational characteristics in clonally expanded B cells from the cerebrospinal fluid of recently diagnosed multiple sclerosis patients. *J Neuroimmunol* 2005;158:170–181. [PubMed: 15589051]
- Monson NL, Dorner T, Lipsky PE. Targeting and selection of mutations in human Vlambda rearrangements. *Eur J Immunol* 2000;30:1597–1605. [PubMed: 10898495]

- O’Riordan JI, Thompson AJ, Kingsley DP, MacManus DG, Kendall BE, Rudge P, McDonald WI, Miller DH. The prognostic value of brain MRI in clinically isolated syndromes of the CNS. A 10-year follow-up. *Brain* 1998;121 (Pt 3):495–503. [PubMed: 9549525]
- Owens GP, Bennett JL, Gilden DH, Burgoon MP. The B cell response in multiple sclerosis. *Neurol Res* 2006;28:236–244. [PubMed: 16687047]
- Owens GP, Kraus H, Burgoon MP, Smith-Jensen T, Devlin ME, Gilden DH. Restricted use of VH4 germline segments in an acute multiple sclerosis brain. *Ann Neurol* 1998;43:236–243. [PubMed: 9485065]
- Owens GP, Ritchie AM, Burgoon MP, Williamson RA, Corboy JR, Gilden DH. Single-cell repertoire analysis demonstrates that clonal expansion is a prominent feature of the B cell response in multiple sclerosis cerebrospinal fluid. *J Immunol* 2003;171:2725–2733. [PubMed: 12928426]
- Owens GP, Wings KM, Ritchie AM, Edwards S, Burgoon MP, Lehnhoff L, Nielsen K, Corboy J, Gilden DH, Bennett JL. VH4 gene segments dominate the intrathecal humoral immune response in multiple sclerosis. *J Immunol* 2007;179:6343–6351. [PubMed: 17947712]
- Paolino E, Fainardi E, Ruppi P, Tola MR, Govoni V, Casetta I, Monetti VC, Granieri E, Carreras M. A prospective study on the predictive value of CSF oligoclonal bands and MRI in acute isolated neurological syndromes for subsequent progression to multiple sclerosis. *J Neurol Neurosurg Psychiatry* 1996;60:572–575. [PubMed: 8778266]
- Polman CH, Reingold SC, Edan G, Filippi M, Hartung HP, Kappos L, Lublin FD, Metz LM, McFarland HF, O’Connor PW, Sandberg-Wollheim M, Thompson AJ, Weinshenker BG, Wolinsky JS. Diagnostic criteria for multiple sclerosis: 2005 revisions to the “McDonald Criteria”. *Ann Neurol* 2005;58:840–846. [PubMed: 16283615]
- Qin Y, Duquette P, Zhang Y, Talbot P, Poole R, Antel J. Clonal expansion and somatic hypermutation of V(H) genes of B cells from cerebrospinal fluid in multiple sclerosis. *J Clin Invest* 1998;102:1045–1050. [PubMed: 9727074]
- Ritchie AM, Gilden DH, Williamson RA, Burgoon MP, Yu X, Helm K, Corboy JR, Owens GP. Comparative analysis of the CD19+ and CD138+ cell antibody repertoires in the cerebrospinal fluid of patients with multiple sclerosis. *J Immunol* 2004;173:649–656. [PubMed: 15210828]
- Siritho S, Freedman MS. The prognostic significance of cerebrospinal fluid in multiple sclerosis. *J Neurol Sci.* 2009
- Soderstrom M, Ya-Ping J, Hillert J, Link H. Optic neuritis: prognosis for multiple sclerosis from MRI, CSF, and HLA findings. *Neurology* 1998;50:708–714. [PubMed: 9521261]
- Stuve O, Bennett JL, Hemmer B, Wiendl H, Racke MK, Bar-Or A, Hu W, Zivadinov R, Weber MS, Zamvil SS, Pacheco MF, Menge T, Hartung HP, Kieseier BC, Frohman EM. Pharmacological treatment of early multiple sclerosis. *Drugs* 2008;68:73–83. [PubMed: 18081373]
- Stuve O, Cepok S, Elias B, Saleh A, Hartung HP, Hemmer B, Kieseier BC. Clinical stabilization and effective B-lymphocyte depletion in the cerebrospinal fluid and peripheral blood of a patient with fulminant relapsing-remitting multiple sclerosis. *Arch Neurol* 2005;62:1620–1623. [PubMed: 16216948]
- Tian C, Luskun GK, Dischert KM, Higginbotham JN, Shepherd BE, Crowe JE Jr. Evidence for preferential Ig gene usage and differential TdT and exonuclease activities in human naive and memory B cells. *Mol Immunol* 2007;44:2173–2183. [PubMed: 17196657]
- Tintore M, Rovira A, Martinez MJ, Rio J, Diaz-Villoslada P, Brieva L, Borrás C, Grive E, Capellades J, Montalban X. Isolated demyelinating syndromes: comparison of different MR imaging criteria to predict conversion to clinically definite multiple sclerosis. *AJNR Am J Neuroradiol* 2000;21:702–706. [PubMed: 10782781]
- Tselis A, Perumal J, Caon C, Hreha S, Ching W, Din M, Van Stavern G, Khan O. Treatment of corticosteroid refractory optic neuritis in multiple sclerosis patients with intravenous immunoglobulin. *Eur J Neurol* 2008;15:1163–1167. [PubMed: 18727675]
- Wardemann H, Yurasov S, Schaefer A, Young JW, Meffre E, Nussenzweig MC. Predominant autoantibody production by early human B cell precursors. *Science* 2003;301:1374–1377. [PubMed: 12920303]

Yurasov S, Wardemann H, Hammersen J, Tsuiji M, Meffre E, Pascual V, Nussenzweig MC. Defective B cell tolerance checkpoints in systemic lupus erythematosus. *J Exp Med* 2005;201:703–711. [PubMed: 15738055]

Codon		24	25	26	27	28	29	30	31	31A	31B	32	33	34	35	36	37
IgH4-30.4	Germline DNA	GTC	TCT	GGT	GGC	TCC	ATC	AGC	AGT	GGT	GAT	TAC	TAC	TGG	AGT	TGG	ATC
	M125-391 DNA	GTC	TCT	GGT	GGC	TCC	ATC	AGC	AGT	GGT	GAT	TAC	CAC	TGG	AGT	TGG	ATC
	Germline Protein	V	S	G	G	S	I	S	S	G	D	Y	Y	W	S	W	I
	M125-391 Changes												H				

Codon		38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53
IgH4-30.4	Germline DNA	CGC	CAG	CCC	CCA	GGG	AAG	GGC	CTG	GAG	TGG	ATT	GGG	TAC	ATC	TAT	TAC
	M125-391 DNA	CGC	CAG	CCC	CCA	GGG	AAG	GGC	CTG	GAG	TGG	ATT	GGG	AAC	ATC	AAT	TAT
	Germline Protein	R	Q	P	P	G	K	G	L	E	W	I	G	Y	I	Y	Y
	M125-391 Changes													N		N	

Codon		54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69
IgH4-30.4	Germline DNA	AGT	GGG	AGC	ACC	TAC	TAC	AAC	CCG	TCC	CTC	AAG	AGT	CGA	GTT	ACC	ATA
	M125-391 DNA	AAT	GGG	GGC	GCG	TAC	CAC	AAT	CCG	TCC	CTC	ACG	AAT	CGA	GTT	ATC	ATG
	Germline Protein	S	G	S	T	Y	Y	N	P	S	L	K	S	R	V	T	I
	M125-391 Changes	N		G	A		H					T	N			I	M

Codon		70	71	72	73	74	75	76	77	78	79	80	81	82	82A	82B	82C
IgH4-30.4	Germline DNA	TCA	GTA	GAC	ACG	TCC	AAG	AAC	CAG	TTC	TCC	CTG	AAG	CTG	AGC	TCT	GTG
	M125-391 DNA	TCA	GTA	GAC	ACG	TCC	AAG	AAT	CAC	TTC	TCC	CTG	AAA	CTG	ACC	TCT	GTG
	Germline Protein	S	V	D	T	S	K	N	Q	F	S	L	K	L	S	S	V
	M125-391 Changes								H						T		

Codon		83	84	85	86	87	88	89	90	91	92	93	94
IgH4-30.4	Germline DNA	ACT	GCC	GCA	GAC	ACG	GCC	GTG	TAT	TAC	TGT	GCC	AGA
	M125-391 DNA	ACT	GCC	GCA	GAC	ACG	GCC	GTG	TAT	TAC	TGT	GCC	AGA
	Germline Protein	T	A	A	D	T	A	V	Y	Y	C	A	R
	M125-391 Changes												

Figure 1. Example of VH4 comparison

A VH4-30.4 sequence is listed as the germline configuration (allele 01) and compared to a patient CD19⁺ B cell sequence. The germline protein conversion and the changes made by replacement mutations in the patient sequence are noted. Signature codons are boxed, with the dashed boxes demarcating cold spots, and the solid boxes demarcating hot spots. CDRs as defined by Kabat (Kabat et al., 1983) are shaded.

Figure 2. Signature score in individual MS and CIS patients

Signature scores were generated by calculating Z-scores for the RF values at the 6 codons within the signature (31B, 40, 56, 57, 81 and 89). Individual z-scores at each of the codon positions were compiled to generate the composite signature z-score. MS patient signature scores are shown as black circles (●), CIS patient signature scores that resulted in prediction of CDMS are black squares (■), and CIS patient signature scores that resulted in prediction of unlikely to convert to definite MS are in open squares (□). The average composite signature score in the MSCSF_{VH4} database was 10.9 ± 2.0 (black line) and so any signature score of an individual CIS patient above 6.8 (average - 2 S.D.; threshold shown as red line) was predicted to convert to CDMS. For reference, ONDCSF_{VH4} group signature score was 4.5, and MSPB_{VH4} signature score was 2.0.

Frequency of VH family usage^a

Table 1

B cell source	Expected Frequency		HCPB ^b		mHCPB ^c		MSCSF ^d		CISCSF ^e		CISCSF ^f		ONDCSF ^g	
	By gene	Frequency ^j	CD19 ⁺	CD19 ⁺	CD19 ⁺ IgD ⁺ CD27 ⁺	CD19 ⁺	CD19 ⁺	CD19 ⁺	CD19 ⁺	CD19 ⁺	CD19 ⁺ CD138 ⁺	CD19 ⁺	CD19 ⁺	CD19 ⁺
VH1	21.6		15.5	10.7		27.1 ^{h,i}	9.9	8.9					20.0	
VH2	5.9		2.0	2.0		1.6	5.3 ^h	7.0					3.1	
VH3	43.1		55.2	62.0		32.4 ^{h,i}	56.3	47.5					46.2 ^j	
VH4	21.6		21.8	16.6		34.3 ^{h,i}	26.2 ^j	35.0					23.1	
Total Sequences	51		348	205		373	302	528					65	
Number of Donors	0		2	6		11	10	11					3	

Abbreviations: VH, variable heavy; HCPB, healthy control peripheral blood; mHCPB, memory healthy control peripheral blood; MSCSF, multiple sclerosis cerebrospinal fluid; CISCSF, clinically isolated syndrome cerebrospinal fluid; ONDCSF, other neurological disease cerebrospinal fluid.

- ^aValues provided in percent
- ^bThe HCPB group includes CD19⁺ B cell antibody sequences from healthy controls BF1 (n=67) and BF2 (n=281).
- ^cThe mHCPB group includes CD19⁺ B cell antibody sequences from healthy controls BF1 (n=18) and BF2 (n=105) with 4 or more mutations (less than 98% homology to germline) and IgD⁺ CD27⁺ B cell antibody sequences from healthy controls HA (n=9), HB (n=44), HC (n=26), and HE (n=3) (Tian et al., 2007).
- ^dThe MSCSF group includes CD19⁺ sequences from MS patients M125 (n=101), M199 (n=19), M354 (n=6), M368 (n=49), M376 (n=8), M484 (n=9), M522 (n=71), M584 (n=85), M875 (n=21), M217 (n=1), and M887 (n=3).
- ^eThe CISCSF group includes CD19⁺ sequences from CIS patients CIS132 (n=19), CIS429 (n=57), CIS3-1 (n=24), ON3-1 (n=23), ON3-3 (n=39), ON3-4 (n=28), ON3-5 (n=35), ON4-7 (n=17), ON4-10 (n=31), and ON5-2 (n=29).
- ^fIn addition to those CD19⁺ sequences listed in ^e, this group includes CD138⁺ sequences from CIS patients CIS3-1 (n=76), ON3-1 (n=45), ON3-3 (n=12), ON3-5 (n=44), ON4-7 (n=20), ON4-8 (n=17), ON5-2 (n=12).
- ^gThe ONDCSF group includes CD19⁺ B cell antibody sequences from OND patients OND341 (n=32), OND758 (n=19), and OND1116 (n=14).
- ^hSignificantly different from HCPB frequency
- ⁱSignificantly different from mHCPB frequency
- ^jExpected frequency from (Cook and Tomlinson, 1995)

Table 2

Percentage of Replacement Mutations in Each Signature Codon

Codon	Location ^b	MSCSF _{VH4}			HCPB _{VH4}			mHCPB _{VH4}		
		RF	RF	p-value ^d	Fold Increase	RF	p-value ^d	Fold Increase	RF	p-value ^d
31B ^e	CDR1	3.5 ⁵	0.5 ⁵	0.001	7.0	0.8	0.001	4.4	0.8	0.001
32	CDR1	2.3	1.5	0.05	1.5	2.1	0.05	1.1	2.1	NS
40 ^e	FR2	2.7	1.0	0.001	2.7	1.1	0.001	2.5	1.1	0.001
56 ^{d,e}	CDR2	5.5	3.0	0.001	1.8	3.2	0.001	1.7	3.2	0.001
57 ^e	CDR2	2.0	1.0	0.005	2.0	0.5	0.005	3.7	0.5	0.001
60	CDR2	2.4	1.5	0.05	1.6	1.1	0.05	2.2	1.1	0.001
81 ^e	FR3	4.7	3.0	0.005	1.5	3.7	0.005	1.3	3.7	NS
89 ^e	FR3	2.0	1.0	0.005	2.0	1.3	0.005	1.5	1.3	NS
Hotspot Total ^c		25.0	12.6	0.001	2.0	13.8	0.001	1.8	13.8	0.001
30	FR1	2.0	4.0	0.005	0.5	2.9	0.005	0.7	2.9	NS
43	FR2	0.9	2.0	0.025	0.5	1.3	0.025	0.7	1.3	NS
77	FR3	1.5	2.5	0.05	0.6	1.6	0.05	0.9	1.6	NS
82	FR3	0.7	2.5	0.001	0.3	1.6	0.001	0.5	1.6	0.05
Coldspot Total ^c		5.1	8.5	0.001	0.5	7.4	0.001	0.7	7.4	0.01

Abbreviations in table: CDR, complementary determining region; FR, framework; RF, replacement frequency; MSCSF, multiple sclerosis cerebrospinal fluid; HCPB, healthy control peripheral blood; mHCPB, memory HCPB; NS, not significant

^aPreviously published replacement hotspot (Dorner et al., 1997; Dorner et al., 1998a)

^bAs defined by Kabat (Kabat et al., 1983)

^c“Hotspot Total” is the total RF within codons 31B, 32, 40, 56, 57, 60, 81 and 89. “Coldspot Total” is the total RF within codons 30, 43, 77 and 82. 199, 337 and 965 replacements respectively were included in this analysis for HCPB_{VH4}, mHCPB_{VH4}, and MSCSF_{VH4}.

^dComparing HCPB_{VH4} or mHCPB_{VH4} to MSCSF_{VH4} RFs at each codon position using χ^2 goodness-of-fit where expected frequency is the RF calculated in HCPB_{VH4} and mHCPB_{VH4} respectively.

^eCodon used in the calculation of signature score

CIS Patient Summary and Signature Score Predictions

Table 3

Subject No. ^a	Time to LP ^b	MRI Brain Lesions	OCB	CD19/CD138 ^c	CD19 VH4 bias ^d	CD138 VH4 bias ^d	Signature Score ^e	Prediction based on signature score	Definite MS	Time to MS Diagnosis ^f
CIS132	1	GD ⁺	Yes	19/NA	No	NA	12.1	CDMS	Clinical	18
CIS429	1	GD ⁺	Yes	56/NA	No	NA	15.0	CDMS	Clinical	3
CIS3-1	2	WML	Yes	24/76	Yes	Yes	15.5	CDMS	MRI	3
ON3-3	4	GD ⁺	Yes	39/13	No	No	11.3	CDMS	Clinical	3
ON3-5	1.75	GD ⁺	Yes	35/44	Yes	Yes	12.8	CDMS	Clinical	2
ON4-7	3	None ^g	Yes	17/20	No	Yes	10.2	CDMS	Clinical	5
ON4-8	1.5	WML	Yes	NA/18	NA	Yes	9.6	CDMS	Clinical	5
ON5-2	1	GD ⁺	Yes	29/12	Yes	Yes	7.9	CDMS	Clinical	3
ON3-1	10	WML	Yes	23/45	No	No	6.4	No	-	NA
ON4-10	1.25	WML	No	31/NA	No	NA	2.2	No	-	NA
ON3-4	1.5	None ^g	No	28/NA	No	NA	11.3	CDMS	-	NA

Abbreviations in table: CIS, Clinically Isolated Syndrome; ON, Optic Neuritis; LP, lumbar puncture; GD, gadolinium enhancing lesion positive; WML, white matter lesions by T2; OCB, oligoclonal bands; CDMS, clinically definite MS; NA, not applicable

^aCIS132 and CIS429 were generated at UTSWMC; the remaining patient CSF B cell repertoires were generated at UCD.

^bmonths from first demyelinating event to LP

^cValues given are number of unique sequences in CD19 repertoire(CD138 repertoire; family usage can be found in (Bennett et al., 2008; Harp et al., 2007); CIS429 was a 62 y.o. male first presenting with optic neuritis and the repertoire had 2% VH1 usage, 0% VH2, 77% VH3, and 19% VH4.

^dBias was considered significantly different from random frequency (Cook and Tomlinson, 1995) or expected frequency in HCPB (Brezinschek et al., 1997; Brezinschek et al., 1998; Dorner et al., 1997; Dorner et al., 1998a; Dorner et al., 1998b; Dorner et al., 1998c; Famer et al., 1999; Hansen et al., 2000; Monson et al., 2000). VH2 bias was also observed in CD19 repertoires from ON4-7, and CD138 repertoires from ON3-3, ON3-5 and ON4-7.

^eSignature score was calculated as outlined in materials and methods, and uses both CD19 and CD138 sequences. The average score among MS patients is 10.9 ±2.0.

^fmonths from first demyelinating event to MS diagnosis.

^gOne spinal cord lesion was observed by T2 weighted MRI