Published in final edited form as:

Synlett. 2009 June 1; 2009(10): 1680–1684.

Bifunctional Asymmetric Catalysis with Hydrogen Chloride: Enantioselective Ring-Opening of Aziridines Catalyzed by a Phosphinothiourea

Tsuyoshi Mita and Eric N. Jacobsen*

Department of Chemistry and Chemical Biology, Harvard University, Cambridge, Massachusetts 02138, USA

Abstract

Ring-opening of aziridines with hydrogen chloride to form β -chloroamine derivatives is catalyzed by a chiral phosphinothiourea derivative in high yields and with high enantioselectivities. On the basis of ^{31}P NMR studies, activation of HCl appears to proceed via quantitative protonation of the catalyst to afford a phosphonium chloride complex.

Keywords

asymmetric catalysis; halogenation; ring-opening; carbocations; aziridines

Hydrogen chloride (HCl) is a commodity chemical used in many industrial processes.¹ HCl is also a convenient and inexpensive reagent for use in laboratory-scale chemical synthesis, such as for the hydrochlorination of carbon-carbon double² and triple³ bonds. However its utility in catalytic enantioselective transformations has thus far been limited.⁴ Indeed, few of the common chiral catalysts are likely to be compatible with HCl given its extremely high acidity.⁵

Over the past several years, chiral thioureas have emerged as a useful class of catalysts for a broad range of asymmetric transformations.⁶ We discovered recently that chiral thiourea derivatives catalyze enantioselective additions to *N*-acyliminium/chloride ion pairs^{7a–c} and α -chloroethers.^{7d} These observations have led to the proposal that thiourea catalysts engage these weakly basic electrophiles by an anion abstraction mechanism that involves a thiourea-chloride hydrogen-bonding interaction.^{7b} We have found that urea and thiourea catalysts are stable to strong mineral acids,⁸ suggesting that these catalysts might be amenable to productive activation of HCl. In particular, we considered the possibility that a bifunctional chiral catalyst⁹ containing a basic group could undergo protonation by HCl with binding of the chloride counterion to the thiourea (Figure 1), thereby activating the molecule to additions to polar organic electrophiles. Although many Lewis or Brønsted basic functional groups are expected to be protonated by HCl, the weak basicity of phosphines ($\text{Ph}_2\text{CyP}^+\text{H}$: $\text{pK}_a = 5.1$ in H_2O),¹⁰ suggested that chiral phosphinothiourea catalysts such as **1**, originally developed in our laboratory for the [3+2] cycloaddition of allenes with imines,¹¹ might be particularly effective in the context of productive HCl activation. Herein, we report the use of chiral phosphinothioureas to catalyze the catalytic enantioselective addition of HCl to *N*-acylaziridines.¹²

Enantioselective desymmetrization of meso epoxides with various chloride sources is a well-studied method for the preparation of optically active chlorohydrins, and many examples using stoichiometric promoters¹³ as well as chiral catalysts¹⁴ have both been reported. Interestingly, the direct use of HCl for the enantioselective variant of this transformation has not been reported.¹⁵ In pioneering studies, Denmark reported that Lewis basic, chiral phosphoramides catalyze enantioselective opening of meso epoxides with SiCl₄.^{14a} Under the reaction conditions developed by Denmark, HCl mediates competitive, unselective epoxide ring-opening, and careful distillation of SiCl₄ was necessary to ensure that reproducible enantioselectivity was obtained. HCl is also very reactive towards aziridines in the absence of any promoters,¹⁶ suggesting that a control over background reaction is necessary for achieving efficient asymmetric catalysis.

We began our study by investigating different thiourea catalysts using benzoyl protected aziridine **2a** as a model substrate (Figure 2).¹⁷ In the absence of a chiral catalyst, hydrochlorination of **2a** is complete within 30 min, even under relatively dilute reaction conditions and low temperatures (0.025 M, -78 °C), indicating that acceleration of a catalytic pathway and/or suppression of background reactivity would be necessary in order to achieve high enantioselectivities. 1,2-Diaminocyclohexane-derived catalysts **1a–1d** (20 mol %) promoted formation of product **3a** with very modest enantioselectivities (3–29% ee). Slow addition of HCl (0.25 M in Et₂O) over 110 min, led only to minor improvements (up to 40% ee for both catalysts **1a** and **1b**), suggesting that background reactivity was only partially responsible for the low ee's. While phosphine oxide catalyst **1e** as well as thioureas **1g** and **1h** also led to product formation with poor enantioselectivity, phosphinothiourea catalyst **1f** promoted hydrochlorination of **2a** within 10 min in 60% ee.¹⁸ In this case, neither slow addition of HCl nor increased catalyst loading had any beneficial effect on enantioselectivity, suggesting that the catalytic pathway was significantly more rapid than the background reaction. This hypothesis was confirmed by *in situ* IR spectroscopic experiments with cycloheptene-derived aziridine **2c**: in these studies, the initial reaction rate of hydrochlorination with catalyst **1f** (20 mol% at -40 °C) was measured to be 140 times greater than in the absence of catalyst.

The α -amino acid component of catalyst **1** was varied systematically in an effort to improve the enantioselectivity in the model reaction (Figure 3). It was found that ee's were responsive to increasing steric demand of the α -substituent (catalysts **1i–1f**). Accordingly, highly hindered unnatural amino acids were prepared, taking advantage of recently-developed thiourea-catalyzed asymmetric Strecker methodology,¹⁹ and incorporated into the new set of phosphinothiourea derivatives **1m–1p**. A measurable improvement in ee was observed with these more hindered catalysts. The amino acid used for the preparation of catalyst **1m** is readily accessible in optically pure form,²⁰ so **1m** was selected as the preferred catalyst for further study.

Absolute concentration has proven to be an important parameter in thiourea-catalyzed reactions, with improvements in both enantioselectivity and rate resulting in certain cases from increased dilution.^{2b,21} This proved to be the case in the aziridine hydrochlorination reaction as well, with highest enantioselectivities obtained under highly dilute conditions (0.0025 M, compare entries 1–4, Table 1). However, further dilution led to significant decreases in reaction rate, necessitating higher reaction temperatures in order to achieve high conversion at the expense of ee (entries 5–6). Given that the racemic uncatalyzed reaction is not competitive with the catalytic pathway (entry 4),²² it appears that the increased enantioselectivity obtained at low concentrations may be ascribed to the existence competing catalyzed pathways of varying kinetic order in catalyst and/or reactants.

The optimal conditions were successfully applied to a variety of aziridines to provide β -chloroamine products with high yields and good enantioselectivities with 10–20 mol% catalyst

loadings (Table 2). Cyclooctene-derived substrate **2d** (entry 4) was unreactive under highly dilute conditions (0.0025 M), but efficient hydrochlorination of this challenging substrate could be achieved at higher concentrations (0.025 M). The scope of enantioselective hydrochlorinations catalyzed by **1m** is noteworthy given that Lewis base-catalyzed variants of this reaction, which generally afford only modest ee's with cyclic substrates.¹⁴

cis-Stilbene-derived aziridine **2g** also underwent the ring-opening in the presence of phosphinothiourea catalyst in reactions carried at 0.025 M (Table 3). Uncatalyzed openings of **2g** lead to mixtures of diastereomeric products, with the ratio dependent on the chloride source used: SiCl₄ leads to selective formation of *syn*-**3g**,²³ the result of retentive ring-opening (entry 1). In contrast, HCl provides 1:1 mixture of *anti*- and *syn*-products (entry 2). Interestingly, *anti*-**3g**²³ was obtained in high diastereo- and enantioselectivity (up to 90% ee) in reactions promoted by chiral phosphinothiourea catalysts **1f** and **1m** (entries 4 and 5). The observation of predominantly invertive ring-opening suggests an S_N2-type pathway in the catalytic reaction. Other chiral thiourea derivatives **1a** and **1h** provided lower levels of selectivity (entries 6 and 7), indicating that a bifunctional phosphinothiourea catalyst was important for this substrate as well.

The absolute configuration of **3a** (71% ee) prepared using **1f** was determined by conversion to the known *cis*-amino alcohol **5**²⁴ in two steps (Scheme 1).

In order to glean insight into the mechanism of the phosphinothiourea-catalyzed additions of HCl to aziridines, we undertook ³¹P NMR studies of the interaction of catalyst **1f** with HCl. An anhydrous, ethereal solution of **1f** was cooled to -78 °C, and excess HCl in Et₂O (10 equiv) was added dropwise via syringe. The solution became cloudy immediately, and was stirred for 30 min at -78 °C. Solvent removal by gradual evaporation under reduced pressure at low temperature (-78 °C to ca. -30 °C) afforded a white solid. Analysis of this material as a CDCl₃ solution by ³¹P NMR spectroscopy revealed a single resonance (δ = 4.6 ppm, J_{P-H} = 508 Hz) distinct from that of **1f** (δ = -8.0 ppm). The chemical shift and P-H coupling constant are similar to those of previously characterized protonated phosphines.^{25,26} Enantioselective hydrochlorination of **2a** (0.0025 M) using the solid, protonated species without added HCl gave the ring-opened product **3a** quantitatively within 1 h in 70% ee,²⁷ comparable to the enantioselectivity obtained under catalytic conditions (77% ee). On the basis of these results, it appears that catalytic hydrochlorinations of benzoyl aziridines such as **2a** catalyzed by phosphinothioureas may involve initial heterolysis of HCl to afford a phosphonium chloride complex as depicted in Figure 1.

A proposed catalytic cycle for the ring-opening of aziridine **2a** is provided in Figure 4. Aliphatic aziridines **2a–2f** undergo ring-opening exclusively with inversion of stereochemistry to provide *trans*-addition products **3a–3f**. Rates of invertive ring-openings of epoxides and aziridines derived from medium-ring cyclic alkenes are typically highly sensitive to the steric environment defined by the ring.²⁸ This is observed in the present case as well, with reactions of five- and six-membered ring substrates **2a** and **2b** being considerably more rapid than those of seven- and eight-membered substrates **2c** and **2d**. The reactivity profiles and the observation of exclusive formation of *trans* products points to an S_N2-type mechanism (Figure 4). We propose that HCl-**1** complex **6** is generated in the first step, and that the phosphonium ion serves to activate substrate **2a** by a general-acid mechanism. Chloride addition is suggested to occur from the thiourea-bound complex, resulting in a highly-ordered enantiodetermining step. Although stilbene-derived aziridines are generally more prone to S_N1-type pathways, the observation of high enantioselectivity and in the predominant invertive pathway suggests a common S_N2 mechanism for all the aziridines described in this study.

In summary, we have developed a catalytic enantioselective ring-opening reaction of meso aziridines with HCl catalyzed by phosphinothioureas. A broad range of cyclic as well as acyclic substrates undergo clean addition to afford the corresponding β -chlorobenzamide derivatives in high yields and with high enantioselectivities. Investigations of other classes of acidic nucleophiles (HX) and potential electrophiles, as well as more detailed mechanistic studies are underway.

Representative Procedure for Catalytic Enantioselective Ring-Opening Reaction with HCl

To a mixture of catalyst **1m** (3.0 mg, 0.005 mmol, 10 mol %) and aziridine **2a** (10.1 mg, 0.050 mmol) in Et₂O (20 mL, 0.0025 M), HCl (1 M stock solution in Et₂O, 60 μ L, 0.06 mmol, 1.2 equiv) was added at -78 °C. After 4 h, the solvent was evaporated under reduced pressure, and the residue was purified by automated column chromatography (silica gel, hexane-AcOEt, 19:1 to 3:2) to afford (1*R*,2*R*)-*N*-(2-chlorocyclohexyl)benzamide (**3a**) (11.6 mg, 0.0488 mmol) in 98% yield as colorless solid. The enantiomeric excess of the product was determined by HPLC analysis to be 83% ee.

IR (neat): 3230, 3059, 2950, 2859, 1635, 1575, 1452, 1330, 1125 cm⁻¹; ¹H NMR (CDCl₃): δ = 7.78 (d, *J* = 7.6 Hz, 2H), 7.50 (t, *J* = 7.6 Hz, 1H), 7.44 (t, *J* = 7.6 Hz, 2H), 6.18 (brd, *J* = 4.5 Hz, 1H), 4.06 (ddd, *J* = 3.8, 9.3, 17.5 Hz, 1H), 3.88 (dt, *J* = 3.8, 10.5 Hz, 1H), 2.34-2.30 (m, 2H), 1.84-1.74 (m, 3H), 1.52-1.44 (m, 1H), 1.40-1.31 (m, 2H); ¹³C NMR (CDCl₃): δ = 167.2, 134.7, 131.4, 128.5, 127.0, 62.5, 55.6, 36.1, 32.8, 25.4, 24.2; HRMS (ESI): *m/z* calcd for C₁₃H₁₆ClNNaO [M+Na⁺]: 260.0813, 262.0783 (chlorine isotope). Found: 260.0810, 262.0782 (chlorine isotope); [α]_D²³ -54.1 (*c* = 0.58, CHCl₃) (83% ee); HPLC (Daicel Chiralpak AD-H, 2-propanol/hexane 1/10, flow rate 1.0 mL/min, detection at both 210 nm and 254 nm): *t*_R 13.8 min (major) and 23.4 min (minor).

Acknowledgments

This research was supported by the NIH (GM-43214). T. M. thanks the Japan Society for the Promotion of Science (JSPS) for scholarship support. We thank Mr. Stephan J. Zuend for helpful discussion.

References and Notes

1. Lacson, J.; Sakuma, Y.; Schlag, S. Chemical Economics Handbook: Hydrochloric Acid. SRI International; Menlo Park: CA: 2006. p. 733.4000A-733.4000F.
2. For the kinetics and mechanism of hydrochlorination of olefins, see: (a) Porker Y, Stevens KD, Champoux JJ. J Am Chem Soc 1969;91:4199. (b) Porker Y, Stevens KD. J Am Chem Soc 1969;91:4205.
3. For the synthesis of vinyl chloride from acetylene, see: Patat F, Weidlich P. Helv Chem Acta 1949;32:783.
4. The kinetic resolutions of terminal epoxides with HCl catalyzed by bimetallic chiral salen complexes were only reported. (a) Thakur SS, Li W, Kim SJ, Kim GJ. Tetrahedron Lett 2005;46:2263. (b) Thakur SS, Li W, Shin CK, Kim GJ. Chirality 2006;18:37. (c) Thakur SS, Chen SW, Li W, Shin CK, Kim SJ, Koo YM, Kim GJ. J Organomet Chem 2006;691:1862.
5. HCl: *pK*_a = 1.8 in DMSO and -8.0 in H₂O.
6. For recent reviews, see: (a) Taylor MS, Jacobsen EN. Angew Chem Int Ed 2006;45:1520. (b) Doyle AG, Jacobsen EN. Chem Rev 2007;107:5713. [PubMed: 18072808]
7. (a) Taylor MS, Jacobsen EN. J Am Chem Soc 2004;126:10558. [PubMed: 15327311] (b) Raheem IT, Thiara PS, Peterson EA, Jacobsen EN. J Am Chem Soc 2007;129:13404. [PubMed: 17941641] (c) Raheem IT, Thiara PS, Jacobsen EN. Org Lett 2008;10:1577. [PubMed: 18341346] (d) Reisman SE, Doyle AG, Jacobsen EN. J Am Chem Soc 2008;130:7198. [PubMed: 18479086]

8. For example, catalyst 1g can be used effectively in reactions catalyzed by triflic acid ($pK_a = 0.3$ in DMSO and -14 in H_2O , respectively). Woll MG, Xu H, Tao Y, Zuend SJ, Jacobsen EN. unpublished results.
9. Kanai M, Kato N, Ichikawa E, Shibasaki M. *Synlett* 2005:1491.
10. Rahman MM, Liu HY, Eriks K, Prock A, Giering WP. *Organometallics* 1989;8:1.
11. (a) Fang YQ, Jacobsen EN. *J Am Chem Soc* 2008;130:5660. [PubMed: 18393504] After our publication, similar type of catalyst was reported by a different research group. (b) Yuan K, Zhang L, Song HL, Hu Y, Wu XY. *Tetrahedron Lett* 2008;49:6262.
12. For examples of catalytic enantioselective ring-opening reactions of meso aziridines, see: (a) Zhang ZD, Scheffold R. *Helv Chim Acta* 1993;76:2602. (b) Hayashi M, Ono K, Hoshimi H, Oguni N. *Tetrahedron* 1996;52:7817. (c) Muller P, Nury P. *Org Lett* 1999;1:439. (d) Li Z, Fernández M, Jacobsen EN. *Org Lett* 1999;1:1611. [PubMed: 10836024] (e) Mita T, Fujimori I, Wada R, Wen J, Kanai M, Shibasaki M. *J Am Chem Soc* 2005;127:11252. [PubMed: 16089448] (f) Fukuda Y, Mita T, Fukuda N, Kanai M, Shibasaki M. *J Am Chem Soc* 2006;128:6312. [PubMed: 16683784] (g) Fujimori I, Mita T, Maki K, Shiro M, Sato A, Furusho S, Kanai M, Shibasaki M. *Tetrahedron* 2007;63:5820. (h) Arai K, Lucarini S, Salter MM, Ohta K, Yamashita Y, Kobayashi S. *J Am Chem Soc* 2007;129:8103. [PubMed: 17567008] (i) Rowland EB, Rowland GB, Rivera-O E, Antilla JC. *J Am Chem Soc* 2007;129:12084. [PubMed: 17877355] (j) Wu B, Gallucci JC, Parquette JR, Rajan-Babu TV. *Angew Chem Int Ed* 2009;48:1126.
13. (a) Joshi NN, Srebnik M, Brown HC. *J Am Chem Soc* 1988;110:6246. (b) Naruse Y, Esaki T, Yamamoto H. *Tetrahedron* 1988;44:4747.
14. (a) Denmark SE, Barsanti PA, Wong KT, Stavenger RA. *J Org Chem* 1998;63:2428. [PubMed: 11672098] (b) Bruns S, Haufe G. *Tetrahedron: Asymmetry* 1999;10:1563. (c) Tao B, Lo MMC, Fu GC. *J Am Chem Soc* 2001;123:353. [PubMed: 11456531] (d) Nakajima M, Saito M, Uemura M, Hashimoto S. *Tetrahedron Lett* 2002;43:8827. (e) Peak SH, Shim SC, Cho CS, Kim TJ. *Synlett* 2003:849. (f) Tokuoka E, Kotani S, Matsunaga H, Ishizuka T, Hashimoto S, Nakajima M. *Tetrahedron: Asymmetry* 2005;16:2391. (g) Denmark SE, Barsanti PA, Beutner GL, Wilson TW. *Adv Synth Catal* 2007;349:567. (h) Takenaka N, Sarangthem RS, Captain B. *Angew Chem Int Ed* 2008;47:9708. (i) Chelucci G, Baldino S, Pinna GA, Benaglia M, Buffa L, Guizzetti S. *Tetrahedron* 2008;64:7574. For chiral bromohydrine synthesis, see: (j) Nugent WA. *J Am Chem Soc* 1998;120:7139.
15. Ring-openings of cyclooctene oxide and *cis*-stilbene oxide with HCl in the presence of phosphinothiourea 1f proceeded to afford corresponding *trans*-chlorohydrines in only 22% ee and 34% ee, respectively.
16. For example, see: Crotti P, Favero L, Gardelli C, Macchia F, Pineschi M. *J Org Chem* 1995;60:2514.
17. Aziridines having different protecting groups such as benzyl, tosyl, benzylcarbamoyl, and also unprotected aziridines gave almost racemic compounds.
18. Diastereomeric phosphinothiourea of 1f previously effective for [3+2] cyclization^{11a} was not promising for this reaction (12% ee).
19. Zuend SJ, Coughlin MP, Lalonde MP, Jacobsen EN. manuscript in preparation.
20. *N*-Boc- α -cyclohexylmethyl glycine was fine solid so that optically active crystal could be easily obtained after the recrystallization from the reaction crude (ca. 90% ee) which was prepared through the enantioselective Strecker reaction and hydrolysis followed by boc-protection. Other boc-protected amino acids are amorphous which need the crystallization after the salt formation with *tert*-butyl amine.
21. Klausen RS, Jacobsen EN. *Org Lett* 2009;11:887. [PubMed: 19178157]
22. Kinetic studies of the uncatalyzed hydrochlorination of 2b and 2c using *in situ* IR spectrometry (Et_2O at $-40^\circ C$) revealed a 1.5-order dependence on [HCl].
23. Authentic samples of anti-3g and syn-3g were prepared through the benzylation of anti- and syn-2,3-diphenyl-2-chloroethylamine hydrochloride. See: Hassner A, Burke SS. *Tetrahedron* 1974;30:2613.
24. Schaus SE, Larrow JF, Jacobsen EN. *J Org Chem* 1997;62:4197. (1R, 2S)-2-amino-cyclohexanol: $[\alpha]_D^{26} -27.9$ ($c = 1.10$, $EtOH$: >99% ee).
25. Dillon KB, Waddington TC, Younger D. *J Chem Soc, Dalton Trans* 1975:790.
26. Pestovsky O, Shuff A, Bakac A. *Organometallic* 2006;25:2894.

27. $\text{PPh}_3\cdot\text{HBr}$ and $\text{PPh}_3\cdot\text{HCl}$ promoted hydrohalogenation of epoxides, see: Afonso CAM, Vieira NML, Mother-well WB. *Synlett* 2000:382.
28. For example, see: Jacobsen EN. *Acc Chem Res* 2000;33:421. [PubMed: 10891060]

Figure 1.
Proposed activation mode of HCl by catalyst **1**.

Figure 2.
Screen of thiourea catalysts

Figure 3.
Optimization of the amino acid component of the catalyst.

Figure 4.
Proposed catalytic cycle for aliphatic aziridine **2a**.

Scheme 1.
Determination of absolute configuration.

Table 1

Concentration Effect

Entry	Conc. (M)	Temp (°C)	Time (h)	Conv. (%)	Ee (%) ^a	
1	0.1	-78	<30 min	>99	35	
2	0.025	-78	<30 min	>99	65	
3	0.0025	-78	4	>99	81	
4 ^b	0.0025	-78	2	<5	–	
5	0.0010	-78 to -50	3/62	99	65	
6	0.0005	-78 to -50	2/45	66	59	

^aEe's were determined by chiral HPLC analysis.

^bBackground reaction carried out in the absence of catalyst **1m**.

Table 2

Substrate Scope

Temp (°C)	Time (h)	Isolated Yield (%) ^d	Ee (%) ^b
-78	4	97	83
-78	6	99	78
-78 to -50	19	96	90

Product

Entry	Temp (°C)	Time (h)	Isolated Yield (%) ^a	Ee (%) ^b
4 ^{c,d}	-78 to -50	84	91	84
5 ^c	-78 to -50	13	94	92

Product

6

-78 1 96 70

^a Isolated yield.

^b Determined by chiral HPLC analysis.

^c 20 mol % of catalyst was used.

^d Reaction was conducted on 0.025 M.

Table 3

Hydrochlorination of *cis*-Stilbene Derived Aziridine **2g**

Entry	Catalyst	Temp (°C)	Time (h)	Conv (%)	Ratio (<i>anti:syn</i>) ^a	Ee of <i>anti</i> (%) ^b
1 ^c	-	-30	1	>99	18:82	-
2	-	-78 to r.t.	19	>99	48:52	-
3	1f	r.t.	2	>99	74:26	50
4	1f	-78 to -30	20	92	94:6	90 ^d
5 ^e	1m	-78 to -30	46	>99 (85 ^f)	93:7	-89 ^d
6	1a	-30	67	95	62:38	39
7	1h	-30	24	ca. 80	39:61	-30

^a Diastereomeric ratios were determined by integration of ¹H NMR resonances.

^b Ee's were determined by chiral HPLC analysis.

^c SiCl₄ in CH₂Cl₂ was used as the chloride source.

^d *Syn-3g* is unstable to silica gel column chromatography and undergoes cyclization to the corresponding *trans*-oxazoline. The ee was determined to be 21% with catalyst **1f** and -21% with catalyst **1m**.

^e [*cis-2g*]₀ = 0.01 M. No reaction was observed under more dilute conditions.

^f Isolated yield of *anti-3g*.