Published in final edited form as:

Curr Atheroscler Rep. 2009 March ; 11(2): 131–138.

Monocyte Chemoattractant Protein (MCP-1)/CCL2 as a biomarker in Acute Coronary Syndromes

Carlos Gonzalez-Quesada, MD and Nikolaos G Frangogiannis, MD

Section of Cardiovascular Sciences, Baylor College of Medicine, Houston, TX.

Abstract

The CC chemokine Monocyte chemoattractant protein (MCP)-1/CCL2 is involved in the formation, progression, and destabilization of atheromatous plaques and plays an essential role in post-infarction remodeling. These properties generated significant interest in the potential significance of MCP-1 as a biomarker in acute coronary syndromes (ACS). Emerging evidence suggests that MCP-1 plasma levels have prognostic value in the acute and chronic phase following ACS, providing information independent of standard clinical variables. The mechanisms responsible for adverse prognosis in patients with elevated plasma MCP-1 following ACS remain unknown. High plasma MCP-1 levels may reflect a higher burden of atherosclerotic disease, may exert prothrombotic effects resulting in recurrent coronary events, or may identify patients who mount a more intense cardiac inflammatory reaction following a coronary event, resulting in enhanced adverse remodeling. Beyond its prognostic significance, the MCP-1 axis may be an attractive target for therapy in patients with ACS.

Introduction

Recent advances in cardiovascular research have provided new insight into the processes involved in the pathogenesis of acute coronary syndromes (ACS). Extensive evidence implicates inflammatory mediators in the development of atherogenesis. In addition, inflammation appears to play a fundamental role in the pathogenesis of plaque rupture. Earlier studies suggested that local inflammation results in rupture of a single vulnerable plaque leading to coronary thrombosis. More recent evidence supported the emerging concept of the “vulnerable patient”, suggesting that unstable coronary syndromes are associated with widespread inflammatory activation of the coronary tree [1], [2]. Thus, both researchers and clinicians have turned to biochemical markers of inflammation as possible noninvasive indicators of the risk of recurrent cardiovascular events following an ACS, and as predictors of the success of therapeutic interventions [3].

High-sensitivity C-Reactive Protein (hsCRP), the prototypical inflammatory biomarker, seems to provide important prognostic information in ACS. In patients with unstable angina or non-Q-wave myocardial infarction enrolled in the Thrombolysis in Myocardial Infarction (TIMI) 11A trial, elevated CRP at presentation correlated with 14-day mortality even in patients with a negative rapid cardiac troponin T assay [4]. In addition, persistent elevation of hsCRP 30 days following an ACS was associated with a significantly increased 2-year mortality rate in the Aggrastat-to-Zocor (A to Z) trial [5]. Patients treated with more aggressive statin therapy were more likely to achieve lower levels of hsCRP. However, although CRP is a reliable marker of inflammation, several other immune and inflammatory mediators are critically involved in the pathogenesis of ACS and may serve as new biomarkers providing additional diagnostic

and prognostic information. These novel molecular markers may serve as a window into the underlying pathophysiologic mechanisms of coronary atherothrombotic disease and may guide a more individualized treatment approach.

The chemokines

The chemokines are a family of small chemotactic cytokines with molecular weights in the range of 8-14kDa and a strikingly similar tertiary structure. There are approximately fifty human chemokines, which can be classified into CC, CXC, C, or CX3C subfamilies based on the number of amino acids between the first two of the four conserved cysteine residues that characterize chemokine structure. From a functional standpoint chemokines can be divided broadly into two categories: homeostatic chemokines are constitutively expressed in certain tissues and may be responsible for basal leukocyte trafficking and formation of the fundamental architecture of lymphoid organs, whereas inducible chemokines are strongly upregulated by inflammatory or immune stimuli and actively participate in the inflammatory reactions by inducing leukocyte recruitment [6]. Although this generalization has been challenged by studies demonstrating that certain chemokines, felt to be homeostatic (such as SDF-1) [7], are inducible upon immune activation, it offers valuable insight into the role of members of the chemokine family in pathological states. Extensive evidence suggests that both CC and CXC chemokines are induced in human atherosclerotic lesions and may play a role in atherogenesis and plaque destabilization.

Monocyte Chemoattractant Protein (MCP)-1/CCL2: an essential mediator in the pathogenesis of atherosclerosis

MCP-1, the most thoroughly characterized CC chemokine, was identified as a monocyte-specific chemoattractant that was later demonstrated to attract T lymphocytes and NK cells, but not neutrophils. Enhanced expression of MCP-1 was demonstrated in a variety of pathologic conditions, associated with inflammation and mononuclear cell infiltration. Extensive experimental evidence suggests that MCP-1 is highly expressed in atherosclerotic plaques [8] and mediates macrophage recruitment in the atheromatous lesion (Figure 1). Fatty streaks, the hallmark of early atherosclerotic disease, are composed of lipid-laden macrophages called foam cells. Oxidized lipids mediate foam cell formation and have long been implicated as important mediators of atherosclerosis. Minimally oxidized Low-Density Lipoproteins (LDLs), but not native LDLs, induce MCP-1 production in vascular endothelial and smooth muscle cells [9]. MCP-1 is tethered to proteoglycans on the luminal side of endothelial cells and interacts with its receptor, CCR2, expressed on rolling monocytes. This interaction initiates firm integrin-mediated adhesive interactions between monocytes and endothelial cells, ultimately resulting in leukocyte diapedesis into the subendothelium. After entering the subendothelial space monocytes differentiate into macrophages, where they continuously ingest lipid to become the foam cells of the fatty streak [10]. Studies using genetically-targeted animals provided strong evidence that MCP-1 plays an essential role in the recruitment of macrophages in atheromatous lesions. Deletion of the MCP-1 gene in LDL receptor deficient mice [11] and in animals overexpressing human apolipoprotein B [12] protected from the development of diet-induced atherosclerosis, dramatically reducing macrophage recruitment in the aortic wall without altering lipoprotein metabolism. In addition, deletion of CCR2, the only functional MCP-1 receptor, attenuated macrophage accumulation in atherosclerotic lesions and protected from the development of atherosclerosis in apoE null mice fed with a high-fat [13] or a regular chow diet [14]. Furthermore, MCP-1 inhibition using transfection with an N-terminal deletion mutant of the MCP-1 gene markedly inhibited formation of new atherosclerotic plaques in hypercholesterolemic mice [15].

The potential role of MCP-1 in progression of atherosclerotic disease and plaque rupture is less established (Figure 1). MCP-1 induces smooth muscle cell proliferation and may exert

angiogenic effects [16] promoting neovessel formation in the plaque; these actions may lead to rapid progression of the lesion [17]. MCP-1 may also play a role in disruption of the atherosclerotic plaque by inducing Matrix Metalloproteinase (MMP) expression and release [18]. In addition, MCP-1 may have procoagulant properties by inducing tissue factor synthesis and activity in smooth muscle cells [19]. Although direct proof of the *in vivo* significance of these mechanisms is lacking, an inhibitory antibody to MCP-1 and MCP-5 administered to ApoE ^{-/-} mice induced a stable plaque phenotype with increased collagen content [20]. In addition, blockade of MCP-1 through gene therapy limited progression of preexisting lesions in the aortic root in hypercholesterolemic mice [21]. This approach also altered the composition of atherosclerotic plaques into a more stable phenotype, containing fewer macrophages and lymphocytes, less lipid, and more smooth muscle cells and collagen [21].

MCP-1 in cardiac injury and repair

In addition to its established role in the pathogenesis of atherosclerotic disease and its potential significance in disease progression and plaque rupture, MCP-1 is also critically involved in the healing response following an acute coronary event [22]. Myocardial infarction triggers a local inflammatory reaction that results in formation of a scar and is closely intertwined with remodeling of the infarcted ventricle (Figure 1). MCP-1 expression is markedly but transiently induced in infarcted hearts [23], [24] and critically regulates the healing response. *In vitro* studies using post-ischemic cardiac lymph collected during canine reperfused infarction suggested that MCP-1 may be a major factor responsible for mononuclear cell recruitment into the ischemic myocardium during the first five hours of reperfusion [25]. In addition, in a rat model of experimental myocardial infarction, administration of a neutralizing antibody to MCP-1 significantly reduced infarct size decreasing adhesion molecule expression and macrophage infiltration [26]. Beyond its leukotactic actions, MCP-1 may have important effects on infarct healing mediated through its direct angiogenic effects on the vascular endothelium [16], or by modulation of fibroblast phenotype and activity [27].

In order to examine the role of endogenous MCP-1 in infarct healing, we studied the effects of MCP-1 gene disruption and antibody neutralization in a mouse model of reperfused myocardial infarction [28]. MCP-1 ^{-/-} mice had decreased and delayed macrophage infiltration in the healing infarct and demonstrated delayed replacement of injured cardiomyocytes with granulation tissue. MCP-1 ^{-/-} infarcts had decreased mRNA expression of the cytokines TNF- α , IL-1 β , Transforming Growth Factor (TGF)- β , and IL-10 and demonstrated defective macrophage differentiation. MCP-1 deficiency diminished myofibroblast accumulation but did not significantly affect infarct angiogenesis. Despite showing delayed phagocytotic removal of dead cardiomyocytes, MCP-1 ^{-/-} mice had attenuated left ventricular remodeling, but similar infarct size, when compared with wild-type animals. MCP-1 antibody inhibition resulted in defects comparable with the pathological findings noted in infarcted MCP-1 ^{-/-} animals without an effect on macrophage recruitment [28].

Our study suggested that the role of MCP-1 extends beyond its monocyte chemoattractant effects: MCP-1 inhibition with a neutralizing antibody resulted in defects comparable with the pathological findings noted in infarcted MCP-1 ^{-/-} animals in the absence of impairment in monocyte recruitment. Suppression of inflammatory cytokine synthesis, decreased macrophage activation, selective recruitment of monocyte subsets with distinct properties, reduced MMP synthesis and activity, and diminished myofibroblast infiltration may be important mechanisms responsible for attenuated left ventricular remodeling in MCP-1 null mice.

The key role of MCP-1 signaling in the pathogenesis of post-infarction remodeling was also suggested by experiments using mice with genetic disruption of CCR2, the primary receptor for MCP-1 [29]. CCR2 null mice had decreased infiltration with macrophages and exhibited

attenuated ventricular dilation following myocardial infarction. CCR2 absence was associated with markedly decreased metalloproteinase expression and lower gelatinolytic activity in the infarcted ventricle. Attenuated matrix degradation may explain, at least in part, the protection from the development of adverse remodeling noted in CCR2 null animals.

During the inflammatory stage of infarct healing, MCP-1 mediates macrophage recruitment and timely clearance of dead cells from the infarct; however, prolonged induction of the chemokine in the infarcted heart may result in extension of granulation tissue formation and adverse remodeling of the ventricle [30]. Thus, timely resolution of the inflammatory response following myocardial infarction is critical for optimal healing.

MCP-1 as a biomarker in ACS

Evidence suggests that MCP-1 may not be a clinically useful tool in diagnosis of subclinical atherosclerosis. In the Dallas Heart Study MCP-1 plasma levels were associated with traditional cardiovascular risk factors and with coronary artery calcium. After adjustment for age MCP-1 levels were no longer independently associated with subclinical atherosclerosis [31]. A few case control studies examined the ability of MCP-1 levels to predict established cardiovascular disease. In the Atherosclerosis Risk in Communities (ARIC) study patients with incident peripheral arterial disease had significantly higher plasma MCP-1 levels in comparison to control patients [32]. However, the association between MCP-1 levels and peripheral arterial disease was no longer statistically significant when CRP was added to model that included multiple inflammation-related variables. The limited value of plasma MCP-1 levels as a predictor of atherosclerotic disease, despite the marked activation of the MCP-1/CCR2 axis in the vascular wall, may reflect the immobilization of the chemokine on the luminal surface of endothelial cells. Binding to glycosaminoglycan chains of cell surface proteoglycans is essential for MCP-1 actions *in vivo*, ensuring high local concentrations of MCP-1 and providing directional signals for monocyte migration [33]. Selective induction and localization of MCP-1 in atheromatous plaques may not result in marked elevations of plasma MCP-1 levels.

The involvement of MCP-1 in atherothrombotic disease and in the pathogenesis of cardiac remodeling suggested that MCP-1 may be a promising biomarker for risk prediction in patients with coronary artery disease [34], [35]. Several small case-control studies indicated that plasma MCP-1 levels were highest among patients with ACS, intermediate among patients with stable coronary disease, and lowest among healthy control subjects [36], [37]. Aukrust and co-workers demonstrated that MCP-1 levels are increased in patients with unstable angina in comparison to gender and age-matched control individuals and to patients presenting with stable effort angina. [38]. However, there was considerable overlap in MCP-1 levels between groups, indicating that MCP-1 will not be useful as a diagnostic marker in ACS.

Emerging evidence suggests that MCP-1 may have value as a prognostic marker in patients with ACS (Table 1). A substudy of the Oral Glycoprotein IIb/IIIa Inhibition with Orbofiban in Patients with Unstable Coronary Syndromes (OPUS-TIMI-16) trial was the first to prospectively assess the value of MCP-1 as a prognostic biomarker in patients with ACS. In this study 279 volunteers and 2270 patients were evaluated [39]. Assessment of MCP-1 levels was performed prior to initiation of treatment with the glycoprotein IIb/IIIa inhibitor and the study population was divided in quartiles according to the levels of MCP-1 in serum. When compared with the first three quartiles, patients with MCP-1 levels above the 75th percentile (> 238 pg/mL, corresponding to the 90th percentile in the control population) exhibited a trend towards increased mortality during the 10-month follow-up period and were more likely to suffer death or myocardial infarction. The association remained after adjustment for several clinical (age, history of diabetes, hypertension, hypercholesterolemia, coronary disease, creatinine clearance, ST deviation) and biochemical (CRP and troponin I elevation) risk predictors. Although the findings supported the role of MCP-1 as a predictive biomarker in

patients with ACS, the lack of serial measurements prevented determination of the optimal timepoint for measuring plasma levels.

The predictive value of MCP-1 plasma levels was supported by a smaller prospective study that enrolled an unselected population of patients with ischemic chest pain representing the whole clinical spectrum of ACS [40]. Blood samples were taken immediately after admission, and patients were followed during 13 months. Plasma MCP-1 level above the median (>144 pg/ml) was a predictor of future coronary events (recurrent angina or acute infarction, coronary revascularization, or cardiac death). In addition, patients who died had the highest mean levels of MCP-1/CCL2 (355 ± 220 pg/mL) suggesting that the severity of a future event increased with increasing MCP-1 levels. Moreover, patients with an early event (≤ 7 days) had higher MCP-1/CCL2 blood levels compared to the ones that presented a nonfatal endpoint later (after 7 days).

Serial measurement of plasma MCP-1 levels may provide additional prognostic information in patients with ACS. A small study demonstrated a sustained increase in MCP-1 levels in patients with complicated myocardial infarction. During the first week of hospitalization peak MCP-1 levels were higher in patients with infarction complicated by heart failure in comparison to patients with an uncomplicated course [41]. One month after the event, patients with severe left ventricular dysfunction (left ventricular ejection fraction $\leq 35\%$) exhibited significantly higher plasma levels of MCP-1 (and several other CC chemokines) than patients with more preserved function (ejection fraction $> 35\%$). A recent prospective study measured MCP-1 levels in a large population of patients with ACS, enrolled in the A to Z (Aggrastat to Zocor) trial, and examined the prognostic value of serial measurement of MCP-1 levels at the time of enrollment to the Z phase (median 3.5 days after symptom onset) and at the 4 and 12-month outpatient visits [42]. The primary outcome was all-cause mortality. The secondary endpoints were myocardial infarction and the composites of death and myocardial infarction; death, MI, and new or worsening heart failure; as well as the phase Z primary endpoint of cardiovascular death, myocardial infarction, readmission for ACS and stroke. The study confirmed that baseline MCP-1 levels provide independent prognostic value in patients with ACS. The incidence of all primary and secondary end points increased across quartiles of baseline MCP-1 plasma levels. In addition, patients with baseline MCP-1 levels above the pre-specified threshold of 238 pg/mL (based on the findings from the OPUS-TIMI16 substudy [39]) were also at increased risk for death and for each of the secondary endpoints. After adjustment for standard risk predictors and levels of CRP and BNP, MCP-1 levels > 238 pg/mL remained associated with mortality and with each of the composite endpoints. Increasing quartiles of MCP-1 plasma levels measured at 4 months were also associated with mortality and with most composite endpoints; however, no association was observed between 4-month MCP-1 quartiles and subsequent myocardial infarction.

The results obtained from the A to Z trial suggest that MCP-1 may have value as a biomarker for risk stratification in both the initial and the chronic phases after an ACS. In contrast to CRP, MCP-1 levels may be an indicator of long-term cardiovascular risk rather than a marker of acute prognosis.

What are the mechanisms responsible for adverse prognosis in patients with ACS who have elevated MCP-1 levels?

Increased cardiovascular risk in ACS patients with elevated MCP-1 levels is not due to increased cardiac injury because the baseline levels of MCP-1 after an acute coronary event do not appear to reflect the extent of cardiomyocyte necrosis [39]. However, the critical involvement of MCP-1 in the pathogenesis of atherosclerosis, plaque destabilization and post-infarction remodeling suggests several distinct mechanisms that may be responsible for adverse prognosis in patients with elevated MCP-1 levels [43]. First, baseline elevation of plasma

MCP-1 may reflect enhanced expression of the chemokine in atheromatous lesions reflecting a higher burden of atherosclerotic disease. Second, enhanced systemic activation of the MCP-1 axis may exert prothrombotic effects resulting in recurrent coronary events. Third, enhanced and prolonged elevation of plasma MCP-1 may identify patients who mount a more intense cardiac inflammatory reaction following a coronary event, or show impaired resolution and defective containment of the post-infarction inflammatory response. Induction of pro-inflammatory mediators and leukocyte infiltration plays an essential role in phagocytotic removal of dead cells and matrix debris from the infarcted myocardium. However, this acute localized inflammatory response is transient, and is followed by resolution of the inflammatory infiltrate and deposition of fibrous tissue [24]. A crucial commitment is made during the late stages of the inflammatory phase to convert the response from phagocytosis and clearance of dead cells and debris, to a mode that promotes tissue repair and scar formation [44]. Inhibition of chemokine and cytokine synthesis after a dramatic early peak is crucial for the repair process, preventing prolonged expression of inflammatory mediators in the healing infarct, and suppressing continuous leukocyte recruitment, extensive matrix degradation and injury. Thus, optimal healing requires mechanisms inhibiting chemokine and cytokine synthesis resulting in resolution of the inflammatory infiltrate and transition to fibrous tissue deposition. Sustained inflammation following the acute event may result in extensive matrix degradation and adverse cardiac remodeling. The findings from the A to Z study are consistent with this hypothesis: prolonged elevation of MCP-1 levels may reflect impaired mechanisms of resolution of inflammation leading to enhanced adverse remodeling and increased mortality, without affecting the incidence of new myocardial infarction. Studies examining the relation between MCP-1 plasma levels and the extent of remodeling following infarction are needed to explore this concept.

Although age, cigarette smoking, triglyceride levels and body mass index are clinical correlates of MCP-1 levels in large populations, genetic variations play a more important role in MCP-1 plasma level variability [45]. In the Framingham Heart Study Offspring Cohort, individuals carrying the *MCP-1-2578G* allele had higher circulating MCP-1 levels and exhibited an increased risk of myocardial infarction [45]. Whether MCP-1 polymorphisms are associated with enhanced serum MCP-1 levels and worse prognosis after an ACS event remains to be investigated.

Can MCP-1 levels guide pharmacotherapy?

MCP-1 may be of potential value in guiding pharmacological treatment in patients with ACS. 3-hydroxy-3-methylglutaryl coenzyme A (HMG-CoA) reductase inhibition decreases MCP-1 expression in atherosclerotic vessels [46] and reduces plasma MCP-1 levels in hypercholesterolemic patients [47]. It has been suggested that the beneficial effects of statins in patients with myocardial infarction may be mediated at least in part through decreased MCP-1 expression. The A to Z trial examined the hypothesis that elevated MCP-1 levels could identify patients more likely to benefit from aggressive statin treatment. However, the findings demonstrated that the effects of early intensive treatment with simvastatin are independent of baseline MCP-1 plasma levels in patients with ACS suggesting that higher MCP-1 levels do not identify patients who derive incremental benefit from statin therapy. The modest effects of aggressive statin treatment on MCP-1 levels do not exclude a significant role of MCP-1 inhibition in mediating the beneficial actions of the statins, but may suggest that therapeutic interventions aiming at further reducing MCP-1 levels in patients with ACS may provide additional benefit. Experimental studies have demonstrated that disruption of the MCP-1 signaling pathway attenuates dilative remodeling following reperfused myocardial infarction [28] suggesting that MCP-1 may be a novel pharmacologic target in patients with acute coronary events [48]. However, a word of caution should be raised regarding the potential consequences of overly aggressive suppression of the MCP-1 axis in the infarcted myocardium:

absence of MCP-1 results in attenuated post-infarction left ventricular remodeling at the expense of impaired phagocytosis and delayed replacement of injured cardiomyocytes with granulation tissue. In patients with acute myocardial infarction, delayed phagocytosis of dead cardiomyocytes may prolong activation of inflammatory pathways resulting in extension of injury and increased matrix degradation. The effects of MCP-1 inhibition in large mammalian models of infarction should be carefully studied before identifying MCP-1 as a potential target for therapeutic intervention.

Conclusions

MCP-1 is critically involved in formation and progression of atheromatous lesions, may contribute to destabilization of the plaque, and appears to be an essential mediator in adverse remodeling following myocardial infarction. Despite its central involvement in the pathogenesis of atherosclerotic disease, MCP-1 may not be a useful biomarker for risk prediction in patients with subclinical atherosclerosis. However, emerging evidence suggests that baseline and sustained elevation of MCP-1 plasma levels may be an independent predictor of mortality following an acute coronary event. Prolonged elevation of systemic MCP-1 levels following ACS may reflect increased atherosclerotic burden, may be associated with “vulnerable lesions”, or may identify patients with impaired resolution of the post-infarction inflammatory response who are likely to exhibit accentuated adverse remodeling. Further research is required to dissect the mechanisms responsible for adverse prognosis in patients with elevated MCP-1 plasma levels and to examine whether MCP-1 is a promising therapeutic target in patients with ACS.

Acknowledgments

This work was supported by NIH grants R01 HL-76246 and R01 HL-85440.

References and recommended reading

Papers of particular interest have been highlighted as:

• Of importance

•• Of major importance

1. Buffon A, Biasucci LM, Liuzzo G, et al. Widespread coronary inflammation in unstable angina. *N Engl J Med* 2002;347:5–12. [PubMed: 12097534]
2. Libby P. Atherosclerosis: disease biology affecting the coronary vasculature. *Am J Cardiol* 2006;98:3Q–9Q.
3. Morrow DA, Braunwald E. Future of biomarkers in acute coronary syndromes: moving toward a multimarker strategy. *Circulation* 2003;108:250–252. [PubMed: 12876133]
4. Morrow DA, Rifai N, Antman EM, et al. C-reactive protein is a potent predictor of mortality independently of and in combination with troponin T in acute coronary syndromes: a TIMI 11A substudy. *Thrombolysis in Myocardial Infarction. J Am Coll Cardiol* 1998;31:1460–1465. [PubMed: 9626820]
5. Morrow DA, de Lemos JA, Sabatine MS, et al. Clinical relevance of C-reactive protein during follow-up of patients with acute coronary syndromes in the Aggrastat-to-Zocor Trial. *Circulation* 2006;114:281–288. [PubMed: 16847150] [This paper showed that persistent elevation of hsCRP levels 30 days or 4 months following an ACS was associated with increased 2-year mortality rate]
6. Gerard C, Rollins BJ. Chemokines and disease. *Nat Immunol* 2001;2:108–115. [PubMed: 11175802]
7. Ceradini DJ, Kulkarni AR, Callaghan MJ, et al. Progenitor cell trafficking is regulated by hypoxic gradients through HIF-1 induction of SDF-1. *Nat Med* 2004;10:858–864. [PubMed: 15235597]

8. Nelken NA, Coughlin SR, Gordon D, Wilcox JN. Monocyte chemoattractant protein-1 in human atheromatous plaques. *J Clin Invest* 1991;88:1121–1127. [PubMed: 1843454]
9. Cushing SD, Berliner JA, Valente AJ, et al. Minimally modified low density lipoprotein induces monocyte chemotactic protein 1 in human endothelial cells and smooth muscle cells. *Proc Natl Acad Sci U S A* 1990;87:5134–5138. [PubMed: 1695010]
10. Charo IF, Taubman MB. Chemokines in the pathogenesis of vascular disease. *Circ Res* 2004;95:858–866. [PubMed: 15514167] [An interesting comprehensive review on the role of chemokines in vascular disease with an emphasis on MCP-1]
11. Gu L, Okada Y, Clinton SK, et al. Absence of monocyte chemoattractant protein-1 reduces atherosclerosis in low density lipoprotein receptor-deficient mice. *Mol Cell* 1998;2:275–281. [PubMed: 9734366]
12. Gosling J, Slaymaker S, Gu L, et al. MCP-1 deficiency reduces susceptibility to atherosclerosis in mice that overexpress human apolipoprotein B. *J Clin Invest* 1999;103:773–778. [PubMed: 10079097]
13. Boring L, Gosling J, Cleary M, Charo IF. Decreased lesion formation in CCR2^{-/-} mice reveals a role for chemokines in the initiation of atherosclerosis. *Nature* 1998;394:894–897. [PubMed: 9732872]
14. Dawson TC, Kuziel WA, Osahar TA, Maeda N. Absence of CC chemokine receptor-2 reduces atherosclerosis in apolipoprotein E-deficient mice. *Atherosclerosis* 1999;143:205–211. [PubMed: 10208497]
15. Ni W, Egashira K, Kitamoto S, et al. New anti-monocyte chemoattractant protein-1 gene therapy attenuates atherosclerosis in apolipoprotein E-knockout mice. *Circulation* 2001;103:2096–2101. [PubMed: 11319201]
16. Salcedo R, Ponce ML, Young HA, et al. Human endothelial cells express CCR2 and respond to MCP-1: direct role of MCP-1 in angiogenesis and tumor progression. *Blood* 2000;96:34–40. [PubMed: 10891427]
17. Doyle B, Caplice N. Plaque neovascularization and antiangiogenic therapy for atherosclerosis. *J Am Coll Cardiol* 2007;49:2073–2080. [PubMed: 17531655]
18. Robinson SC, Scott KA, Balkwill FR. Chemokine stimulation of monocyte matrix metalloproteinase-9 requires endogenous TNF-alpha. *Eur J Immunol* 2002;32:404–412. [PubMed: 11813159]
19. Schecter AD, Rollins BJ, Zhang YJ, et al. Tissue factor is induced by monocyte chemoattractant protein-1 in human aortic smooth muscle and THP-1 cells. *J Biol Chem* 1997;272:28568–28573. [PubMed: 9353321]
20. Lutgens E, Faber B, Schapira K, et al. Gene profiling in atherosclerosis reveals a key role for small inducible cytokines: validation using a novel monocyte chemoattractant protein monoclonal antibody. *Circulation* 2005;111:3443–3452. [PubMed: 15967845]
21. Inoue S, Egashira K, Ni W, et al. Anti-monocyte chemoattractant protein-1 gene therapy limits progression and destabilization of established atherosclerosis in apolipoprotein E-knockout mice. *Circulation* 2002;106:2700–2706. [PubMed: 12438296]
22. Frangogiannis NG. The mechanistic basis of infarct healing. *Antioxid Redox Signal* 2006;8:1907–1939. [PubMed: 17034340]
23. Frangogiannis NG. Chemokines in ischemia and reperfusion. *Thromb Haemost* 2007;97:738–747. [PubMed: 17479184]
24. Dewald O, Ren G, Duerr GD, et al. Of mice and dogs: species-specific differences in the inflammatory response following myocardial infarction. *Am J Pathol* 2004;164:665–677. [PubMed: 14742270]
25. Birdsall HH, Green DM, Trial J, et al. Complement C5a, TGF-beta 1, and MCP-1, in sequence, induce migration of monocytes into ischemic canine myocardium within the first one to five hours after reperfusion. *Circulation* 1997;95:684–692. [PubMed: 9024158]
26. Ono K, Matsumori A, Furukawa Y, et al. Prevention of myocardial reperfusion injury in rats by an antibody against monocyte chemotactic and activating factor/monocyte chemoattractant protein-1. *Lab Invest* 1999;79:195–203. [PubMed: 10068207]
27. Gharaee-Kermani M, Denholm EM, Phan SH. Costimulation of fibroblast collagen and transforming growth factor beta1 gene expression by monocyte chemoattractant protein-1 via specific receptors. *J Biol Chem* 1996;271:17779–17784. [PubMed: 8663511]

- 28•. Dewald O, Zymek P, Winkelmann K, et al. CCL2/Monocyte Chemoattractant Protein-1 regulates inflammatory responses critical to healing myocardial infarcts. *Circ Res* 2005;96:881–889. [PubMed: 15774854] [This study demonstrates that MCP-1 regulates macrophage recruitment in the infarcted myocardium and plays an important role in the pathogenesis of adverse remodeling]
29. Kaikita K, Hayasaki T, Okuma T, et al. Targeted deletion of CC chemokine receptor 2 attenuates left ventricular remodeling after experimental myocardial infarction. *Am J Pathol* 2004;165:439–447. [PubMed: 15277218]
30. Frangogiannis NG, Ren G, Dewald O, et al. The critical role of endogenous Thrombospondin (TSP)-1 in preventing expansion of healing myocardial infarcts. *Circulation* 2005;111:2935–2942. [PubMed: 15927970]
31. Deo R, Khera A, McGuire DK, et al. Association among plasma levels of monocyte chemoattractant protein-1, traditional cardiovascular risk factors, and subclinical atherosclerosis. *J Am Coll Cardiol* 2004;44:1812–1818. [PubMed: 15519012]
32. Hoogeveen RC, Morrison A, Boerwinkle E, et al. Plasma MCP-1 level and risk for peripheral arterial disease and incident coronary heart disease: Atherosclerosis Risk in Communities study. *Atherosclerosis* 2005;183:301–307. [PubMed: 16285993]
33. Lau EK, Paavola CD, Johnson Z, et al. Identification of the glycosaminoglycan binding site of the CC chemokine, MCP-1: implications for structure and function in vivo. *J Biol Chem* 2004;279:22294–22305. [PubMed: 15033992]
- 34•. Aukrust P, Halvorsen B, Yndestad A, et al. Chemokines and Cardiovascular Risk. *Arterioscler Thromb Vasc Biol*. Jul 31;2008 [Epub ahead of print, DOI:10.1161/ATVBAHA.107.161240] [This recently published comprehensive review manuscript discusses the role of the chemokines in cardiovascular risk prediction]
35. Aukrust P, Yndestad A, Smith C, et al. Chemokines in cardiovascular risk prediction. *Thromb Haemost* 2007;97:748–754. [PubMed: 17479185]
36. Matsumori A, Furukawa Y, Hashimoto T, et al. Plasma levels of the monocyte chemotactic and activating factor/monocyte chemoattractant protein-1 are elevated in patients with acute myocardial infarction. *J Mol Cell Cardiol* 1997;29:419–423. [PubMed: 9040055]
37. Nishiyama K, Ogawa H, Yasue H, et al. Simultaneous elevation of the levels of circulating monocyte chemoattractant protein-1 and tissue factor in acute coronary syndromes. *Jpn Circ J* 1998;62:710–712. [PubMed: 9766714]
38. Aukrust P, Berge RK, Ueland T, et al. Interaction between chemokines and oxidative stress: possible pathogenic role in acute coronary syndromes. *J Am Coll Cardiol* 2001;37:485–491. [PubMed: 11216967]
39. de Lemos JA, Morrow DA, Sabatine MS, et al. Association between plasma levels of monocyte chemoattractant protein-1 and long-term clinical outcomes in patients with acute coronary syndromes. *Circulation* 2003;107:690–695. [PubMed: 12578870]
40. Kervinen H, Manttari M, Kaartinen M, et al. Prognostic usefulness of plasma monocyte/macrophage and T-lymphocyte activation markers in patients with acute coronary syndromes. *Am J Cardiol* 2004;94:993–996. [PubMed: 15476610]
41. Parissis JT, Adamopoulos S, Venetsanou KF, et al. Serum profiles of C-C chemokines in acute myocardial infarction: possible implication in postinfarction left ventricular remodeling. *J Interferon Cytokine Res* 2002;22:223–229. [PubMed: 11911805]
- 42•. de Lemos JA, Morrow DA, Blazing MA, et al. Serial measurement of monocyte chemoattractant protein-1 after acute coronary syndromes: results from the A to Z trial. *J Am Coll Cardiol* 2007;50:2117–2124. [PubMed: 18036447] [This study demonstrates that patients with persistently elevated MCP-1 plasma levels have adverse outcome after an ACS event]
43. Frangogiannis NG. The prognostic value of monocyte chemoattractant protein-1/CCL2 in acute coronary syndromes. *J Am Coll Cardiol* 2007;50:2125–2127. [PubMed: 18036448]
44. Nathan C. Points of control in inflammation. *Nature* 2002;420:846–852. [PubMed: 12490957]
45. McDermott DH, Yang Q, Kathiresan S, et al. CCL2 polymorphisms are associated with serum monocyte chemoattractant protein-1 levels and myocardial infarction in the Framingham Heart Study. *Circulation* 2005;112:1113–1120. [PubMed: 16116069]

46. Bustos C, Hernandez-Presa MA, Ortego M, et al. HMG-CoA reductase inhibition by atorvastatin reduces neointimal inflammation in a rabbit model of atherosclerosis. *J Am Coll Cardiol* 1998;32:2057–2064. [PubMed: 9857893]
47. Rezaie-Majd A, Maca T, Bucek RA, et al. Simvastatin reduces expression of cytokines interleukin-6, interleukin-8, and monocyte chemoattractant protein-1 in circulating monocytes from hypercholesterolemic patients. *Arterioscler Thromb Vasc Biol* 2002;22:1194–1199. [PubMed: 12117737]
48. Frangogiannis NG. Targeting the inflammatory response in healing myocardial infarcts. *Curr Med Chem* 2006;13:1877–1893. [PubMed: 16842199]

Figure 1. The role of MCP-1 in atherothrombotic disease and myocardial infarction

MCP-1 is markedly induced in early atherosclerotic lesions and is immobilized on the luminal surface of the endothelium through binding with proteoglycans. Interactions between MCP-1 and its receptor CCR2, expressed on mononuclear cells, mediate mononuclear cell (Mo) recruitment into the subendothelial space. Monocytes differentiate into macrophages (Ma) and ingest lipids to form foam cells (FC). MCP-1 is involved in foam cell formation, one of the earliest manifestations of atherosclerosis. (upper left) Enhanced expression of MCP-1 in the vascular wall may contribute to progression of atherosclerotic disease through foam cell activation, stimulation of smooth muscle cell (SMC) proliferation and induction of plaque neovascularization (v, neovessel). MCP-1 may also induce Matrix Metalloproteinase (MMP) synthesis promoting matrix degradation and plaque disruption, and is capable of upregulating tissue factor (TF) expression exerting procoagulant effects. (upper right) MCP-1 is also involved in infarct healing and cardiac remodeling. During the pro-inflammatory phase of healing, MCP-1 is upregulated in the infarcted myocardium and mediates mononuclear cell recruitment. As the infarct heals, MCP-1 synthesis is suppressed leading to resolution of inflammation and formation of a scar. Prolonged increase of plasma MCP-1 levels following ACS may identify patients exhibiting an enhanced inflammatory response in the healing myocardium. Timely repression of MCP-1 synthesis is crucial for optimal healing; sustained inflammation may result in extensive matrix degradation and adverse remodeling of the infarcted ventricle. Additional symbols: EC, endothelial cells.

Table 1

Studies supporting the prognostic value of MCP-1 in ACS

Name of study/Author	Number of patients enrolled	Follow-up	Endpoints	Study design	Outcome
OPUS-TIMI-16 / de Lemos et al. [39]	2,270 patients with ACS 279 healthy volunteers	10 months	All-cause death, nonfatal MI, recurrent ischemia requiring urgent revascularization congestive heart failure and composite (death or MI)	Prospective	Higher levels of MCP-1 were associated with death or MI
Kervinen et al. [40]	183	13 months	Recurrent unstable angina pectoris or MI, coronary revascularization (PCI or CABG), and cardiac death	Prospective	High MCP-1 levels predicted recurrent coronary event
A to Z trial/de Lemos et al. [42]	4,244	12 months	Death (primary) and secondary composite endpoints (death or MI; death, MI or heart failure; cardiovascular death, MI, readmission for ACS or stroke)	Prospective (serial MCP-1 measurements)	High MCP-1 levels predicted death and composite endpoints

ACS, Acute coronary syndrome; MCP, monocyte chemoattractant protein; OPUS-TIMI, Oral glycoprotein IIb/IIIa inhibition with orbofiban in patients with unstable coronary syndromes- Thrombolysis In Myocardial Infarction; PCI, percutaneous coronary intervention; CABG, coronary artery bypass grafting; A to Z, Aggrastat to Zocor, MI, myocardial infarction.