Published in final edited form as:

Tetrahedron Lett. 2010 ; 51(1): 197–200. doi:10.1016/j.tetlet.2009.10.124.

Synthesis of labeled meropenem for the analysis of *M. tuberculosis* transpeptidases

David B. Kastrinsky and Clifton E. Barry III*

The National Institutes of Health, National Institute of Allergy and Infectious Diseases, Tuberculosis Research Section, Bethesda MD 20892

Abstract

A concise synthesis of ^{14}C labeled meropenem prepared from ^{14}C dimethylamine hydrochloride is described. Using a similar reaction sequence, the meropenem nucleus was also attached to biotin providing a probe for protein interaction studies.

The mycobacterial cell wall is a complex, multi-layered heteropolymer composed of a peptidoglycan core that is covalently bound to arabinogalactan and mycolic acids.^{1–3} Although the size and hydrophobic nature of this complex contribute to the impermeability of mycobacteria to many antibiotics,^{4,5} other features likely contribute to resistance to specific classes of antibiotics. The β -lactams (penicillins, cephalosporins, and carbapenems) are the most widely prescribed anti-infectives and derive their biological activity through the acylation and deactivation of the transpeptidases involved in peptidoglycan crosslinking.^{6–8} *Mycobacterium tuberculosis* (*Mtb*) contains at least one chromosomal β -lactamase, Rv2068c, a Class A, extended spectrum β -lactamase.⁹ Rv2068c can be inhibited by clavulanic acid and thus combinations of antibiotic and β -lactamase inhibitor or newer classes of β -lactamase resistant antibiotics should prove effective for treating tuberculosis.

Historically the β -lactams have not been used in treating tuberculosis despite the fact that the peptidoglycan of mycobacteria is extensively crosslinked and the β -lactams penetrate the cell wall and inhibit transpeptidase targets in *Mtb*.^{11,12} A recent explanation for this paradox suggested that the mycobacterial cell wall actually contains two distinct types of crosslink.¹³ The more intensively studied 4-3 crosslinks (Figure 1) are synthesized by a class of D,D transpeptidases called penicillin binding proteins that are inhibited by the β -lactams. In several species of bacteria including *M. tuberculosis*, a different type of 3-3 crosslink (Figure 1) has been observed. These crosslinks are formed by L,D transpeptidases, a new class of cysteine transpeptidase previously thought to be insensitive to β -lactams but now known to be inhibited by the carbapenem class of β -lactam antibiotics. Conceivably, the 3-3 crosslinks provide an alternative architectural modification and some evidence suggests that these linkages play a role in stationary phase rigidification providing benefits for the long term survival of non-replicating bacilli.^{14–20}

© 2009 Elsevier Science. All rights reserved

*Corresponding author. Tel.: 301-435-7509; fax: 301-402-0993; Clifton E. Barry, III; 33 North Dr.; Bldg. 33; Rm. 2W20D; Bethesda, MD 20892 USA; cbarry@mail.nih.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Supplementary Material

Supplementary material including ^1H NMR spectra and experimental information is available and can be found at ().

The carbapenems, a class of four FDA approved antibiotics (imipenem, meropenem, ertapenem and doripenem) are β -lactams with a structure derived from the natural product thienamycin.²¹ This class received much attention due to its broad spectrum potency notably towards gram negative and anaerobic bacteria, its stability to clinically significant β -lactamases, and its rapid, bactericidal activity.²² The penicillins and cephalosporins contain fused bicyclic structures, and are suicide substrates reminiscent of D-Ala-D-Ala, the substrate for the 4-3 transpeptidation reaction, as originally suggested by Tipper and Strominger (Figure 2).⁶ The saturated five membered thiazolidine ring of the penicillins and the saturated six membered dihydrothiazoline ring of the cephalosporins reveal common precursors in valine and cysteine. While the differences in core structure and *N*-acyl substitution account for differences in reactivity, β -lactamase stability, and spectrum of activity, the putative targets of these compounds are still D,D transpeptidases. The structural similarity of the carbapenem core, the hydroxyethyl and sulfide linked proline sidechains excepted, suggested that the carbapenems also inhibited the D,D transpeptidases. This fact was corroborated in several studies,^{23–25} however recent evidence suggests the L,D transpeptidases are another possible target.

The recent discovery that the peptidoglycan of stationary phase cultures of *M. tuberculosis* contains up to 80% 3-3 crosslinks contrasted strikingly with the traditional view of mycobacterial peptidoglycan and strongly supported the notion that the L,D transpeptidases contribute to resistance to the β -lactam antibiotics.²⁶ More intriguingly, a putative L,D transpeptidase, *Mtb* Rv0116c, (Ldt_{Mtb}), was inhibited by meropenem, and from studies conducted in our laboratory, meropenem demonstrated efficacy against extensively drug resistant (XDR)-TB when co-administered with the β -lactamase inhibitor clavulanic acid.²⁷ This combination was also active against non-replicating bacilli, suggesting an essential remodeling or recycling function of these enzymes in metabolically static bacteria. With the immediate goal of identifying the protein targets of the carbapenems in whole cells of *M. tuberculosis*, we undertook the synthesis of two labeled forms of meropenem for use as probes.

The synthesis of ¹⁴C labeled meropenem (**1**, Scheme 1) utilizes bicyclic intermediate **2** (prepared in 2 steps from *trans* 4-hydroxy-L-proline)^{28,29} which is synthesized in kilogram quantities for the production of Merrem®. From **2**, the synthesis was optimized at milligram scale for introduction of the radiolabel. Unlabeled intermediates were prepared as TLC standards and spectral data is contained in the supporting information. A 250 μ Ci sample of ¹⁴C labeled Me₂NH-HCl (4.0 mg) was obtained from American Radiolabeled Chemicals specially prepared in acetonitrile. This reagent was treated with DIPEA (18 μ L, 107 μ moles) and **2** (16 mg, 52 μ moles) for 2 h at 0 °C. The solution was subsequently treated with unlabeled Me₂NH-HCl (4.4 mg, 54 μ moles), DIPEA (18 μ L, 107 μ moles) and stirred for an additional 2 h at 0 °C. While there are reports of in situ opening of **2** and coupling to **4**, we purified the thiol intermediate **3** by flash chromatography (SiO₂, 5% MeOH-CH₂Cl₂). Labeled **3** was coupled to the carbapenem enol phosphate **4** (35 mg, 59 μ moles, Bosche Scientific) with DIPEA (20 μ L, 112 μ moles) in acetonitrile (0.5 mL, 3h, 25 °C) to provide **5** (19 mg, 56% two steps). Hydrogenolysis of **5** (10 mg, H₂, 10% Pd/C, 1 atm, 2 h), in a biphasic (1:1, 2 mL) solvent system of ethyl acetate:aqueous potassium phosphate buffer (0.050 M, pH = 7) followed by removal of the catalyst by filtration through Celite, wash of the aqueous layer with ethyl acetate (2 $\times$ 1 mL) provided an aqueous solution of labeled meropenem **1** which was pure by non radiographic TLC (*R_f* = 0.25) 7:3 MeOH-CH₂Cl₂), radiographic TLC is contained in the supporting info. The aqueous phase containing the pure, labeled antibiotic was adjusted with additional phosphate buffer to a volume of 1.0 mL (29.3 μ Ci/mL, 22% radioactive yield) and was stored in 100 μ L aliquots and frozen at -30 °C for future use.

We also considered the use of an alternative, non-radioactive label for utilization in a protein pull-down strategy (Scheme 2). Since the radiolabeled meropenem synthesis relied on the introduction of a labeled amine, it would be advantageous to introduce an alternative label in

a similar fashion with the only constraints that the label be compatible with the carbapenem nucleus and not impart any significant steric demand. Biotin does not contain an amine handle, but previous reports documented its introduction by Curtius rearrangement of the carboxylic acid.³⁰ Treatment of biotin (1.0 g, 4.2 mmol) in *t*-BuOH (15 mL, distilled from CaH₂) with DPPA (1.0 mL, 4.6 mmol) and Et₃N (0.64 mL, 4.6 mmol) at 95 °C for 14 h provides *N*-Boc norbiotinamine (1.0 g, 74%). Deprotection of this compound (1.6 g, 5.1 mmol) in neat TFA (3.0 mL) at 25 °C provides the salt **7** isolated as a white solid (1.2 g, 87%). Insoluble in acetonitrile, the reaction of **7** (0.56 g, 1.7 mmol) with **2** (0.50 g, 1.6 mmol) in DMF (8.0 mL) with DIPEA (310 µL, 1.8 mmol) at 25 °C for 2 h provides the thiol **8** (0.450 g, 80%). Coupling of **8** (0.181 g, 0.347 mmol) to **4** (0.206 g, 0.347 mmol) with DIPEA (63 µL, 0.364 mmol) provided **9** (0.215 g, 71%). Hydrogenolysis (H₂, 10% Pd/C, 45 psi, 2 h) of **9** (0.233 g, 0.268 mmol) in the biphasic solvent system detailed above (5 mL ethyl acetate: 5 mL phosphate buffer) removed the *p*-nitrobenzyl ester and carbamate. In this case, the aqueous layer was collected, washed with ethyl acetate (2 × 5 mL), and concentrated in vacuo. The crude residue was dissolved in methanol, filtered to remove the precipitated salts, and concentrated again to provide the biotin labeled meropenem **9** (111 mg, 75%) as a yellow solid, pure by ¹H NMR (supporting info). Studies to identify the targets of meropenem in *M. tuberculosis* are underway. It is worth noting that ¹⁴C and biotin labeled meropenems offer tools to identify and characterize the targets of the carbapenems in other organisms.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge Dr. Kriti Arora and Dr. Pradeep Kumar for help with scintillation counts and biological studies. We would like to thank Dr. Bong-Jin Kim of the Korean Research Institute for Chemical Technology (KRICT) for helpful suggestions and Dr. Noel Whittaker of the National Institute of Diabetes, Digestive and Kidney Diseases (NIDDK) for high resolution mass spectral analysis. Funding was provided by the Intramural Research Program of the National Institutes of Health, National Institute of Allergy and Infectious Diseases.

References

1. Brennan PJ, Nikaido H. Annu Rev Biochem 1995;64:29–63. [PubMed: 7574484]
2. Daffe M, Draper P. Adv Microb Physiol 1998;39:131–203. [PubMed: 9328647]
3. Brennan PJ, Crick DC. Curr Top Med Chem 2007;7:475–488. [PubMed: 17346193]
4. Nikaido H, Jarlier V. Res Microbiol 1991;142:437–443. [PubMed: 1871430]
5. Jarlier V, Nikaido H. FEMS Microbiol Lett 1994;123:11–18. [PubMed: 7988876]
6. Tipper DJ, Strominger JL. Proc Natl Acad Sci U S A 1965;54:1133–1141. [PubMed: 5219821]
7. Kelly JA, Moews PC, Knox JR, Frere JM, Ghuyssen JM. Science 1982;218:479–481. [PubMed: 7123246]
8. Lee W, McDonough MA, Kotra L, Li ZH, Silvaggi NR, Takeda Y, Kelly JA, Mobashery S. Proc Natl Acad Sci U S A 2001;98:1427–1431. [PubMed: 11171967]
9. Hugonnet JE, Blanchard JS. Biochemistry 2007;46:11998–12004. [PubMed: 17915954]
10. Dincer I, Ergin A, Kocagoz T. Int J Antimicrob Agents 2004;23:408–411. [PubMed: 15081094]
11. Goffin C, Ghuyssen JM. Microbiol Mol Biol Rev 2002;66:702–738. table of contents. [PubMed: 12456788]
12. Chambers HF, Moreau D, Yajko D, Miick C, Wagner C, Hackbarth C, Kocagoz S, Rosenberg E, Hadley WK, Nikaido H. Antimicrob Agents Chemother 1995;39:2620–2624. [PubMed: 8592990]
13. Wietzerbin J, Das BC, Petit JF, Lederer E, Leyh-Bouille M, Ghuyssen JM. Biochemistry 1974;13:3471–3476. [PubMed: 4210702]

14. Mainardi JL, Legrand R, Arthur M, Schoot B, van Heijenoort J, Gutmann L. *J Biol Chem* 2000;275:16490–16496. [PubMed: 10748168]
15. Mainardi JL, Morel V, Fourgeaud M, Cremniter J, Blanot D, Legrand R, Frehel C, Arthur M, Van Heijenoort J, Gutmann L. *J Biol Chem* 2002;277:35801–35807. [PubMed: 12077139]
16. Mainardi JL, Fourgeaud M, Hugonnet JE, Dubost L, Brouard JP, Ouazzani J, Rice LB, Gutmann L, Arthur M. *J Biol Chem* 2005;280:38146–38152. [PubMed: 16144833]
17. Biarrotte-Sorin S, Hugonnet JE, Delfosse V, Mainardi JL, Gutmann L, Arthur M, Mayer C. *J Mol Biol* 2006;359:533–538. [PubMed: 16647082]
18. Cremniter J, Mainardi JL, Josseaume N, Quincampoix JC, Dubost L, Hugonnet JE, Marie A, Gutmann L, Rice LB, Arthur M. *J Biol Chem* 2006;281:32254–32262. [PubMed: 16943188]
19. Mainardi JL, Hugonnet JE, Rusconi F, Fourgeaud M, Dubost L, Moumi AN, Delfosse V, Mayer C, Gutmann L, Rice LB, Arthur M. *J Biol Chem* 2007;282:30414–30422. [PubMed: 17646161]
20. Mainardi JL, Villet R, Bugg TD, Mayer C, Arthur M. *FEMS Microbiol Rev* 2008;32:386–408. [PubMed: 18266857]
21. Kahan FM, Kropp H, Sundelof JG, Birnbaum J. *J Antimicrob Chemother* 1983;12:1–35. (including references cited therein). [PubMed: 6365872]
22. Nicolau DP. *Expert Opin Pharmacother* 2008;9:23–37. (including references cited therein). [PubMed: 18076336]
23. Edwards JR. *J Antimicrob Chemother* 1995;36:1–17. [PubMed: 8543486]
24. Hujer AM, Kania M, Gerken T, Anderson VE, Buynak JD, Ge X, Caspers P, Page MG, Rice LB, Bonomo RA. *Antimicrob Agents Chemother* 2005;49:612–618. [PubMed: 15673741]
25. Ubukata K, Shibasaki Y, Yamamoto K, Chiba N, Hasegawa K, Takeuchi Y, Sunakawa K, Inoue M, Konno M. *Antimicrob Agents Chemother* 2001;45:1693–1699. [PubMed: 11353613]
26. Lavollay M, Arthur M, Fourgeaud M, Dubost L, Marie A, Veziris N, Blanot D, Gutmann L, Mainardi JL. *J Bacteriol* 2008;190:4360–4366. [PubMed: 18408028]
27. Hugonnet JE, Tremblay LW, Boshoff HI, Barry CE 3rd, Blanchard JS. *Science* 2009;323:1215–1218. [PubMed: 19251630]
28. Matsumura H, Bando T, Sunagawa M. *Heterocycles* 1995;41:147–159.
29. Sunagawa M, Matsumura H, Inoue T, Fukasawa M, Kato M. *J Antibiot (Tokyo)* 1990;43:519–532. [PubMed: 2358404]
30. Foulon CF, Alston KL, Zalutsky MR. *Bioconjug Chem* 1997;8:179–186. [PubMed: 9095358]

Figure 1.
Two Types of Peptidoglycan Crosslink in *M. tuberculosis*

Meropenem

Penicillins

Cephalosporins

Peptidoglycan D-Ala-D-Ala

Figure 2.
Common β -lactam Structures

Scheme 1.
¹⁴C Labeled Meropenem Synthesis

Scheme 2.
Synthesis of a Biotinylated Carbapenem