Published in final edited form as:

Neuroimage. 2010 January 15; 49(2): 1741–1749. doi:10.1016/j.neuroimage.2009.09.058.

Where syntax meets math: Right Intraparietal Sulcus activation in response to grammatical number agreement violations

Manuel Carreiras^{1,2,3}, Lindsay Carr⁴, Horacio A. Barber³, and Arturo Hernandez⁴

¹ Basque Center on Cognition Brain and Language. Donostia-San Sebastián, Spain

² IKERBASQUE. Basque Foundation for Science. Bilbao, Spain

³ Universidad de La Laguna, Tenerife, Spain

⁴ University of Houston, Texas, USA

Abstract

Previous research has shown that the processing of words referring to actions activated motor areas. Here we show activation of the right intraparietal sulcus, an area that has been associated with quantity processing, when participants are asked to read pairs of words with number agreement violations as opposed to phrases with gender agreement violations or with no violation. In addition, we show activation in the left premotor and left inferior frontal areas when either gender or number agreement is violated. We argue that number violation automatically activates processes linked to quantity processing which are not directly related to language mechanisms.

Keywords

fMRI; Agreement; Gender and number; Quantity processing

Introduction

Models of phylogenetic human brain development assume that language emerged from neural structures dedicated to more basic cognitive functions. From this point of view, it is not surprising to find growing evidence from neuroimaging studies that language comprehension and production is not limited to left perisylvian areas (e.g., left inferior frontal and left superior temporal areas), as is commonly viewed in classic neuropsychology texts. Recent research has shown that areas associated with non-linguistic processing may reveal increased activity during language tasks. For instance, word processing related to actions involving different body parts, such as “pick” and “kick”, activate motor and premotor cortex in a somatotopic fashion (Hauk & Pulvermuller, 2004; Hauk et al., 2004; Shtyrov et al., 2004), reading words with strong olfactory associations in their meaning activated olfactory regions of the brain (Gonzalez et al., 2006), and generating colour words activated a region near brain areas involved in the perception of colour (Martin et al., 1995). Thus, evidence has begun to accrue that action and perceptual information associated with the reference of a word contributes in some way to its

Corresponding author: Manuel Carreiras, Basque Center on Cognition Brain and Language, Paseo Mikeletegi, 69. 20009. Donostia-San Sebastian (Spain). m.carreiras@bcbl.eu, Url: www.bcbl.eu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

neural representation (see review in Vigliocco, Barber, Vinson & Druks, submitted). Furthermore, these studies suggest a bridge between the semantic properties of words and their neural activity. To date no study has shown a link between grammatical processing and underlying conceptual representations. Here we show that grammatical processing activates quantity processing areas in phrases with number agreement violations compared to phrases with gender agreement violations or that agree in both number and gender.

Agreement is an important mechanism for language comprehension and production. This is especially noticeable in romance languages, which are richly inflected. In Spanish, nouns are marked as either masculine or feminine, with a natural criterion for assigning gender to human beings (based on the biological sex of the referent) and an arbitrary criterion for objects, roles and abstract entities (Corbett, 1991). They also carry explicit plural markers signalling one aspect of quantity of the semantic referent. The gender and number of determiners, adjectives, pronouns and past participles must always agree with the entity to which they refer. Thus, agreement is a syntactic operation that reflects conceptual features (e.g., the sex of a referent, or the number of objects) and a syntactic relation that is realized through morpho-phonological markers. For instance, in *The woman saw the films last month. She remembers them well now*, the personal pronoun *She* agrees in gender and number with the antecedent and unambiguously co-refers to *the woman*, while the pronoun *them* agrees in number with the antecedent and unambiguously co-refers to *the films*. In romance languages such as in Spanish *them* and *films* should also agree in gender – morphosyntactic gender. Furthermore, this gender is not always directly related to masculine and feminine gender concepts. For example, the word for nose is masculine in Spanish (*la nariz*) but feminine in Portuguese (*o nariz*). Hence, gender is used as an agreement marker which aids the recognition of gender marked items. The alignment of the different agreement cues facilitates the computation of the co-referential relationship necessary to comprehend the sentence introduced by the personal pronoun contributing to discourse cohesion, as well as for computing the relations among elements within a sentence. In fact, empirical behavioural evidence showed that readers are sensitive to the use of gender and number cues in on-line agreement computation (e.g. Cacciari, Carreiras, Barbolini, 1997; Garnham, Oakhill, Ehrlich & Carreiras, 1995; Carreiras, Garnham and Oakhill, 1993).

The mechanisms of agreement computation have been also investigated in the last few years using electrophysiological and neuroimaging techniques. Event-related potentials (ERPs) studies have found effects of agreement violation mainly in two components: the Left Anterior Negativity (LAN) and the P600 (e.g. Barber & Carreiras, 2003, 2005; Barber, Salillas, & Carreiras, 2004; Deutsch & Bentin, 2001; Gunter, Friederici, & Schriefers, 2000; Kutas & Hilliard, 1983; Münte & Heinze, 1994; Osterhout & Mobley, 1995; Silva-Pereyra & Carreiras, 2007). In particular, Barber & Carreiras (2005) studied the agreement of gender and number mechanisms in Spanish between word pairs presented in isolation or embedded in sentences. Disagreement in word pairs formed by a noun and an adjective (e.g., *faro–alto* [lighthouse–high]) produced an N400-type effect, while word pairs formed by a determiner and a noun (e.g., *el–piano* [the-piano]) showed an additional left anterior negativity effect (LAN). Agreement violations with the same words inserted in sentences (e.g., *El piano estaba viejo y desafinado* [the_{masc-sing} piano_{masc-sing} was old_{masc-sing} and off-key]) resulted in a pattern of LAN–P600.

Some recent fMRI investigations (e.g., Miceli et al., 2002; Hernández et al., 2004) indicate that gender features are processed in a network that involves frontal (BA 45 and BA9) and temporal (BA20/21 and BA21) areas of the left hemisphere. For instance, Hernández et al. (2004) investigated the neural correlates of grammatical gender decisions in Spanish. Participants were visually presented with transparent and opaque words and were required to make a gender decision while being scanned. The gender of the word can be easily derived

from the ending in transparent words (e.g., masculine when the word ends in –o, such as *faro* “lighthouse” and feminine if they end in –a, such as *mesa* “table”), but it cannot be derived from the ending in opaque words (e.g., *reloj* “watch”). The comparison between opaque and transparent words revealed increased activity in the left inferior frontal gyrus in BA 44/45, a more superior locus near BA 44/6 and, bilaterally in the insula (near BA 47) and the anterior cingulate gyrus for opaque items with respect to transparent ones. The authors argued that the gender decision for opaque words requires additional morphological processing as compared to transparent ones and confirmed the critical role played by the left inferior frontal gyrus in processing grammatical information. Many fMRI studies, using different tasks, stimuli, languages and paradigms have found activation of left inferior frontal areas in syntactic processing (e.g., Fiebach et al. 2001; Frederici et al., 2003; Heim et al., 2002; Indefrey et al. 2001, 2004; Moro et al., 2001, Ni et al., 2000). In particular, Hammer et al. (2007) found increased activation in left inferior frontal areas when morphosyntactic agreement between pronouns and antecedents was violated, as compared to when it was not; i.e., when antecedents were things, which only carries morphosyntactic gender, and therefore only a syntactic violation is involved. However, it is unclear whether different brain areas will be recruited when agreement involves not only a grammatical relationship, but broken agreement has conceptual consequences, such as in the case of number agreement.

The “triple-code model” of number processing assumes three distinct systems of representation: a quantity system –a nonverbal analogical-semantic representation of the size and distance relations between numbers–, a verbal system –where numerals are represented lexically, phonologically, and syntactically, much like any other type of word–, and a visual system –in which numbers can be encoded as Arabic numerals– (Dehaene, 1992; Dehaene & Cohen, 1995; Dehaene, et al., 2003). These authors proposed that the three circuits coexist in the parietal lobe and capture most of the observed differences between numerical processing tasks: a bilateral intraparietal system associated with a core quantity system, a region of the left angular gyrus associated with verbal processing of numbers, and a posterior superior parietal system of spatial and non spatial attention, strongly engaged in visual attention processes that may contribute to the visual processing of numbers. Thus, the “triple-code model” assumes that there are modality-specific symbolic codes in the visual Arabic and auditory verbal domain, but that there is also a supramodal abstract “number sense” that conveys semantic, i.e., magnitude, information and contributes to mathematical performance (Dehaene, 1992). Thus if, as suggested by Dehaene and Cohen (1995) a region of intraparietal cortex underlies this abstract “number sense”, then it should respond to numbers even when the task itself does not explicitly require processing of numerical magnitude. Interestingly, Eger et al., (2003) have found bilateral activation in the intraparietal sulcus in the absence of explicit magnitude processing when they compared non-arithmetical processing of number with that of letters and colors using a target-detection task across the visual and auditory modalities. They proposed that these intraparietal responses to numbers reflect access to an abstract supramodal representation of numbers that codes magnitude (see however, Cohen Kadosh et al., 2007; Cohen Kadosh & Walsh, in press, for a proposal that numerical representation is notation dependent and thus includes non-abstract representations). Here we asked whether numerical processing can be activated by the processing of grammatical number in a reading task in which agreement of grammatical gender and number is manipulated.

Some previous research has investigated relationships between number sense and quantifiers understanding in natural language. It has been shown that patients with corticobasal degeneration performed significantly worse on quantifier processing than Alzheimer disease patients (Clark & Grossman, 2007). This is thought to be due to the fact that corticobasal degeneration patients have a specific impairment of number sense associated with right hemisphere parietal lobe disease (Halpern, et al., 2004). In addition, McMillan et al. (2005) carried out an fMRI study in which they presented participants two consecutive events. In the

first event they presented sentences containing quantifiers expressions (e.g., some, less than half, etc.) and in the second event the sentences were presented accompanied by a stimulus array which contained eight randomly distributed familiar objects (women, balls, flowers, cars and dinosaurs). The participants were asked to judge the truth of the sentences with respect to the visual array. They found that first-order (some, all) and high-order (less than half, odd) quantifiers both activated the right inferior parietal cortex, and suggested that processing of numerosity is involved in quantifier comprehension. In addition, they found that only high-order quantifiers activated the right dorsolateral prefrontal cortex, and suggested that this could be caused by the contribution of executive resources such as working memory to understand these more complex quantifiers. Since the right intraparietal sulcus has been shown to be involved in number comparison, a task calling for internal manipulation of numerical quantity (Chochon et al, 1999), these results are consistent with the view that numerosity is involved in quantifier comprehension. However, since the present results are based on the contrasts between activation for sentences plus array plus response minus activation of sentences alone, the effects may be caused not necessarily by processes occurring at the comprehension stage but by later processes triggered by the need to compare the two representations (sentences and arrays) as well as by the need to give an explicit response.

The aim of the present fMRI study was to identify the cortical areas involved in agreement processing during comprehension when this is a pure grammatical operation and when it has conceptual consequences. To that end we presented the determiner-noun and noun-adjective pairs used by Barber & Carreiras (2005) in three conditions: agreement in number and gender, number agreement violation and gender agreement violation. If agreement in gender and number involves a stage of syntactic processing irrespective of what cues (purely syntactic or syntactic and semantic) are intervening in the processing (as suggested by Barber & Carreiras, 2005) we should be able to find activations in similar regions when gender and number agreement was violated. In particular, we should expect activation of the left inferior frontal areas based on gender-related previous studies. However, taking into account the “triple code model” as well as previous findings (e.g., Chochon et al, 1999; Eger et al. 2003; Halpern, et al., 2004; McMillan et al. 2005) showing activation of the intraparietal sulcus for quantity processing, we should expect activation in the intraparietal sulcus for number disagreement.

Methods

Participants

A total of fifteen right handed volunteers (7 women) undergraduate students, native speakers of Spanish, aged between 20 and 35 years participated in the study. Each participant received \$45 for participating. Language proficiency was assessed with the Boston Naming Test-Spanish. Participants all received a score of at least 80% (48/60) correct on the Boston Naming Test. They were all either monolingual or had not learned English until adulthood (later than 16 years of age). Participants were assessed for handedness with an abridged Spanish version of the Edinburgh Handedness Inventory (Oldfield, 1971), and completed a language history questionnaire. The participants were also asked about claustrophobia, or any other criteria which could exclude them from participating in an fMRI experiment. All subjects had normal or corrected vision with no history of neurological disorders or reading impairments. This project was approved by the Institutional Review Boards responsible for the treatment of experimental subjects at both Baylor College of Medicine and the University of Houston. Informed consent was obtained from each participant in accordance with the approval of both institutions. One subject was discarded for the data analysis because of movement artifacts.

Design and Task

The 2×3 factorial design included four different types of word pairs (determinant-noun pairs vs. noun adjective pairs) and three disagreement conditions (gender agreement violation vs. number agreement violation vs. baseline: agreement). In each trial participants were presented with two words one after the other. The pairs of words could be determinant-noun (el carro) or noun-adjective pairs (carro-negro), which could agree or disagree in gender or number. Specifically, each trial began with a “+” as a visual cue, presented for 100 msec, followed by the first word (either a determiner or a noun) for 300 msec, followed by a blank screen for 200 msec, followed by a second word (either a noun or an adjective) for 300 msec. They were instructed to judge each pair for grammatical congruency using the buttons in their hands, pressing one of two buttons whether the two words agree (i.e., were grammatically correct) or disagree in gender or number (i.e., were grammatically incorrect). Participants completed the grammatical task, while undergoing simultaneous and continuous whole brain functional imaging. They were given a button-press device for each hand, and instructed which hand should press for grammatically correct and incorrect. The assignment of hands was counterbalanced across participants. During the task, stimuli were presented visually to the participant via a rear projection video screen and a series of mirrors, which allowed the participant to view the screen placed behind their head. Jython software (Reston, Virginia) was used to present the stimuli to the participant every 6.0 – 8.0 seconds (average inter-stimulus interval = 7.01 seconds, SD = 0.72 seconds). The interstimulus interval was varied in order to counteract expectation effects which might diminish or change participants’ strategies. In addition, varying these times also helped to insure that brain activity was sampled at different points of the BOLD response. An event related design was used in which a different randomization of trials was used for each participant.

Stimuli

The stimuli consisted of 266 word pairs used by Barber and Carreiras (2005) of which 192 word pairs were experimental pairs (96 determiner-noun pairs, and 96 noun-adjective pairs) and the other 64 were fillers. The 192 experimental trials were divided in six experimental conditions (see below): determiner-noun agreement phrases; determiner-number violation phrases; determiner-noun gender violation phrases; noun-adjective agreement phrases; noun-adjective number violation phrases; noun-adjective gender violation phrases. Stimuli in the remaining 64 trials were fillers (determiner-noun and noun-adjective word pairs) in which the nouns and adjectives used were opaque; that is, their gender was not identifiable by looking at the word ending. There were thirty-two word-pairs in each of 6 conditions, plus the 64 fillers. Thus, the contrasts involved the 6 phrase types described above, divided first by word condition (e.g., determiner-noun or noun-adjective), and then by violation type (e.g., congruent, gender, or number).

Determiner-noun word pairs

- a. Agreement, e.g. *El piano (the m-s piano m-s)*.
- b. Gender disagreement, e.g. *La piano (the f-s piano m-s)*.
- c. Number disagreement, e.g. *Los piano (the m-p piano m-s)*.

Noun-adjective word pairs

- (d) Agreement, e.g. *Faro alto (lighthouse m-s high m-s)*.
- (e) Gender disagreement, e.g. *faro alta (lighthouse m-s high f-s)*.
- (f) Number disagreement, e.g. *faro altos (lighthouse m-s high m-p)*.

Three lists of 266 experimental word pairs each were generated. Assignment of word pairs to conditions in each list was counterbalanced across participants. Thus, each pair occurred three times across subjects, once in each condition, so that each subject only saw one form of each pair during the experiment. In Spanish it is mandatory that determiners, nouns and adjectives agree in gender and in number. In the present stimuli, gender was always a strictly morphosyntactic feature without semantic significance. All nouns and adjectives were morphologically marked in gender and number, this is, they ended with the canonical suffixes in Spanish for gender (“-o” for masculine and “-a” for feminine) and number (“-s” or “-es” for plural). This way, both gender and number agreement were similar in terms of transparency cues for agreement. Although there are some exceptions (e.g. tesis “thesis”) in which ending does not change for singular and plural, number is almost always transparent, while gender is not. All words were of medium lexical frequency according to the Spanish database (Sebastián-Gallés et al., 2000) and of 3 to 7 letters long. In the noun-adjective pairs the mean frequency for adjectives was 23.27 per million, and the differences of frequency for the masculine, feminine, singular and plural forms were balanced across the counterbalancing conditions. Length was also balanced across conditions. In the article noun pairs, the target (mean frequency 20.19 per million) was always the same in the three conditions (agreement, gender violation and number violation). (See Appendix with the list of experimental materials).

In addition, a list of 64 filler trials was introduced. Some fillers had opaque gender (e.g. the word “reloj” [clock] lacks any explicit morphological marking) and some other fillers were irregular words (e.g. “mano” [hand] ends with the letter “-o” but is feminine). This type of filler was included to prevent participants from using a superficial strategy for solving the task such as, for example, attending only to the suffixes or tracking whether there were the same number of grammatically correct and incorrect sentences. This way, participants had to respond *yes* and *no* the same number of times. All the filler word pairs agreed in gender and number. Each subject received a different randomization order of the trials.

Data Acquisition

Participants were scanned using a 3.0 Tesla Siemens Allegra scanner at Baylor College of Medicine. Anatomical imaging parameters were: 192 transversal slices at 0.89 mm, voxel size = 0.96 mm × 0.96 mm × 0.89 mm, TR = 1200 ms, TE = 2.93 ms, FoV = 245mm × 214mm; in-plane resolution = 256 × 208, flip angle = 12 degrees. The functional images were EPI gradient echo, using 26 transversal slices at 4 mm thickness (0 mm overlap), voxel size 3.44 mm × 3.44 mm × 4.00 mm, TR = 2000 ms, TE = 40 ms, in-lane resolution = 64 × 64, flip angle = 90°. A total of 1200 volume images were taken. The first 2 (dummy) volumes of each run were discarded to allow for T1 equilibration effects.

Data Analysis

Data were analyzed with statistical parametric mapping (SPM2: Wellcome Department of Imaging Neuroscience, London, UK. <http://www.fil.ion.ucl.ac.uk>), running under Matlab 6.5.1 (Mathworks, Sherbon, MA, USA). All volumes from each participant were realigned and unwarped (Jesper et al., 2001), adjusting for residual motion-related signal changes. The average of the motion-corrected images was co-registered to each individual’s structural MRI using a 12-parameter affine transformation. The images were then spatially normalized to the Montreal Neurological Institute (MNI) template by applying a 12-parameter affine transformation, followed by a nonlinear warping using basis functions following the method of Friston, Ashburner, et al., (1995). Functional data were spatially smoothed with a 8-mm full width half maximum isotropic Gaussian kernel, to compensate for residual variability after spatial normalization and to permit application of Gaussian random-field theory for corrected statistical inference (Friston, Holmes, et al., 1995). At the first level, data were analyzed in a participant-specific fashion, with each of the six experimental conditions (gender violation,

number violation and agreement for both the determiner-noun and the noun-adjective pairs) and the filler condition modelled separately and convolved with the canonical hemodynamic response function (HRF). The data were high-pass filtered using a set of discrete cosine basis functions with a cut-off period of 128 seconds. The significance threshold was set at $p < 0.001$ uncorrected. Clusters smaller than 20 contiguous voxels were omitted from the results.

Statistical Analysis

The contrasts of interest were each of the 4 violation conditions [2 type of violation: (gender violation vs. number violation) $\times$ 2 type of word pair: (determiner-noun vs. noun adjective)] relative to its corresponding specific baseline condition (agreement in the determiner-noun pairs and agreement in the noun-adjective pairs). This would help us determine whether differences between the four experimental violation conditions were due to increases or decreases in activation –i.e., of the BOLD signal with respect to the baseline condition. These contrast images were then entered into a second level ANOVA to permit inferences about condition effects across participants (i.e., a random-effects analysis; Holmes & Friston, 1998). Only correctly responded trials were included in the fMRI and in the behavioural analyses. In addition, responses that exceeded 4 sec beyond the presentation of the second word were eliminated from both analyses.

The second level analysis focused on the effects of disagreement vs. agreement (i) over number and gender, and for each type of phrase separately (ii) over determiner-noun pairs and noun-adjective pairs and for each pair separately. Details of the relevant interactions are reported in the tables. Activation over and above the baseline conditions is illustrated in the figures.

Results

Behavioural data

Percentage of errors and mean response times for agreement, gender violation and number violation in determiner-noun and noun-adjective pairs are presented in Table 1. ANOVAs on mean response latencies and error rates were conducted based on the 2 (word condition: determiner-noun vs. noun-adjective pairs) by 3 (type of violation: agreement vs. gender violation vs. number violation) design.

Reaction times—Incorrect responses were excluded from the latency analysis. In addition, in order to avoid the influence of outliers, reaction times more than 2.0 standard deviations above or below the mean for that participant in each condition were excluded.

The ANOVA showed a main effect of word condition [$F(1,13) = 40.3$, $p < .0001$], indicating that reaction times were slower in the noun adjective pairs than in the determiner noun pairs. The main effect of violation type was also significant [$F(2,26) = 9.3$, $p < .001$]. The interaction was not significant [$F < 1$]. Pairwise comparisons showed that there were no differences between the gender and number violation conditions. However, both gender violation and number violation were slower than the agreement condition.

Error rates—The ANOVA showed a main effect of word condition [$F(1,13) = 24.5$, $p < .0001$], indicating that the percentage of errors was larger in the noun adjective pairs than in the determiner noun pairs. The main effect of violation type was not significant [$F(2,26) = 2.7$, $p < .09$]. Importantly, the interaction was significant [$F(2,26) = 3.8$, $p < .05$]. Pairwise comparisons showed that there were no differences between the three violation type conditions in the determiner-noun pairs. However, in the noun adjective conditions the number of errors was larger in the number violation pairs than in the agreement and in the gender violation conditions.

fMRI data

A main effect of increase of activation for number and gender violation agreement relative to the agreement baseline was observed in the left premotor cortex and in the left inferior frontal areas (see Table 2 and Figure 1 for details). In addition, an increase of activation of number violation relative to the agreement baseline and to the gender violation was observed in the right parietal areas (right intraparietal sulcus and right superior parietal gyrus) (see Table 2). Importantly, this effect was mainly driven by the determiner-noun pairs conditions which showed an increase of activation in number violation as compared to gender violation and as compared to agreement in the right intraparietal sulcus and the right superior parietal gyrus (see Table 3 and Figure 2). Interestingly, the same contrast in the left homologue areas of the parietal cortex did not show any effect significant even at $p < .05$ uncorrected threshold. Laterality analyses were performed for the two parietal regions (intraparietal sulcus and superior parietal gyrus) extracting the signal for the two main coordinates $+40, -48, 40$ and $+24, -70, 46$ within a 6mm sphere. The ANOVA including Region and Hemisphere as factors on the BOLD signal for number violation relative to agreement showed a main effect of Hemisphere $F(1,13)=4.96, p < .05$. Neither the main effect of Region nor the interaction between Region and Hemisphere were significant (both $F_s < 1$).

No further increases of activation were observed when agreement was compared to the violation conditions or when gender violation was compared to number violation or to agreement. In addition, no effects were observed for the contrasts relative to the type of word (determiner-noun vs. noun-adjective and viceversa).

Finally, singular vs plural contrasts were performed for the two parietal regions extracting the signal of the coordinates within a 6mm sphere. No significant effects were found either in the singular > plural or in the plural > singular contrasts.

Discussion

Grammatical violations of gender and number agreement produced similar and distinct effects on regional brain activation. Activation in the left premotor and left inferior frontal regions was higher for the two ungrammatical conditions as compared to the grammatical condition, both in the case of determiner-noun and noun-adjective pairs. However, the activation in the right parietal areas (intraparietal sulcus and superior parietal gyrus) was higher for number disagreement than for gender disagreement and agreement conditions, but this effect was mainly driven by the determiner-noun pairs. Thus, we have identified brain areas that are modulated by both gender and number agreement violations. Furthermore, we have identified neural activity unique to number agreement violation. Finally, the behavioural data showed that it was easy and faster to judge the determiner noun pairs than the noun adjective pairs, and the agreement than the disagreement conditions.

Increased activation for number and gender agreement violations over agreement

The left premotor and left inferior frontal areas that we found to be more active for number and gender violations relative to congruent pairs are very close to the areas other authors have previously associated with gender computation (Hammer et al., 2007; Hernández et al., 2004; Miceli et al., 2002) as well as to areas involved in different types of grammatical processing of different structures in different languages (Fiebach et al. 2001; Frederici et al., 2003; Heim et al., 2002; Indefrey et al. 2004; Moro et al., 2001, Ni et al., 2000). These results suggest that agreement violation processing during visual word processing tasks modulates activation in regions that are also engaged during syntactic processing. The similarity in the effects of agreement violation and syntactic processing is also consistent with the results obtained in ERPs showing that both gender and number violation modulated the LAN and the

P600 (Barber & Carreiras, 2005). These two components were found to be sensitive to other grammatical violations and did not differ in early stages of processing. Thus, the present results add evidence to consider that these left frontal areas are involved in morphosyntactic processing. Number and gender violation seem to require additional morphosyntactic processing in order to determine the nature and cause when two words do not agree in morphosyntactic features.

Increased of activation for number violation relative to gender violation and agreement

Interestingly, number agreement violation activated the right intraparietal sulcus and the right superior parietal gyrus, this effect being mainly driven by the determiner-noun pairs. The “triple code model” and several fMRI studies have suggested a crucial role of regions situated along the horizontal segment of the intraparietal sulcus of both hemispheres for the representation of numerical quantities (e.g., Dehaene et al., 2003). The present results only showed right but not left IPS activation, what seems to be partially inconsistent with the “triple code model” predictions. However, recent studies have suggested that left and right IPS seem to play a different role. Briefly, the right intraparietal sulcus has been implicated in several tasks related to quantity processing (e.g., Dehaene et al., 2003; Piazza et al., 2002; 2006; 2007; Pinel et al., 2004). Independent lines of research have pointed to parietal cortex as crucial for coding numerical quantity both when it is conveyed by number symbols (digits, number words; Eger et al., 2003; Pinel et al., 2001) or by nonsymbolic displays of dot patterns (Ansari et al., 2006; Cantlon et al., 2006; Piazza et al., 2004), which suggests that is involved in computing an abstract code for numerical magnitude (see however, Cohen Kadosh et al., 2007; Cohen Kadosh & Walsh, in press, who recently challenged the idea of abstract representation and suggested that numerical representation in the intra-parietal sulcus (IPS) is notation dependent). Activation of the more posterior dorsal parietal areas has been observed whenever subjects count, which relates also to movements of spatial attention (Dehaene et al., 2003). However, we did not observe activation in the left angular gyrus, the most likely candidate a priori in case number violation triggered some verbal numerical code of numerical computation. Nonetheless, the lack of activation in the left angular gyrus could be due to the fact that our control condition is also a linguistic task. All three conditions (number disagreement, gender disagreement and agreement) involve verbal stimuli, so effects of verbal decoding and comprehension would be cancelled out since they are present in all three conditions. If expectations about several or a single entity is what makes number agreement (specially in determiner-noun pairs) different from the other two conditions, and other results involving numerical comparison or magnitude estimations have found mainly right IPS activation, it is quite likely instead that we have only found right IPS activation.

A comparison between correct plurals and singulars was performed because linguistic and psycholinguistic theories have debated about differences between singulars and plurals in terms of markedness. Although the most common view in the psycholinguistic literature assumes that singular is unmarked and plural marked (Eberhard 1997) the linguistics literature is very much more nuanced. Thus, Greenberg’s (1966) Number Hierarchy (singular before plural before dual before trial) establishes the typological unmarkedness of the singular based on the fact that (a) more languages have singulars than plurals, (b) more languages use the singular more than the plural, and (c) whenever languages have both numbers, there will be more nouns in the singular than in the plural. However, that ordering does not cover the case of facultative number (Corbett 2000) and in fact it does not even apply to the full to languages with obligatory singular/plural contrasts in the sense that, for specific nouns (like arms), the plural may actually be the unmarked term (Tiersma 1982). Additionally, plurality interacts closely with (in) definiteness and this in turn with whether NPs have specific determiners, and whether such NPs project kinds or properties (see Dayal 2004). Note also Chierchia’s idea (1998) that nouns in numeral classifier languages (like Chinese or Japanese) will exhibit neutralization of the

singular/plural distinction. Since we have neither manipulated (in)definiteness nor the inherent specificity of unmarked plurals like arms in any way, we prefer to remain agnostic on the issue of plurality. For the linguistic view that plural number is the unmarked term, see Krifka (1989) and Sauerland et al. (2005). We looked for differences of activation between correct singulars and plurals in the regions of interest in which we have found an increase of activation for number violation, but we did not find differences (either increase of activation for singulars or for plurals).

Importantly, the effect of number violation relative to gender violation and agreement was mainly specific to determiner-noun pairs. Figure 2 shows that number violation in the determiner-noun pairs led to increased activation of the right intraparietal sulcus and the superior parietal gyrus compared to the agreement baseline and to the gender violation conditions. It is quite unlikely that the right intraparietal sulcus and the right superior parietal gyrus are brain regions specifically involved in language processing. In contrast, the increased right parietal activation for number violation is more likely to reflect the automatic activation of a quantity computation while participants were performing a grammatical judgment task: is this pair of words grammatically correct or incorrect? When reading the determiner, participants trigger expectations about whether the noun would be singular or plural, that is, whether it should refer to only one or many entities. Such expectations about quantity could be fulfilled or not when they read the noun. Therefore, the number mismatch leads to an increase of activation in areas involved in computing quantity. The fact that the effect is negligible for the noun-adjective pairs alone may be accounted for by the fact that plural or singular nouns do not create such important expectations (see Barber & Carreiras, 2005). Determiners must be followed by nouns, but nouns may be followed by many other possible constituents (verbs, adjectives, prepositions, adverbs etc.) some of which do not need to agree in number with the noun. In addition, in the case of noun-adjective pairs, the noun creates a concrete representation with a particular number and the adjective qualifies the noun, however, the determiner creates an expectation of number that will be confirmed with the representation of the noun that follows.

The human intraparietal sulcus is systematically activated in all number tasks and could host a representation of quantity (see however, Gobel et al., 2004; Shuman and Kanwisher, 2004). There is considerable evidence from neuroimaging studies to indicate that the parietal lobes are involved in the processing of symbolic and non symbolic numerical information (e.g., Cappelletti et al., in press; Castelli, Glaser, & Butterworth, 2006; Dehaene et al., 2003; Dehaene, Molko, & Cohen, 2004; Zago et al., 2001; Piazza et al., 2006; 2007). However, the left and right intraparietal sulcus may play slightly different roles. The left parietal region is critical for most calculation deficits, it is involved in tasks requiring access to verbal memory, such as verbal coding of numbers, in exact computation, comes into play as a result of education and symbolic acquisition and provides the interconnection of the quantity representation with the linguistic code (see Dehaene et al., 2003; Dehaene & Cohen, 1995). In contrast, the right intraparietal sulcus is engaged when recognizing numerosity (the number of objects in a set), which provides us with a basic intuition that guides the acquisition of formal arithmetic, and it is present from infancy/early childhood (see Cantlon et al., 2006; Dehaene 1997; Dehaene et al., 2003). In fact, activation of the right intraparietal sulcus only or activation higher than the left intraparietal sulcus has been observed when subjects estimate the numerosity of a set of concrete visual or auditory objects (Piazza et al., 2002, 2006, 2007). Approximate judgments (in both the visual and auditory domain) correlate with stronger activation in the right than in the left IPS, while exact judgments correlate with more activation in the left versus right IPS (Piazza et al., 2006). For instance, Piazza, et al., (2002) demonstrated right lateralized activation of inferior parietal cortex associated with determining and manipulating the numerical quantities of a set. Chochon et al. (1999) demonstrated greater right hemisphere activation in a magnitude comparison task, but greater left hemisphere activation in a multiplication task.

Cantlon et al. (2006) reported that the only region showing overlapping activation in preschool children and adults during processing of numerosity was the right IPS. In addition, Cappelletti et al (in press) found that right parietal activation was number-selective. They compared activation to numbers and object names during the same conceptual and perceptual tasks while factoring out activations correlating with response times. Activation was higher for conceptual decisions of numbers relative to the same tasks on object names only in right parietal areas.

Interestingly, recruitment of the right inferior parietal cortex was also observed when assessing quantifiers (Clark & Grossman, 2007; McMillan et al., (2005). The fact that processing of quantifiers involves the right inferior parietal cortex, an area associated with number sense which is not typically associated with language, suggests an involvement of number knowledge in quantifiers processing. This link between number processing and quantifiers processing is further supported by the finding that patients with corticobasal degeneration performed significantly worse on quantifier processing than Alzheimer disease patients (Clark & Grossman, 2007). This is thought to be due to the fact that corticobasal degeneration patients have a specific impairment of number sense associated with right hemisphere parietal lobe disease (Halpern, et al., 2004). In addition, this finding of right parietal activation and damage associated with quantifiers' comprehension is consistent with the previously mentioned neuroimaging literature investigating the neural basis of number knowledge. Finally, Cohen Kadosh et al., (2007) applied TMS to disrupt left- or right-IPS activation clusters in order to induce dyscalculic-like behavioral deficits in healthy volunteers. They found that automatic magnitude processing was impaired only during disruption of right-IPS activity. Thus, it seems that the right, but not the left, IPS is functionally necessary for automatic magnitude processing. It might well be that the right IPS is used for coding magnitude in general (Walsh, 2003), whether numerical or nonnumerical.

While we cannot completely assure that the intraparietal activation reported here is due to numerical processing, there are two facts that seem to suggest that this is the case (1) differences between the two disagreement conditions are limited to agreement in number or in gender. The same determiners and nouns are used across the two conditions (gender and number agreement). Since both gender and number lead to increased activity in the left inferior frontal and left premotor regions, but only number disagreement led to activity in the intraparietal sulcus, it can be inferred that the purely grammatical effects common to the two disagreement conditions are leading to the common activation in left frontal and premotor regions, whereas the conceptual number effects due to the number disagreement condition are leading to the intraparietal activation. Gender and number conditions are equated in syntactic complexity (in fact they showed equivalent response times in the syntactic judgement task), but they differ mainly at the conceptual level in the semantic representation of numerosity. In addition (2) the coordinates revealed in the present study overlap with those reported in other studies looking at numerosity processing. The coordinates in the present paper (40, -48, 40) fall within the area that is associated with numerical processing (37, -46, 42, in a meta-analysis of 14 fMRI studies of magnitude comparisons (Cohen-Cadosh, Lammertyn, & Izard, 2008). Thus, the overlap of coordinates with numerosity processing and the fact that gender and number disagreement most likely differ only in numerosity have led us to conclude that grammatical number disagreement involves numerosity computation which in turn leads to increased activity in the right intraparietal sulcus.

In sum, the results of the present experiment add evidence to the suggestion that the quantity processing would be a function implemented in intraparietal regions of the right hemisphere and reinforces the assumption of a non-verbal system for numerosity judgements. It is striking that the right intraparietal sulcus has been activated by a verbal task for which quantity processing, at least on the surface, is not required. In addition, the results are in line with the general proposal that claim that language processing is not restricted to the classical "languages

areas". In this regard, motor and sensory areas may be involved in some aspects of semantic language processing. The present study provides evidence that syntactic processes can also involve cerebral areas not directly related to the language processing.

Conclusion

The present results converge on the conclusion that the human brain is equipped with a shared mechanism to compare numbers and other magnitudes. In addition, the present research suggests that semantic representation is distributed in a large number of different networks which extend beyond perisylvian areas. Results show that number violation specially when processing determiner-noun phrases correlates with increased activity of a right lateralized fronto-parietal cortical network, which is recruited during tasks which involve quantity processing. In sum, the current findings show the involvement of the right parietal lobe for automatic activation of magnitude processing during language processing. These results add evidence to the growing view of language as a complex process that is not restricted to the classical cerebral areas of language.

Acknowledgments

MC was funded by grants (CONSOLIDER INGENIO2010 CSD2008-00048 and SEJ2006-09238) from the Spanish Ministry of Science and Innovation, and (MTKD-CT-2005-029639) from the European Commission. HAB is funded by the "Ramón y Cajal" program and the grant SEJ2007-67364 from the Spanish Ministry of Education and Science. AH work on the project was supported by the National Institute of Child Health and Development grant (1R03HD50313-01). We would like to thank three anonymous reviewers for comments to a previous version of the manuscript.

References

- Ansari D, Dhital B, Siong SC. Parametric effects of numerical distance on the intraparietal sulcus during passive viewing of rapid numerosity changes. *Brain Res* 2006;1067:181–188. [PubMed: 16359648]
- Barber H, Carreiras M. Integrating gender and number information in Spanish word pairs: An ERP study. *Cortex* 2003;39:465–482. [PubMed: 12870822]
- Barber H, Carreiras M. Grammatical gender and number agreement in Spanish: An ERP comparison. *Journal of Cognitive Neuroscience* 2005;17:137–156. [PubMed: 15701245]
- Barber, H.; Salillas, E.; Carreiras, M. Gender or genders agreement?. In: Carreiras, M.; Clifton, C., editors. *On-line study of sentence comprehension; eye-tracking, ERP and beyond*. Brighton, UK: Psychology Press; 2004.
- Cacciari C, Carreiras M, Barbolini Cionini C. When words have two genders: Anaphor resolution for Italian functionally ambiguous words. *Journal of Memory and Language* 1997;37:517–532.
- Cappelletti M, Lee HL, Elliot D, Freeman ED, Price CJ. The Role of Right and Left Parietal Lobes in the Conceptual Processing of Numbers. *Journal of Cognitive Neuroscience*. In press
- Carreiras M, Garnham A, Oakhill JV. The use of superficial and meaning-based representations in interpreting pronouns: Evidence from Spanish. *European Journal of Cognitive Psychology* 1993;5:93–116.
- Cantlon JF, Brannon EM, Carter EJ, Pelphrey KA. Functional imaging of numerical processing in adults and 4-y-old children. *PLoS Biol* 2006;4:e125.10.1371/journal.pbio.0040125 [PubMed: 16594732]
- Castelli F, Glaser DE, Butterworth B. Discrete and analogue quantity processing in the parietal lobe: A functional MRI study. *Proceedings of the National Academy of Science* 2006;103:4693–4698.
- Chierchia, Gennaro. Reference to kinds across languages. *Natural Language Semantics* 1998b;6:339–405.
- Chochon F, Cohen L, van de Moortele PF, Dehaene S. Differential contributions of the left and right inferior parietal lobules to number processing. *Journal of Cognitive Neuroscience* 1999;11:617–630. [PubMed: 10601743]
- Clark R, Grossman M. Number sense and quantifier interpretation. *Topoi* 2007;26:51–62.

- Cohen Kadosh R, Walsh V. Numerical representation in the parietal lobes: Abstract or not abstract? Behavioral and Brain Sciences. In press
- Cohen Kadosh R, Cohen Kadosh K, Kaas A, Henik A, Goebel R. Notation-dependent and -independent representations of numbers in the parietal lobes. *Neuron* 2007;53:307–314. [PubMed: 17224410]
- Cohen Kadosh R, Cohen Kadosh K, Schuhmann T, Kaas A, Goebel R, Henik A, Sack AT. Virtual dyscalculia induced by parietal-lobe TMS impairs automatic magnitude processing. *Curr Biol* 2007;17:689–693. [PubMed: 17379521]
- Cohen Kadosh R, Lammertyn J, Izard V. Are numbers special? An overview of chronometric, neuroimaging, developmental and comparative studies of magnitude representation. *Progress in Neurobiology* 2008;84:132–147. [PubMed: 18155348]
- Corbett, GG. Gender. Cambridge: Cambridge University Press; 1991.
- Corbett Greville, G. Number. Cambridge, UK: Cambridge University Press; 2000.
- Dayal V. Linguistics and Philosophy 2004;27:393–450.
- Dehaene S. Varieties of numerical abilities. *Cognition* 1992;44:1–42. [PubMed: 1511583]
- Dehaene, S. The Number Sense: How the Mind Creates Mathematics. Oxford University Press; Oxford, UK: 1997.
- Dehaene S, Cohen L. Towards an anatomical and functional model of number processing. *Mathematical Cognition* 1995;1:83–120.
- Dehaene S, Piazza M, Pinel P, Cohen L. Three parietal circuits for number processing. *Cogn Neuropsychol* 2003;20:487–506.
- Dehaene S, Molko N, Cohen L. Arithmetic and the brain. *Current Opinion in Neurobiology* 2004;14:218–224. [PubMed: 15082328]
- Deutsch A, Bentin S. Syntactic and semantic factors in processing gender Grammatical Agreement in Hebrew: Evidence from ERPs and Eye Movements. *Journal of Memory and Language* 2001;45:200–224.
- Eberhard KM. The marked effect of number on subject-verb agreement. *Journal of Memory and Language* 1997;36:147–164.
- Eger E, Sterzer P, Russ MO, Giraud AL, Kleinschmidt A. A supramodal number representation in human intraparietal cortex. *Neuron* 2003;37:719–725. [PubMed: 12597867]
- Fiebach CJ, Schlesewsky M, Friederici AD. Syntactic working memory and the establishment of filler-gap dependencies: insights from ERPs and fMRI. *J Psycholinguist Res* 2001;30:321–338. [PubMed: 11523277]
- Friederici AD, Ruschemeyer SA, Hahne A, Fiebach CJ. The role of left inferior frontal and superior temporal cortex in sentence comprehension: localizing syntactic and semantic processes. *Cerebral Cortex* 2003;13:170–177. [PubMed: 12507948]
- Friston KJ, Ashburner J, Frith CD, Poline JB, Heather JD, Frackowiak RSJ. Spatial registration and normalization of images. *Hum Brain Map* 1995;2:1–25.
- Garnham A, Oakhill JV, Ehrlich MF, Carreiras M. Representations and processes in the interpretation of pronouns: New evidence from Spanish. *Journal of Memory and Language* 1995;34:41–62.
- Göbel SM, Johansen-Berg H, Behrens T, Rushworth MFS. Response-selection-related parietal activation during number comparison. *J Cogn Neurosci* 2004;16:1536–1551. [PubMed: 15601517]
- Gonzalez J, Barros-Loscertales A, Pulvermuller F, Meseguer V, Sanjuán A, Belloch V, Avila C. Reading cinnamon activates olfactory brain regions. *Neuroimage* 2006;32:906–012. [PubMed: 16651007]
- Greenberg, JH. Some universals of grammar with particular reference to the order of meaningful elements. In: Greenberg, JH., editor. *Universals of grammar*. Cambridge, Mass: M.I.T; 1966. p. 73–113.
- Gunter TC, Friederici AD, Schriefers H. Syntactic gender and semantic expectancy: ERPs reveal early autonomy and late interaction. *Journal of Cognitive Neuroscience* 2000;12:556–568. [PubMed: 10936910]
- Halpern C, Glosser G, Clark R, Gee J, Moore P, Dennis K, McMillan C, Colcher A, Grossman M. Dissociation of numbers and objects in corticobasal degeneration and semantic dementia. *Neurology* 2004;62:1163–1169. [PubMed: 15079017]

- Hammer A, Goebel R, Schwarzbach J, Münte TF, Jansma BM. When sex meets syntactic gender on a neural basis during pronoun processing. *Brain Research* 2007;1146:185–198. [PubMed: 16904083]
- Hauk O, Pulvermüller F. Neurophysiological distinction of action words in the fronto-central cortex. *Hum Brain Mapp* 2004;21:191–201. [PubMed: 14755838]
- Hauk O, Johnsrude I, Pulvermüller F. Somatotopic representation of action words in the motor and premotor cortex. *Neuron* 2004;41:301–307. [PubMed: 14741110]
- Hernandez AE, Kotz SA, Hofmann J, Valentin VV, Dapretto M, Bookheimer SY. The neural correlates of grammatical gender decisions in Spanish. *Neuroreport* 2004;15:863–866. [PubMed: 15073532]
- Heim S, Opitz B, Friederici AD. Broca's area in the human brain is involved in the selection of grammatical gender for language production: Evidence from event-related functional magnetic resonance imaging. *Neurosci Lett* 2002;328:101–104. [PubMed: 12133565]
- Holmes AP, Friston KJ. Generalisability, Random Effects and Population Inference. *Neuroimage* 1998;7:S754.
- Indefrey P, Hagoort P, Herzog H, Seitz RJ, Brown CM. Syntactic processing in left prefrontal cortex is independent of lexical meaning. *Neuroimage* 2001;14:546–555. [PubMed: 11506529]
- Indefrey P, Hellwig F, Herzog H, Seitz RJ, Hagoort P. Neural responses to the production and comprehension of syntax in identical utterances. *Brain Lang* 2004;89:312–319. [PubMed: 15068913]
- Jesper LR, Andersson JLR, Hutton C, Ashburner J, Turner R, Friston K. Modeling Geometric Deformations in EPI Time Series. *Neuroimage* 2001;13:903–919. [PubMed: 11304086]
- Krifka, Manfred. Nominal Reference, Temporal Constitution and Quantification in Event Semantics. In: Bartsch, Renate; van Benthem, Johan; van Emde Boas, Peter, editors. *Semantics and Contextual Expressions*. Dordrecht: Foris Publications; 1989. p. 75–115.
- Kutas M, Hillyard SA. Event-related brain potentials to grammatical errors and semantic anomalies. *Memory and Cognition* 1983;11:539–550.
- Martin A, Haxby JV, Lalonde FM, Wiggs CL, Ungerleider LG. Discrete cortical regions associated with knowledge of color and knowledge of action. *Science* 1995;270:102–105. [PubMed: 7569934]
- McMillan CT, Clark R, Moore P, Devita C, Grossman M. Neural basis for generalized quantifier comprehension. *Neuropsychologia* 2005;43:1729–1737. [PubMed: 16154448]
- Miceli G, Turriziani P, Caltagirone G, Capasso R, Tomaiuolo F, Caramazza A. The neural correlates of grammatical gender: an fMRI investigation. *J Cogn Neurosci* 2002;14:618–628. [PubMed: 12126502]
- Moro A, Tettamani M, Perani D, Donati C, Cappa S, Fazio F. Syntax and the brain: disentangling grammar by selective anomalies. *Neuroimage* 2001;13:110–118. [PubMed: 11133314]
- Münte, TF.; Heinze, HJ. ERP negativities during syntactic processing of written words. In: Heinze, H-J.; Münte, TF.; Mangun, GR., editors. *Cognitive electrophysiology*. Boston: Birkhäuser; 1994. p. 211–238.
- Ni W, Constable RT, Mencl WE, Pugh KR, Fulbright RK, Shaywitz SE, Shaywitz BA, Gore JC, Shankweiler D. An event-related neuroimaging study distinguishing form and content in sentence processing. *J Cogn Neurosci* 2000;12:120–133. [PubMed: 10769310]
- Oldfield RC. The assessment and analysis of handedness: the Edinburgh inventory. *Neuropsychologia* 1971;9:97–113. [PubMed: 5146491]
- Osterhout L, Mobley LA. Event-related potentials elicited by failure to agree. *Journal of Memory and Language* 1995;34:739–773.
- Piazza M, Mechelli A, Price C, Butterworth B. Exact and approximate judgements of visual and auditory numerosity: an fMRI study. *Brain Res* 2006;1106:177–188. [PubMed: 16828717]
- Piazza M, Izard V, Pinel P, Le Bihan D, Dehaene S. Tuning curves for approximate numerosity in the human intraparietal sulcus. *Neuron* 2004;44:547–555. [PubMed: 15504333]
- Piazza M, Izard V, Pinel P, Le Bihan D, Dehaene S. Tuning curves for approximate numerosity in the human intraparietal sulcus. *Neuron* 2004;44:547–555. [PubMed: 15504333]
- Piazza M, Mechelli A, Price C, Butterworth B. Exact and approximate judgements of visual and auditory numerosity: an fMRI study. *Brain Res* 2006;1106:177–188. [PubMed: 16828717]
- Piazza M, Pinel P, Le Bihan D, Dehaene S. A magnitude code common to numerosities and number symbols in human intraparietal cortex. *Neuron* 2007;53:293–305. [PubMed: 17224409]

- Pinel P, Dehaene S, Riviere D, LeBihan D. Modulation of parietal activation by semantic distance in a number comparison task. *Neuroimage* 2001;14:1013–1026. [PubMed: 11697933]
- Pinel P, Piazza M, Le Bihan D, Dehaene S. Distributed and overlapping cerebral representations of number, size, and luminance during comparative judgments. *Neuron* 2004;41:983–993. [PubMed: 15046729]
- Sauerland, U.; Jan, A.; Kazuko, Y. The plural involves comparison. In: Kesper, S.; Reis, M., editors. *Linguistic evidence*. Berlin: Mouton de Gruyter; 2005.
- Sebastián, N.; Martí, MA.; Carreiras, M.; Cuetos, F. *LEXESP: Una base de datos informatizada del español*. Barcelona: Servicio de Publicaciones de la Universitat de Barcelona; 2000.
- Shtyrov Y, Hauk O, Pulvermüller F. Distributed neuronal networks for encoding category-specific semantic information: the mismatch negativity to action words. *Eur J Neurosci* 2004;19:1083–1092. [PubMed: 15009156]
- Shuman M, Kanwisher N. Numerical magnitude in the human parietal lobe: tests of representational generality and domain specificity. *Neuron* 2004;44:557–569. [PubMed: 15504334]
- Silva-Pereyra JF, Carreiras M. An ERP study of agreement features in Spanish. *Brain Research* 2007;1185:201–211. [PubMed: 17963736]
- Tiersma P. Local and general markedness. *Language* 1982;58:832–489.
- Walsh V. A theory of magnitude: Common cortical metrics of time, space and quantity. *Trends Cogn Sci* 2003;7:483–488. [PubMed: 14585444]
- Zago L, Pesenti M, Mellet E, Crivello F, Mazoyer B, Tzourio-Mazoyer N. Neural correlates of simple and complex mental calculation. *Neuroimage* 2001;13:314–327. [PubMed: 11162272]

Appendix

Determiner-noun pairs

El techo, el puerto, el saco, el hierro, el sábado, el premio, el pozo, el gobierno, los dedos, los lagos, los arros, los labios, los lavabos, los litros, los metros, los hornos, la costa, la saliva, la capa, la sopa, la rama, la cola, la pierna, la danza, las risas, las orejas, las letras, las navajas, las páginas, las espadas, las sierras, las reglas, la receta, la boda, la colcha, la plata, la luna, la aguja, la taza, la selva, las perlas, las abejas, las flautas, las gomas, las palas, las cuerdas, las iglesias, las alarmas, el pico, el banco, el piano, el circo, el martillo, el rebaño, el campo, el espejo, los hielos, los rayos, los besos, los torneos, los clavos, los frenos, los cocos, los votos, los hígados, los daños, los hombros, los arcos, los rollos, los caldos, los cercos, los sótanos, el trueno, el cráneo, el engaño, el mérito, el ahorro, el vidrio, el código, el trofeo, las vueltas, las gotas, las pausas, las ondas, las aldeas, las burlas, las minas, las retinas, la deuda, la pasta, la prisa, la misa, la duda, la urna, la rueda, la bandera.

Noun-Adjective pairs

cuerpo moreno, número mínimo, mundo justo, caso cierto, pelo rizado, fruto ácido, fuego flojo, llanto amargo, tiempos pasados, grupos serios, tipos listos, ídolos sabios, hilos finos, quesos agrios, huevos fritos, caminos anchos, cabeza menuda, lucha épica, vida sana, basura nociva, zona rara, suela gorda, piedra áspera, visita rápida, puertas seguras, mañanas futuras, cartas vacías, casas ajenas, ropas caras, citas médicas, blusas viejas, novelas baratas, pueblo bélico, estado sereno, libro único, lado turbio, ruido brusco, museo bello, vuelo largo, dibujo bonito, rostros pálidos, cambios súbitos, brazos tensos, juegos lícitos, autos rojos, discos buenos, barcos lentos, sonidos lejanos, guerra mítica, lengua humana, bolsa usada, droga tóxica, cena mala, tarea ardua, sábana blanca, sombra oscura, alturas idóneas, mantas raídas, cosas opacas, hojas pegadas, telas sosas, tumbas hondas, maderas blandas, plantas quietas, verano cálido, juicio severo, deseo mutuo, estilo maduro, piso pagado, cielo claro, plato llano, equipo entero, asuntos íntimos, cuellos rígidos, suelos planos, relatos amenos, muros duros, gestos lindos, anillos dorados, abrigos gruesos, fiesta loca, dama culta, copa tapada, mesa baja, tabla curva,

falda negra, papaya fresca, tierra remota, semanas previas, medidas válidas, cintas atadas, aguas tibias, comidas jugosas, etapas cortas, jarras llenas, figuras famosas.

Figure 1.
Left part: Axial, coronal and sagittal sections (left part) for the contrast violation > agreement. All contrasts depicted at $p < 0.001$ uncorrected. **Right part:** Graphs of contrast estimates and 90% confidence intervals. Number: Number violation > agreement; Gender: Gender violation > agreement.

Figure 2.
Left part: Axial, coronal and sagittal sections for the contrast Number violation > Gender violation in Determiner-Noun pairs. All contrasts depicted at $p < 0.001$ uncorrected. **Right part:** Graphs of contrast estimates and 90% confidence intervals for a right frontal middle area. Number: Number violation > agreement; Gender: Gender violation > agreement.

Table 1

Percentage of correct responses and mean decision times (in ms) (standard deviations within parentheses) for agreement, gender violation and number violation conditions in determiner-noun and noun-adjective pairs.

	Agreement	Gender	Number
Determiner-Noun	1113 (306) .99 (.02)	1276 (369) .98 (.03)	1282 (378) .97 (.02)
Noun-Adjective	1284 (353) .94 (.05)	1416 (429) .96 (.06)	1466(417) .90 (.08)

Table 2

Brain activation for type of violation (number and gender) and type of word pairs (determiner-noun and noun-adjective).

Region & Effect	x	y	z	Violation > Agreement		Number Violation > Agreement		Gender Violation > Agreement		Number > Gender		Gender > Number		Interaction type of violation by type of word	
				Z	Vx	Z	Vx	Z	Vx	Z	Vx	Z	Vx	Z	Vx
Premotor cortex	-40	4	36	4.2		3.2	121	4.6	166	n.s.		2.0		n.s.	
	-48	4	22	3.3		3.8	**	3.1	**	2.6		n.s.		n.s.	
Inferior frontal gyrus	-58	12	22	4.1		3.8	**	3.9	17	2.6		n.s.		n.s.	
Right intraparietal sulcus	40	-48	40	3.1	I	3.6	306	2.1		3.4	91	n.s.		3.4	11
	42	-50	48	n.s.		3.5	**	n.s.		3.9	**	n.s.		2.4	
Superior parietal gyrus	24	-70	46	2.6		4.4	**	n.s.		3.4	17	n.s.		3.0	

x,y,z = Coordinates of local maxima. **Z** = Z scores. **Vx** = Number of Voxels at $p < 0.001$ uncorrected. Z scores and cluster size are reported in **bold** if they are significant in height or extent at $p < .05$ after family wise correction for multiple comparisons across the whole brain. Those in *italics* were reported at $p < .05$ uncorrected. All others are significant at $p < .001$ uncorrected. When activation foci are part of the same cluster to that of the coordinates reported with the highest peak for the cluster in the upper cell, the corresponding cells below contain two asterisks (**). When the Z scores do not surpass the threshold of $p > .0001$ uncorrected, the corresponding cells for number of voxels are empty.

Table 3

Brain activation for number agreement violations only in determiner-noun pairs.

Region & Effect	x	y	z	Number Violation > Agreement Z	Number > Gender Z	Voxels
Right intraparietal sulcus	40	-48	40	3.9	4.7	727
	42	-50	48	3.5	4.6	***
Superior parietal gyrus	24	-70	46	3.9	4.4	***

x,y,z = Coordinates of local maxima. **Z** = Z scores. **Voxels** = Number of Voxels at $p < 0.001$ uncorrected. Z scores and cluster size are reported in **bold** if they are significant in height or extent at $p < .05$ after family wise correction for multiple comparisons across the whole brain. All others are significant at $p < .001$ uncorrected. When activation foci are part of the same cluster to that of the coordinates reported with the highest peak for the cluster in the upper cell, the corresponding cells below contain two asterisks (**). When the Z scores do not surpass the threshold of $p > .0001$ uncorrected, the corresponding cells for number of voxels are empty.