Published in final edited form as:

Nat Rev Microbiol. 2009 September ; 7(9): 677–683. doi:10.1038/nrmicro2184.

Transport of lipopolysaccharide across the cell envelope: the long road of discovery

Natividad Ruiz^{*}, Daniel Kahne^{‡,§}, and Thomas J. Silhavy^{*}

^{*}Department of Molecular Biology, Princeton University, Princeton, NJ 08544

[‡]Department of Chemistry and Chemical Biology, Harvard University, Cambridge, MA 02138

[§]Department of Biological Chemistry and Molecular Pharmacology, Harvard Medical School, Boston, MA 02115

Abstract

Intracellular lipid transport is poorly understood. Genetic studies to identify lipid-transport factors are complicated by the essentiality of many lipids, while biochemical and cell biology approaches aiming to determine localization and mechanisms of lipid transport are often challenged by the lack of adequate technology. Here we review the epic history of how different approaches, technological advances, and ingenuity contributed to the recent discovery of a multi-protein pathway that transports lipopolysaccharide across the envelope of Gram-negative bacteria.

One hundred and twenty-five years ago, Christian Gram developed a staining method that allows the differentiation of two major classes of bacteria, the Gram positives and the Gram negatives¹. This classification is widely used by microbiologists and the procedure still remains an important primary diagnostic tool in the clinic. Although it was assumed that the Gram stain reflected differences in the cellular envelope of these bacterial types, it took almost 100 years to understand the basis for the Gram reaction^{2,3}. This delay occurred mostly because discovering the composition and structure of the prototypical Gram -positive and -negative cell envelopes depended on the development of new technologies.

Although most progress in determining the composition, structure, and function of the different constituents of the bacterial envelope occurred during the 1940s–1970s, many researchers have been fascinated since the 1920s by a glycolipid that is present at the outer membrane (OM) of Gram-negative bacteria. This glycolipid, known as lipopolysaccharide (LPS) or endotoxin, is a potent inducer of the immune system and the causative agent of septic shock⁴. LPS is asymmetrically located at the outer leaflet of the OM^{5,6} and typically the molecule has three distinct regions: lipid A, a core oligosaccharide, and a long polysaccharide called the O-antigen⁷ (Fig. 1). Its location at the cell surface and its physicochemical properties make LPS primarily responsible for the barrier quality of the OM, which is crucial for the survival of many Gram-negative bacteria in many environments and the reason why it has been so hard to develop antibiotics against these organisms.⁸ Furthermore, LPS is thought to be essential in most Gram-negative bacteria.

What now might seem simple LPS biogenesis facts, such as its chemical composition, cellular localization and site of synthesis actually took decades to elucidate. As we describe in this *Timeline*, answers to these questions required years of study and the application of different approaches. This great body of work has recently led to the identification of the proteins that

are required to transport LPS from its site of synthesis, the inner membrane (IM), to the cell surface. The discovery of this transport pathway poses new questions and raises novel technological challenges that will guide future research in the field of bacterial envelope biogenesis.

Uncovering envelope structure

The realization in the 1920s that Gram-negative bacteria produce an endotoxin drove much of the biochemical work done on the bacterial envelope during the 1920s–50s. This early work was both facilitated and motivated by the potent immunogenic and toxic properties of LPS. Crude estimates of the chemical composition of the bacterial envelope were emerging but our understanding of its ultrastructure and the location of its constituents could not advance until the development of the electron microscope (EM) and ultracentrifugation. Likewise, although the general composition of LPS was known early on, LPS structure (Fig. 1) was not resolved until nuclear magnetic resonance was used in 1983 to elucidate the structure of lipid A.^{9–12}

An excellent review by Glauert and Thornley¹³ summarizes early work that probed the physical structure of the bacterial envelope by examining thin sections with EM. Using a variety of staining methods, the cell envelope of Gram-positive bacteria appeared to contain a cytoplasmic membrane surrounded by a thick but uniformly dense layer, which we now know is the cell-wall polymer known as peptidoglycan. In contrast, in Gram-negative bacteria the cytoplasmic membrane was surrounded with what appeared to be a triple-layered structure¹⁴ that in most of these early papers was referred to as the *cell wall*, a term that is very confusing to modern readers.

The difficulties encountered in resolving these outer layers likely reflect the fact that the OM is covalently attached to the peptidoglycan. However, as EM techniques improved, a clearer view of the Gram-negative cell envelope emerged. One of the most revealing EM studies was done by Bladen and Mergenhagen¹⁵ with the dental pathogen *Veillonella*. Their micrographs (Fig. 2) showed that the cell envelope contains three distinct layers: the *plasma membrane* that confines the cytoplasm, a structure the authors called a *solid membrane*, and an outermost unit membrane, which they called the *outer membrane*. The *solid membrane* was degraded by lysozyme and thus was the peptidoglycan layer. They also demonstrated that treatment with phenol-water extracted the OM and that this preparation contained LPS. In hindsight, this is a remarkable paper not only for the clarity of the EM images but also because it is the first to use the term *outer membrane*, although, in fairness, Murray¹⁶ reported similar observations for *E. coli* and referred to the outermost layer as the *outer unit membrane*. Nevertheless, as pointed out by Hiroshi Nikaido,¹⁷ the term *outer membrane* did not really come into common use until Mary Jane Osborn proved the membrane-like nature of this layer by analyzing its components.¹⁸

The lack of electron density between the IM and OM observed in EM micrographs suggested the existence of an aqueous, extracytoplasmic compartment. Evidence that this compartment contains a unique set of proteins came from Leon Heppel and colleagues, who developed methods such as cold osmotic shock and spheroplasting, which release proteins from this compartment without contamination by cytoplasmic proteins.¹⁹ Their work revealed that this extracytoplasmic compartment sequesters hydrolytic enzymes such as ribonuclease and alkaline phosphatase, and contains soluble substrate-binding proteins that are required for the active transport of certain small molecules. The concentration of macromolecules in this compartment is high, and it is, in fact, more viscous than the cytoplasm.²⁰ This compartment is now called the *periplasm*, a term apparently coined by Peter Mitchell in 1961, although it did not come into common use until much later. In fact, in the aforementioned review,¹⁹ Heppel does not use this word. Together, these pioneering works paved the way to many other studies

that have provided us with the current detailed picture of the structure and composition of the Gram-negative cell envelope (Fig. 3).

LPS localization

Armed with the knowledge derived from EM of the physical structure of the Gram-negative cell envelope, Miura and Mizushima developed a method to separate the OM from the IM using sucrose density-gradient centrifugation.²¹ This technological advance allowed Mary Jane Osborn and collaborators to determine in 1972 the composition of the OM and to conclude that LPS is indeed found in this organelle¹⁸. Although this method was also crucial for those studies discussed below probing the localization of LPS synthesis and the transport of LPS across the cell envelope, it did not resolve how OM components are arranged.

Indeed, demonstration that LPS is localized at the outer leaflet of the OM was accomplished in 1975 by Muhlradt and Golecki⁵ using immune-EM. A year later, the study by Kamio and Nikaido showing that the outer leaflet of the OM of *Salmonella enteritica* is devoid of its most abundant phospholipid, phosphatidylethanolamine,⁶ further confirmed the lipid asymmetry of the OM. These findings, together with the research described in the following section, were the foundation for our understanding of the barrier quality of the OM.

Function of LPS

Like most lipid bilayers, the OM is impermeant to large hydrophilic molecules, but it also has the unusual property of preventing the rapid diffusion of small hydrophobic molecules.²² This is why Gram-negative bacteria are generally more resistant to hydrophobic antibiotics and detergents than their Gram-positive counterparts. Of course, the OM still allows the exchange of solutes with its environment. Small hydrophilic molecules of up to approximately 600 Da diffuse through non-specific OM protein channels called porins, and certain large molecules such as vitamins and iron chelates can pass through specialized channels.

Probing the barrier function of the OM became possible when in 1958 MacGregor and Elliker realized that ethylenediaminetetraacetic acid (EDTA) increases the sensitivity of *Pseudomonas* to quaternary amines.²³ In the 1960–70s, Loretta Leive demonstrated that this was because EDTA promotes the entry of several compounds into Gram-negative cells.^{24, 25} However, understanding how this compound increases permeability was not possible until biochemical and genetic studies proved, respectively, that EDTA releases LPS from the cell surface²⁶ and that certain mutants producing truncated forms of LPS (known as rough mutants) are more permeable than wild type.^{27, 28}

Most recent physical studies on the barrier quality of the OM have been conducted in enteric bacteria and are discussed in detail by Nikaido.²² They have revealed that the main feature responsible for the impermeability of the OM is the low fluidity conferred by the high degree of packing of LPS molecules into the outer leaflet. This is accomplished by the following properties of LPS. LPS has a higher number of fatty acid chains per molecule than phospholipids, resulting in a higher number of hydrophobic interactions between neighboring molecules. In the *Enterobacteriaceae*, lateral interactions between LPS molecules are also facilitated by hydrogen bonds and by cations (Mg^{2+} and Ca^{2+}) that neutralize the negative charges of the phosphate groups present in lipid A. Furthermore, in the enteric LPS, the fatty acid chains in lipid A are saturated, which allows tighter packing. Consequently, altering either the structure or the cellular location of LPS compromises the barrier function of the OM. We now understand that the aforementioned EDTA treatments chelate the cations that are intercalated between LPS molecules, resulting in the repulsion and release of LPS molecules into the medium. Somehow, LPS release causes the translocation of phospholipids to the outer leaflet, which increases the permeability of the OM as it now contains phospholipid bilayers.

Where is LPS synthesized?

Separation of the inner and outer membranes using sucrose density-gradient ultracentrifugation allowed Osborn and collaborators to prove not only that LPS is located at the OM, but also that LPS is synthesized in the IM and that translocation of LPS to the OM is unidirectional.^{18, 29} Those studies were seminal in the field of LPS biogenesis, since they began sketching the pathway of LPS transport across the cell envelope.

During the next decades, the biosynthetic pathway of LPS (often referred to as the *Raetz pathway*), its participating enzymes and their location were mainly elucidated by Christian Raetz and collaborators and those findings are discussed in an excellent review by Raetz and Whitfield.⁷ From those studies we know that all components of LPS (lipid A, core, and O-antigen) are synthesized at the inner leaflet of the IM where lipid A and core are ligated to form what is known as rough LPS. Rough LPS and the O-antigen are independently flipped across the IM and ligated at the outer leaflet of the IM. Notably, *E. coli* K-12 strains only synthesize rough LPS because they carry a mutation in the O-antigen biosynthetic pathway³⁰.

Once we knew that LPS is synthesized at the IM, it became clear that LPS molecules are first translocated across the IM, then transported across the periplasm, and ultimately delivered to the cell surface. At the same time, many questions emerged. Were there specific proteins required for LPS transport? Was LPS transport to the OM coupled to the transport of OM proteins? Were the controversial zones of contact between the IM and the OM bridges that mediated LPS transport across the cell envelope? What is the source of energy for LPS assembly at the OM, an environment devoid of ATP? We now know the answer to some of these questions. For example, Rothfield and Pearlman-Hothencz determined early on that inhibition of protein synthesis did not prevent either LPS synthesis or transport to the OM for hours,³¹ implying that OMPs and LPS were not transported to and assembled at the OM as a unit. Most importantly, employing a variety of genetic, biochemical, and bioinformatic approaches, a number of laboratories have recently discovered the proteins that transport LPS across the cell envelope, and a model for LPS biogenesis has emerged.

Current model for LPS biogenesis

For clarity, we will first summarize the current model for LPS biogenesis (Figure 4). This model was based on the fact that every envelope compartment has at least one factor required for LPS transport and from the genetic, biochemical, and bioinformatic studies described below. After its synthesis, rough LPS is translocated across the IM by the ABC (ATP-binding cassette) transporter MsbA. The O-antigen is independently translocated across the IM usually by the Wzx flippase as an undecaprenol-phosphate-linked molecule; however, in some *E. coli* strains, the Wzm-Wzt ABC transporter translocates their undecaprenol-linked O-antigen. The WaaL ligase then adds the O-antigen to rough LPS at the periplasmic face of the IM.⁷ Next, LPS (or rough LPS in the case of *E. coli* K-12 strains) interacts with the Lpt (LPS transport) pathway where LPS is first extracted from the IM by an ABC transporter composed of LptBFG and likely the bitopic membrane protein LptC. For LPS to cross the periplasm and be assembled at the OM, the periplasmic protein LptA, the OM lipoprotein LptE, and the β -barrel OM protein LptD are required. As we discuss below, although we believe that every essential Lpt factor has been identified, we still lack a fundamental understanding of the mechanism of LPS transport.

Flipping LPS across the IM

MsbA translocates LPS across the IM. Perhaps owing to its early identification and that it belongs to the ABC transporter superfamily, MsbA has been the subject of many biophysical and protein biochemical studies during the last decade; as a result, MsbA is the best understood

protein involved LPS transport. We even have four x-ray structures of MsbA trapped in different conformations!³² However, it all started with a simple genetic selection.

Karow and Georgopoulos identified *msbA* in 1993 as a multicopy suppressor of the temperature sensitivity conferred by *htrB* null alleles.³³ Primary sequence and genetic analyses revealed that MsbA likely was an essential ABC transporter in *E. coli* but the lack of knowledge of HtrB (later renamed LpxL) function obscured the identity of the transported substrate. Indeed, the demonstration in 1996 by Raetz and collaborators that LpxL is a lipid A laurate acetyltransferase³⁴ led Polissi and Georgopoulos³⁵ to show that same year that LPS transport to the OM is defective in an *htrB* mutant under non-permissive conditions. More importantly, after demonstrating that in multicopy *msbA* alleviates this defect and that loss of MsbA causes accumulation of LPS species at the IM, they proposed that MsbA was a translocator of either LPS or one of its precursors.

In an effort to demonstrate a role for MsbA in the transport of LPS across the IM, Raetz and collaborators made an important contribution by developing a biochemical assay to probe LPS transport system *in vivo*.³⁶ In this assay, lipid A and its phosphorylated precursors are radiolabeled and these species are then extracted and analyzed by thin-layer chromatography. Using this assay in combination with sucrose-gradient fractionations to separate the IM and OM, Zhou *et al.*³⁶ showed that under non-permissive conditions, *htrB* mutants accumulate tetra-acylated lipid A (lipid IV_A) species at the IM and that overexpressing MsbA does not correct this defect in the maturation of LPS; instead, overexpressing MsbA suppresses by increasing the *transport* of the immature lipid IV_A species to the OM. They also demonstrated that cells depleted of MsbA accumulate LPS species containing mature hexa-acylated lipid A at the IM.

Proof that this accumulation of the hexa-acylated lipid A occurs at the inner leaflet of the IM was accomplished by exploiting the fact that LPS undergoes chemical modification at different locations in the cell.³⁷ Specifically, Doerrler *et al.*³⁸ showed that in cells deficient in MsbA function, the LPS that accumulates at the IM is not accessible to modifications by ArnT and EptA/PmrC, two enzymes that catalyze respectively the transfer of aminoarabinose and phosphoethanolamine to lipid A at the periplasmic side of the IM.^{37, 39–41} Later we will explain a similar strategy used to determine whether LPS reaches the cell surface.

Thus, genetic and biochemical *in vivo* studies agree that MsbA translocates LPS across the IM. Crystal structure analyses reveal that MsbA forms a dimer where each monomer provides six transmembrane (TM) helices and half of a cytoplasmic nucleotide-binding domain (NBD)³². The dimer hydrolyzes ATP and undergoes large conformational changes thought to be required for transport. Although reconstitution of MsbA's flippase activity awaits, *in vitro* biochemical studies have shown that hexa-acylated lipid A species and lipophilic drugs stimulate MsbA's ATPase activity.^{42, 43} In fact, the Sharom laboratory has recently shown that the MsbA dimer contains two high-affinity substrate-binding sites capable of binding lipid A and amphipathic drugs⁴⁴ and has suggested that MsbA may function as both a lipid flippase and a multidrug transporter.^{43, 44}

From the IM to the surface

The Lpt pathway transports LPS from the IM to the cell surface (Fig. 4). Most of the Lpt factors have been identified in the last three years but the first *lpt* mutations were isolated in *lptD* (formerly *imp* or *ostA*) in 1989. Benson and collaborators designed a genetic selection for mutants which allowed maltodextrins to pass across their OM in the absence of the maltodextrin-specific channel LamB.⁴⁵ The *imp* mutations increased membrane permeability not only to maltodextrins but also to many hydrophobic and hydrophilic antibiotics, implying that the overall barrier function of the OM was compromised.

Although those early studies suggested the essential nature of *lptD* (*imp*), it took more than a decade to uncover LptD's role in OM biogenesis. New attention to *lptD* was sparked by two findings. First, *lptD* is located immediately upstream of *surA*, a gene that encodes a major periplasmic chaperone involved in the assembly of OMPs.⁴⁶ Second, both *lptD* and *surA* are in an operon regulated by σ^E envelope stress response.⁴⁷ The σ^E response monitors the status of the cell envelope and, when activated, increases the production of envelope-biogenesis factors.⁴⁸ With this information in hand, Braun and Silhavy showed that LptD is an essential OMP required for the biogenesis of the *E. coli* cell envelope.⁴⁹ Proof of its role in LPS assembly at the surface came later from work in *Neisseria meningitidis* by Tommassen and collaborators,⁵⁰ who showed that in mutants lacking the neisserial LptD ortholog, LPS does not reach the cell surface since it is not accessible to either cleavage by extracellular neuraminidase or modification by the OM enzyme PagL. Notably, in *N. meningitidis* neither LptD nor LPS are essential^{50, 51}; likewise, LptD is also not essential in *Helicobacter pylori*.⁵²

To identify additional Lpt factors, Kahne and collaborators undertook a biochemical approach. Using affinity purification of an epitope-tagged LptD, they discovered that in *E. coli*, LptD is in a complex with the essential OM lipoprotein LptE (formerly RlpB).⁵³ They also showed LPS molecules synthesized after depletion of either LptD or LptE are not accessible to modification by PagP, an OM enzyme that palmitoylates LPS at the outer leaflet of the OM.^{37, 54} Therefore, both LptD and LptE are required for LPS transport to the cell surface in *E. coli*.

The remaining Lpt factors were identified by different approaches. Polissi and collaborators found *lptA* and *lptB* (formerly *yhbN* and *yhbG*, respectively) in a genetic screen searching for essential genes.⁵⁵ Clues that these genes could be involved in LPS biogenesis first came from the realization that they are part of a highly conserved locus that includes genes specifying enzymes involved in the synthesis of LPS core.⁵⁶ This led Polissi and collaborators to demonstrate using sucrose density-gradient centrifugation that depletion of either LptA or LptB halts the transport of LPS across the cell envelope, causing the accumulation of LPS at the IM.⁵⁷ Because sequence analysis predicts that LptB is an IM-associated⁵⁸ cytoplasmic protein with an NBD typical of ABC transporters and LptA is a periplasmic proteins, they proposed the following model. LptB hydrolyzes ATP to provide energy to an ABC transporter that extracts LPS from the IM, while LptA facilitates LPS transport across the periplasm.⁵⁷ ABC transporters require an IM component, therefore, they also suggested that the IM protein encoded by the essential gene *yrbK*⁵⁶ might play this role since the 3' end of *yrbK* overlaps with the 5' end of *lptA*. Such chromosomal arrangement is usually indicative of genes specifying related functions.

Since then, Polissi and collaborators have proven that YrbK (renamed LptC) is required for the extraction of LPS from the IM.⁵⁹ However, LptC is a bitopic protein and typically the integral components of ABC transporters are composed of either one IM protein with 12 TM domains or two with 6-TM domains each.⁶⁰ Therefore, it was likely that we were missing at least one more Lpt factor. Indeed, using a reductionist bioinformatic approach designed to identify OM-biogenesis factors, Ruiz and collaborators recently found the 6-TM domain IM proteins LptF and LptG (formerly YjgP and YjgQ).⁶¹ Their approach took advantage of the high degree of conservation of OM-biogenesis factors among Gram negative bacteria, including endosymbionts. They reasoned that even though endosymbionts have lost many genes during their adaptation to intracellular life, these intracellular bacteria must still possess those encoding factors essential for OM-biogenesis. Thereby, they identified LptF and LptG as the missing components of the ABC transporter that together with LptB and, possibly LptC, extracts LPS from the outer leaflet of the IM so that it can be transported to the OM. This bioinformatic analysis also suggests that all essential Lpt factors of *E. coli* have been identified.

Lpt pathway order

In *E. coli*, every cellular compartment contains proteins required for LPS transport. Phenotypes exhibited by cells depleted of MsbA are distinct from those observed upon depletion of Lpt factors: LPS is not flipped across the IM in the absence of MsbA but it is in the absence of Lpt factors. However, disruption of the pathway at any step after MsbA leads to indistinguishable phenotypes: in all cases, LPS accumulates in the outer leaflet of the IM.^{59, 61, 62} Therefore, we cannot conduct epistasis analysis to determine exact pathway order. We can only rely on the information about the cellular location of these proteins to predict temporal order in the pathway as follows: MsbA, LptBCFG, LptA, and finally LptDE.

Future directions

The central question of LPS biogenesis at the moment is how mechanistically LPS is transported across the aqueous periplasmic compartment and inserted in functional form in its correct spatial location, the outer leaflet of the OM. In a very real sense, the mechanism by which the Lpt proteins accomplish transport is a stereochemical problem of the highest order. Since it involves proteins in two different membranes that are separated by an aqueous compartment, it is not even clear what kinds of tools will be needed to study this problem.

LPS cannot traverse the aqueous periplasm by diffusion owing to its amphipathic nature. Thus, LPS is somehow assisted across the periplasm and through the peptidoglycan layer. Two general models have been proposed for how LPS crosses the periplasm to reach the OM (Fig. 4).⁴⁶ One model posits that LPS is bound by a soluble periplasmic chaperone protein that diffuses across the periplasmic space to a docking site at the inner leaflet of the OM. Specifically, according to this model, LPS transport would resemble OM lipoprotein transport in that a soluble chaperone, LolA, binds lipoproteins after they have been extracted from the IM by the ABC transporter LolCDE and carries them to an assembly site, LolB, in the OM.⁶³ In fact, LptA has recently been shown to bind LPS *in vitro*, an observation that the authors suggest is consistent with the chaperone model.⁶⁴ An alternate model posits that there is some sort of proteinaceous bridge that physically connects the IM and OM and allows for direct efflux of LPS to the cell surface. It has been demonstrated that spheroplasts, which are effectively drained of periplasmic contents, are still able to transport newly-synthesized LPS to the outer leaflet of the OM, implying that all components required for LPS transport remain stably associated with the spheroplasts.⁶⁵ In addition, the recently solved crystal structure of LptA suggests that the protein is capable of forming fibrils composed of LptA molecules oriented in a head-to-tail fashion, and thus could physically bridge the gap between the two membranes.⁶⁶ It is important to note, however, that none of the evidence presented for one model can rule out the other model entirely. More biochemical experiments need to be done to discern between these two transport models.

Even though we still do not understand how the Lpt system functions, we have clearly made great advances in understanding LPS assembly in the recent years. We hope that the scientific history presented here illustrates how this endeavor was only possible because of the development of new technologies and the use of different approaches throughout decades. We foresee that the same level of creativity and diversity of scientific approaches will be needed to explore and elucidate the molecular mechanisms of the multi-component, inter-membrane Lpt transport system.

TIMELINE

1964: Envelope composition and structure:

EM micrographs clearly showing Gram-negative envelope structure and first use of the term *outer* membrane.¹⁵

1965: Demonstration of the barrier function of the OM and role of LPS.^{24–26}

1967: Envelope composition and structure:

Demonstration that the periplasm contains proteins.¹⁹

1968: Envelope composition and structure:

Development of density gradient centrifugation procedure to separate the IM and OM.²¹

1972: LPS synthesis:

Demonstration that LPS is synthesized at the IM.²⁹

Envelope composition and structure:

Analysis of the composition of the IM and OM and demonstration that LPS is at the OM.¹⁸

1975–1976: Envelope composition and structure:

Demonstration of OM lipid asymmetry and localization of LPS at the cell surface.^{5, 6}

1972–1990s: LPS synthesis:

Discovery and characterization of enzymes involved in LPS synthesis.⁷

1981–1983: LPS synthesis:

LPS precursor lipid X is discovered and its structure is solved, providing the pivotal clue to the biosynthetic pathway.^{69, 70}

1983: Complete LPS structure is resolved.^{9–12}

1993: LPS transport:

MsbA is discovered but function remains unknown.³³

1996–98: LPS transport:

Demonstration that MsbA is required for flipping LPS across the IM.^{35, 36}

1989: LPS transport:

Identification of the first *lptD* mutants.⁴⁵

2002: LPS transport:

Demonstration that LptD is an essential OM biogenesis factor in *E. coli*.⁴⁹

2004: LPS transport:

Demonstration that neisserial Imp (LptD ortholog) is required for LPS transport.⁵⁰

Identification of *lptAB* (*yhbNG*) as essential genes.⁵⁵

2006: LPS transport:

LptE is identified and shown to be in a complex with LptD that is required for LPS transport.⁵³

2007: LPS transport:

Demonstration that LptAB are required for LPS transport.⁵⁷

MsbA crystal structure is solved.³²

2008: LPS transport:

LptFG are identified and shown to be required for LPS transport.⁶¹

Demonstration that LptC is required for LPS transport.⁵⁹

LptA is crystallized⁶⁶ and shown to bind LPS.⁶⁴

Acknowledgments

This work was supported by National Institute of General Medical Sciences Grants GM34821 (to T.J.S.) and GM66174 (to D.K.).

REFERENCES

1. Gram HCJ. Über die isolirte Färbung der Schizomyceten in Schnitt- und Trockenpräparaten. Fortschritte der Medizin 1884;2:185–189.
2. Beveridge TJ, Davies JA. Cellular responses of *Bacillus subtilis* and *Escherichia coli* to the Gram stain. J Bacteriol 1983;156:846–858. [PubMed: 6195148]
3. Davies JA, Anderson GK, Beveridge TJ, Clark HC. Chemical mechanism of the Gram stain and synthesis of a new electron-opaque marker for electron microscopy which replaces the iodine mordant of the stain. J Bacteriol 1983;156:837–845. [PubMed: 6195147]
4. Miller SI, Ernst RK, Bader MW. LPS, TLR4 and infectious disease diversity. Nat Rev Microbiol 2005;3:36–46. [PubMed: 15608698]
5. Muhlradt PF, Golecki JR. Asymmetrical distribution and artifactual reorientation of lipopolysaccharide in the outer membrane bilayer of *Salmonella typhimurium*. Eur J Biochem 1975;51:343–352. [PubMed: 807474]
6. Kamio Y, Nikaido H. Outer membrane of *Salmonella typhimurium*: accessibility of phospholipid head groups to phospholipase c and cyanogen bromide activated dextran in the external medium. Biochemistry 1976;15:2561–2570. [PubMed: 820368]
7. Raetz CR, Whitfield C. Lipopolysaccharide endotoxins. Annu Rev Biochem 2002;71:635–700. [PubMed: 12045108]
8. Delcour AH. Outer membrane permeability and antibiotic resistance. Biochim Biophys Acta. 2008
9. Qureshi N, Takayama K, Heller D, Fenselau C. Position of ester groups in the lipid A backbone of lipopolysaccharides obtained from *Salmonella typhimurium*. J Biol Chem 1983;258:12947–12951. [PubMed: 6355099]
10. Strain SM, Fesik SW, Armitage IM. Structure and metal-binding properties of lipopolysaccharides from heptoseless mutants of *Escherichia coli* studied by ¹³C and ³¹P nuclear magnetic resonance. J Biol Chem 1983;258:13466–13477. [PubMed: 6358205]
11. Strain SM, Fesik SW, Armitage IM. Characterization of lipopolysaccharide from a heptoseless mutant of *Escherichia coli* by carbon 13 nuclear magnetic resonance. J Biol Chem 1983;258:2906–2910. [PubMed: 6338007]
12. Takayama K, Qureshi N, Mascagni P. Complete structure of lipid A obtained from the lipopolysaccharides of the heptoseless mutant of *Salmonella typhimurium*. J Biol Chem 1983;258:12801–12803. [PubMed: 6355095]
13. Glauert AM, Thornley MJ. The topography of the bacterial cell wall. Annu Rev Microbiol 1969;23:159–198. [PubMed: 4186529]
14. Kellenberger E, Ryter A. Cell wall and cytoplasmic membrane of *Escherichia coli*. J Biophys Biochem Cytol 1958;4:323–326. [PubMed: 13549505]
15. Bladen HA, Mergenhagen SE. Ultrastructure of *Veillonella* and Morphological Correlation of an Outer Membrane with Particles Associated with Endotoxic Activity. J Bacteriol 1964;88:1482–1492. [PubMed: 14234809]
16. Murray RG. On the cell wall structure of *Spirillum serpens*. Can J Microbiol 1963;9:381–392.
17. Nikaido, H. The periplasm. Ehrmann, M., editor. Washington, D.C.: ASM Press; 2007.

18. Osborn MJ, Gander JE, Parisi E, Carson J. Mechanism of assembly of the outer membrane of *Salmonella typhimurium*. Isolation and characterization of cytoplasmic and outer membrane. *J Biol Chem* 1972;247:3962–3972. [PubMed: 4555955]
19. Heppel LA. Selective release of enzymes from bacteria. *Science* 1967;156:1451–1455. [PubMed: 4304946]
20. Mullineaux CW, Nenninger A, Ray N, Robinson C. Diffusion of green fluorescent protein in three cell environments in *Escherichia coli*. *J Bacteriol* 2006;188:3442–3448. [PubMed: 16672597]
21. Miura T, Mizushima S. Separation by density gradient centrifugation of two types of membranes from spheroplast membrane of *Escherichia coli* K12. *Biochim Biophys Acta* 1968;150:159–161. [PubMed: 4868245]
22. Nikaido H. Molecular basis of bacterial outer membrane permeability revisited. *Microbiol Mol Biol Rev* 2003;67:593–656. [PubMed: 14665678]
23. Macgregor DR, Elliker PR. A comparison of some properties of strains of *Pseudomonas aeruginosa* sensitive and resistant to quaternary ammonium compounds. *Can J Microbiol* 1958;4:499–503. [PubMed: 13573235]
24. Leive L. A Nonspecific Increase in Permeability in *Escherichia Coli* Produced by Edta. *Proc Natl Acad Sci U S A* 1965;53:745–750. [PubMed: 14324530]
25. Leive L. Actinomycin Sensitivity in *Escherichia Coli* Produced by Edta. *Biochem Biophys Res Commun* 1965;18:13–17. [PubMed: 14265747]
26. Leive L. Release of lipopolysaccharide by EDTA treatment of *E. coli*. *Biochem Biophys Res Commun* 1965;21:290–296. [PubMed: 4159978]
27. Leive L. The barrier function of the gram-negative envelope. *Ann N Y Acad Sci* 1974;235:109–129. [PubMed: 4212391]
28. Tamaki S, Sato T, Matsushashi M. Role of lipopolysaccharides in antibiotic resistance and bacteriophage adsorption of *Escherichia coli* K-12. *J Bacteriol* 1971;105:968–975. [PubMed: 4926688]
29. Osborn MJ, Gander JE, Parisi E. Mechanism of assembly of the outer membrane of *Salmonella typhimurium*. Site of synthesis of lipopolysaccharide. *J Biol Chem* 1972;247:3973–3986. [PubMed: 4624447]
30. Liu D, Reeves PR. *Escherichia coli* K12 regains its O antigen. *Microbiology* 1994;140(Pt 1):49–57. [PubMed: 7512872]
31. Rothfield L, Pearlman-Kothencz M. Synthesis and assembly of bacterial membrane components. A lipopolysaccharide-phospholipid-protein complex excreted by living bacteria. *J Mol Biol* 1969;44:477–492. [PubMed: 4899474]
32. Ward A, Reyes CL, Yu J, Roth CB, Chang G. Flexibility in the ABC transporter MsbA: Alternating access with a twist. *Proc Natl Acad Sci U S A* 2007;104:19005–19010. [PubMed: 18024585]
33. Karow M, Georgopoulos C. The essential *Escherichia coli* *msbA* gene, a multicopy suppressor of null mutations in the *htrB* gene, is related to the universally conserved family of ATP-dependent translocators. *Mol Microbiol* 1993;7:69–79. [PubMed: 8094880]
34. Clementz T, Bednarski JJ, Raetz CR. Function of the *htrB* high temperature requirement gene of *Escherichia coli* in the acylation of lipid A: *HtrB* catalyzed incorporation of laurate. *J Biol Chem* 1996;271:12095–12102. [PubMed: 8662613]
35. Polissi A, Georgopoulos C. Mutational analysis and properties of the *msbA* gene of *Escherichia coli*, coding for an essential ABC family transporter. *Mol Microbiol* 1996;20:1221–1233. [PubMed: 8809774]
36. Zhou Z, White KA, Polissi A, Georgopoulos C, Raetz CR. Function of *Escherichia coli* MsbA, an essential ABC family transporter, in lipid A and phospholipid biosynthesis. *J Biol Chem* 1998;273:12466–12475. [PubMed: 9575204]
37. Raetz CR, Reynolds CM, Trent MS, Bishop RE. Lipid A modification systems in gram-negative bacteria. *Annu Rev Biochem* 2007;76:295–329. [PubMed: 17362200]
38. Doerrler WT, Gibbons HS, Raetz CR. MsbA-dependent translocation of lipids across the inner membrane of *Escherichia coli*. *J Biol Chem* 2004;279:45102–45109. [PubMed: 15304478]

39. Lee H, Hsu FF, Turk J, Groisman EA. The PmrA-regulated pmrC gene mediates phosphoethanolamine modification of lipid A and polymyxin resistance in *Salmonella enterica*. *J Bacteriol* 2004;186:4124–4133. [PubMed: 15205413]
40. Trent MS, et al. Accumulation of a polyisoprene-linked amino sugar in polymyxin-resistant *Salmonella typhimurium* and *Escherichia coli*: structural characterization and transfer to lipid A in the periplasm. *J Biol Chem* 2001;276:43132–43144. [PubMed: 11535605]
41. Trent MS, Ribeiro AA, Lin S, Cotter RJ, Raetz CR. An inner membrane enzyme in *Salmonella* and *Escherichia coli* that transfers 4-amino-4-deoxy-L-arabinose to lipid A: induction on polymyxin-resistant mutants and role of a novel lipid-linked donor. *J Biol Chem* 2001;276:43122–43131. [PubMed: 11535604]
42. Doerrler WT, Raetz CR. ATPase activity of the MsbA lipid flippase of *Escherichia coli*. *J Biol Chem* 2002;277:36697–36705. [PubMed: 12119303]
43. Eckford PD, Sharom FJ. Functional characterization of *Escherichia coli* MsbA: interaction with nucleotides and substrates. *J Biol Chem* 2008;283:12840–12850. [PubMed: 18344567]
44. Siarheyeva A, Sharom FJ. The ABC transporter MsbA interacts with lipid A and amphipathic drugs at different sites. *Biochem J*. 2009
45. Sampson BA, Misra R, Benson SA. Identification and characterization of a new gene of *Escherichia coli* K-12 involved in outer membrane permeability. *Genetics* 1989;122:491–501. [PubMed: 2547691]
46. Bos MP, Robert V, Tommassen J. Biogenesis of the gram-negative bacterial outer membrane. *Annu Rev Microbiol* 2007;61:191–214. [PubMed: 17506684]
47. Dartigalongue C, Missiakas D, Raina S. Characterization of the *Escherichia coli* sigma E regulon. *J Biol Chem* 2001;276:20866–20875. [PubMed: 11274153]
48. MacRitchie DM, Buelow DR, Price NL, Raivio TL. Two-component signaling and gram negative envelope stress response systems. *Adv Exp Med Biol* 2008;631:80–110. [PubMed: 18792683]
49. Braun M, Silhavy TJ. Imp/OstA is required for cell envelope biogenesis in *Escherichia coli*. *Mol Microbiol* 2002;45:1289–1302. [PubMed: 12207697]
50. Bos MP, Tefsen B, Geurtsen J, Tommassen J. Identification of an outer membrane protein required for the transport of lipopolysaccharide to the bacterial cell surface. *Proc Natl Acad Sci U S A* 2004;101:9417–9422. [PubMed: 15192148]
51. Steeghs L, et al. Meningitis bacterium is viable without endotoxin. *Nature* 1998;392:449–450. [PubMed: 9548250]
52. Chiu HC, Lin TL, Wang JT. Identification and characterization of an organic solvent tolerance gene in *Helicobacter pylori*. *Helicobacter* 2007;12:74–81. [PubMed: 17241305]
53. Wu T, et al. Identification of a protein complex that assembles lipopolysaccharide in the outer membrane of *Escherichia coli*. *Proc Natl Acad Sci U S A* 2006;103:11754–11759. [PubMed: 16861298]
54. Bishop RE, et al. Transfer of palmitate from phospholipids to lipid A in outer membranes of gram-negative bacteria. *Embo J* 2000;19:5071–5080. [PubMed: 11013210]
55. Serina S, et al. Scanning the *Escherichia coli* chromosome by random transposon mutagenesis and multiple phenotypic screening. *Res Microbiol* 2004;155:692–701. [PubMed: 15380559]
56. Sperandio P, Pozzi C, Deho G, Polissi A. Non-essential KDO biosynthesis and new essential cell envelope biogenesis genes in the *Escherichia coli* yrbG-yhbG locus. *Res Microbiol* 2006;157:547–558. [PubMed: 16765569]
57. Sperandio P, et al. Characterization of lptA and lptB, two essential genes implicated in lipopolysaccharide transport to the outer membrane of *Escherichia coli*. *J Bacteriol* 2007;189:244–253. [PubMed: 17056748]
58. Stenberg F, et al. Protein complexes of the *Escherichia coli* cell envelope. *J Biol Chem* 2005;280:34409–34419. [PubMed: 16079137]
59. Sperandio P, et al. Functional analysis of the protein machinery required for transport of lipopolysaccharide to the outer membrane of *Escherichia coli*. *J Bacteriol* 2008;190:4460–4469. [PubMed: 18424520]
60. Linton KJ, Higgins CF. Structure and function of ABC transporters: the ATP switch provides flexible control. *Pflugers Arch* 2007;453:555–567. [PubMed: 16937116]

61. Ruiz N, Gronenberg LS, Kahne D, Silhavy TJ. Identification of two inner-membrane proteins required for the transport of lipopolysaccharide to the outer membrane of *Escherichia coli*. *Proc Natl Acad Sci U S A* 105;2008:5537–5542.
62. Ma B, Reynolds CM, Raetz CR. Periplasmic orientation of nascent lipid A in the inner membrane of an *Escherichia coli* LptA mutant. *Proc Natl Acad Sci U S A* 2008;105:13823–13828. [PubMed: 18768814]
63. Tokuda H, Matsuyama S. Sorting of lipoproteins to the outer membrane in *E. coli*. *Biochim Biophys Acta* 2004;1694:IN1–IN9. [PubMed: 15672528]
64. Tran AX, Trent MS, Whitfield C. The LptA protein of *Escherichia coli* is a periplasmic lipid A-binding protein involved in the lipopolysaccharide export pathway. *J Biol Chem* 2008;283:20342–20349. [PubMed: 18480051]
65. Tefsén B, Geurtsen J, Beckers F, Tommassen J, de Cock H. Lipopolysaccharide transport to the bacterial outer membrane in spheroplasts. *J Biol Chem* 2005;280:4504–4509. [PubMed: 15576375]
66. Suits MD, Sperandio P, Deho G, Polissi A, Jia Z. Novel structure of the conserved gram-negative lipopolysaccharide transport protein A and mutagenesis analysis. *J Mol Biol* 2008;380:476–488. [PubMed: 18534617]
67. Trent MS, Stead CM, Tran AX, Hankins JV. Diversity of endotoxin and its impact on pathogenesis. *J Endotoxin Res* 2006;12:205–223. [PubMed: 16953973]
68. Meredith TC, et al. Modification of lipopolysaccharide with colanic acid (M-antigen) repeats in *Escherichia coli*. *J Biol Chem* 2007;282:7790–7798. [PubMed: 17227761]
69. Nishijima M, Raetz CR. Characterization of two membrane-associated glycolipids from an *Escherichia coli* mutant deficient in phosphatidylglycerol. *J Biol Chem* 1981;256:10690–10696. [PubMed: 7026561]
70. Takayama K, et al. Fatty acyl derivatives of glucosamine 1-phosphate in *Escherichia coli* and their relation to lipid A. Complete structure of A diacyl GlcN-1-P found in a phosphatidylglycerol-deficient mutant. *J Biol Chem* 1983;258:7379–7385. [PubMed: 6345522]

Figure 1. Structure of *E. coli* LPS

The structure of *E. coli* LPS with a K-12 core region is shown. The structure of LPS of several species has been solved and it is clear that it is very diverse.⁶⁷ Moreover, one single species can further modify its typical LPS structure in response to environmental signals and defects in envelope conditions.^{37, 59, 61, 68} Abbreviations: Kdo, 3-deoxy-D-manno-oct-2-ulonic acid; Hep, L-glycero-D-manno-heptose; Glu, D-glucose; Gal, D-galactose; EtN, ethanolamine; P, phosphate.

Figure 2. Electron micrograph of *Veillonella* illustrating the ultrastructure of the Gram-negative cell envelope

Bladen and Mergenhagen published this micrograph in 1964¹⁵ and they labeled the different structures as the outer membrane (OM), the solid membrane (SM), which is known nowadays as the peptidoglycan, and the plasma membrane (PM), which we refer to as the inner or cytoplasmic membrane. This figure was reproduced with the permission of the *Journal of Bacteriology* (pending).

Figure 3. Structure of the Gram-negative cell envelope

In Gram-negative bacteria, the cytoplasm is surrounded by the IM, a phospholipid (PL) bilayer that also contains proteins. There are two types of proteins in the IM: integral IM proteins, which span membrane through α -helical TM domains, and IM lipoproteins, which are anchored to the outer leaflet through a lipid moiety. The periplasm is the aqueous compartment bounded by the IM and the OM; it contains soluble proteins and the peptidoglycan layer. The OM is anchored to the rest of the cell via proteins that are covalently attached to the peptidoglycan. The OM is asymmetric, as it contains phospholipids in the inner leaflet and LPS in the outer leaflet. In addition, there are two types of proteins, integral OM proteins (OMPs) and lipoproteins.

Figure 4. Current models for LPS transport across the cell envelope

After being synthesized, rough LPS molecules are flipped across the IM by the ABC transporter MsbA. If the O-antigen is made, it is then ligated to the rest of the molecule by WaaL (step not shown). Then, LPS is extracted from the IM by LptC and the LptFGB ABC transporter. In the soluble-intermediate model (shown on the left), LptA serves as a soluble periplasmic chaperone that transports LPS from the IM to the LptDE assembly site. In the trans-envelope complex model (shown on the right), the Lpt factors constitute a multiprotein complex that spans the cell envelope. Because LptA crystallizes as a fiber in the presence of LPS, it is possible that it serves as a bridge between the OM and IM components. Note that the only physical interaction between Lpt factors that has been conclusively demonstrated to date is the one that exists between LptD and LptE.⁵³