Published in final edited form as:

Invest Ophthalmol Vis Sci. 2009 December ; 50(12): 6015–6022. doi:10.1167/iovs.09-3643.

The Reduction of Retinal Autofluorescence Caused by Light Exposure

Jessica I. W. Morgan^{1,*}, Jennifer J. Hunter¹, William H. Merigan¹, and David R. Williams¹

¹University of Rochester, Center for Visual Science, Rochester, New York

*J.I.W.M. is currently at the University of Pennsylvania, Department of Ophthalmology, Philadelphia, Pennsylvania

Abstract

Purpose—We have previously shown that long exposure to 568 nm light at levels below the maximum permissible exposure safety limit produces retinal damage preceded by a transient reduction in the autofluorescence of retinal pigment epithelial (RPE) cells *in vivo*. Here, we determine how the effects of exposure power and duration combine to produce this autofluorescence reduction and find the minimum exposure causing a detectable autofluorescence reduction.

Methods—Macaque retinas were imaged using a fluorescence adaptive optics scanning laser ophthalmoscope to resolve individual RPE cells *in vivo*. The retina was exposed to 568 nm light over a square subtending 0.5° with energies ranging from 1 J/cm² to 788 J/cm², where power and duration were independently varied.

Results—*In vivo* exposures of 5 J/cm² and higher caused an immediate decrease in autofluorescence followed by either full autofluorescence recovery (exposures ≤ 210 J/cm²) or permanent RPE cell damage (exposures ≥ 247 J/cm²). No significant autofluorescence reduction was observed for exposures of 2 J/cm² and lower. Reciprocity of exposure power and duration held for the exposures tested, implying that the total energy delivered to the retina, rather than its distribution in time, determines the amount of autofluorescence reduction.

Conclusions—That reciprocity holds is consistent with a photochemical origin, which may or may not cause retinal degeneration. The implementation of safe methods for delivering light to the retina requires a better understanding of the mechanism causing autofluorescence reduction. Finally, RPE imaging was demonstrated using light levels that do not cause a detectable reduction in autofluorescence.

Introduction

It has long been known that the retina can be harmed by bright light, and that it is important to understand the limits and mechanisms of retinal phototoxicity in order to incorporate safe practices of retinal light exposures. In particular, phototoxicity is important because several clinical procedures (such as slit-lamp examination, fundus photography, fluorescein angiography and retinal surgery) are often performed at light levels close to the limits imposed by current safety standards.^{1–3} In addition, those procedures are used to examine retinas affected by eye diseases, and the extent to which patients have increased susceptibility to retinal light damage remains unknown. Indeed, it has been shown that certain genetic mutations can cause higher susceptibility to light in animal models,⁴ and that phototoxicity increases with age even in the normal human retina⁵ as is observed by the accumulation of lipofuscin in the retinal pigment epithelial cells.⁶ Previous studies have shown that A2E, a primary component of lipofuscin, is one of the phototoxic materials in the retina; in cultured ARPE-19 cells the combination of A2E and blue light resulted in cell death⁷ by a photooxidative mechanism.⁸

The role of phototoxicity and photooxidation in the retina has led to the theory that light exposure plays a role in some retinal diseases including age-related macular degeneration,^{5, 9-12} although this association is controversial.^{13, 14} Regardless, until phototoxic mechanisms are fully understood, studies of retinal toxicity remain important for the implementation of safe practices in ophthalmic procedures such as retinal surgery and ophthalmic imaging applications including fundus photography, fluorescein angiography, and lipofuscin autofluorescence imaging.

Lipofuscin autofluorescence imaging has been used to observe features of the RPE layer in both normal and diseased retinas *in vivo*.¹⁵⁻²⁰ This imaging modality takes advantage of the autofluorescent properties of lipofuscin, a conglomeration of materials that accumulate in the cytoplasm of the RPE cells as byproducts of the visual cycle and phagocytosis.²¹⁻²³

There has been considerable effort to develop *in vivo* retinal imaging modalities with cellular resolution. The use of adaptive optics (AO), which involves measuring the higher-order optical aberrations of the eye with a wavefront sensor and correcting these aberrations with a wavefront corrector – usually a deformable mirror, has allowed improvements in contrast and resolution such that it has become possible to routinely characterize the photoreceptor mosaic in both normal and diseased eyes *in vivo*.²⁴⁻³¹ Recently, the combination of autofluorescence and AO with the scanning laser ophthalmoscope (SLO) has allowed the visualization of the individual cells of the RPE mosaic in human and macaque retina *in vivo*.^{32, 33}

With the ability to image the RPE mosaic *in vivo*, we previously reported observations of retinal phototoxicity in macaque retina caused by exposure to the 568 nm light that was used for autofluorescence excitation.³⁴ Indeed, this study led to the unexpected finding that exposures to 568 nm light at levels below the American National Standards Institute (ANSI) Safe Use of Lasers³⁵ maximum permissible exposure (MPE) produced retinal damage. In all cases, this retinal damage was preceded by a reduction in the autofluorescence intensity from the RPE cells. In addition, dependent upon the retinal radiant exposure, some exposures caused the immediate reduction in autofluorescence but subsequently the autofluorescence recovered in full and no long-term retinal damage was detected in the RPE and cone mosaics at the site of the exposure.³⁴

It is known that retinal damage can occur by either thermal or photochemical mechanisms, depending on exposure conditions such as wavelength and duration.³⁶⁻³⁸ Thermal damage is thought to occur when the temperature of the retina rises 10° above its ambient temperature,^{36, 39, 40} whereas photochemical effects cause retinal damage when the incoming light interacts with molecules to cause a chemical change.^{36, 38, 40-42} Retinal lesions exhibit different characteristics depending on whether they are caused by thermal or photochemical mechanisms. For example, thermal lesions are in general immediately visible, while photochemical lesions can take up to 48 hours to appear.^{38, 43} In addition, thermal lesions can spatially expand beyond the exposed retinal region because of lateral heat diffusion, while photochemical lesions are usually confined to the irradiated retinal area.³⁶ Also, photochemical lesions will exhibit reciprocity of time and irradiance, while thermal lesions do not.⁴² The concept of reciprocity is that exposures of equivalent energies cause equivalent effects regardless of how the total exposure is distributed in time.⁴²

Morgan et al. showed that the retinal damage observed with exposures to 568 nm light is caused by a photochemical mechanism.³⁴ However, it is currently unknown whether the observed reduction in lipofuscin autofluorescence and long-term retinal damage are caused by the same mechanism and represent stages of a phototoxic continuum or whether the two phenomena are caused by separate mechanisms. It is important to understand the mechanism causing the reduction in autofluorescence to determine if this phenomenon is detrimental to retinal health,

and if reduced autofluorescence represents a pathway to phototoxicity in the retina. In the present study, we test the mechanism causing autofluorescence reduction by examining reciprocity of time and irradiance for 568 nm light exposures.

Regardless of the mechanism causing retinal damage and autofluorescence reduction, it is important to determine the limits of phototoxicity in order to establish exposure guidelines that allow the safe practice of procedures requiring retinal illumination such as retinal imaging. In the present study, the effects of exposures over a range of almost four log units of radiant exposure were measured to determine the boundaries between permanent retinal damage, reduced autofluorescence, and no detectable change. Following establishing those limits, we demonstrate successful imaging of the RPE cell mosaic using illumination conditions more than an order of magnitude below the exposure level causing a detectable reduction in lipofuscin autofluorescence.

Methods

Macaque Preparation

Six eyes of 4 macaques were used for these experiments. Table 1 characterizes the parameters for each macaque. During each imaging session, macaques were anesthetized with isoflurane (1.0-3.0%), body temperature was monitored, and pupils were dilated and cyclopleged with one drop each of Phenylephrine Hydrochloride (2.5%) and Tropicamide (1%). A lid speculum held the eye open for imaging, and a rigid gas permeable contact lens was used to protect the cornea.⁴⁴ The animal's head and pupil of the eye were aligned with the imaging system by a head post rotation mount and a three-axis translation stage. Axial lengths were used to determine the dimensions of the retinal images by linearly scaling the LeGrand model eye.⁴⁵ For the macaques, axial length was measured by averaging 10 B-scan ultrasound measurements. The University of Rochester review board approved all experiments and the study adhered to the ARVO Statement for the use of animals in research.

Autofluorescence Imaging of the RPE Cells

The fluorescence adaptive optics scanning laser ophthalmoscope (AOSLO) used in this experiment has been previously described.^{32, 33} Three lasers were simultaneously scanned across the retinal area – 568 nm light for the autofluorescence excitation, 830 nm or 794 nm radiation for reflectance imaging, and 904 nm radiation for wavefront sensing. Autofluorescence images were taken by exciting the retina with 568 nm light and collecting the emission over a 40 nm bandwidth centered at 624 nm. The simultaneous imaging and registration scheme was employed as previously described.³² In this scheme, the reflectance video frames were registered to determine interframe eye motion, the motion was corrected by shifting the frames of the simultaneously recorded autofluorescence video, and the shifted autofluorescence frames were averaged to give the final RPE mosaic image.

Laser Modulation

As previously described,³⁴ the 568 nm light was modulated with an acousto-optic modulator relative to the position of the two scanners. Under normal conditions of operation, the 568 nm laser light illuminated the retina 37% of the time (ON-ratio = 0.37). To increase the average power for some exposures, the 568 nm laser light remained on continuously (ON-ratio = 1.00). The 794 nm SLD and the 830 nm and 904 nm lasers always had an ON-ratio of 1.00.

Exposures

Exposures were delivered to the retina using the same general method as previously described.³⁴ Pre-exposure images of the RPE cells and cone photoreceptors were taken of a square 2° in

visual angle per side using an average power of 20 μW (ON-ratio = 0.37, instantaneous power of 55 μW) of 568 nm for autofluorescence imaging, 50 μW of 904 nm for wavefront sensing, and approximately 300 μW of 794 nm for reflectance imaging. Some exposures used 830 nm radiation for reflectance imaging, but the majority of exposures performed in this study used 794 nm for reflectance imaging. No differences were observed between exposures using the two different reflectance sources. The same powers were used for wavefront sensing and for reflectance imaging during the exposures. The power of all radiation entering the cornea was measured using a power meter with a silicon detector (power meter model 1930-C, detector model Phto-918-SL, Newport Corporation, Irvine, CA, USA). The power meter and detector have an accuracy of $\pm 1\%$ and were calibrated prior to the experiments. Retinal radiant exposure was then calculated from the measured power by assuming an ocular focal length of 15 mm.

Exposures of 568 nm light were delivered to the retina over a square 0.5° in visual angle per side (approximately 120 μm per side) using the AOSLO. Radiant exposures ranged from 1 to 788 J/cm^2 of the 568 nm light. All exposures were obtained using the normal mode of operation for the AOSLO (ON-ratio = 0.37) except for the 788 J/cm^2 exposures, which used an ON-ratio of 1.00. The exposure power and duration were independently varied to test whether reciprocity of radiant exposure holds for those exposure conditions. Tables 2 and 3 provide the exposure powers and durations that were tested as well as the number of trials tested for each radiant exposure and each combination of exposure power and duration.

During the exposures, eye motion was monitored and the retina was stabilized as previously described.³⁴ If the eye motion could not be stabilized within approximately 0.1° in any direction, the exposure location was abandoned. Immediately after each exposure, a post-exposure image was taken of the RPE cells and cone photoreceptors over the same 2° square area and under the same conditions as the pre-exposure image. Thus, the immediately post-exposure image contains retinal area both exposed and unexposed. Additional images of the RPE cells and cone photoreceptors were taken at the same location between 5 and 63 days post-exposure (average $\pm$ standard deviation: 33 ± 13 days). Pre-images for each exposure did not overlap with each other, and each exposure location was used only once. All exposures were contained within the vascular arcade. Results from some exposures previously published are included in the current analysis.³⁴

Quantification of the Autofluorescence Intensity Decrease

The amount of autofluorescence reduction was quantified by the “AF-ratio” as previously described.³⁴ The AF-ratio is the ratio of autofluorescence intensity inside to outside the exposure area for a post-exposure image normalized by the ratio of the autofluorescence intensity inside to outside the exposure area for the pre-exposure image.

$$AF - ratio = \frac{\left(\frac{\text{mean}(\text{autofluorescence}_{\text{inside-post}})}{\text{mean}(\text{autofluorescence}_{\text{outside-post}})} \right)}{\left(\frac{\text{mean}(\text{autofluorescence}_{\text{inside-pre}})}{\text{mean}(\text{autofluorescence}_{\text{outside-pre}})} \right)}$$

AF-ratios were averaged for exposures of the same radiant exposure. Paired t-tests between the pre-, immediately post-, and several days post-exposure AF-ratios were performed. Those differences with $p < 0.05$ were considered significant. To test reciprocity of radiant exposure, the AF-ratios were averaged for exposures of the same retinal irradiance and an analysis of variance (ANOVA) was conducted between the power of the exposures and the immediately post-exposure AF-ratios. Again, the results were considered significant when $p < 0.05$.

Results

Effect of Radiant Exposure on Autofluorescence Reduction

Figure 1 shows three series of images for exposures of 788, 39 and 1 J/cm². For the 788 and 39 J/cm² exposures, a reduction in lipofuscin autofluorescence was observed immediately following the exposure. Long-term imaging showed disruption in the RPE mosaic at the site of the 788 J/cm². No long-term disruption was observed at the site of the 39 J/cm² exposure, and instead the autofluorescence intensity fully recovered. No changes were observed in the autofluorescence intensity immediately post-exposure or in long term follow up imaging for the 1 J/cm² exposure.

Figure 2 is a graph of the mean and standard error of the AF-ratios immediately post-exposure versus the radiant exposures. Exposures of 247 J/cm² and higher caused permanent retinal damage.³⁴ Exposures of 5 J/cm² and higher caused a significant immediate reduction in autofluorescence at the site of the exposure, where the magnitude of autofluorescence reduction increased as radiant exposure increased (ANOVA, $p < 0.001$, Observed Power = 1). No significant autofluorescence reduction was observed immediately post-exposure for exposures of 1 J/cm² (paired t-test, $p = 0.074$) or 2 J/cm² (paired t-test, $p = 0.177$).

Table 2 shows the mean and standard error of the AF-ratios immediately post-exposure and several days post-exposure for exposures of various radiant exposures. Autofluorescence images several days post-exposure showed either full recovery of autofluorescence or permanent RPE cell damage at the site of the exposure. For exposures of 210 J/cm² and lower,³⁴ the several days post-exposure AF-ratio was not significantly different from the pre-exposure AF-ratio, (paired t-test, $p = 0.233$) indicating full recovery of autofluorescence.

Reciprocity over Radiant Exposure

Table 3 shows the various combinations of exposure power and duration tested for four different radiant exposures, and the resulting mean and standard error of the AF-ratios for those exposures. No significant changes between the pre-exposure and the immediately post-exposure AF-ratio were observed for any of the four combinations of exposure powers and durations yielding a radiant exposure of 1 J/cm² (ANOVA, $p = 0.407$, Observed Power = 0.217). Figure 3 plots the AF-ratio immediately post-exposure for the combinations of exposure power and duration tested for the four radiant exposures. The two combinations of exposure power and duration tested for exposures of 5 J/cm² were not significantly different from each other (ANOVA, $p = 0.752$, Observed Power = 0.059), but did cause a significant decrease in the AF-ratio immediately post-exposure (paired t-test, $p < 0.001$). Likewise, the three combinations of exposure power and duration tested for 14 J/cm², and the four combinations tested for 39 J/cm² were not significantly different from one another (ANOVA, $p = 0.727$, Observed Power = 0.110, $p = 0.081$, Observed Power = 0.536, respectively). Overall, reciprocity of exposure power and duration held for the exposures tested, however there is a possibility that the variability and small sample size of this study masked an effect of radiant exposure on AF-ratio.

Discussion

The two primary objectives of this study were to determine the maximum exposure that does not produce a detectable reduction in autofluorescence, and to establish the nature of the reduction in autofluorescence. No measurable change in the autofluorescence was detected after exposures to 2 J/cm² or less but a reduction in autofluorescence was observed after radiant exposures of 5 J/cm² and higher. This range (2 – 5 J/cm²) represents the sensitivity limit for detecting the autofluorescence reduction with the AOSLO imaging technique rather than

defining a true threshold for the phenomenon. The photochemical nature of this phenomenon leads to the expectation that any light exposure would result in some amount of autofluorescence reduction.

Varying the exposure duration and power while maintaining a constant radiant exposure produced the same amount of autofluorescence reduction, demonstrating that the total radiant exposure delivered to the retina, rather than its distribution in time, determined the amount of autofluorescence reduction. That implies that reciprocity of radiant exposure holds over the tested exposure durations of 15 to 900 s. However, there is a possibility that the variability and small sample size of this study masked an effect of radiant exposure on AF-ratio, and thus did not permit the finding of a break-down in reciprocity for these exposures.

Our previous study describing the long-term retinal damage³⁴ detailed some reasons supporting the idea that the retinal damage was caused by a photochemical mechanism. These reasons include the delayed appearance of structural damage in the RPE cell mosaic, and the lack of a damaging temperature rise in the retina. In addition, and as expected for a photochemical mechanism, the autofluorescence reduction is confined to a square area of the retina limited by the spatial extent of the radiant exposure (Figure 1a, 1b). Likewise, the observation from this study that reciprocity holds is consistent with a photochemical origin^{40, 43, 46} for the autofluorescence reduction phenomenon as well.

The current study tested and found reciprocity of radiant exposure using exposure durations from 15 to 900 s. However, reciprocity is expected to fail under certain conditions on both ends of the time scale. For example, Ham *et al.*⁴⁰ found reciprocity held down to 10 s, below which the exposures did not show reciprocity and the observed retinal lesions were thus presumed to be caused by thermal rather than photochemical mechanisms. A study by Lund *et al.*⁴³ shows conditions in which reciprocity holds down to 5 s. On the other side of the scale, Ham *et al.*⁴⁰ found reciprocity broke down after extremely long durations. After finding the radiant exposure that produced a minimal lesion, Ham *et al.*⁴⁰ exposed the retina four separate times to ¼ of the radiant exposure causing a minimal lesion, where each exposure was separated by 48 hours. In this case, no lesion was observed, and thus those exposure conditions exemplified a failure of reciprocity. A study of the additivity of radiant exposure by Griess and Blankenstein⁴⁶ found that additivity could be described by an exponential function with a time constant of 4 days.

As previously discussed by Morgan *et al.*,³⁴ possible mechanisms for the autofluorescence reduction include A2E photoisomerization^{47, 48} or photooxidation.^{8, 49} Understanding the mechanism causing the decrease in autofluorescence remains an important question because it would help to determine whether the observed reduction in autofluorescence is detrimental to retinal health. For example, if the autofluorescence reduction is caused by the photooxidation of A2E or another component of lipofuscin, the phenomenon may be toxic to the retina. However, if the autofluorescence reduction is caused by the photoisomerization of one or more of the components of lipofuscin, the observed retinal changes may be benign. Regardless, the long-term full recovery of the autofluorescence and the breakdown of reciprocity described by Ham *et al.*⁴⁰ and Griess and Blankenstein⁴⁶ point to the existence of at least one retinal repair mechanism, by which the retina can to some extent counterbalance the deleterious effects of light exposures. Given that no long-term changes can be detected in the AOSLO images of the RPE cells or cone photoreceptors at exposure sites following autofluorescence recovery in this study, it is likely that the autofluorescence reduction phenomenon is either intrinsically benign or is counteracted by a retinal repair mechanism that happens to restore autofluorescence. Of course, we cannot eliminate the possibility that retinal damage is caused by the process of autofluorescence reduction and recovery and the permanent damage just remains undetected by the AOSLO imaging methods used in this study. Histological studies including electron

microscopy may help determine if any toxic retinal changes following autofluorescence reduction and recovery exist.

Effect of Multiple Wavelengths

In this study, the retina was simultaneously exposed to multiple wavelengths for wavefront sensing (904 nm), reflectance imaging (830 or 794 nm), and autofluorescence imaging or test exposures (568 nm). We have previously shown that the expected temperature rise in the retina from the infrared (IR) and 568 nm exposures is minimal.³⁴ In addition, the photochemical mechanism of autofluorescence reduction is likely caused by a molecular change in one or more components of lipofuscin, which minimally absorbs IR radiation.⁵⁰ In current safety standards, photochemical and thermal mechanisms are treated independently,⁵¹ and thus the minimal temperature rise and absorption by lipofuscin of the IR radiation used in this study would have no effect on the observation of autofluorescence reduction. However, we cannot rule out the possibility that thermal and photochemical mechanisms interact in the retina. If this is the case, then theoretically a minimal rise in retinal temperature could lead to enhanced photochemical effects, including the long-term retinal damage and autofluorescence reduction phenomena in this study. Indeed Han *et al.* show that multiple photon absorption in synthesized melanin does not follow an independent model of thermal and photochemical effects, and therefore that thresholds for damage in multiple wavelength exposures may be underestimated.⁵² Further experiments will be necessary to determine the full effect of multi-wavelength simultaneous exposures and any resulting interactions between thermal and photochemical effects in the retina.

Light Exposures and Safety Standards

Because the photochemical molecular mechanism causing the reduction and recovery of autofluorescence is unresolved, it is important to examine the exposures in the context of light safety standards.^{35, 53-55} Figure 4 shows the tested exposures in comparison with the ANSI standard³⁵ MPE for exposures of 568 nm over a square 0.5° in visual angle per side for the monkey eye of 15 mm focal length. As described in the present study and previously by Morgan *et al.*,³⁴ we have found long-term damage at light levels below the ANSI standard and previously thought to be safe. As well, we have observed autofluorescence reduction with exposures (5 J/cm²) 31 times less than the ANSI MPE (155 J/cm², thermal limit). In light of that new data, the ANSI standard,³⁵ and other light safety standards,⁵³⁻⁵⁵ need to be modified to ensure adequate retinal protection from light exposures. However, the extent to which those standards must be decreased will depend on whether the mechanism causing the reduction in autofluorescence is found to be harmful.

Safe RPE Mosaic Imaging

The ability to image the RPE cells on an individual cellular level *in vivo* provides numerous advantages for studying the normal and diseased retina because quantitative metrics such as cell density, size and regularity can be applied to advancing the understanding of retinal disease progression and treatment efficacy.³² However, the imaging techniques used on the living human eye need to be completely non-invasive. Until it is known whether or not autofluorescence reduction is benign, light exposures causing a reduction in autofluorescence should be avoided whenever possible. Figure 5 shows an example image of the macaque RPE mosaic taken with an exposure of only 0.15 J/cm² of 568 nm and using a broader emission filter for autofluorescence detection. This light exposure is more than an order of magnitude below the 2 J/cm² exposure that caused no detectable autofluorescence reduction. Thus, we have demonstrated safe and successful imaging of the individual cells in the RPE mosaic despite the reduced light intensity for autofluorescence excitation.

Conclusion

This study investigated the nature of and limits to the reduction and recovery of autofluorescence following exposure to light as was described by Morgan et al.³⁴ The sensitivity of this method yielded a limit to detecting reduced autofluorescence with the AOSLO following a light exposure between 2 and 5 J/cm². The magnitude of the autofluorescence reduction increased as the radiant exposure increased, and reciprocity held for all exposures tested from 15 to 900 s, showing that the phenomenon is a result of photochemical changes in the retina. It is still unknown whether autofluorescence reduction poses a safety risk to the retina. Future studies to describe the action spectrum of autofluorescence reduction and histological characteristics of retinal locations experiencing both autofluorescence reduction with recovery or long-term disruption could help elucidate the retinal health risk and the mechanisms involved. Until these phenomena are completely understood, it is prudent to ensure exposures to the retina remain below those that cause any retinal changes, including the reduction in autofluorescence, unless absolutely necessary. To meet this goal, this study demonstrates RPE cellular imaging using light exposures more than an order of magnitude below the autofluorescence reduction detection limit. In general, the implementation of safe methods, not only to image the RPE cell mosaic *in vivo*, but also to deliver high intensity light to the retina, requires a better understanding of the cause and effects of autofluorescence reduction.

Acknowledgments

The authors thank Robert Wolfe and Benjamin Masella. This research was supported by the National Institute for Health, Bethesda, Maryland (NIH EY014375, NIH EY01319, NIH EY07125), and Research to Prevent Blindness. This work has been supported in part by the National Science Foundation Science and Technology Center for Adaptive Optics (Santa Cruz, California), managed by the University of California at Santa Cruz (cooperative agreement no.: AST-9876783). Jessica I. W. Morgan is currently at the University of Pennsylvania.

References

1. Delori FC, Parker JS, Mainster MA. Light levels in fundus photography and fluorescein angiography. *Vision Research* 1980;20:1099–104. [PubMed: 7269267]
2. Delori FC, Pomerantzeff O, Mainster MA. Light levels in ophthalmic diagnostic instruments. *Society of Photo-Optical Instrumentation Engineers* 1980;229:154–60.
3. Stiller H, Rassow B. Light hazards to the patient's retina from ophthalmic instruments. *Applied Optics* 1991;30(16):2187–96.
4. Cideciyan AV, Jacobson SG, Aleman TS, et al. In vivo dynamics of retinal injury and repair in the rhodopsin mutant dog model of human retinitis pigmentosa. *Proceedings National Academy of Sciences (USA)* 2005;102(14):5233–8.
5. Margrain TH, Boulton M, Marshall J, Sliney DH. Do blue light filters confer protection against age-related macular degeneration? *Progress in Retinal and Eye Research* 2004;23:523–31. [PubMed: 15302349]
6. Delori FC, Goger DG, Dorey CK. Age-related accumulation and spatial distribution of lipofuscin in RPE of normal subjects. *Investigative Ophthalmology & Visual Science* 2001;42(8):1855–66. [PubMed: 11431454]
7. Sparrow JR, Nakanishi K, Parish CA. The lipofuscin fluorophore A2E mediates blue light-induced damage to retinal pigmented epithelial cells. *Invest. Ophthalmol. Vis. Sci* 2000;41:1981–9. [PubMed: 10845625]
8. Sparrow JR, Zhou J, Ben-Shabat S, et al. Involvement of oxidative mechanisms in blue-light-induced damage to A2E-laden RPE. *Invest. Ophthalmol. Vis. Sci* 2002;43:1222–7. [PubMed: 11923269]
9. Taylor HR, Munoz B, West S, et al. Visible light and risk of age-related macular degeneration. *Trans Am Ophth Soc* 1990;88:163–78.

10. Taylor HR, West S, Munoz B, et al. The long-term effects of visible-light on the eye. *Archives of Ophthalmology* 1992;110:99–104. [PubMed: 1731731]
11. Cruickshanks KJ, Klein R, Klein BEK. Sunlight and age-related macular degeneration - the Beaver Dam Eye Study. *Archives of Ophthalmology* 1993;111:514–8. [PubMed: 8470986]
12. Tomany SC, Cruickshanks KJ, Klein R, et al. Sunlight and the 10-year incidence of age related maculopathy. *Archives of Ophthalmology* 2004;122:750–7. [PubMed: 15136324]
13. Darzins P, Mitchell P, Heller RF. Sun exposure and age-related macular degeneration - an Australian case-control study. *Ophthalmology* 1997;104:770–6. [PubMed: 9160021]
14. Group EDCCS. Risk factors for neovascular age-related macular degeneration. *Archives of Ophthalmology* 1992;110:1701–8. [PubMed: 1281403]
15. von Ruckmann A, Fitzke FW, Bird AC. Distribution of fundus autofluorescence with a scanning laser ophthalmoscope. *British Journal of Ophthalmology* 1995;79:407–12. [PubMed: 7612549]
16. von Ruckmann A, Fitzke FW, Bird AC. Fundus autofluorescence in age-related macular disease imaged with a laser scanning ophthalmoscope. *Invest. Ophthalmol. Vis. Sci* 1997;38:478–86. [PubMed: 9040481]
17. Lois N, Halfyard AS, Bird AC, Fitzke FW. Quantitative evaluation of fundus autofluorescence imaged “in vivo” in eyes with retinal disease. *British Journal of Ophthalmology* 2000;84:741–5. [PubMed: 10873986]
18. Holz FG, Bellman C, Staudt S, et al. Fundus autofluorescence and development of geographic atrophy in age-related macular degeneration. *Investigative Ophthalmology & Visual Science* 2001;42(5):1051–6. [PubMed: 11274085]
19. Lois N, Halfyard AS, Bird AC, et al. Fundus autofluorescence in Stargardt macular dystrophy-fundus flavimaculatus. *American Journal of Ophthalmology* 2004;138(1):55–63. [PubMed: 15234282]
20. Bindewald A, Jorzik JJ, Loesch A, et al. Visualization of retinal pigment epithelial cells in vivo using digital high-resolution confocal scanning laser ophthalmoscopy. *American Journal of Ophthalmology* 2004;137(3):556–8. [PubMed: 15013882]
21. Eldred GE, Katz ML. Fluorophores of the human retinal pigment epithelium: separation and spectral characterization. *Experimental Eye Research* 1988;47:71–86. [PubMed: 3409988]
22. Sparrow JR, Fishkin N, Zhou J, et al. A2E, a byproduct of the visual cycle. *Vision Research* 2003;43:2983–90. [PubMed: 14611934]
23. Fishkin NE, Sparrow JR, Allikmets R, Nakanishi K. Isolation and characterization of a retinal pigment epithelial cell fluorophore: An all-trans-retinal dimer conjugate. *Proceedings National Academy of Sciences (USA)* 2005;102(20):7091–6.
24. Liang J, Williams DR, Miller DT. Supernormal vision and high-resolution retinal imaging through adaptive optics. *Journal of the Optical Society of America A* 1997;14(11):2884–92.
25. Hofer H, Chen L, Yoon GY, et al. Improvement in retinal image quality with dynamic correction of the eye's aberrations. *Optics Express* 2001;8(11):631–43. [PubMed: 19421252]
26. Roorda A, Williams DR. The arrangement of the three cone classes in the living human eye. *Nature (London)* 1999;397:520–2. [PubMed: 10028967]
27. Roorda A, Romero-Borja F, Donnelly WJ III, et al. Adaptive optics scanning laser ophthalmoscopy. *Optics Express* 2002;10(9):405–12. [PubMed: 19436374]
28. Hermann B, Fernandez EJ, Unterhuber A, et al. Adaptive-optics ultrahigh-resolution optical coherence tomography. *Optics Letters* 2004;29(18):2142–4. [PubMed: 15460883]
29. Carroll J, Neitz M, Hofer H, et al. Functional photoreceptor loss revealed with adaptive optics: An alternate cause for color blindness. *Proceedings of the National Academy of Science, USA* 2004;101(22):8461–6.
30. Wolfing JI, Chung M, Carroll J, et al. High-resolution retinal imaging of cone-rod dystrophy. *Ophthalmology* 2006;113:1014–9.
31. Choi SS, Doble N, Hardy JL, et al. In vivo imaging of the photoreceptor mosaic in retinal dystrophies and correlations with visual function. *Invest. Ophthalmol. Vis. Sci* 2006;47:2080–92. [PubMed: 16639019]

32. Morgan JIW, Dubra A, Wolfe R, et al. In vivo autofluorescence imaging of the human and macaque retinal pigment epithelial cell mosaic. *Invest. Ophthalmol. Vis. Sci* 2009;50(3):1350–9. [PubMed: 18952914]
33. Gray DC, Merigan W, Wolfing JI, et al. In vivo fluorescence imaging of primate retinal ganglion cells and retinal pigment epithelial cells. *Optics Express* 2006;14:7144–58. [PubMed: 19529085]
34. Morgan JIW, Hunter JJ, Masella B, et al. Light-induced retinal changes observed using high-resolution autofluorescence imaging of the retinal pigment epithelium. *Invest. Ophthalmol. Vis. Sci* 2008;49:3715–29. [PubMed: 18408191]
35. ANSI. American National Standard for Safe Use of Lasers ANSI Z136.1-2007. Laser Institute of America; 2007.
36. Glickman RD. Phototoxicity to the retina: mechanisms of damage. *International Journal of Toxicology* 2002;21(6):473–90. [PubMed: 12537644]
37. Lund, DJ. Action spectrum for retinal thermal injury.. In: Matthes, R.; Sliney, DH., editors. *Measurements of Optical Radiation Hazards*. Maerkl-Druck; Munchen: 1998.
38. Stuck, BE. The Retinal and Action Spectrum for Photoreinitis (“Blue-Light Hazard”).. In: Matthes, R.; Sliney, DH., editors. *Measurements of Optical Radiation Hazards*. Maerkl-Druck; Munchen: 1998.
39. Clarke AM, Geeraets WJ, Ham WTJ. An equilibrium thermal model for retinal injury from optical sources. *Appl. Opt* 1969;8(5):1051–4.
40. Ham WTJ, Mueller HA, Ruffolo JJ, Clarke AM. Sensitivity of the retina to radiation damage as a function of wavelength. *Photochemistry and Photobiology* 1979;29:735–43. [PubMed: 109869]
41. Boulton M, Rozanowska M, Rozanowski B. Retinal Photodamage. *Journal of Photochemistry and Photobiology* 2001;64:144–61.
42. Wu JW, Seregard S, Algvere PV. Photochemical Damage of the Retina. *Survey of Ophthalmology* 2006;51(5):461–81. [PubMed: 16950247]
43. Lund DJ, Stuck BE, Edsall P. Retinal injury thresholds for blue wavelength lasers. *Health Physics* 2006;90(5):477–84. [PubMed: 16607179]
44. Metha AB, Crane AM, Rylander HGI, et al. Maintaining the cornea and the general physiological environment in visual neurophysiology experiments. *Journal of Neuroscience Methods* 2001;109:153–66. [PubMed: 11513950]
45. Wyszecki, G.; Stiles, WS. *Color Science: Concepts and Methods, Quantitative Data and Formulae*. Vol. Second Edition. John Wiley & Sons; New York: 1982.
46. Griess GA, Blankenstein MF. Additivity and repair of actinic retinal lesions. *Invest. Ophthalmol. Vis. Sci* 1981;20(6):803–7. [PubMed: 7195384]
47. Parish CA, Hashimoto M, Nakanishi K, et al. Isolation and one-step preparation of A2E and iso-A2E, fluorophores from human retinal pigment epithelium. *Proceedings National Academy of Sciences (USA)* 1998;95:14609–13.
48. Lamb LE, Simon JD. A2E: a component of ocular lipofuscin. *Photochemistry and Photobiology* 2004;79(2):127–36. [PubMed: 15068025]
49. Zhou J, Jang YP, Kim SR, Sparrow JR. Complement activation by photooxidation products of A2E, a lipofuscin constituent of the retinal pigment epithelium. *Proceedings National Academy of Sciences (USA)* 2006;103(44):16182–7.
50. Delori FC, Dorey CK, Staurenghi G, et al. In vivo fluorescence of the ocular fundus exhibits retinal pigment epithelium lipofuscin characteristics. *Investigative Ophthalmology & Visual Science* 1995;36(3):718–29. [PubMed: 7890502]
51. Delori FC, Webb RH, Sliney DH. Maximum permissible exposures for ocular safety (ANSI 2000), with emphasis on ophthalmic devices. *Journal of the Optical Society of America A* 2007;24(5):1250–65.
52. Han Y, Rosenshein JS, Wu ZL, et al. Abnormal photon absorption behavior in melanin detected by photothermal lensing due to simultaneous irradiation with multiple laser wavelengths. *Optical Engineering* 2001;40(2):303–9.
53. ACGIH. American Conference of Governmental Industrial Hygienists; TLVs and BEIs.; ACGIH. Cincinnati, Ohio. 2005;

54. ICNIRP. International Commission on Non-Ionizing Radiation Protection: Guidelines on limits of exposure to laser radiation of wavelengths between 180nm and 1,000 microns. *Health Physics* 1996;71:804–19. [PubMed: 8887534]
55. ICNIRP. Revision of guidelines on limits of exposure to laser radiation of wavelengths between 400 nm and 1.4 micron. International Commission on Non-Ionizing Radiation Protection. *Health Physics* 2000;79(4):431–40. [PubMed: 11007467]

Figure 1.

Time series of autofluorescence images of the RPE mosaic for locations exposed to (a) 788, (b) 39 and (c) 1 J/cm². The 788 J/cm² exposure (a) caused an immediate decrease in autofluorescence intensity followed by long-term disruption in the RPE mosaic at the site of the exposure. The 39 J/cm² exposure (b) caused an immediate decrease in autofluorescence intensity at the site of the exposure followed by long-term full recovery of the autofluorescence. The 1 J/cm² exposure (c) caused no change in autofluorescence intensity. The white squares show the exposure locations. Scale bar: 50 μm.

Figure 2.

Mean and standard error of AF-ratios immediately post-exposure for a range of radiant exposures. Data points labeled with * and † mark exposures that resulted in permanent damage and no significant autofluorescence reduction, respectively. Ranges of radiant exposures causing no change, transient autofluorescence reduction, or permanent damage are shaded in white, light gray, and dark gray, respectively.

Figure 3.

Data points plot the mean and standard error of the AF-ratio immediately post-exposure for various combinations of exposure power and duration resulting in radiant exposures of 1, 5, 14, and 39 J/cm². Although the AF-ratio appears to decrease with exposure power for the 39 J/cm² trials, this was not a significant trend. No significant reduction in autofluorescence was observed for exposures of 1 J/cm². Black diamonds: 1 J/cm²; gray squares: 5 J/cm²; black X's: 14 J/cm²; gray circles: 39 J/cm².

Figure 4.

The exposures tested in this study are compared to the ANSI standard.³⁵ The solid line shows the thermal MPE exposure limit calculated from the ANSI standard for exposures to 568 nm light over a square 0.5° in visual angle per side for an eye with a focal length of 15 mm. The dashed line shows the photochemical MPE for the same conditions. Black triangles show exposures that caused long-term permanent damage to the RPE. Black circles show exposures that caused a significant reduction in autofluorescence. Open squares show exposures in which no significant change in the RPE autofluorescence was observed. The radiant exposures tested in this study are labeled.

Figure 5. Images of the RPE mosaic taken with an exposure of only 0.15 J/cm² of 568 nm light. A 90 nm bandpass emission filter centered at 630 nm was used for the detection of autofluorescence emission. Scale bar: 50 μ m.

Table 1

Macaque Parameters

Macaque Number	Species	Age (years)	Gender	Eye	Axial Length (mm)
320	Macaca fascicularis	8	M	OD	17.85
				OS	17.59
526	Macaca mulatta	7	F	OS	20.22
620	Macaca fascicularis	3	M	OS	18.07
903	Macaca nemestrina	11	M	OD	20.97
				OS	20.95

Table 2

AF-ratio and Radiant Exposure

Radiant Exposure (J/cm ²)	Number of Trials	AF-Ratio (mean $\pm$ standard error)	
		Immed. Post-Expo	Days Post-Expo
1	16	0.987 $\pm$ 0.007	1.024 $\pm$ 0.008
2	4	0.984 $\pm$ 0.009	0.992 $\pm$ 0.013
5	9	0.922 $\pm$ 0.006	1.000 $\pm$ 0.017
14	13	0.903 $\pm$ 0.011	0.996 $\pm$ 0.007
39	16	0.829 $\pm$ 0.011	1.003 $\pm$ 0.014
105	4	0.765 $\pm$ 0.004	0.993 $\pm$ 0.051 [‡]
289	3	0.709 $\pm$ 0.022	N/A [§]
788 [*]	4	0.577 $\pm$ 0.028	N/A [§]

[†] The AF-ratio was significantly reduced (paired t-test, $p < 0.05$).

[‡] Only 3 of the 4 exposures were followed several days post-exposure.

[§] The AF-ratio was not measured days post-exposure for those locations because the RPE exhibited permanent damage at the site of the exposures.

^{*} Exposures of 788 J/cm² used the 568 nm light with an ON-ratio of 1.00. All other exposures used the normal mode of AOSLO operation with an ON-ratio of 0.37.

Table 3

AF-ratios for Exposures of Various Combinations of Power and Duration

Energy (J/cm ²)	Exposure Average Power (μW)	Exposure Duration (seconds)	Number of Trials	AF-Ratio (mean ± standard error)	
				Immed. Post-Expo	Days Post-Expo
1	1.9	90	4	0.987 ± 0.017	1.034 ± 0.018
	2.9	60	4	0.968 ± 0.008	1.040 ± 0.016
	5.7	30	4	1.000 ± 0.019	1.002 ± 0.015
	11	15	4	0.992 ± 0.004	1.018 ± 0.017
5	7.4	120	4	0.920 ± 0.013	0.996 ± 0.009
	20	45	5	0.924 ± 0.004	1.003 ± 0.017
	7.4	330	4	0.894 ± 0.016	1.003 ± 0.011
	20	120	4	0.921 ± 0.033	0.978 ± 0.016
39	45	53	5	0.896 ± 0.009	1.005 ± 0.006
	7.4	900	4	0.863 ± 0.013 *	1.004 ± 0.012
	20	330	4	0.847 ± 0.014 *	0.997 ± 0.028
	36	184	4	0.810 ± 0.005 *	1.031 ± 0.020
	45	147	4	0.795 ± 0.032 *	0.978 ± 0.045

* Although the AF-ratio appears to decrease with exposure power for the 39 J/cm² trials, this was not a significant trend.