Comprehensive Cancer Center Annual Meeting, Durham, NC, March 10, 2008. Funded through an Outcomes Research service agreement with Pfizer. Pfizer has no access to individual data or authority to prohibit publication results.

Authors' Disclosures of Potential Conflicts of Interest

Although all authors completed the disclosure declaration, the following author(s) indicated a financial or other interest that is relevant to the subject matter under consideration in this article. Certain relationships marked with a "U" are those for which no compensation was received; those relationships marked with a "C" were compensated. For a detailed description of the disclosure categories, or for more information about ASCO's conflict of interest policy, please refer to the Author Disclosure Declaration and the Disclosures of Potential Conflicts of Interest section in Information for Contributors.

References

- de Gramont A, Figer A, Seymour M, et al: Leucovorin and fluorouracil with or without oxaliplatin as first-line treatment in advanced colorectal cancer. *J Clin Oncol* 18:2938-2947, 2000
- Douillard JY, Cunningham D, Roth AD, et al: Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: A multicentre randomised trial. *Lancet* 355:1041-1047, 2000
- Maindrault-Goebel F, Louvet C, André T, et al: Oxaliplatin added to the simplified bimonthly leucovorin and 5-fluorouracil regimen as second-line therapy for metastatic colorectal cancer (FOLFOX6): GERCOR. *Eur J Cancer* 35:1338-1342, 1999
- André T, Louvet C, Maindrault-Goebel F, et al: CPT-11 (irinotecan) addition to bimonthly, high-dose leucovorin and bolus and continuous-infusion 5-fluorouracil (FOLFIRI) for pretreated metastatic colorectal cancer: GERCOR. *Eur J Cancer* 35:1343-1347, 1999
- Goldberg RM, Rothenberg ML, Van Cutsem E, et al: The continuum of care: A paradigm for the management of metastatic colorectal cancer. *Oncologist* 12:38-50, 2007
- Tournigand C, André T, Achille E, et al: FOLFIRI followed by FOLFOX6 or the reverse sequence in advanced colorectal cancer: A randomized GERCOR study. *J Clin Oncol* 22:229-237, 2004
- World Medical Association: Declaration of Helsinki. <http://www.wma.net/ethics/b3.htm>
- André T, Boni C, Mounedji-Boudiaf L, et al: Oxaliplatin, fluorouracil, and leucovorin as adjuvant treatment for colon cancer. *N Engl J Med* 350:2343-2351, 2004
- Saltz LB, Niedzwiecki D, Hollis D, et al: Irinotecan fluorouracil plus leucovorin is not superior to fluorouracil plus leucovorin alone as adjuvant treatment for stage III colon cancer: Results of CALGB 89803. *J Clin Oncol* 25:3456-3461, 2007
- Goldberg RM, Sargent DJ, Morton RF, et al: A randomized controlled trial of fluorouracil plus leucovorin, irinotecan, and oxaliplatin combinations in patients with previously untreated metastatic colorectal cancer. *J Clin Oncol* 22:23-30, 2004
- Schrag D, Cramer LD, Bach PB, et al: Age and adjuvant chemotherapy use after surgery for stage III colon cancer. *J Natl Cancer Inst* 93:850-7, 2001
- Ayanian JZ, Zaslavsky AM, Fuchs CS, et al: Use of adjuvant chemotherapy and radiation therapy for colorectal cancer in a population-based cohort. *J Clin Oncol* 21:1293-1300, 2003
- Schrag D, Gelfand SE, Bach PB, et al: Who gets adjuvant treatment for stage II and III rectal cancer? Insight from surveillance, epidemiology, and end results—Medicare. *J Clin Oncol* 19:3712-3718, 2001
- Goldberg RM, Tabah-Fisch I, Bleiberg H, et al: Pooled analysis of safety and efficacy of oxaliplatin plus fluorouracil/leucovorin administered bimonthly in elderly patients with colorectal cancer. *J Clin Oncol* 24:4085-4091, 2006
- Townsend C, Pond GR, Pelozo B, et al: Analysis of treatment practices for elderly cancer patients in Ontario, Canada. *J Clin Oncol* 23:3802-3810, 2005
- Popescu RA, Norman A, Ross PJ, et al: Adjuvant or palliative chemotherapy for colorectal cancer in patients 70 years or older. *J Clin Oncol* 17:2412-2418, 1999
- Fuchs CS, Marshall J, Mitchell E, et al: Randomized, controlled trial of irinotecan plus infusional, bolus, or oral fluoropyrimidines in first-line treatment of metastatic colorectal cancer: Results from the BICC-C Study. *J Clin Oncol* 25:4779-4786, 2007
- Saltz LB, Clarke S, Díaz-Rubio E, et al: Bevacizumab in combination with oxaliplatin-based chemotherapy as first-line therapy in metastatic colorectal cancer: A randomized phase III study. *J Clin Oncol* 26:2013-2019, 2008

Employment or Leadership Position: None **Consultant or Advisory Role:** Michael A. Morse, Genentech (C); Amy P. Abernethy, Pfizer (C) **Stock Ownership:** None **Honoraria:** S. Yousuf Zafar, Genentech **Research Funding:** Amy P. Abernethy, Pfizer **Expert Testimony:** None **Other Remuneration:** None

Corresponding author: S. Yousuf Zafar, MD, MHS, Division of Medical Oncology, Department of Medicine, Duke University Medical Center, DUMC Box 3505, Durham, NC 27710; e-mail: yousuf.zafar@duke.edu.

DOI: 10.1200/JOP.091010; posted online ahead of print at <http://jop.ascopubs.org> on August 20, 2009.

Commentary: Practice Patterns and Potential Impact on Quality Measures for a Practicing Physician

By Cathy Eng, MD, FACP, and Scott Kopetz, MD

Practicing physicians make multiple difficult treatment decisions daily. Several influences may affect choice of therapy, including consensus recommendations of the practice, ASCO or National Comprehensive Cancer Network guidelines, medical insurance coverage, geographic location of the practice, data from recently published or presented clinical trials, and familiarity with chemotherapeutic or biologic agent. Given the US Food and Drug Administration (FDA) approval of seven therapeutic agents in metastatic

colorectal cancer, various treatment combinations may be considered. Practice patterns vary significantly within individual practices and may be more pronounced when comparing community versus academic settings.

Zafar et al¹ have approached capturing community practice patterns in the population of patients with metastatic colorectal cancer, a challenging subject on which, to our knowledge, few data have been published. The objective of

this study was to analyze longitudinal patterns of chemotherapy use and how recently published or presented scientific data translate into the real world of community practice. A retrospective medical record review of patients with metastatic colorectal cancer at the Duke University Medical Center (Durham, NC) and nine Duke University–affiliated community oncology practices was completed, inclusive of patients diagnosed between June 2003 and June 2006. All patients were treated with either fluorouracil (FU)-based chemotherapy with oxaliplatin (FOLFOX) or irinotecan (FOLFIRI). Patients participating in clinical trials were excluded. Of the 400 patients initially eligible, only 110 patients (28%) were included for the purpose of the analysis. Zafar et al noted that the majority of physicians chose oxaliplatin-based regimens for first-line therapy and irinotecan-based treatment for subsequent lines of therapy. Approximately two thirds of the medical records reviewed included no documentation to support the treatment decisions made. The authors suggested that age bias was present, because the majority of patients were younger than age 70 years. The authors encouraged prospective studies to identify physician intent in chemotherapy treatment.

Given the frequency of colorectal cancer in the United States, the number of recently approved therapeutics, and the concerns of pharmacoeconomics, practice patterns will continue to evolve as major discussion points. The objectives of such practice pattern studies must be clearly defined and critical to the primary study aim. In the study by Zafar et al,¹ despite the stated aims of assessing longitudinal management representative of community practice, regimens other than FOLFOX and FOLFIRI were excluded, including irinotecan, bolus FU, and leucovorin (LV; IFL) at a time when this regimen was commonly used. A recent community-based practice pattern study demonstrated that one half of patients received a regimen other than FOLFOX, FOLFIRI, and IFL.²

Defining lines of therapy for metastatic colorectal cancer is also difficult because of variability in practice patterns. Reasons for changing therapy on the basis of clinical or radiographic progression, deferred treatment, rising carcinoembryonic antigen levels, or toxicity were not well documented. Does restarting oxaliplatin after progression while a patient is receiving maintenance FU/LV or on a prolonged treatment break constitute a new line of therapy? Given that 87% of patients in the study by Zafar et al¹ received first-line oxaliplatin-based therapy, it is likely patients developed the dose limiting toxicity of peripheral neuropathy, which emphasizes the importance of transparency in documentation, because oxaliplatin may be reconsidered for future treatment if discontinued as a result of toxicity rather than progression of disease. Future studies investigating practice patterns should provide greater insight into the decisions physicians make.

From the data derived, the analysis by Zafar et al¹ also confirms the tendency of physicians to make treatment choices on the basis of familiarity with therapy. As mirrored in practices across the country, oxaliplatin was preferentially chosen for first-line therapy on the basis of the superiority of FOLFOX versus IFL (NCCTG [North Central Cancer Treatment Group] 9741 trial, which did not include bevacizumab).³ After FDA approval of oxaliplatin, the FOLFOX regimen was widely utilized. Yet our European colleagues completed two phase III trials of FOLFIRI and FOLFOX as first-line therapy and noted equivalent response and overall survival and excellent tolerance of therapy.^{4,5} The impact of familiarity with drug on preferentially accepted use is additionally exemplified by the fact that the combination of FOLFOX and bevacizumab continues to be the most widely accepted first-line regimen among practicing physicians. Bevacizumab was originally approved by the FDA in combination with irinotecan. Hence, familiarity with the drug of interest appears to have precedence in practice patterns over recent evidence-based literature, as exemplified in the observational analysis by Zafar et al. The combination of FOLFIRI and bevacizumab continues to be underutilized in the United States as first-line therapy and should be considered in first-line therapy for any patient who is receiving palliative therapy, given its relatively few toxicities.⁶

Zafar et al¹ suggested the existence of age bias because only 14% of patients were age 70 years or older. Yet the median age and percentage of patients age 70 years or older among those originally screened is unknown. Age bias has been reported in earlier studies but is neither supported nor refuted in the data presented here. A recent pooled analysis suggested that elderly patients age 70 years or older do not derive benefit from adjuvant FU-based therapies, contradicting prior studies suggesting equivalent benefit in elderly patients.^{7,8} Stratifying patients on the basis of comorbidities and functional status (ie, assessing biologic age in lieu of chronologic age) is required to define optimal chemotherapy utilization.

Whether the findings of the observation analysis by Zafar et al¹ truly reflect the practice patterns in the community is difficult to ascertain, because the nine community practices are affiliated with Duke University, and this may have affected their current practice patterns. Although this small observational analysis was limited in its interpretation of practice patterns generalizable to the community setting, it emphasizes the importance and value of observational analyses and databases in improving quality measures for any practicing physician.

Accepted for publication on June 30, 2009.

Authors' Disclosures of Potential Conflicts of Interest

Although all authors completed the disclosure declaration, the following author(s) indicated a financial or other interest that is relevant to the subject matter under consideration in this article. Certain relationships

marked with a "U" are those for which no compensation was received; those relationships marked with a "C" were compensated. For a detailed description of the disclosure categories, or for more information about ASCO's conflict of interest policy, please refer to the Author Disclosure Declaration and the Disclosures of Potential Conflicts of Interest section in Information for Contributors.

Employment or Leadership Position: None **Consultant or Advisory Role:** None **Stock Ownership:** None **Honoraria:** Cathy Eng, Bristol-Myers Squibb, Pfizer, Roche **Research Funding:** Cathy Eng, Amgen, sanofi-aventis, Bristol-Myers Squibb, Novartis, Genentech **Expert Testimony:** None **Other Remuneration:** Scott Kopetz, Roche, sanofi-aventis, Genentech

Cathy Eng, MD, FACP, is an associate professor at The University of Texas M. D. Anderson Cancer Center in Houston, TX. Contact her at ceng@mdanderson.org.

Scott Kopetz, MD, is an assistant professor at The University of Texas M. D. Anderson Cancer Center in Houston, TX. Contact him at skopetz@mdanderson.org.

DOI: 10.1200/JOP.091013; posted online ahead of print at <http://jop.ascopubs.org> on August 20, 2009.

References

1. Zafar SY, Marcello JE, Wheeler JL, et al: Longitudinal patterns of chemotherapy use in metastatic colorectal cancer. *J Oncol Pract* 5:228-233, 2009
2. Ferro SA, Myer BS, Wolff DA, et al: Variation in the cost of medications for the treatment of colorectal cancer. *Am J Manag Care* 14:717-725, 2008
3. Goldberg RM, Sargent DJ, Morton RF, et al: A randomized controlled trial of fluorouracil plus leucovorin, irinotecan, and oxaliplatin combinations in patients with previously untreated metastatic colorectal cancer. *J Clin Oncol* 22:23-30, 2004
4. Colucci G, Gebbia V, Paoletti G, et al: Phase III randomized trial of FOLFIRI versus FOLFOX4 in the treatment of advanced colorectal cancer: A multicenter study of the Gruppo Oncologico Dell'Italia Meridionale. *J Clin Oncol* 23:4866-4875, 2005
5. Tournigand C, André T, Achille E, et al: FOLFIRI followed by FOLFOX6 or the reverse sequence in advanced colorectal cancer: A randomized GERCOR study. *J Clin Oncol* 22:229-237, 2004
6. Douillard JY, Cunningham D, Roth AD, et al: Irinotecan combined with fluorouracil compared with fluorouracil alone as first-line treatment for metastatic colorectal cancer: A multicentre randomised trial. *Lancet* 355:1041-1047, 2000
7. Jackson McCleary NA, Meyerhardt J, Green E, et al: Impact of older age on the efficacy of newer adjuvant therapies in > 12,500 patients (pts) with stage II/III colon cancer: Findings from the ACCENT Database. *J Clin Oncol* 27:170s, 2009 (suppl; abstr 4010)
8. Köhne CH, Folprecht G, Goldberg RM, et al: Chemotherapy in elderly patients with colorectal cancer. *Oncologist* 13:390-402, 2008

Shape Your Future and Ours: Join ASCO's Leadership Development Program

If you completed your final subspecialty training between 2000 and 2005 and are interested in becoming a future leader in both ASCO and the practice of oncology, ASCO's Leadership Development Program is for you. Participants will:

- Network with ASCO leadership at the 2010 Annual Meeting in Chicago, IL
- Gain proficiency with key ASCO research activities
- Participate in a leadership boot camp at ASCO Headquarters
- Receive first-hand advocacy experience on Capitol Hill in Washington, DC

To apply or learn more about ASCO's Leadership Development Program, please visit www.asco.org/leadership. All applications must be received by October 13, 2009.