Published in final edited form as:

Clin Psychol (New York). 2009 December 1; 16(4): 449–469. doi:10.1111/j.1468-2850.2009.01184.x.

Psychosocial Interventions for Bipolar Disorder: Perspective from the Behavioral Approach System (BAS) Dysregulation Theory

Robin Nusslock and Lyn Y. Abramson

University of Wisconsin-Madison

Eddie Harmon-Jones

Texas A&M University

Lauren B. Alloy

Temple University

James A. Coan

University of Virginia

Abstract

Research has emerged providing consistent support for the behavioral approach system (BAS) dysregulation theory of bipolar disorder. The objective of the current article is to examine the extent to which findings from the BAS dysregulation theory can inform psychosocial interventions for bipolar disorder. Towards this end, we first provide an overview of the BAS dysregulation theory. Second, we review extant research on psychosocial interventions for bipolar disorder. And, third, we discuss means by which research and theory in line with the BAS dysregulation model can inform psychosocial interventions for bipolar disorder. Particular attention is given to the clinical implications of research suggesting that bipolar disorder is characterized by high drive/incentive motivation, ambitious goal-setting, and perfectionism in the achievement domain.

Keywords

bipolar disorder; treatment; BAS

Approximately 4.4% of the U.S. population experience a bipolar spectrum disorder (Merikangas, Akiskal, Angst, Greenberg, et al., 2007). The consequences of this disorder are striking, often creating significant impairment such as erratic work history, alcohol abuse, and poor academic achievement (Goodwin & Jamison, 1990; Nusslock, Alloy, Abramson, Harmon-Jones, & Harmon-Jones, 2008; Lagace, Kutcher, Robertson, 2003). Indeed, the World Health Organization (WHO) has consistently ranked bipolar disorder as one of the top ten leading causes of disability world-wide among adults (Ayuso-Mateos, 2006; Murray & Lopez, 1996).

The quality of outpatient treatment for bipolar disorder advanced considerably with the introduction of lithium carbonate in the 1960s and the anticonvulsants in the 1980s. Whereas patients with bipolar disorder tended to follow deteriorating courses in the pre-pharmacological era (Cutler & Post, 1982), lithium or anticonvulsant treated patients often remain out of the hospital for extended periods (Goodwin & Jamison, 1990; Patel, DelBello, Bryan, Adler, et

al., 2006). Nonetheless, there is increasing recognition that pharmacotherapy alone forestalls but does not prevent relapses of bipolar episodes (Miklowitz, Simoneau, George, Richards, et al., 2000). Despite the use of mood-stabilizing agents, data suggest relapse rates as high as 40% in 1 year, 60% in 2 years, and 73% in 5 or more years (Gitlin, Swendsen, Heller, & Hammen, 1995). Moreover, the efficacy of medications is limited by the fact that poor medication compliance occurs in 50–67% of individuals with bipolar disorder within the first year of treatment (Keck, McElroy, Strakowski, West, et al., 1998).

Recognizing the limitations of pharmacotherapy alone, a 1996 report by the National Institute of Mental Health (Prien & Rush, 1996) recommended developing adjunctive psychosocial interventions as a central focus for research on bipolar disorder. The importance of psychosocial interventions is also highlighted by the fact that environmental variables play an important role in determining whether an individual at risk develops bipolar disorder, and the timing, frequency, and polarity (depressive vs. hypomanic/manic) of his/her bipolar episodes (Ellicott, Hammen, Gitlin, Brown, & Jamison, 1990; Nusslock, Abramson, Harmon-Jones, Alloy, & Hogan, 2007). Over the past decade, researchers have begun to answer this call, and currently there are three psychosocial interventions for bipolar disorder that have shown promise as adjuncts to pharmacotherapy – Cognitive-Behavioral Therapy, Psychoeducational Interventions, and Interpersonal and Social Rhythm Therapy. Growing evidence highlights the efficacy of these interventions, as indicated in a recent meta-analysis (Scott, Colom, & Vieta, 2007) which reported a significant reduction in relapse rates (~ 40%) for individuals with bipolar disorder engaged in psychosocial treatment. Moreover, findings from the multisite Systematic Treatment Enhancement Program for Bipolar Disorder (STEP-BD) indicate that each of the three existing psychosocial interventions for bipolar disorder enhance life functioning (Miklowitz, Otto, Frank, Reilly-Harrington, Kogan, et al., 2007) and hasten recovery from a bipolar depressive episode (Miklowitz, et al., 2007).

All three of the psychosocial interventions for bipolar disorder were developed by researchers and clinicians translating basic science on bipolar disorder into the clinical arena. For example, Cognitive-Behavioral Therapy has been applied to bipolar disorder given research suggesting that the logic of cognitive vulnerability-stress theories of unipolar depression (Abramson, Metalksy, & Alloy, 1989; Beck, 1967) extends to understanding bipolar episodes and symptoms (Alloy, Reilly-Harrington, Fresco, Whitehouse, & Zechmeister, 1999; Reilly-Harrington, Alloy, Fresco, & Whitehouse, 1999). Psychoeducational interventions developed from findings that life events such as levels of familial expressed emotion (i.e., criticism, hostility, and/or emotional over-involvement [Miklowitz, et al., 1988]), low parental warmth (Geller, Craney, Bolhofner, Nickelsburg, Williams, & Zimmerman, 2002), and social support (Johnson, Winett, Meyer, Greenhouse, & Miller, 1999) predict the course of bipolar disorder. Interpersonal and Social Rhythm Therapy is grounded in the circadian rhythm and zeitgeber theory of bipolar disorder (Ehlers, Frank, & Kupfer, 1988; Malkoff-Schwartz, et al., 1998).

Over the past few decades, exciting work has emerged providing support for another model of bipolar disorder – the behavioral approach system (BAS) dysregulation theory. This theory proposes that weak regulation of the behavioral approach system (BAS), a system involved in approach to reward, might be involved in hypomanic/manic highs and depressive lows that characterize bipolar spectrum disorders (Depue & Iacono, 1989; Depue, Krauss, & Spoont, 1987; Urosevic, Abramson, Harmon-Jones, & Alloy, 2008). The BAS dysregulation theory is compelling insofar as it proposes specific psychosocial factors relevant to understanding the etiology and course of bipolar disorder, as well as provides a unified model of both poles of the disorder – depression and hypomania/mania. Recently, Lam and colleagues have begun to incorporate findings in line with the BAS dysregulation theory into their treatment protocols (Lam, Watkins, Hayward, Bright, et al., 2003). The objective of the current article is to bring attention to this important development and to examine the extent to which the BAS

dysregulation theory can inform the treatment of bipolar disorder. The article is organized as follows: First, we provide an overview of the BAS dysregulation theory. Second, we review theory and research on the three psychosocial interventions for bipolar disorder and discuss how findings from the BAS dysregulation theory can help inform these interventions.

Behavioral Approach System Dysregulation Theory

Researchers suggest that two psychobiological systems are critical in regulating behavior (Gray, 1981). One of these systems, the behavioral approach system (BAS), is hypothesized to regulate approach behavior to attain rewards and goals, whereas the other, the behavioral inhibition system (BIS), regulates inhibition of behavior in response to threat and punishment. We focus on the BAS because theory and research underscore its importance in bipolar disorder. The BAS is activated by signals of reward as well as punishment avoidance cues (Fowles, 1988). These signals can be either external (e.g., presence of a goal/reward) or internal (expectancies of goal-attainment). The objective of the BAS is to regulate appetitive motivation and goal-directed behavior in order to obtain rewards and/or avoid punishment. The BAS has been implicated in the generation of positive emotions (Depue & Iacono, 1989; Gray, 1994), and, in particular, goal-striving related positive affect and drive/incentive motivation (Fowles, 1988). Activation of the BAS causes the person to increase cognitive activity aimed at promoting goal attainment (e.g., hope, self-efficacy, planning [Depue & Iacono, 1989]). Also, certain negative emotions sometimes accompany approach behavior, as research has revealed that anger is associated with heightened BAS activity (see Harmon-Jones, 2004 for review). Finally, research has implicated dopaminergic projections of the Ventral Tegmental Area (Depue & Iacono, 1989), as well as relative left frontal cerebral activity (Harmon-Jones & Allen, 1997; Sutton & Davidson, 1997), as neurobiological processes important in BAS function.¹

The BAS dysregulation theory (Depue & Iacono, 1989; Depue et al., 1987; Urosevic et al., 2008) proposes that weak regulation of the behavioral approach system is involved in the rollercoaster of hypomanic/manic highs and depressive lows that characterize bipolar spectrum disorders. Individuals with bipolar disorder experience excessive variability in BAS activity (e.g., extreme fluctuations in approach motivation). Consistent with theory, research demonstrates that 1) individuals with a bipolar spectrum disorder experience excessive within day, and between day, variability on a variety of indices of BAS (Depue, Slater, Wolfstetter-Kausch, Klein, Goplerud, & Farr, 1981), 2) this variability is a temperamental quality of bipolar disorder (i.e., state-independent; Depue et al., 1987; 1989), and 3) this variability predicts relapse over time (Depue et al., 1981). A source of this variability, according to the theory, is that individuals with bipolar disorder have an overly-sensitive BAS that is hyper-responsive to BAS relevant cues. This BAS hyper-responsivity can lead to excessive BAS activity in response to events involving themes of goal-striving and attainment, reward incentive, and anger-evocation. This excessive increase in BAS activity in vulnerable individuals is hypothesized to be reflected in hypomanic/manic symptoms such as euphoria, excessive goal-seeking behavior, decreased need for sleep, irritability, excessive self-confidence, etc. (Depue et al., 1987; 1989). In contrast, depressive symptoms such as sadness, low energy, anhedonia, psychomotor retardation, and hopelessness reflect a shutdown or an excessive decrease in BAS

¹Although there is some evidence for specific brain areas associated with BAS functioning (see Depue & Collins, 1999, for review), there are still many unexamined questions regarding the neural circuitry of this system. Thus, at this point in time, the BAS is more of a theoretical construct than a well defined neurobiological pathway. Throughout the present article we occasionally use the term BAS activation and BAS deactivation. With regards to BAS activation, we are not referring to the activation of specific neural structures, but rather more generally to the activation of an approach oriented motivational state. Similarly, with regards to the term BAS deactivation, we are not referring to the deactivation of specific neural structures, but rather to the deactivation or “shutdown” of an approach oriented state.

activity in response to BAS deactivation-relevant events such as definite failure and non-attainment of a desired goal (Depue et al., 1987).

The empirical testing of the BAS dysregulation theory has been aided by the development of a self-report measure of BAS sensitivity by Carver and White (1994), the BAS scale of the BIS/BAS scales. The BAS subscales tap individual differences in BAS sensitivity (i.e., BAS responsiveness to relevant stimuli). Compared to relevant control groups, individuals with bipolar I disorder (Meyer, Johnson, & Winters, 2001) and people prone to hypomanic symptoms (Meyer, Johnson, & Carver, 1999) show elevated BAS scores. In a bipolar I sample, high BAS sensitivity at recovery predicted an increase in manic symptoms over six-months (Meyer et al., 2001). In addition, high BAS sensitivity prospectively predicted a greater likelihood and shorter time to onset of bipolar episodes over a 3.5 year follow-up period among individuals with a bipolar II and/or cyclothymia diagnosis (Alloy, Abramson, Walshaw, Cogswell, Hughes, et al., 2008). Research also has demonstrated that individuals with a bipolar spectrum disorder evidence elevated responses on psychophysiological indices of BAS sensitivity (Harmon-Jones, et al. 2002; Harmon-Jones, et al., 2008). We now review psychosocial interventions for bipolar disorder and examine the extent to which the BAS dysregulation model can inform these interventions.

Cognitive-Behavioral Therapy for Bipolar Disorder

Cognitive-Behavioral Therapy (CBT) emerged from highly successful cognitive models of unipolar depression. These models hypothesize that maladaptive cognitive patterns [negative styles for inferring causes, consequences, and self-worth implications in hopelessness theory (Abramson et al., 1989) and negative self-schemata and dysfunctional attitudes in Beck's (1967) theory] act as vulnerabilities for depression when individuals experience stressful life events. These maladaptive cognitive styles increase the likelihood of negative appraisals of negative life events, thereby leading to hopelessness and negative views of one's self and personal world, and ultimately, depressive symptoms. Cognitive theories of depression have garnered considerable empirical support from cross-sectional studies, prospective studies, and behavioral high risk studies (See Abramson, Alloy, Hankin, Haefel, MacCoon, & Gibb, 2002 for review).

CBT for unipolar depression identifies, challenges, and ultimately aims to modify cognitive styles to generate a less depressogenic information processing scheme (Hollon, 2006). CBT is effective for treating unipolar depression (for review, see Deckersbach, Gershuny, & Otto, 2000), with research indicating that this efficacy extends to severe and treatment resistant cases (Fava, Grandi, Zielesny, Rafanelli, et al., 1996). CBT also appears to protect against relapse in patients with unipolar depression and CBT offers relapse protection in the same range as maintenance pharmacotherapy (Evans, Hollon, DeRubeis, Piasecki, Grove, et al., 1992).

Over the past two decades, growing evidence indicates that the cognitive processes involved in unipolar depression also play a role in the bipolar spectrum disorders. Recent reviews of research on cognition in bipolar disorder (See Alloy, Abramson, Neeren, Walshaw, Urosevic, & Nusslock, 2006 for review; Alloy, Abramson, Urosevic, Bender, & Wagner, 2009) concluded that individuals with bipolar spectrum disorders exhibit underlying cognitive patterns as negative as those of unipolar depressed persons overall, but with certain unique characteristics (see below). In addition, there is growing evidence that cognitive styles also predict the course of bipolar disorder, alone or in combination with relevant life events. For example, Scott and Pope (2003) reported that cognitive profiles were the most robust predictor of relapse at 12-month follow-up in patients with hypomania. Moreover, Francis-Raniere, Alloy, and Abramson (2006) found that cognitive styles interacted with congruent negative events to predict increases in depressive symptoms and with congruent positive events to

predict increases in hypomanic symptoms over follow-up. Accordingly, cognitive-behavioral therapy is beginning to be applied to bipolar spectrum disorders as an adjunctive treatment to pharmacotherapy.

Although there is variability in how CBT is implemented for bipolar disorder, most protocols address at least the three following themes: medication adherence, biased or dysfunctional thinking, and psychosocial difficulties (Basco & Rush, 1996; Otto, Reilly-Harrington, & Sachs, 2003; Fava, Bartolucci, Rafanelli, & Mangelli, 2001). Regarding, medication adherence, clients are informed of the importance of strict treatment adherence and educated on pharmacology, toxicity, side effects, and the positive effects that mood-stabilizers can have on the course of bipolar disorder. Clients are provided behavioral strategies to help increase adherence, and receive psychoeducation on the nature of bipolar disorder and its corresponding symptoms. Regarding, biased or dysfunctional thinking, clients are first introduced to the concept that negative thinking can influence the interpretation of events and subsequent emotions. Clients are encouraged to monitor their thoughts, provide evidence for and against negative cognitions, and to replace maladaptive interpretations of events with more adaptive explanations. The challenge for both the clinician and client, however, is that they not only need to address the client's depressogenic cognitions, but also cognitions associated with hypomanic and/or manic episodes. Clients are educated in identifying increased self-esteem and "positively biased" cognitions as indicators of possible hypomania/mania. By challenging and keeping these cognitions "in check", the client, ideally, can manage or preempt a bipolar episode. Lastly, CBT therapists typically addresses psychosocial difficulties associated with bipolar disorder. The therapist helps the client identify strengths that facilitate psychosocial adjustment and weaknesses that may interfere with functioning. Clients are then taught specific problem-solving and behavioral skills to help manage these psychosocial stressors.

The small number of studies conducted to date provides reasonably consistent evidence that CBT has a positive prophylactic effect on bipolar disorder and preliminary evidence that it is also effective in managing acute episodes of bipolar depression. Two uncontrolled studies (Fava, Bartolucci, Rafanelli, & Mangelli, 2001; Patelis-Siotis, et al., 2001) found that CBT reduced depressive and manic episode relapse rates over a 30-month follow-up compared to the 30 months prior to beginning CBT, and improved psychosocial functioning. Fava and colleagues (2001) further reported that CBT was associated with a significant reduction in subsyndromal symptoms during remission, which is of import given that subsyndromal symptoms increase risk of relapse (Benazzi, 2001). Four randomized controlled trials of CBT for individuals with bipolar disorder receiving maintenance medication provide further evidence of the prophylactic effects of CBT. In a study involving individuals with both bipolar I and bipolar II, Scott et al. (2001) reported that compared to a wait list control, their CBT group exhibited 60% fewer relapses and fewer hospitalizations during an 18-month follow-up, significant improvements in global functioning and some symptoms (particularly depressive symptoms), and greater medication adherence. Cochran (1984) also found that their CBT plus lithium group showed greater medication compliance than a lithium-only group, as well as fewer hospitalizations and mood episodes. Lam and colleagues (2000) reported that compared to a "treatment as usual" control group, a CBT group had significantly fewer bipolar episodes and hospitalizations throughout a 12-month follow-up, and lower depressive and manic symptoms. In a follow-up, Lam and colleagues (2003) found that the CBT group had significantly fewer bipolar episodes and hospitalizations relative to a "treatment as usual" control group. The proportion of patients in the control group who relapsed during follow-up was 75% compared with 44% in the CBT group. Additionally, the CBT group had fewer days in episode, less residual depressive symptoms, and less manic symptom fluctuation during the 12-month follow-up. One study (Zaretsky, Segal, & Gemar, 1999) examined the efficacy of CBT for managing an acute episode of depression in individuals with bipolar I and bipolar II to standard CBT for unipolar depression. The authors reported that the two forms of CBT were

equally effective. Additional support for the efficacy of CBT in managing acute bipolar depression comes from the STEP-BD study by Miklowitz and colleagues (2007) mentioned earlier in which all three of the psychosocial interventions (CBT, Family-Focused Treatment, and Interpersonal and Social Rhythm Therapy) for bipolar disorder hastened the recovery from a depressive episode (Miklowitz, Otto, et al., 2007). Lastly, work by Scott and colleagues (2006) suggest that severity of course may moderate client's responsiveness to CBT. The authors report that CBT was more effective than treatment as usual in those with fewer than 12 previous episodes, but less effective in those with more episodes.

CBT for Bipolar Disorder: Perspective from the BAS Dysregulation Theory

CBT was developed from cognitive models of unipolar depression and was subsequently applied to bipolar disorder. Research indicates, however, that, although negative cognitive styles characterize unipolar depression and bipolar disorder alike (Alloy et al., 2005; Johnson & Kizer, 2002), individuals with bipolar disorder exhibit cognitive profiles with unique characteristics that are in line with the high drive/incentive motivation associated with high BAS sensitivity (Alloy et al., 2005; 2006; Urosevic et al., 2008). This is in contrast to the dependency and attachment attitudes typically observed among unipolar depressed individuals (see Hammen et al., 1992, for alternative findings). This next section examines the implications of research on the cognitive profiles of individuals with bipolar disorder in line with the BAS dysregulation theory for CBT for bipolar disorder. As outlined above, CBT for bipolar disorder typically addresses medication adherence, biased or dysfunctional thinking, and psychosocial difficulties. We argue that research in line with the BAS dysregulation theory may be particularly helpful in informing the biased or dysfunctional thinking component of CBT for bipolar disorder. Specifically, clinicians and scientists informed by research suggesting that cognitive profiles of individuals with bipolar disorder are in line with high drive/incentive motivation may be able to more effectively identify and address the cognitive distortions of their clients.

In traditional CBT for unipolar depression and bipolar disorder, there are two levels of cognitions that a CBT therapist typically addresses with a client (Basco & Rush, 1996; Greenberger & Padesky, 1995; Lam, Jones, Hayward, & Bright, 1999). The first level are state-dependent 'automatic thoughts' that typically occur prior to or during a mood disorder episode. Next, the clinician addresses the underlying state-independent world view, schema, or assumptions that an individual has about reality.

In addressing state-dependent automatic thoughts, there is growing consensus that CBT therapists should target the cognitive prodromes of bipolar episodes as opposed to automatic thoughts experienced during full-blown episodes (Lam et al., 2003). In medicine, prodromes are defined as the early signs and symptoms that herald a full episode (Molnar, Feeney, & Fava, 1988). Research indicates that individuals with bipolar disorder, as well as their relatives, are able to report prodromes reliably (Keitner, Solomon, Ryan, Miller, Mallinger, et al., 1996; Lam, Wong, & Sham, 2001). Two advantages in targeting prodromes in the treatment of bipolar disorder are: 1) full-blown bipolar episodes may overwhelm the coping strategies of patients, where these strategies may be highly effective in managing prodromes of bipolar episodes; and 2) the prodromal period precedes the full bipolar syndrome (Smith & Tarrier, 1992), representing a window in which intervention might protect patients from relapse. In line with this view, good coping skills during prodromal periods have been associated with higher social functioning (Lam & Wong, 1997), which predicts longer intervals between episodes (Gitlin et al., 1995).

Growing evidence suggests that the cognitive prodromes of manic episodes are characterized by extreme-goal setting and heightened expectations of success in the achievement domain.

This is consistent with the BAS's involvement in regulating appetitive motivation and goal-directed activity (Gray, 1981; 1994). Meyers, Beevers, and Johnson (2004) noted that current symptoms of hypomania/mania positively correlated with higher expectations of attaining upcoming goals. The relationship between hypomanic/manic symptoms and success expectancies is most pronounced following a proposed BAS activation relevant event (i.e., goal – attainment [Johnson, Ruggero, & Carver, 2005]). Individuals with bipolar disorder seem to over-interpret this initial success and become unrealistically confident about achieving the next goal (Johnson, et al., 2005). They may express beliefs such as 'there is nothing that I can not do', 'I have to obtain the goal', 'I am invincible.' This unrealistic confidence may fuel excessive goal-striving behaviors and even higher goal-setting (Johnson, 2005). Indeed, increased goal-directed activity is one of the two most common behavioral prodromes of mania and has been independently associated with increased rates of manic episodes (Lam, Wong, & Sham, 2001). In line with this research, certain clinicians have begun to take a BAS dysregulation perspective and are giving particular attention to the importance of helping patients understand the relationship between ambitious goal-setting and the onset of manic episodes (Lam et al., 2003). These clinicians argue that the therapist and patient can develop a treatment plan in which surges of ambitious goal-setting and confidence are identified and challenged during the prodromal period. An important cognitive strategy for dealing with prodromal mania is reframing the ambitious goal-setting and surge of confidence as early symptoms of mania as opposed to thoughts to be considered in their own right (Lam, Jones, Hayward, & Bright, 1999). This allows the client to more objectively identify what is a manic prodrome versus simply a good mood. Lam and Wong (1997) report that the optimal coping strategies for prodromes of mania involve behavioral deactivation strategies including 'modifying high activities', 'restraining oneself', and 'engaging in calming activities.' Accordingly, an extreme surge in ambition (i.e., BAS hyper-activation) should serve as a cue to the individual with bipolar disorder to reduce goal-striving behavior and normalize social and sleep routines. We argue that CBT as a whole for bipolar disorder would benefit from the perspective taken by Lam and colleagues of giving particularly close attention to ambitious goal-setting and increased goal-directed activity during the prodromal period.

Where manic prodromes of extreme confidence and ambitious goal-setting typically occur following success in the achievement domain (Johnson, 2005), depressive prodromes of low self-esteem and decreased goal-setting typically occur in response to failure in the achievement domain (Lam, et al., 2000). According to Urosevic and colleagues' (2008) expanded BAS dysregulation theory, low efficacy expectancy resulting from failure in the achievement domain leads to BAS deactivation. This BAS deactivation is characterized by decreased goal-setting and expectancy of success, followed by reduced motivation and goal-oriented behaviors. As noted by an individual with bipolar disorder, "I know that I am becoming depressed when I think there is no point in pursuing goals." Accordingly, a CBT therapist taking a BAS dysregulation perspective should be particularly primed to target cognitive-behavioral prodromes of depression in the achievement domain. Once a client's prodromes have been identified, the clinician can assist the client in monitoring and modifying cognitive prodromes of depression to reduce the risk of relapse. Lam and Wong (1997) report that the coping strategy most often employed among individuals with bipolar disorder in the "Good Coping Group" involved 'getting myself organized and keeping busy' (i.e., increased BAS activity). Accordingly, behavioral activation strategies such as exercise, goal-striving, and pleasurable activities can be employed in order to target the BAS deactivation often associated with bipolar depression (Depue et al., 1987; 1989).

Once the clinician and client have addressed state-dependent automatic thoughts – which we argue are best addressed during the prodromal period (i.e., cognitive prodromes) –, the next step in traditional CBT is to address the underlying assumptions or cognitive style that the bipolar patient has about his/her self and world (Basco & Rush, 1996; Lam et al., 1999;

2003). These assumptions are state-independent cognitions from which automatic thoughts arise in response to relevant life events (Lam et al., 1999). Assumptions can be considered rules for living or conditional statements that incorporate an 'if....then' component (e.g., "if I am not the best, I am nothing"). Once identified, strategies that have been outlined to address dysfunctional assumptions among individuals with bipolar disorder include, but are not limited to, the use of thought records, collaborative empiricism, and behavioral experiments (Basco & Rush, 1996; Lam et al., 1999; 2003; Otto et al., 1999).

In line with the BAS dysregulation theory's focus on high drive/incentive motivation, as well as the research just outlined on state-dependent cognitions in manic and depressive episodes, the cognitive style of individuals with bipolar disorder is marked by ambitious goal-setting, autonomy, and perfectionism in the achievement domain (Johnson, 2005; Lam, Wright, & Smith, 2004). Clinical observations report high goal setting, drive, and work motivation among individuals with a history of mania that are "inappropriately high" (Peven & Shulman, 1983, p.13). Researchers have also found that a history of bipolar disorder is associated with stronger emphasis on goal-pursuit and goal-attainment. For example, two studies found that individuals with a history of mania are more likely to endorse items reflecting perfectionism and the need to achieve goals on the Dysfunctional Attitudes Scale relative to individuals with no mood disorder (Lam, et al., 2004; Scott, Stanton, Garland, & Ferrier, 2000). Alloy and colleagues (2009) found that euthymic bipolar spectrum participants scored higher on BAS-relevant cognitive dimensions of perfectionism, autonomy, and self-criticism than did normal controls, but not on non-BAS-relevant dimensions of sociotropy, dependency, or approval from others. Moreover, in this study, only BAS-relevant cognitive dimensions predicted the likelihood of onset of depressive and hypomanic/manic episodes among individuals with bipolar disorder over a 3.2 year follow-up. Further, a study by Spielberger and colleagues (1963) reported that 93% of individuals with a history of bipolar disorder endorsed the item "I nearly always strive hard for personal achievement." Ambitious goal-setting and perfectionism in the achievement domain among individuals with bipolar disorder appears to be a state-independent feature of the disorder given: 1) studies of goal-oriented cognitive styles in bipolar disorder have been conducted when individuals are in a remitted/euthymic state (Lam, et al., 2004; Scott et al., 2000), 2) the effect holds even after controlling for baseline mood state (Alloy et al., 2008; Johnson, Ruggero, & Carver, 2005), and 3) current symptoms of bipolar disorder do not correlate with ambitious goal-setting (Lozano & Johnson, 2001). Researchers have extended these findings to individuals with subsyndromal bipolar disorder (Johnson & Carver, 2006) and non-affected family members of individuals with bipolar disorder (see Johnson, 2005, for review). These findings suggest that ambitious goal setting and perfectionism in the achievement domain may index a temperamental/cognitive vulnerability to bipolar disorder. In line with this view, high scores on the Neuroticism-Extraversion-Openness (NEO) Inventory achievement-striving scale predicted increases in bipolar symptoms over 6 months (Lozano & Johnson, 2001).

What is the incremental value of a BAS dysregulation perspective in the application of CBT to bipolar disorder? We argue that research indicating that ambitious goal-setting and perfectionism in the achievement domain characterize the cognitive style of bipolar disorder suggest that CBT therapists may benefit from being particularly sensitive to cognitive distortions in the achievement domain of their clients with bipolar disorder. To date, however, only one of the six studies outlined above on the efficacy of CBT for bipolar disorder systematically targeted goal-oriented cognitions. In this study, Lam and colleagues (2003) randomly assigned 103 individuals with bipolar disorder to CBT or medication management. In the CBT group, the therapists specifically targeted extreme striving attitudes. As reported above, the authors found evidence that CBT was protective for both bipolar depression and mania, with the CBT group experiencing significantly fewer episodes and hospitalizations relative to the control group. Importantly, highly driven and extreme goal attainment beliefs

were identified as vulnerability factors, and at 6 month follow-up, the CBT group scored significantly lower than a comparison control group regarding such goal-striving attitudes.

In summary, the BAS dysregulation theory proposes high drive/incentive motivation and high BAS sensitivity characterize bipolar disorder (Depue & Iacono, 1989; Fowles, 1988). Consistent with theory, growing evidence suggests that the cognitive profiles of bipolar disorder are characterized by extreme goal-striving tendencies and perfectionism in the achievement domain, rather than the sociotropic, dependent features often exhibited by unipolar individuals. Accordingly, we propose that an important contribution of the BAS dysregulation theory to CBT for bipolar disorder is highlighting the potential value of targeting achievement-oriented cognitive profiles in managing both the depressive and manic phases of bipolar disorder. Preliminary research supports this proposal (Lam et al., 2003). Further work, however, is needed to test the nuances of this issue. Although research indicates that bipolar-spectrum individuals have higher mean levels on BAS-relevant cognitive dimensions (perfectionism, autonomy, self-criticism), there are likely notable individual differences in such cognitive profiles. It will be important for researchers and clinicians to examine these individual differences, and adjust their treatment plan accordingly.

Psychoeducational Interventions

A second prominent psychosocial intervention for bipolar disorder is psychoeducation. Models of psychoeducation for bipolar disorder have been informed by the literature on psychoeducational treatments for schizophrenia (Miklowitz & Goldstein, 1997). Early research demonstrated that family-based psychoeducation reduced relapse rates and improved psychosocial functioning in individuals with schizophrenia over a 2 year period (Falloon, Boyd, McGill, Williamson, et al., 1985; Hogarty, Anderson, Reiss, Kornblith, et al., 1986). Given the fact that bipolar disorder shares many of the clinical characteristics of schizophrenia (e.g., relapse-remission course, need for maintenance medication), clinicians and researchers began examining psychoeducation as a treatment for bipolar disorder.

Central to psychoeducational models for bipolar disorder is research indicating that environmental variables influence the timing, frequency, and polarity (depressive vs. hypomanic/manic) of bipolar episodes (see Craighead & Miklowitz, 2000 for review). Environmental variables in bipolar disorder have been examined in two domains: life events and family/marital discord. Regarding the former domain, research indicates that individuals with bipolar disorder experience an elevated rate of life events in the period preceding the onset of an episode relative to other periods in their life (Alloy et al., 2005; Ambelas, 1979, 1987; Johnson & Roberts, 1995). For example, a prospective study that followed bipolar individuals across a 2-year period found that individuals who received the highest total life event scores had 4.53 times the risk of relapse compared to individuals who did not experience such events (Ellicott, et al., 1990). Moreover, Perris (1984) found that a greater number of negative events preceded the onset of bipolar depression than the onset of unipolar depression.

The second domain – family discord as a trigger for recurrences- has focused on “expressed emotion” (EE) attitudes among care-giving relatives. EE reflects the extent to which relatives of patients express critical, hostile, or emotionally over-involved attitudes towards their disturbed family member (Miklowitz, et al., 1988). High EE in family members of individuals with bipolar disorder predicts a more pernicious course relative to bipolar individuals whose family members express low-EE attitudes (Miklowitz, et al., 1988; 2000). Furthermore, high-EE relative/patient pairs have more conflict in laboratory assessments during the post-episode stabilization period than low-EE relative/patient pairs (Simoneau, Miklowitz, & Saleem, 1998).

Psychoeducation for bipolar disorder has been implemented in different forms. One commonly used approach is a 21 session Family-Focused Treatment (FFT) program (Miklowitz & Goldstein, 1990; Miklowitz & Goldstein, 1997). This protocol involves three primary components: psychoeducation about bipolar disorder, communication enhancement, and problem-solving skills training. The *Psychoeducation* component includes information about the course, causes, and treatment of bipolar disorder. Clients and their family members are educated on the important role that life events and stressors play in the course of bipolar disorder and life events are viewed as “occasion setters” for the onset of bipolar episodes. Similar to CBT, clients and their family members are educated on the importance of identifying early warning signs or prodromes of bipolar episodes given that full-blown episodes often overwhelm a person's coping strategies. The *Communication Enhancement* component is concerned with enhancing the quality and efficiency of the family's or couple's communication. Family members are informed about the role of the family in the course of bipolar disorder, and that family tension, when combined with ineffective or negative communication, can result in increased risk of relapse. Family members are acquainted with new communication strategies for negotiating family tension, including expressing positive feelings, active listening, making positive requests for change, and expressing negative feelings. Finally, the *Problem-Solving Skills Training* involves identifying specific problems faced by families during the aftermath of an episode and strategies for managing these problem areas. Family members are assisted in defining the problem areas, brainstorming solutions, and implementing these solutions.

Psychoeducation programs have also been implemented in group and marital format. Group psychoeducation for bipolar disorder addresses illness awareness, treatment compliance, early detection of symptoms, and life style regularity (Colom, et al., 2003). Marital psychoeducation addresses these issues in the context of an intimate relationship (Clarkin, Carpenter, Hull, Wilner, & Glick, 1998).

Research indicates that psychoeducational interventions for bipolar disorder have a positive prophylactic effect for a number of clinically relevant indices. Most of the studies used a randomized controlled trial in which clients were assigned to either a psychoeducational intervention or Crisis Management (CM). All clients received concomitant pharmacotherapy. Both FFT (Miklowitz, George, Richards, et al., 2003; Miklowitz, et al., 2000; Rea, Tompson, Miklowitz, Goldstein, et al., 2003) and group psychoeducation (Colom, et al., 2003) were associated with reduced relapse rates relative to CM. Miklowitz and colleagues (2000; 2003) reported that clients undergoing FFT had longer survival intervals (i.e., time to relapse) and a greater decrease in affective symptoms than clients undergoing CM at follow-up. FFT (Rea, et al., 2003) and group psychoeducation (Colom et al., 2003) have also been associated with reduced hospitalization, and a psychoeducation intervention for clients and their spouses has been associated with increased medication compliance (Clarkin et al., 1998). However, these studies indicated that psychoeducation is typically more effective in managing depression than mania. In Colom et al. (2003), at the end of follow-up, group psychoeducation clients had a lower number of all types of recurrences than CM patients, except for mania.

Two studies have examined the effect of FFT on expressed emotion and the interaction patterns of individuals with bipolar disorder and their relatives. One of these studies found that following treatment, clients and relatives who received FFT showed a greater amount of positive interactional behavior as compared to those in the CM condition (Simoneau, Miklowitz, Richards, Saleem, et al., 1999). Additionally, Honig and colleagues (1997) found that FFT lowered EE and that clients with key relatives who had low EE had a better course (lower hospital admissions) than clients with high EE key relatives.

Over the past decade, clinicians have begun to apply family-based psychoeducation for children and adolescents with bipolar disorder. Regarding children, family-based psychoeducation is associated with an increase among parents in their knowledge of bipolar disorder (Fristad, Arnett, & Gavazzi, 1998; Fristad, Goldberg-Arnold, & Gavazzi, 2003), high consumer satisfaction (Fristad, Gavazzi, & Soldano, 1998), greater social/parental support (Fristad, Goldberg-Arnold, & Gavazzi, 2002; 2003), and a decrease in negative expressed emotion (Fristad, Arnett, & Gavazzi, 1998). Miklowitz and colleagues (2004) recently examined the efficacy of FFT for adolescents with bipolar disorder and found that FFT was associated with improvements in both depression and manic symptoms and behavior over a 1-year period.

Psychoeducation for Bipolar Disorder: Perspective from the BAS Dysregulation Theory

Although psychoeducation interventions for bipolar disorder typically address a number of topics, including information about symptoms, medication adherence, and etiology, one important objective is to help clients and their family members identify triggers of bipolar episodes (Morris, Miklowitz, & Waxmonsky, 2007). Bipolar disorder is presented in the context of a vulnerability-stress model in which genetic, biological, and social factors interact in bringing about episodes of mood disorder or in protecting against their occurrence (Miklowitz and Goldstein, 1997). Clients and their family members/spouses are educated about the relationship between life stress and affective episodes, encouraged to anticipate future stressors, and encouraged to identify low stress activities. However, aside from work on Expressed Emotion and circadian rhythm disruption, many of the psychoeducational manuals take a more generalist approach to life stress and do not explicitly discuss specific types of life events that have been found to precipitate bipolar episodes in the research literature. In many respects, this is understandable given that much of the research on life events and bipolar disorder has been largely atheoretical, with many studies relying on global classifications of events such as “negative events”, “stressful events”, and “severe events.” However, an advantage of the BAS dysregulation theory is that it makes specific predictions about the types of life events relevant to the onset of both manic and depressive episodes. Accordingly, as we outline below, we propose that research in line with the BAS dysregulation theory may be helpful to psychoeducation interventions in identifying specific types of life events relevant to the course of bipolar disorder.

BAS activation and BAS deactivation relevant life events

According to the BAS dysregulation theory, there are two prerequisites for the development of bipolar symptoms/episodes – an occurrence of an event that is appraised as BAS relevant and a hypersensitivity to BAS relevant events (Urosevic et al., 2008). The BAS hypersensitivity characterizing individuals at risk for bipolar disorder “...transforms the normally mild effects of events into periods of dysregulation. That is, the biobehavioral systems of bipolar prone individuals will be more perturbed by stimuli of both positive and negative valence” (Depue et al., 1987, pp. 118-119). In essence, individuals vulnerable to bipolar disorder are unable to effectively regulate their emotions and behavior because their proneness to BAS dysregulation renders them excessively responsive to BAS relevant events. In support, Alloy, Abramson, Whitehouse, Sylvia, et al. (2009) found that BAS-relevant events prospectively predicted increases in bipolar symptoms among bipolar spectrum participants over a one year follow-up.

Two types of BAS relevant events have been proposed – BAS activation and deactivation relevant events. BAS activation-relevant events involve an opportunity for an individual to attain a highly valued reward/goal. Depue and Collins (1999) have proposed a variety of

rewards to be BAS activating – ranging from food, sex, social rewards, money, to long term goals. Alloy et al. (2009) noted that self-reported BAS sensitivities, especially the BAS-Drive subscale, interacted with BAS-activation relevant events to prospectively predict increases in hypomanic symptoms. High BAS participants who experienced more BAS activation relevant events showed the largest increase in hypomanic symptoms. Johnson et al. (2000) found that life events involving high goal-attainment were significantly related to higher levels of follow-up manic, but not depressive, symptoms, even when controlling for baseline manic symptoms. In line with the BAS dysregulation theory's focus on high drive/incentive motivation among individuals with bipolar disorder, positive events that did not involve the attainment of a valued reward were not associated with an increase in manic symptoms. These results are consistent with research showing that the relationship between hypomanic/manic symptoms and success expectancies is most pronounced following a proposed BAS activation relevant event (i.e., goal-attainment [Johnson, Ruggero, & Carver, 2005]). Nusslock and colleagues (2007) extended this work in reporting that a full 42% of bipolar spectrum individuals developed a new episode of hypomania in response to a goal-striving life event (Nusslock, Abramson, Harmon-Jones, Alloy, & Hogan, 2007). In contrast, only 4% of bipolar individuals not exposed to this event developed a new hypomanic episode during the same time. Taken together, the findings of these two studies suggest that goal-striving and goal-attainment life events are two BAS activation relevant events particularly important to the course of bipolar disorder.

BAS deactivation-relevant events involve definite failure to obtain or loss of pertinent reward/goals (Depue et al., 1987; 1989; Urosevic et al., 2008). The lower the expectations of a positive outcome of one's actions, or the more hopeless the individual becomes, the greater the subsequent BAS deactivation will be (Abramson et al., 2002). BAS deactivation-relevant events have been shown to prospectively predict increases in depressive symptoms among individuals with a bipolar spectrum disorder (Alloy, Abramson, Whitehouse, Sylvia, Hafner et al., 2009). Given the BAS dysregulation theory's focus on drive/incentive motivation, and the fact that the cognitive profiles of bipolar individuals are characterized by extreme goal-striving tendencies, perfectionism, and autonomy, we predict that bipolar individuals will be particularly vulnerable to depression in response to failure or loss in the achievement domain.

As indicated, psychoeducational interventions tend to focus on the role that more global life stress plays in the course of bipolar disorder. Research and theory on BAS activation and deactivation events, however, may be able to help provide better resolution on the specific types of life events likely to precipitate bipolar episodes. We propose that events in the achievement domain are particularly salient to the course of bipolar disorder. It may appear counterintuitive to bipolar individuals and their families that events involving the pursuit and attainment of achievement goals can act as risk factors for bipolar episodes. Indeed, bipolar individuals highly value these goal-oriented events (Johnson, 2005). We are not suggesting that bipolar individuals refrain from pursuing or attaining goals. We are suggesting, however, that bipolar individuals enter into these events with an understanding that they may be at heightened risk of relapse. With this understanding, psychoeducational interventions can help bipolar individuals develop problem-solving and coping strategies to regulate the affect these events likely invoke.

Self-generated BAS relevant life events

Clinicians long have noted that individuals with bipolar disorder, often, by their own behavior, create life events that worsen the course of their illness. This “stress generation” effect indicates that depressed or bipolar individuals often experience an increased rate of life events that are dependent on their behavior (Hammen, 1991; Daley, Hammen, Burge, Davila, Paley, et al., 1997). This suggests a “2-hit model” in which individuals with BAS hypersensitivity not only react more strongly to BAS-relevant events, but also are exposed to such events more frequently

through event generation or selection. That is, the stimulus seeking, hyper-driven, and workaholic traits characteristic of individuals with bipolar disorder may create or select the very life events likely to trigger bipolar episodes. In line with this view, Urosevic and colleagues (2009) reported that bipolar spectrum individuals experienced both more BAS activation-relevant (e.g., took on new tasks) and more BAS deactivation-relevant events (e.g., failed at a task) than normal participants.

One example of this might be an individual with bipolar disorder who has recently attained a goal for which she has strived. According to Johnson et al. (2005), this goal-attainment is likely to induce a surge in confidence in the individual and an expectancy of future success in the achievement domain (i.e., cognitive prodrome). This success expectancy leads the individual to excessive goal-striving behaviors and grandiose plans. Given the suggestion that manic prodromes are associated with decreased awareness of danger signs (Lembke & Ketter, 2002), the individual begins making risky financial decisions, takes on more work than she can handle, and engages in excessive pleasurable activities. These self-generated events exacerbate the individual's prodromal symptoms, leading to further dysregulation, and eventually to a full manic episode. Over time, it becomes clear to the person that she is unable to manage her current state. Because she is so overextended, she fails to meet professional/familial obligations; she becomes aware of the financial and/or personal implications of her impulsive decisions, and, as a result, experiences significant failure in the achievement domain. This failure serves as a BAS deactivation relevant event, which may cause a subsequent depressive episode.

This sequence of events could potentially have been preempted had the bipolar individual been aware of the concept of stress generation. There is growing awareness among clinicians of the importance of informing individuals with bipolar disorder and their families that the events that have the most notable impact on the course of bipolar disorder are often generated by the bipolar individual (Lam et al., 1999). However, to date, the majority of psychoeducational manuals do not devote much discussion to self-generated events, particularly as it pertains to the achievement domain. BAS relevant events that seem positive and proactive can, in excess, create chaos and significant stress. The goal is to strike a balance so that the individual with bipolar disorder can enjoy goal-striving, but moderate it so that such behaviors do not result in dysregulation. This is especially true during a prodromal period in which the bipolar individual is hypersensitive to BAS relevant events (Urosevic et al., 2008). Thus, psychoeducational programs may benefit from allocating more attention to the impact of self-generated life events in the achievement domain on the course of bipolar disorder.

This sequence of events also highlights the bi-directional relationship between life events and motivational state. That is, under certain circumstances, the presence of a BAS relevant life event may serve as an initial trigger for an excessive increase or decrease in approach motivation, which, ultimately, may result in a bipolar episode. However, under other circumstances, a person's motivational or affective state may cause them to actually generate or select the particular types of life events that may lead to a more severe bipolar course. Clinicians can help patients understand this process and to monitor both their response and exposure to relevant life events, particularly during prodromal periods.

Expressed Emotion in Achievement Domain

Although less direct, research in line with the BAS dysregulation theory may also have implications for the *communication enhancement training* module of family-focused treatment programs (i.e., FFT). On this topic, elevated drive, ambition, and achievement motivation have not only been observed in individuals with bipolar disorder, but also are characteristic of the family members of individuals with a history of bipolar disorder (Spielberger, et al., 1963).² Family members of bipolar probands have higher lifetime occupational and educational

attainment than the general population (Tsuchiya, Agerbo, Byrne, & Mortensen, 2004). Growing up in such families, individuals with bipolar disorder are likely expected to attain an equivalent degree of professional success. Indeed, 2/3 of individuals with bipolar disorder had academic performance in the good to excellent range prior to the onset of their first bipolar episode (Kutcher, Robertson, & Bird, 1998). These expectations may result in the individual with bipolar disorder engaging in excessive goal-striving behaviors in the presence of possible academic/professional reward, thus putting themselves at risk for a hypomanic/manic episode. These expectations, however, may also result in family members being particularly critical, hostile, and/or over involved when the individual with bipolar disorder experiences failure in the achievement domain. Accordingly, family members of individuals with bipolar disorder may benefit from being particularly attuned to negative “expressed emotion” attitudes in the achievement domain. Presumably, this attention would have benefits for managing both mania and depression. With respect to mania, decreased EE in the achievement domain may help individuals with bipolar disorder moderate extreme goal striving attitudes/behaviors associated with mania tendencies. In terms of depression, decreased EE in the achievement domain may help reduce catastrophic responses to academic/professional failure and increase a sense of social support. Further research is needed to test this prediction.

In summary, we propose that research in line with the BAS dysregulation theory has important implications for psychoeducational interventions for bipolar disorder. Family members and clients may benefit from being educated on the role that BAS activation and deactivation relevant life events have on triggering bipolar episodes. Research indicates that goal-striving and goal-attainment life events are particularly salient BAS activation relevant events (Johnson et al., 2000; Nusslock et al., 2007). We hypothesize that definite failure in the achievement domain is a salient BAS deactivation relevant event, although future research is needed to test this prediction. Second, clinicians should inform their clients of the importance of minimizing excessive goal-striving behaviors, particularly during manic prodromes, in order to minimize exposure to self-generated BAS relevant events. As will be discussed in the next section, self-generated events that may be particularly important to manage are disruptions in social and circadian rhythms. Lastly, with respect to *Communication Enhancement* in family-focused treatment programs, research on BAS dysregulation theory indicates that bipolar disorder is characterized by elevated drive, ambition, and achievement motivation. Thus, negative Expressed Emotion in the achievement domain may be a particularly relevant stressor for bipolar individuals.

Interpersonal and Social Rhythm Therapy (IPSRT)

The Social Rhythm or Zeitgeber Theory (Ehlers, Frank, & Kupfer, 1988; Ehlers, Kupfer, Frank, & Monk, 1993) postulates that life events that disrupt daily rhythms or schedules will be especially likely to precipitate bipolar symptoms and episodes. This theory suggests that individuals with bipolar disorder have a predisposition to circadian rhythm and sleep-wake cycle abnormalities that may be responsible, in part, for the symptomatic manifestations of the illness. In this model, a *social Zeitgeber* is defined as a personal relationship, social demand, or life task, that entrains biological rhythms such as circadian rhythms or the sleep-wake cycle. It is hypothesized that life events (both positive and negative) that involve the disruption or loss of a social Zeitgeber can trigger bipolar episodes by causing a dysregulation of biological rhythms (Ehlers et al., 1988). In line with this view, Wehr and colleagues (1987) have demonstrated that sleep reduction can lead to mania in individuals with bipolar disorder and

²Preliminary evidence suggesting that the similarity in levels of achievement motivation and creativity between bipolar individuals and their non-affected family members is driven, at least in part, by genetic factors comes from two sources: (1) An early study by McNeil (1971) demonstrating that biological, but not adoptive, parents of bipolar individuals displayed high levels of creativity, and (2) research estimating the overall heritability of bipolar disorder to be approximately (70%; Edvardsson, Torgersen, Roysamb, et al., 2008).

sleep deprivation has significant antidepressant effects in individuals with both unipolar and bipolar depression (Leibenluft et al., 1993; Leibenluft and Wehr, 1992). Furthermore, Malkoff-Schwartz et al. (1998; 2000) found that manic patients had significantly more pre-onset stressors characterized by social rhythm disruption (e.g., change in sleep-wake cycle) than did depressed patients with bipolar disorder. Wehr and colleagues (1987) argued that sleep loss may be a common causal pathway in the genesis of mania. Other research has shown that social rhythm disruption precipitates depressive symptoms and episodes among bipolar individuals (Sylvia, et al., in press).

Based on such research, Frank and colleagues (1997; 1999) developed Interpersonal and Social Rhythm Therapy (IPSRT; Frank, Hlastala, Ritenour, Houck, et al., 1997). This therapy integrates interpersonal psychotherapy for unipolar depression (Klerman et al., 1984) with behavioral and environmental strategies to help stabilize irregularities of the sleep-wake cycle among bipolar individuals. Techniques are drawn from interpersonal psychotherapy given the hypothesis that social or interpersonal events play an important role in entraining biological rhythms (Ehlers et al., 1988; 1993). It is proposed that teaching individuals with bipolar disorder skills to navigate interpersonal stressors will help them have more regularity in their biological and circadian rhythms.

With regards social rhythm maintenance, IPSRT focuses on 1) the reciprocal relationships between life stress and the onset of mood disorder symptoms, 2) the importance of maintaining regular daily rhythms and sleep/wake cycles, and 3) the identification and management of potential precipitants of rhythm dysregulation with special attention to interpersonal triggers. Clients are first instructed to monitor their social routines and rhythms using the Social Rhythm Metric (SRM; Monk et al., 1991), a self-report instrument designed to quantify daily life rhythms. Using the SRM, the client ideally begins to see the interplay among instabilities in daily routines, patterns of social stimulation, sleep-wake times, and mood fluctuations. IPSRT then implements behavioral strategies to help the client alter those activities that promote rhythm irregularities (minimizing overstimulation, monitoring the frequency and intensity of social interactions). The client is encouraged to make significant life changes in order to protect the integrity of his or her circadian rhythms and sleep-wake cycle.

The interpersonal techniques employed in IPSRT are similar to those described in Klerman and colleagues description of Interpersonal Psychotherapy for unipolar depression. The therapist determines the important individuals in the client's life, known as the interpersonal inventory. In addition to outlining the "cast of characters" in the client's life, the therapist probes the quality of those relationships and aspects of the relationships that the client would like to change. The therapist also identifies an interpersonal problem area during the initial phase of treatment that will serve as the interpersonal treatment focus. These problem areas include grief, role disputes, role transitions, and interpersonal deficits. Thus, IPSRT is a truly integrated therapy that allows interpersonal and social rhythm strategies to function synergistically.

Preliminary evidence suggests that IPSRT has a positive prophylactic effect as an adjunct to long-term maintenance pharmacotherapy. Frank and colleagues (2005) reported that individuals with bipolar disorder assigned to IPSRT went longer without a new affective episode. Furthermore, IPSRT was associated with higher regularity of social rhythms and this increased regularity was associated with a reduced likelihood of recurrence. In an earlier study, individuals with bipolar disorder who received IPSRT had fewer depressive episodes over the course of a year relative to patients in a comparative clinical management group (Frank, 1999). No differences, however, were observed for rates of manic episodes. Another study compared the efficacy of IPSRT versus clinical management (CM) on rates of suicide attempts among individuals with bipolar I disorder who were followed for an average of 1.4 years (Rucci, Frank, Kostelnik, Fagiolini, et al., 2002). Both IPSRT and CM were associated with a

reduction in suicide attempt risk. Given the low number of suicide attempts, the authors were unable to compare the efficacy of IPSRT to CM. In an attempt to examine the mechanisms through which IPSRT works, Frank and colleagues (1997) examined the extent to which initial exposure to IPSRT leads to lifestyle regularity. IPSRT was efficacious in regularizing the daily lifestyles of recovering bipolar patients, whereas this lifestyle regularity was not observed in the CM group. Research also indicates that changing the treatment protocol (from IPSRT to CM, and vice versa) for individuals with bipolar disorder is associated with higher rates of recurrence and levels of symptomatology (Frank, Swartz, Mallinger, Thase, et al., 1999). The authors note that “it appears that a constant treatment regimen contributes to enhanced stability. By contrast, changing treatment parameters may represent yet another destabilizing pathway to recurrence in bipolar disorder” (p. 585). Lastly, a recent treatment outcome study integrated the central tenets of FFT and IPSRT (Miklowitz et al., 2003), reporting that clients given this combined therapy along with medication showed longer time to relapse than those given CM. Consistent with existing FFT and IPSRT studies, however, the combined treatment had a greater impact on depressive than manic symptoms.

IPSRT for Bipolar Disorder: Perspective from the BAS Dysregulation Theory

A central component of IPSRT is identifying and managing potential precipitants of rhythm dysregulation (Frank, Swartz, & Kupfer, 2000). As outlined above, IPSRT has given particular attention to the disruptive effects that interpersonal events have on circadian rhythms (Frank, et al., 2000). This attention is highly warranted given the important role that such events (childcare, marriage, etc) play in entraining biological rhythms (see Frank, 2007 for review). However, the research outlined in the current article indicating that bipolar disorder is characterized by high drive/incentive motivation indicates that individuals with bipolar disorder may also be especially sensitive to events in the achievement domain. Moreover, this hypersensitivity may put individuals with bipolar disorder at particular risk for social/circadian rhythm disruption in response to such events. Accordingly, we argue that it will be helpful for the IPSRT therapist to be particularly attuned to the social and circadian rhythm disruption associated with both interpersonal and achievement oriented life events.

IPSRT also focuses primarily on the disruptive effects that independent life events (i.e., caused by external factors and/or not related to the individual's behavior) have on circadian rhythms. As noted by Frank and colleagues (2000), “the search for triggers of rhythm disruption leads the IPSRT therapist to comb the patient's history in search of *external* sources of rhythm disruption” (pp. 599). However, the stress-generation hypothesis proposes that individuals with bipolar disorder often create or select the very events that trigger bipolar episodes (Hammen, 1991; Daley et al., 1997). Accordingly, the social/circadian rhythm disruption typically observed prior to a bipolar episode is often generated by the patient rather than imposed on her by an external source (Akiskal et al., 1996; Lam et al., 2003). The BAS dysregulation theory expands on this view, proposing that a common source of circadian rhythm disruption is self-generated, goal-striving behaviors. The drive to succeed can often lead to working excessively long hours and neglecting important social routines (Lam et al., 1999). In extreme cases, this can involve sleep disruption and total disruption to social routines, which can lead to disruption of circadian rhythms. In line with this view, the two most common prodromes of a manic episode are increased goal-directed activity and decreased need for sleep (Lam & Wong, 1997). Indeed, Molnar et al. (1988) reported that 100% of bipolar participants reported increased goal-directed activity during a manic prodrome and 90% reported decreased sleep. This highlights the important relationship between self-generated goal-oriented events and social/circadian rhythm disruption in the onset and/or exacerbation of bipolar episodes. It also highlights the importance of regulating excessive goal-striving related behaviors during manic prodromal periods.

We are not suggesting that IPSRT reduce its focus from the disruptive effects of independent interpersonal events. We are suggesting, however, that it may be helpful for the IPSRT therapist to be particularly sensitive to the disruptive effects of both independent and self-generated triggers of rhythm disruption. Following the BAS dysregulation theory (Depue et al., 1987), we propose that self-generated events in the achievement domain may be particularly relevant triggers for social/circadian rhythm disruption. Many of the therapeutic strategies designed to address the effect of independent interpersonal events on circadian/social rhythms could be applied to both independent and self-generated goal-relevant events. For example, the interpersonal inventory used in IPSRT is designed to identify and assess the major relationships in an individual's life. This logic could be extended to the development of a "goal-relevant inventory" in which the therapist and client could identify the meaning and value of relevant goals. Using the Social Rhythm Metric (Monk, et al., 1991), the therapist and client could then examine the extent to which these events are likely to induce social/circadian rhythm disruption and identify goal-pursuit strategies that would minimize such rhythm disruption.

Conclusion and Future Directions

Evidence suggests that psychosocial interventions have a positive prophylactic effect on bipolar disorder. However, a limited number of studies have been conducted and further work is needed. Accordingly, the objective of the current article was to examine whether recent findings in line with the BAS dysregulation theory can inform the three prominent psychosocial interventions for bipolar disorder – Cognitive-Behavioral Therapy, Psychoeducation, and Interpersonal and Social Rhythm Therapy. The research presented in the current article suggests that the biopsychosocial vulnerability for bipolar disorder may be particularly expressed in the achievement domain. We propose that clinicians and researchers may benefit from being particularly sensitive to the role of achievement relevant themes in the treatment and management of bipolar disorder. Accordingly, we propose the following recommendations: First, research indicates that the cognitive profiles of bipolar disorder are characterized by extreme goal striving tendencies and perfectionism in the achievement domain, rather than the sociotropic, dependent features often exhibited by unipolar depressed individuals. In line with Lam and colleagues (1999, 2003), clinicians working with bipolar individuals should be primed to address these perfectionistic tendencies in the achievement domain. Second, bipolar patients and their family members should be educated on the role that goal-relevant events play in precipitating bipolar episodes. In line with the stress-generation hypothesis, we argue that clinicians should address both independent and self-generated BAS relevant events. Third, the fact that bipolar disorder seems to be characterized by elevated drive, ambition, and achievement motivation suggests that negative expressed emotion in the achievement domain may be a particularly relevant stressor for individuals with bipolar disorder and should be addressed by FFT therapists. Fourth, IPSRT may benefit from being attuned to the possible disruptive effect that independent and self-generated goal-relevant events have on social/circadian rhythms.

The majority of research on psychosocial interventions for bipolar disorder has focused on adults. However, epidemiological findings (Weissman, et al., 1996) suggest that bipolar disorder first peaks in rates between ages 15–19. This transition from adolescence to young adulthood has been referred to as an "age of risk" (Weissman, et al., 1996) during which bipolar conditions "consolidate" and sometimes progress to a more severe condition. Accordingly, researchers and clinicians have begun the important task of developing age appropriate psychosocial interventions for children and adolescents with bipolar disorder (Fristad et al., 1998; 2003; Miklowitz et al., 2004; Post & Leverich, 2006). Adolescence is a period of significant transition in which individuals encounter a number of BAS activation (graduating from high-school, applying to college, dating) and BAS deactivation (academic failure, social

rejection) relevant events. Helping individuals with bipolar disorder navigate these events may help prevent a worsening of course.

References

- Abramson, LY.; Alloy, LB.; Hankin, BL.; Haeffel, GJ.; MacCoon, DG.; Gibb, BE. Cognitive vulnerability-stress models of depression in a self-regulatory and psychobiological context. In: Gotlib, IH.; Hammen, CL., editors. *Handbook of depression*. Guilford; New York: 2002. p. 368-294.
- Abramson LY, Metalsky GI, Alloy LB. Hopelessness depression: A theory-based subtype of depression. *Psychological Review* 1989;96:358–372.
- Akiskal, HS. The temperamental foundations of affective disorders. In: Mundt, C.; Goldstein, MJ.; Hahlweg, K.; Fielder, P., editors. *Interpersonal Factors in the Origin and Course of Affective Disorders*. Gaskell; London: 1996. p. 3-30.
- Alloy, LB.; Abramson, LY.; Whitehouse, WG.; Sylvia, LG.; Hafner, J.; Bender, RE.; Urosevic, S.; Nusslock, R.; Hogan, ME.; Harmon-Jones. Prospective prediction of hypomanic and depressive symptoms by Behavioral Approach System (BAS) activation-relevant and deactivation-relevant life events. 2009. Manuscript in preparation
- Alloy, LB.; Abramson, LY.; Whitehouse, WG.; Sylvia, LG.; Walshaw, PD.; Gerstein, RK.; Urosevic, S.; Nusslock, R.; Hogan, ME.; Harmon-Jones, E. Behavioral Approach System (BAS) and Behavioral Inhibition System (BIS) sensitivities, BAS-relevant life events, and mood symptoms: A prospective study. 2009. Manuscript in preparation
- Alloy, LB.; Abramson, LY.; Neeren, AM.; Walshaw, PD.; Urosevic, S.; Nusslock, R. Psychosocial risk factors for bipolar disorder: Current and early environment and cognitive styles. In: Jones, S.; Bentall, R., editors. *The psychology of bipolar disorder – New developments and research strategies*. Oxford University Press; Oxford, UK: 2006. p. 11-46.
- Alloy LB, Abramson LY, Walshaw PD, Cogswell A, Hughes M, Iacoviello B, Urosevic S, Nusslock R, Whitehouse WG, Hogan ME. Behavioral Approach System (BAS) and Behavioral Inhibition System (BIS) sensitivities and bipolar spectrum disorders: Prospective prediction of bipolar mood episodes. *Bipolar Disorders* 2008;10:310–322. [PubMed: 18271911]
- Alloy LB, Abramson LY, Walshaw PD, Gerstein RK, Keyser JD, Whitehouse WG, Urosevic S, Nusslock R, Hogan ME, Harmon-Jones E. Behavioral approach system (BAS) – relevant cognitive styles and bipolar spectrum disorders: Concurrent and prospective associations. *Journal of Abnormal Psychology* 2009;118:459–471. [PubMed: 19685944]
- Alloy LB, Abramson LY, Urosevic S, Bender RE, Wagner CA. Longitudinal predictors of bipolar spectrum disorders: A Behavioral Approach System (BAS) perspective. *Clinical Psychology: Science and Practice* 2009;16:206–226. [PubMed: 20161008]
- Alloy LB, Abramson LY, Urosevic S, Walshaw PD, Nusslock R, Neeren AM. The psychosocial context of bipolar disorder: Environmental, cognitive, and developmental risk factors. *Clinical Psychology Review* 2005;25:1043–1075. [PubMed: 16140445]
- Alloy LB, Reilly-Harrington NA, Fresco DM, Whitehouse WG, Zechmeister JS. Cognitive styles and life events in subsyndromal unipolar and bipolar mood disorders: Stability and prospective prediction of depressive and hypomanic mood swings. *Journal of Cognitive Psychotherapy: An International Quarterly* 1999;13:235–240.
- Ambelas A. Psychologically stressful events in the precipitation of manic episodes. *British Journal of Psychiatry* 1979;135:15–21. [PubMed: 497617]
- Ambelas A. Life events and mania: A special relationship? *British Journal of Psychiatry* 1987;150:235–240. [PubMed: 3651678]
- Ayuso-Mateos, JL. Global Burden of Bipolar Disorder in the Year 2000. 2006. Draft 21-06-06. http://www.who.int/healthinfo/statistics/bod_bipolar.pdf
- Basco, MR.; Rush, JA. *Cognitive-behavioral therapy for bipolar disorder*. The Guilford Press; New York, NY: 1996.
- Beck, AT. *Depression: Clinical, experimental, and theoretical aspects*. Harper & Row; New York: 1967.
- Benazzi F. Prevalence and clinical correlates of residual depressive symptoms in bipolar II disorder. *Psychotherapy and Psychosomatic* 2001;70:232–238.

- Carver CS, White TL. Behavioral inhibition, behavioral activation, and affective responses to impending reward and punishment: The BIS/BAS scales. *Journal of Personality and Social Psychology* 1994;67:319–333.
- Clarkin JF, Carpenter D, Hull J, et al. Effects of psychoeducational intervention for married patients with bipolar disorder and their spouses. *Psychiatric Services* 1998;49:531–533. [PubMed: 9550248]
- Cochran SD. Preventing medical non-compliance in the outpatient treatment of bipolar affective disorders. *Journal of Consulting and Clinical Psychology* 1984;52:873–878. [PubMed: 6501672]
- Colom F, Vieta E, Martinez-Aran A, Reinares M, Goikolea JM, Benabarre A, Torrent C, Comes M, Corbella B, Parramon G, Corominas J. A randomized trial on the efficacy of group psychoeducation in the prophylaxis of recurrences in bipolar patients whose disease is in remission. *Archives of General Psychiatry* 2003;60:402–407. [PubMed: 12695318]
- Craighead WE, Miklowitz DJ. Psychosocial interventions for bipolar disorder. *Journal of Clinical Psychiatry* 2000;61(Suppl 13):58–64. [PubMed: 11153813]
- Cutler NR, Post RM. Life course of illness in untreated manic-depressive patients. *Comprehensive Psychiatry* 1982;23:101–115. [PubMed: 7067425]
- Daley SE, Hammen C, Burge D, Davila J, Paley B, Lindberg N, Herzberg DS. Predictors of the generation of episodic stress: A longitudinal study of late adolescent women. *Journal of Abnormal Psychology* 1997;106:251–259. [PubMed: 9131845]
- Deeckersbach T, Gershuny BS, Otto MW. Cognitive-behavioral therapy for depression: applications and outcome. *Psychiatric Clinics of North America* 2000;23:795–809. [PubMed: 11147248]
- Depue RA, Collins PF. Neurobiology of the structure of personality: Dopamine, facilitation of incentive motivation, and extraversion. *Behavioral and Brain Sciences* 1999;22:491–569. [PubMed: 11301519]
- Depue RA, Iacono WG. Neurobehavioral aspects of affective disorders. *Annual Review of Psychology* 1989;40:457–492.
- Depue, RA.; Krauss, SP.; Spoont, MR. A two-dimensional threshold model of seasonal bipolar affective disorder. In: Maggnusson, D.; Ohman, A., editors. *Psychopathology: An interactional perspective*. Academic Press; New York: 1987. p. 95-123.
- Depue RA, Slater JF, Wolfstetter-Kausch H, Klein D, Goplerud E, Farr D. A behavioral paradigm for identifying persons at risk for bipolar depressive disorder: A conceptual framework and five validation studies. *Journal of Abnormal Psychology* 1981;90:381–437. [PubMed: 7298991]
- Ehlers CL, Frank E, Kupfer DJ. Social zeitgebers and biological rhythms: A unified approach to understanding the etiology of depression. *Archives of General Psychiatry* 1988;45:948–952. [PubMed: 3048226]
- Ehlers CL, Kupfer DJ, Frank E, Monk TH. Biological rhythms and depression: The role of zeitgebers and zeitstorsers. *Depression* 1993;1:285–293.
- Elicott A, Hammen C, Gitlin M, Brown G, Jamison K. Life Events and the course of bipolar disorder. *American Journal of Psychiatry* 1990;147:1194–1198. [PubMed: 1974746]
- Evans M, Hollon S, DeRubeis R, Piasecki J, Grove W, Garvey M, Tuason V. Differential relapse following cognitive therapy and pharmacotherapy for depression. *Archives of General Psychiatry* 1992;49:802–808. [PubMed: 1417433]
- Falloon IRH, Boyd JL, McGill CW, Williamson M, Razani J, Moss HB, et al. Family management in the prevention of morbidity of schizophrenia. *Archives of General Psychiatry* 1985;42:887–896. [PubMed: 2864032]
- Fava GA, Bartolucci G, Rafanelli C, Mangelli L. Cognitive-behavioral management of patients with bipolar disorder who relapsed while on lithium prophylaxis. *Journal of Clinical Psychiatry* 2001;62:556–559. [PubMed: 11488368]
- Fava GA, Grandi S, Zielezny M, Rafanelli C, et al. Four year outcome for cognitive behavioral treatment of residual symptoms in major depression. *American Journal of Psychiatry* 1996;153:945–947. [PubMed: 8659620]
- Fowles DC. Psychophysiology and psychopathology: A motivational approach. *Psychophysiology* 1988;25:373–391. [PubMed: 3051073]

- Francis-Raniere E, Alloy LB, Abramson LY. Depressive personality styles and bipolar spectrum disorders: Prospective tests of the event congruency hypothesis. *Bipolar Disorders* 2006;8:382–399. [PubMed: 16879139]
- Frank E. Interpersonal and social rhythm therapy prevents depressive symptomatology in bipolar I patients [abstract]. *Bipolar Disorders* 1999;1(suppl 1)
- Frank E. Interpersonal and social rhythm therapy: A means of improving depression and preventing relapse in bipolar disorder. *Journal of Clinical Psychology* 2007;63:463–473. [PubMed: 17417811]
- Frank E, Hlastala S, Ritenour A, Houck P, Tu XM, Monk TH, Mallinger AG, Kupfer DJ. Inducing lifestyle regularity in recovering bipolar disorder patients: Results from the maintenance therapies of bipolar disorder protocol. *Biological Psychiatry* 1997;41:1165–1173. [PubMed: 9171907]
- Frank E, Kupfer DJ, Thase ME, Mallinger AG, Swartz HA, Fagioli AM, Grochocinski V, Houck P, Scott J, Thompson W, Monk T. Two-year outcomes for interpersonal and social rhythm therapy in individuals with bipolar I disorder. *Archives of General Psychiatry* 2005;62:996–1004. [PubMed: 16143731]
- Frank E, Swartz HA, Kupfer DJ. Interpersonal and social rhythm therapy: Managing the chaos of bipolar disorder. *Biological Psychiatry* 2000;48:593–604. [PubMed: 11018230]
- Frank E, Swartz HA, Mallinger AG, Thase ME, Weaver EV, Kupfer DJ. Adjunctive psychotherapy for bipolar disorder: Effects of changing treatment modality. *Journal of Abnormal Psychology* 1999;108:579–587. [PubMed: 10609422]
- Fristad MA, Arnett MM, Gavazzi SM. The impact of psychoeducational workshops on families of mood-disordered children. *Family Therapy* 1998;25:151–159.
- Fristad MA, Gavazzi SM, Soldano KW. Multi-family psychoeducation groups for childhood mood disorders: A program description and preliminary efficacy data. *Contemporary Family Therapy: An International Journal* 1998;20:385–402.
- Fristad MA, Goldberg-Arnold JS, Gavazzi SM. Multifamily psychoeducation groups (MFPG) for families of children with bipolar disorder. *Bipolar Disorders* 2002;2002:254–262. [PubMed: 12190715]
- Fristad MA, Goldberg-Arnold JS, Gavazzi SM. Multi-family psychoeducation groups in the treatment of children with mood disorders. *Journal of Marital & Family Therapy* 2003;29:491–504. [PubMed: 14593691]
- Geller B, Craney JL, Bolhofner K, Nickelsburg MJ, Williams M, Zimmerman B. Two-year prospective follow-up of children with a prepubertal and early adolescent bipolar disorder phenotype. *American Journal of Psychiatry* 2002;159:927–933. [PubMed: 12042179]
- Gitlin MJ, Swendsen J, Heller TL, Hammen C. Relapse and impairment in bipolar disorder. *American Journal of Psychiatry* 1995;152:1635–1640. [PubMed: 7485627]
- Goodwin, FK.; Jamison, KR. Manic depressive illness. Oxford University Press; New York: 1990.
- Gray, JA. A critique of Eysenck's theory of personality. In: Eysenck, HJ., editor. *A model for personality*. Springer-Verlag; Berlin: 1981. p. 246-276.
- Gray, JA. Three fundamental emotion systems. In: Eckman, P.; Davidson, RJ., editors. *The nature of emotion: Fundamental questions*. Oxford University Press; New York: 1994. p. 243-247.
- Greenberger, D.; Padesky, CA. *Mind over mood: A cognitive therapy treatment manual for clients*. Guilford; New York: 1995.
- Hammen C. Generation of stress in the course of unipolar depression. *Journal of Abnormal Psychology* 1991;100:555–561. [PubMed: 1757669]
- Hammen C, Ellicott A, Gitlin M. Stressors and sociotropy/autonomy: A longitudinal study of their relationship to the course of bipolar disorder. *Cognitive Therapy and Research* 1992;16:409–418.
- Harmon-Jones E. On the relationship of anterior brain activity and anger: Examining the role of attitude toward anger. *Cognition and Emotion* 2004;18:337–361.
- Harmon-Jones E, Abramson LY, Nusslock R, Sigelman JD, Urosevic S, Turonie L, Alloy LB, Fearn M. Effect of bipolar disorder on left frontal cortical responses to goals differing in valence and task difficulty. *Biological Psychiatry* 2008;63:693–698. [PubMed: 17919457]
- Harmon-Jones E, Abramson LY, Sigelman J, Bohlig A, Hogan ME, Harmon-Jones C. Proneness to hypomania/mania symptoms or depression symptoms and asymmetrical frontal cortical responses to

- an anger-evoking event. *Journal of Personality and Social Psychology* 2002;82:610–618. [PubMed: 11999927]
- Harmon-Jones E, Allen JJB. Behavioral activation sensitivity and resting frontal EEG asymmetry: Covariation of putative indicators related to risk for mood disorders. *Journal of Abnormal Psychology* 1997;106:159–163. [PubMed: 9103728]
- Hogarty GE, Anderson CM, Reiss DJ, Kornblith SJ, Greenwald DP, Javna CD, et al. Family psychoeducation, social skills training, and maintenance chemotherapy in the aftercare treatment of schizophrenia: One-year effects of a controlled study on relapse and expressed emotion. *Archives of General Psychiatry* 1986;43:633–642. [PubMed: 2872870]
- Hollon, SD. Cognitive therapy in the treatment and prevention of depression. In: Joiner, TE.; Brown, JS.; Kistner, J., editors. *The interpersonal, cognitive, and social nature of depression*. Lawrence Erlbaum Associates Publishers; Mahwah, NJ: 2006. p. 131-151.
- Honig A, Hofman A, Rozendaal N, Dingemans P. Psycho-education in bipolar disorder: Effect on expressed emotion. *Psychiatry Research* 1997;72:17–22. [PubMed: 9355815]
- Johnson SL. Mania and dysregulation in goal pursuit: A review. *Clinical Psychology Review* 2005;25:241–262. [PubMed: 15642648]
- Johnson SL, Carver CS. Extreme goal-setting and vulnerability to mania among undiagnosed young adults. *Cognitive Therapy and Research* 2006;30:377–395.
- Johnson, SL.; Kizer, A. Bipolar and unipolar depression: A comparison of clinical phenomenology and psychosocial predictors. In: Gotlib, IH.; Hammen, CL., editors. *Handbook of Depression*. Guilford Press; New York, NY, US: 2002. p. 141-165.
- Johnson SL, Roberts JE. Life events and bipolar disorder: Implications from biological theories. *Psychological Bulletin* 1995;117:434–449. [PubMed: 7777648]
- Johnson SL, Ruggero C, Carver CS. Cognitive, behavioral and affective responses to reward: Links with hypomanic symptoms. *Journal of Social and Clinical Psychology* 2005;24:894–906.
- Johnson SL, Sandrow D, Meyer B, Winters R, Miller I, Solomon D, Keitner G. Increases in manic symptoms after life events involving goal attainment. *Journal of Abnormal Psychology* 2000;109:721–727. [PubMed: 11195996]
- Johnson SL, Winett CA, Meyer B, Greenhouse WJ, Miller I. Social support and the course of bipolar disorder. *Journal of Abnormal Psychology* 1999;108:558–566. [PubMed: 10609420]
- Keck PE, McElroy SL, Strakowski SM, West SA, Sax KW, Hawkins JM, Bourne ML, Haggard P. 12-Month outcome of patients with bipolar disorder following hospitalization for a manic or mixed episode. *American Journal of Psychiatry* 1998;155:646–652. [PubMed: 9585716]
- Keitner GI, Solomon DA, Ryan CE, Miller IW, Mallinger A, Kupfer DJ, Frank E. Prodromal and residual symptoms in bipolar I disorder. *Comprehensive Psychiatry* 1996;37:363–367.
- Klerman, GL.; Weissman, MM.; Rounsaville, BJ.; Chevron, ES. *Interpersonal Psychotherapy of Depression*. Basic Books; New York: 1984.
- Kutcher S, Robertson HA, Bird D. Premorbid functioning in adolescent onset bipolar I disorder. A preliminary report from an ongoing study. *Journal of Affective Disorders* 1998;51:137–144. [PubMed: 10743846]
- Lagace DC, Kutcher SP, Robertson HA. Mathematics deficits in adolescents with bipolar I disorder. *American Journal of Psychiatry* 2003;160:100–104. [PubMed: 12505807]
- Lam DH, Bright J, Jones S, Hayward P, Schuck N, Chisholm D, Sham P. Cognitive therapy for bipolar illness: A pilot study of relapse prevention. *Cognitive Therapy and Research* 2000;24:503–520.
- Lam, DH.; Jones, SH.; Hayward, P.; Bright, JA. *Cognitive therapy for bipolar disorder: A therapist's guide to concepts, methods, and practice*. John Wiley & Sons; New York, NY: 1999.
- Lam DH, Watkins ER, Hayward P, Bright J, Wright K, Kerr N, Parr-Davis G, Sham P. A randomized controlled study of cognitive therapy for relapse prevention for bipolar affective disorder: Outcome of the First Year. *Archives of General Psychiatry* 2003;60:145–152. [PubMed: 12578431]
- Lam D, Wong G. Prodromes, coping strategies, insight, and social functioning in bipolar affective disorders. *Psychological Medicine* 1997;27:1091–1100. [PubMed: 9300514]
- Lam D, Wong G, Sham P. Prodromes, coping strategies and course of illness in bipolar affective disorder – a naturalistic study. *Psychological Medicine* 2001;31:1387–1402.

- Lam D, Wright K, Smith N. Dysfunctional assumptions in bipolar disorder. *Journal of Affective Disorders* 2004;79:193–199. [PubMed: 15023494]
- Lembke A, Ketter TA. Impaired recognition of facial emotion in mania. *American Journal of Psychiatry* 2002;159:302–304. [PubMed: 11823275]
- Leibenluft E, Moul DE, Schwartz PJ, Madden PA, Wehr TA. A clinical trial of sleep deprivation in combination with antidepressant medication. *Psychiatry Research* 1993;46:213–227. [PubMed: 8493292]
- Leibenluft E, Wehr TA. Is sleep deprivation useful in the treatment of depression? *American Journal of Psychiatry* 1992;149:159–168. [PubMed: 1734735]
- Lozano BE, Johnson SL. Can personality traits predict increases in manic and depressive symptoms? *Journal of Affective Disorders* 2001;63:103–111. [PubMed: 11246086]
- Malkoff-Schwartz S, Frank E, Anderson BP, Hlastala SA, Luther JF, Sherrill JT, Houck PR, Kupfer DJ. Social rhythm disruption and stressful life events in the onset of bipolar and unipolar episodes. *Psychological Medicine* 2000;30:1005–1010. [PubMed: 12027038]
- Malkoff-Schwartz S, Frank E, Anderson B, Sherrill JT, Siegel L, Patterson D, Kupfer DJ. Stressful life events and social rhythm disruption in the onset of manic and depressive bipolar episodes. *Archives of General Psychiatry* 1998;55:702–707. [PubMed: 9707380]
- Merikangas KR, Akiskal HS, Angst J, Greenberg PE, Hirschfeld RM, Petukhova M, Kessler RC. Lifetime and 12-month prevalence of bipolar spectrum disorder in the National Comorbidity Survey replication. *Archives of General Psychiatry* 2007;64:543–552. [PubMed: 17485606]
- Meyer B, Beevers CG, Johnson SL. Goal appraisals and vulnerability to bipolar disorder: A personal projects analysis. *Cognitive Therapy and Research* 2004;28:173–182.
- Meyer B, Johnson SL, Carver CS. Exploring behavioral activation and inhibition sensitivities among college students at risk for bipolar spectrum symptomatology. *Journal of Psychopathology and Behavioral Assessment* 1999;21:275–292.
- Meyer B, Johnson SL, Winters R. Responsiveness to threat and incentive in bipolar disorder: Relations of the BIS/BAS scales with symptoms. *Journal of Psychopathology and Behavioral Assessment* 2001;23:133–143.
- Miklowitz DJ, George EL, Axelson DA, Kim EY, Birmaher B, Schneck C, Beresford C, Craighead WE, Brent DA. Family-focused treatment for adolescents with bipolar disorder. *Journal of Affective Disorders* 2004;82(Suppl 1):S113–128. [PubMed: 15571785]
- Miklowitz DJ, George EL, Richards JA, Simoneau TL, Suddath RL. A randomized study of family-focused psychoeducation and pharmacotherapy in the outpatient management of bipolar disorder. *Archives of General Psychiatry* 2003;60:904–911. [PubMed: 12963672]
- Miklowitz DJ, Goldstein MJ. Behavioral family treatment for patients with bipolar affective disorder. *Behavioral Modification* 1990;14:457–489.
- Miklowitz, DJ.; Goldstein, MJ. *Bipolar Disorder: A Family-Focused Treatment Approach*. Guilford; New York: 1997.
- Miklowitz DJ, Goldstein MJ, Nuechterlein KH, Snyder KS, Mintz J. Family Factors and the course of bipolar affective disorder. *Archives of General Psychiatry* 1988;45:225–231. [PubMed: 3341878]
- Miklowitz DJ, Otto MW, Frank E, Reilly-Harrington NA, Kogan JN, Sachs GS, Thase ME, Calabrese JR, Marangell LB, Ostacher MJ. Intensive psychosocial intervention enhances functioning in patients with bipolar depression: Results from a 9-month randomized controlled trial. *American Journal of Psychiatry* 2007;164:1340–1347. [PubMed: 17728418]
- Miklowitz DJ, Otto MW, Frank E, Reilly-Harrington N, Wisniewski SR, Kogan JN, Nierenberg AA, Calabrese JR, Marangell LB, Gyulai L, Araga M, Gonzalez JM, Shirley ER, Thase ME, Sachs GS. Psychosocial treatments for bipolar depression: A 1-year randomized trial from the systematic treatment enhancement program. *Archives of General Psychiatry* 2007;64:419–427. [PubMed: 17404119]
- Miklowitz DJ, Richards JA, George EL, Frank E, Suddath RL, Powell KB, Sacher JA. Integrated family therapy and individual therapy for bipolar disorder: Results of a treatment development study. *Journal of Clinical Psychiatry* 2003;64:182–191. [PubMed: 12633127]

- Miklowitz DJ, Simoneau TL, George EL, Richards JA, Kalbag A, Sachs-Ericsson N, Suddath R. Family-focused treatment of bipolar disorder: 1-year effects of a psychoeducational program in conjunction with pharmacotherapy. *Biological Psychiatry* 2000;48:582–592. [PubMed: 11018229]
- Molnar GJ, Feeney MG, Fava GA. Duration and symptoms of bipolar prodromes. *American Journal of Psychiatry* 1988;145:1576–1578. [PubMed: 3195679]
- Monk TH, Kupfer DJ, Frank E, Ritenour AM. The Social Rhythm Metric (SRM): Measuring daily social rhythms over 12 weeks. *Psychiatry Research* 1991;36:195–207. [PubMed: 2017534]
- Morris CD, Miklowitz DJ, Waxmonsky JA. Family-focused treatment for bipolar disorder in adults and youth. *Journal of Clinical Psychology: In Session* 2007;63:433–445.
- Murray, CJL.; Lopez, AD. The global burden of disease: A comprehensive assessment of mortality and disability from diseases, injuries, and risk factors in 1990 and projected to 2020. Harvard University Press; Boston: 1996.
- Nusslock R, Abramson LY, Harmon-Jones E, Alloy LB, Hogan ME. A goal-striving life event and the onset of bipolar episodes: Perspective from the Behavioral Approach System (BAS) dysregulation theory. *Journal of Abnormal Psychology* 2007;116:105–115. [PubMed: 17324021]
- Nusslock R, Alloy LB, Abramson LY, Harmon-Jones E, Hogan ME. Impairment in the achievement domain in bipolar spectrum disorders: Role of behavioral approach system hypersensitivity and impulsivity. *Minerva Pediatrica* 2008;59:41–50. [PubMed: 18277364]
- Otto, MW.; Reilly-Harrington, N.; Kogan, JN.; Henin, A.; Knausz, RO. Cognitive-Behavior Therapy for Bipolar Disorder: Treatment Manual. Massachusetts General Hospital; Boston, M.A.: 1999. Unpublished manual
- Otto MW, Reilly-Harrington N, Sachs GS. Psychoeducational and cognitive-behavioral strategies in the management of bipolar disorder. *Journal of Affective Disorders* 2003;73:171–181. [PubMed: 12507750]
- Patel NC, Delbello MP, Bryan HS, Adler CM, Kowatch RA, Stanford K, Strakowski SM. Open-label lithium for the treatment of adolescents with bipolar depression. *Journal of the American Academy of Child & Adolescent Psychiatry* 2006;45:289–297. [PubMed: 16540813]
- Patelis-Siotis I, Young LT, Robb JC, Marriott M, Bieling PJ, Cox LC, Joffe RT. Group cognitive behavioral therapy for bipolar disorder: A feasibility and effectiveness study. *Journal of Affective Disorders* 2001;65:145–153. [PubMed: 11356238]
- Perris H. Life events and depression: Part 2. Results in diagnostic subgroups, and in relation to the recurrence of depression. *Journal of Affective Disorders* 1984;7:25–36. [PubMed: 6236241]
- Peven DE, Shulman BH. The psychodynamics of bipolar affective disorder: Some empirical findings and their implications for cognitive theory. *Individual Psychology: Journal of Adlerian Theory, Research, and Practice* 1983;39:2–16.
- Post RM, Leverich GS. The role of psychosocial stress in the onset and progression of bipolar disorder and its comorbidities: The need for earlier and alternative modes of therapeutic intervention. *Development and Psychopathology* 2006;18:1181–1211. [PubMed: 17064434]
- Prien RF, Rush AJ. National Institute of Mental Health workshop report on the treatment of bipolar disorder. *Biological Psychiatry* 1996;40:215–220. [PubMed: 8830955]
- Rea MM, Tompson MC, Miklowitz DJ, Goldstein MJ, Hwang S, Mintz J. Family-focused treatment versus individualized treatment for bipolar disorder: Results of a randomized clinical trial. *Journal of Consulting and Clinical Psychology* 2003;71:482–492. [PubMed: 12795572]
- Reilly-Harrington NA, Alloy LB, Fresco DM, Whitehouse WG. Cognitive styles and life events interact to predict bipolar and unipolar symptomatology. *Journal of Abnormal Psychology* 1999;108:567–578. [PubMed: 10609421]
- Rucci P, Frank E, Kostelnik B, Fagioli A, Mallinger A, Swartz HA, Thase ME, Siegel L, Wilson D, Kupfer DJ. Suicide attempts in patients with bipolar I disorder during acute and maintenance phases of intensive treatment with pharmacotherapy and adjunctive psychotherapy. *American Journal of Psychiatry* 2002;159:1160–1164. [PubMed: 12091194]
- Scott J, Paykel E, Morriss R, Bentall R, Kinderman P, Johnson T, Abbott R, Hayhurst H. Cognitive-behavioural therapy for severe and recurrent bipolar disorders: Randomized controlled trial. *British Journal of Psychiatry* 2006;188:313–320. [PubMed: 16582056]

- Scott J, Pope M. Cognitive styles in individuals with bipolar disorders. *Psychological Medicine* 2003;33:1081–1088. [PubMed: 12946092]
- Scott J, Colom F, Vieta E. A meta-analysis of relapse rates with adjunctive psychological therapies compared to usual psychiatric treatment for bipolar disorders. *International Journal of Neuropsychopharmacology* 2007;10:123–129. [PubMed: 16787554]
- Scott J, Garland A, Moorhead S. A pilot study of cognitive therapy in bipolar disorders. *Psychological Medicine* 2001;31:459–467. [PubMed: 11305854]
- Scott J, Stanton B, Garland A, Ferrier IN. Cognitive vulnerability in patients with bipolar disorder. *Psychological Medicine* 2000;30:467–472. [PubMed: 10824667]
- Simoneau TL, Miklowitz DJ, Richards JA, Saleem R, George EL. Bipolar disorder and family communication: Effects of a psychoeducation treatment program. *Journal of Abnormal Psychology* 1999;108:588–597. [PubMed: 10609423]
- Simoneau TL, Miklowitz DJ, Saleem R. Expressed emotion and interaccitonal patterns in the families of bipolar patients. *Journal of Abnormal Psychology* 1998;107:497–507. [PubMed: 9715584]
- Smith JA, Tarrier N. Prodromal symptoms in manic depressive psychosis. *Social Psychiatry and Psychiatric Epidemiology* 1992;27:245–248. [PubMed: 1439998]
- Spielberger CD, Parker JB, Becker J. Conformity and achievement in remitted manic-depressive patients. *Journal of Nervous and Mental Disorders* 1963;137:162–172.
- Sutton SK, Davidson RJ. Prefrontal brain asymmetry: A biological substrate if the behavioral approach and inhibition systems. *Psychological Science* 1997;8:204–210.
- Sylvia LG, Alloy LB, Hafner JA, Gauger MC, Verdon K, Abramson LY. Life events and social rhythms in bipolar spectrum disorders: A prospective study. *Behavior Therapy*. (in press).
- Tsuchiya KJ, Agerbo E, Byrne M, Mortensen PB. Higher socio-economic status of parents may increase risk for bipolar disorder in offspring. *Psychological Medicine* 2004;34:787–793. [PubMed: 15500299]
- Urosevic, S.; Abramson, LY.; Alloy, LB.; Bender, RE.; Hogan, ME.; Nusslock, R. Bipolar disorders are related to increased Behavioral Approach System (BAS) activating and deactivating events. 2009. Manuscript in preparation
- Urosevic S, Abramson LY, Harmon-Jones E, Alloy LB. Dysregulation of the Behavioral Approach System (BAS) in bipolar spectrum disorders: Review of theory and evidence. *Clinical Psychology Review* 2008;28:1188–1205. [PubMed: 18565633]
- Wampold, B. *The Great Psychotherapy Debate: Models, Methods, and Findings*. Lawrence Erlbaum Associates; New Jersey: 2001.
- Wehr TA, Sack DA, Rosenthal NE. Sleep reduction as a final common pathway in the genesis of mania. *American Journal of Psychiatry* 1987;144:201–204. [PubMed: 3812788]
- Weissman MM, Bland RC, Canino GJ, Faravelli C, Greenwald S, Hwu HG, et al. Cross-national epidemiology of major depression and bipolar disorder. *Journal of the American Medical Association* 1996;276:293–299. [PubMed: 8656541]
- Zaretsky AE, Segal ZV, Gemar M. Cognitive therapy of bipolar depression: A pilot study. *Canadian Journal of Psychiatry* 1999;44:491–49.