Published in final edited form as:

J Am Chem Soc. 2009 October 21; 131(41): 15061–15074. doi:10.1021/ja906056v.

Total Synthesis and Stereochemical Assignment of (–)-Ushikulide A

Barry M. Trost^{*}, Brendan M. O’Boyle, and Daniel Hund

Department of Chemistry, Stanford University, Stanford, California, 94305-5080

Abstract

We report the determination of the full stereostructure of (–)-ushikulide A (**1**), a spiroketal containing macrolide by total synthesis. Ushikulide A (**1**) was isolated from a culture broth of *Streptomyces* sp. IUK-102 and exhibits potent immunosuppressant activity ($IC_{50} = 70$ nM). To embark upon an ushikulide A synthesis, a tentative assignment was made based on analogy to cytovaricin (**2**), a related macrolide isolated from a culture of *Streptomyces diastatochromogenes* whose full structure was previously established *via* synthesis and X-ray crystallography. This report delineates studies on several key steps, namely a direct aldol reaction catalyzed by the dinuclear zinc ProPhenol complex, a metal catalyzed spiroketalization, as well as application of an unprecedented asymmetric alkynylation of a simple saturated aldehyde with methyl propiolate to prepare the nucleophilic partner for a Marshall-Tamaru propargylation. These studies culminated in the first total synthesis and stereochemical assignment of (–)-ushikulide A and significantly extended the scope of the above-mentioned methodologies.

Keywords

Ushikulide A; cytovaricin; spiroketal; macrolactone; dinuclear zinc aldol; Suzuki cross-coupling; Marshall/Tamaru propargylation

Introduction

In spite of the tremendous advances in modern spectroscopic methods, organic synthesis continues to play a pivotal role in the elucidating the full structure of complex natural products.¹ This method has the advantage that even in the absence of a firm structural assignment, a combination of logic and spectroscopic comparison can arrive at the correct structure. Previous work in our group led to the assignment of amphidinolide A after a total of eleven diastereomers were prepared synthetically.² Thus we were encouraged to consider the synthesis of other natural products of uncertain stereochemistry.

Ushikulide A (**1**), a newly isolated³ and stereochemically undefined member of the oligomycinrutamycin family was chosen out of many possible synthetic targets based on several considerations. Due to the stereochemical complexity of such natural products randomly preparing diastereomers would lead to an exceedingly tedious task (ushikulide A contains 14 stereogenic centers - leading to $2^{14} = 16384$ stereoisomers). Therefore, a reasonable amount of likely stereochemical information was required before synthetic studies could be initiated. Ushikulide A (**1**) has remarkable resemblance to cytovaricin (**2**), a natural product

bmtrost@stanford.edu.

Supporting Information Available: Detailed experimental procedures, full characterization of all products, and comparison NMR spectra. This material is available free of charge on the Internet at <http://pubs.acs.org>.

isolated from the same species of bacteria, whose full structure was established through X-ray crystallography⁴ and confirmed *via* synthesis.⁵ Furthermore, careful comparison of the NMR spectra of these structures revealed close similarity between the spiroketal regions of **1** and **2** (Figure 2). While this was clearly insufficient evidence to firmly assign the structure of ushikulide, it appeared at the outset that eight of the stereocenters (carbons 16, 19, 20, 21, 23, 26, 27 and 29) were either clearly analogous to cytovaricin or their configuration should be apparent after the first synthetic diastereomer was prepared. Specifically, stereochemistry at C₁₆ could be lost through base catalyzed isomerization of the β,γ -unsaturated ketone yielding ushikulide B (which contains an *E* olefin between C₁₅ and C₁₆), while the stereochemistry at C₂₉ (which is isolated from the macrolactone) could be assigned so long as the spiroketal portion of the synthetic material had the correct configuration. Only six centers remained, a task roughly comparable to our successful synthesis of amphidinolide A.² In the absence of significant quantities of a natural product for degradation studies, synthesis is the only method to reliably assign structure in the event that X-ray crystallography proves untenable, as was the case for **1**.⁶ We hope that this full account⁷ will provide a deeper understanding of the chemistry employed in this study while also outlining the power of synthesis for the determination of structure.

Synthetic Planning

A flexible approach that could be amended to prepare any diastereomer of **1** was envisioned (Scheme 1). As stereochemistry of the spiroketal was pivotal, it seemed well advised to split the spiroketal (**3**) away from the aliphatic portion (**4**) *via* Suzuki coupling and esterification. While sp^2 - sp^2 Suzuki coupling is a well established disconnection for diene containing members of this family,⁸ our retrosynthesis relied on a less common sp^2 - sp^3 Suzuki reaction⁹ to access the C₁₄-C₁₅ olefin of ushikulide A. For spiroketal fragment **3**, a Marshall/Tamaru propargylation would provide the homopropargylic alcohol that could be elaborated to the C₁₅-C₁₄ olefin. The metal catalyzed spiroketalization, pioneered by Utimoto¹⁰ represents a highly atom-economical and redox-neutral¹¹ approach to the desired spiroketal ring system. Utilization of an alkyne as an orthogonally reactive surrogate¹² for the desired spiroketal is strategically advantageous as it enabled two carbon-carbon bond forming reactions without requiring the use of additional protecting group manipulations or activating groups, which certainly would have been required if more conventional carbonyl chemistry had been employed. Either **5** or **6** could potentially lead to the desired spiroketal however, only the former case appeared likely to give the desired regioselectivity (See **6** to **7**; Scheme 1). Application of our dinuclear zinc aldol methodology¹³ was expected to simplify the aliphatic fragment into ketone **9** and aldehyde **10**. Due to the similar steric bulk of the methyl and allyl groups at C₁₀ conventional aldol methodologies would not be expected to give acceptable diastereoselectivity for this reaction. However, we envisioned that through catalyst control this highly convergent strategy would become tenable.

Synthesis of the Spiroketal Fragment

Our studies began with the preparation of several aldehydes to fulfill the role of synthon **8** (**16ad**, Scheme 2). To accomplish this, a Crimmins¹⁴ aldol reaction between thiazolidinethione **14** and three known aldehydes (**13a-c**)¹⁵ was employed. Reaction of the (*Z*)-titanium enolate derived from **14** with aldehydes **13a-c** resulted in essentially complete diastereoselectivity and quantitative yield for each case. Reduction of aldol adduct **15a** to the β -hydroxy aldehyde was complicated by facile epimerization of the α -stereogenic center on silica gel (not shown). However, after protection of the secondary alcohol, mono-reduction proceeded uneventfully to yield the desired aldehydes (**16a-d**) in good yield over three steps.

Preparation of an alkyne fragment to fulfill the role of synthon **11** began with a known Noyori reduction of methyl-3-oxopentanoate (Scheme 3).¹⁶ Formation of the *p*-methoxybenzyl ether and mono-reduction with diisobutylaluminum hydride gave aldehyde **17** in good yield. Addition of the lithium anion of TMS acetylene in the presence of a chelating Lewis acid (Et₂AlCl) gave a mixture of propargyl alcohols **18a** and **18b** in low diastereoselectivity and yield (~2:1 at C₂₇; Scheme 3). Separation of these epimers on a preparative scale was prohibitively difficult, particularly at this early stage in the synthesis. As an alternative, a number of crotylations were surveyed, starting with the conditions of Keck,¹⁷ which afforded a mixture of predominantly two diastereomers (2:1 - **19a** : **19b**) that were also inseparable by flash chromatography on silica gel. Varying the Lewis acid (SnCl₄, CH₂Cl₂, Entry 2) led to only a negligible increase in diastereoselectivity. Fortunately, ample precedent suggested that this problem could be overcome by reagent-control.¹⁸ In our case, using either antipode of Brown's¹⁹ (Z)-crotyl-B(Ipc)₂ reagent led to excellent control of stereochemistry for the preparation of **19a** or **19b** in >20:1 d.r. by ¹H NMR analysis for both cases. Preparation of alkyne **20** was completed in three additional steps (silylation, hydroboration-iodination and nucleophilic substitution). Because no chromatographic separation of diastereomers was required, this route proved to be highly scalable and more than 10 grams of **20** were ultimately prepared.

The stereochemistry of **19a** was also confirmed *via* formation of a cyclic *para*-methoxybenzyl acetal (Equation 1). Exposure of **19a** to DDQ in the presence of molecular sieves²⁰ furnished acetal **21**, which could be cleanly separated from an initial mixture of diastereomers in 60% isolated yield. NOE irradiation of the benzylic acetal proton showed strong enhancements at C₂₆ and C₂₉, consistent with the indicated chair form and demonstrating the 1,3 diol to be of an *anti* stereochemical relationship.

With the two fragments in hand, our attention turned to nucleophilic addition of alkyne **20** to the aldehyde fragment **16c** (Equation 2). The addition proceeded with low diastereoselectivity; however, separation of the C₂₁ epimers was readily accomplished by silica gel chromatography. Desilylation with fluoride (HF•pyridine or *n*-Bu₄NF) proceeded only very slowly at one of the two silyl groups, but exposure of either **22a** or **22b** to hydrochloric acid yielded either epimeric diol in good yield (**23a** or **23b**).

(2)

The stereochemistry of each triol was assigned by the method of Rychnovsky (Equation 3).²¹ Protection of 1,3-diols **23a** and **23b** was accomplished with 2-methoxypropene and PPTS, yielding acetonides **24** and **25**. gHSQC analysis of either acetonide confirmed the relative stereochemistry with *syn* acetonide **24** displaying large differences in the acetonide ¹³C NMR shifts, consistent with a chair conformation, while *anti* acetonide **25** displayed nearly equivalent ¹³C NMR shifts consistent with a half-chair conformation.

(3)

Attempts were also made to improve the diastereoselectivity of the alkyne addition reaction (Scheme 4). Moderate Felkin-Anh selectivity could be obtained in the presence of lithium bromide and molecular sieves (6:1 d.r., Entry 2). However, obtaining the chelation controlled product (**22b**) proved not to be feasible under the reaction conditions examined. Switching the C₁₉ *tert*-butyldimethylsilyl group to a methoxymethyl ether (**16d**, Entry 4) gave some improvement of the diastereoselectivity; unfortunately the resulting epimers (**27a** and **27b**) were very recalcitrant to chromatographic separation causing this route to be impractical.

The low selectivity of the alkynylation led us to consider two possible alternatives. The first would be addition of the alkyne to a Weinreb amide, followed by diastereoselective Noyori reduction,²² while the second would be inversion of the *syn* 1,3-diol to converge both epimers to the *anti* stereochemistry. Because of the ease of chromatographic separation, we ultimately elected to pursue the latter course (Scheme 5). Alternatives to the 1,3-dithiane protecting group were also examined as the sulfur atoms it contained might cause catalyst poisoning during the key spiroketalization reaction. Three aldehydes (**16a-c**) were subjected to alkyne addition with the lithium acetylene derived from **20**. After separation of the resulting epimers the *anti* product was esterified with retention (BzCl, pyridine), while the *syn* product was esterified under conditions leading to inversion (PPh₃, DEAD, BzOH, toluene). All three pairs of epimers converged, yielding propargyl benzoates **28a-c**. The TBDPS group (**28a**) proved too acid-labile and yielded predominantly triol **29a** when subjected to the deprotection conditions developed previously (HCl, H₂O, THF). Fortunately, both **28b** and **28c** could be cleanly desilylated yielding diols **29b** and **29c** respectively.

With the triol **23b** in hand, we examined application of a metal catalyzed spiroketalization for the preparation of spiroketal **26**. Coordination of a carbophilic Lewis acid (Pd^{II}) to the alkyne present in the starting material should lead to oxypalladation, generating an enol-ether and ultimately producing the desired spiroketal. Unfortunately, Utimoto's conditions (Pd [CH₃CN]₂Cl₂, THF, CH₃CN, reflux)¹⁰ failed entirely for the preparation of **26** (Equation 4). When acetone was substituted as solvent with the same catalyst, a mixture of acetonide **25**

along with an unidentified product was obtained. All other Pd^{II} catalyzed conditions that were examined afforded only decomposition after prolonged heating.

(4)

Further investigation of the spiroketalization reaction focused on triol **30b** and propargyl benzoate **29b** as precursors to spiroketal **31a** (Scheme 6). Palladium catalysis again provided only limited reactivity (Entry 1), prompting us to examine platinum²³ and gold.²⁴ While platinum also failed (Entry 2), gold(I)-chloride gave complete conversion at room temperature, but almost none of the desired spiroketal (**31a**) was observed. Instead elimination product **32** was isolated in 61% yield (Entry 3). Examination of other gold catalysts (such as AuClPPH₃ with or without added silver salts) failed to resolve this difficulty. Under nearly identical conditions propargyl benzoate **29b** gave a 17% yield of spiroketal **31b** along with 47% of elimination product **32** (Entry 4). Examination of several Brønsted acid additives showed that the distribution of products could be biased toward the formation of **31b** in the presence of PPTS (Entry 5). In the presence of stronger acid (CSA, Entry 6) the elimination problem was significantly exacerbated and the overall yield decreased significantly. Variation of the solvent caused significant perturbation to the ratio of products. Moving from MTBE to THF (Entry 7) significantly improved the yield of **31b**, but continuing to an even more polar solvent, such as methanol increased the elimination problem relative to THF. Out of the conditions surveyed, the best were those outlined in Entry 7 and these were used for all further spiroketalization reactions.

Aponick²⁵ reported the preparation of unsaturated 1,7-dioxaspiro[5.5]undec-4-ene spiroketals *via* a gold catalyzed dehydrative cyclization of monopropargylic triols. Thus in retrospect, it is not surprising that triol **30b** was so prone to elimination. However, the leaving group ability of benzoate appears to be greater than a hydroxyl group at least in a thermodynamic sense (pK_a 15.7 vs. 4.3 in water).²⁶ At first, we were somewhat puzzled by this counterintuitive observation. This issue is also convoluted by the fact that it is unclear whether the gold cation is lost *via* protonation prior to or following the elimination. The mechanism below (Scheme 7) assumes loss of gold cation prior to formation of the spiroketal. While we cannot rule out that enol ether **33** or **34** could induce elimination directly after protonolysis of the gold cation, the heightened propensity of the hydroxy group to eliminate suggests that gold is involved in the elimination step prior to protonolysis. This premise is consistent with the fact that weak acid disfavors elimination, as acid should favor protonolysis of the gold cation before elimination can occur. The length the reaction time did not alter the ratio of **31b** and **32**, implying that **31b** does not convert to **32** directly *via* reionization and solvolysis. The first report of a gold catalyzed spiroketalization²⁴ presented several reactions in which 5-*exo*-dig cyclization dominated over a possible 6-*endo*-dig pathway. Therefore, it seems more likely that the C₂₇ hydroxyl group attacks first *via* a 6-*exo* dig pathway to form intermediate **33** or **34**, although

we cannot rule out a 6-*endo* cyclization of the C₁₉ hydroxyl group, as it would lead to the same product. Nucleophilic addition to C₂₃ in preference to C₂₂ is enforced by the benzoyloxy or hydroxy group at C₂₁, which through its inductive electron withdrawing nature disfavors a cationic intermediate at C₂₂.²⁷ Calculations have indicated that this oxy-metalation consists of a *syn* addition across the alkyne, resulting from prior coordination of the hydroxy group to the gold Lewis acid (see intermediates **33** and **34**).²⁸ One possible rationalization for the benzoyl groups lessened propensity toward elimination would be formation of a 6-membered chelate (**35**), in which the σ^* orbital of the C₂₁-O bond is orthogonal to both the π -system and the Au-C σ bond, leading to a kinetically inert conformation. Intermediate **34** instead encourages elimination through coordination of the hydroxyl group to the gold center, resulting in formation allenyl enol ether **37**, which could undergo reversible protonation to form oxonium **38**, explaining the formation of the *cis* olefin geometry required for **32**.

Because we had assumed that triol **30b** would be a superior spiroketalization substrate, it had not been anticipated that the benzoyl group would be present in the product. This led to unexpected protecting group difficulties as the benzoyl group was not compatible with conditions known to selectively cleave benzyl groups in the presence of *p*-methoxybenzyl groups (lithium 4,4'-di-*tert*-butylbiphenyl, THF, -78 °C).²⁹ Hydrogenolysis was unsatisfactory, and after surveying a fair number of conditions only a 40% yield of the desired alcohol (**39**) could be isolated (Equation 5).

As a result, the 1,3 dithiane group was reinvestigated and to our delight it proved to be an effective substrate for the spiroketalization (Scheme 8). In analogy to the spiroketalization with the benzyl ether, employing triol **30c** in the spiroketalization led almost exclusively to elimination product **42**. In contrast, benzoate **29c** led at slightly elevated temperature and increased catalyst loading (50 °C, 10 mol% AuCl, THF) to spiroketal **41** in 63% isolated yield. Discrimination of the protecting groups was now trivial and methyl iodide in buffered aqueous acetonitrile gave aldehyde **45** directly.

Our need to access mesylate **44** prompted us to examine the asymmetric alkynylation of isovaleraldehyde with methyl propiolate, a very challenging transformation³⁰ that was unprecedented before our communication.⁷ Implementation of our zinc alkynylation method³¹ led to the isolation of **43** in high yield and excellent enantioselectivity, the first such transformation reported for a simple aliphatic aldehyde. Under identical conditions, TMS acetylene fails entirely due to its increased basicity, which causes destructive polymerization of the aldehyde by a competing aldol pathway. Methyl propiolate serves as a very effective acetylene equivalent as saponification and decarboxylation were easily accomplished,³² yielding **44** after mesylation. Propargylation of **45** with **44** under the conditions of Marshall³³ afforded a 3.7:1 mixture of separable propargyl epimers (C₁₇) in a combined 90% yield. The stereochemistry of **46a** is assigned by Marshall's model in which propargyl mesylate **44** undergoes oxidative addition with palladium(0) with inversion, followed by

transmetallation to diethyl zinc with retention. The resulting allenylzinc undergoes nucleophilic addition to aldehyde **45** via coordination of the aldehyde to zinc. Thus, stereochemistry at C₁₆ is completely controlled, whereas C₁₇ is formed as a mixture favoring **46a** through a Zimmerman-Traxler transition state in which the aldehyde group occupies a pseudo-equatorial position. Both epimers were independently carried forward to the complete spiroketal fragment, as the C₁₇ stereochemistry is ultimately irrelevant and will be removed by oxidation later in the synthesis. After some experimentation, it proved best to incorporate the sensitive vinyl iodide last, giving the completed spiroketal fragment (**47**) in 16 linear steps from methyl 3-oxopentanoate.

Spiroketal product **41** was examined extensively by various NMR experiments (gCOSY, gHSQC, gHMBC, ROESY, Figure 3). The expected NOE correlations (H₁₉ to H₂₇ and H₁₉ to H₂₁) and coupling constants ($J_{C26-C27} = 2.5$ Hz [equatorial-axial], $J_{C21-C22} = 4.5$ Hz [axial-equatorial] 12.0 Hz [axial-axial], $J_{C19-C20} = 2.5$ Hz [axial-equatorial]) were observed within the bicyclic framework of spiroketal **41** consistent with the indicated structure. The strong resemblance of these data to the NMR spectrum of authentic ushikulide A (Figure 2) was taken to be a very positive sign and improved the likelihood that our synthetic spiroketal had the same relative stereochemistry as the natural product.

Synthesis of the Aliphatic Fragment

A first generation approach toward an aliphatic fragment to fulfill the role of synthon **4** began from the known dibromide **48** (available in three steps from the (*S*)-Roche ester, Scheme 9).^{20,34} Exposing dibromide **48** to *n*-BuLi led to the expected Fritsch-Buttenberg-Wiechell rearrangement³⁵ and the resulting lithium acetylide was quenched with *N*-methoxy-*N*-methylacetamide to yield ynone **49** in 76% yield. This ketone gave good reactivity in the direct zinc aldol reaction¹³ with aldehyde **50**^{13d} to yield aldol adduct **51** as a single diastereomer by ¹H NMR analysis. Unfortunately, hydrosilylation³⁶ of this product under the standard conditions (Cp*Ru(CH₃CN)₃PF₆, BDMSH, acetone) caused deprotection of the diethoxyketal. Changing the solvent to dichloromethane failed to remedy the problem. Therefore, aldehyde **53** was prepared in enantiopure form via an allylation reported by Mukaiyama.³⁷ This aldehyde also gave good yield and diastereoselectivity in the zinc aldol reaction to afford aldol adduct **54**, again as a single diastereomer. The alkyne moiety of **54** was reduced in a *syn* fashion via a palladium-catalyzed reduction,³⁸ yielding enone **55**. Lindlar reduction was also attempted on ynone **54** or the corresponding *syn* 1,3-diol after reduction of the ketone function. Both substrates gave preferential reduction of the terminal double bond, demonstrating the superior selectivity of homogeneous palladium catalysis in this case. Enone **55** was then reduced to the *syn* diol and protected as a cyclic acetonide (*syn* stereochemistry of **56** and **64** was consistent with Rychnovsky's ¹³C shift analysis).²¹ Epoxidation of **56** gave no diastereoselectivity and modest chemoselectivity for the more substituted olefin to yield epoxides **57** and **58** as a separable but unassigned mixture of diastereomers. Before any attempt was made to remedy this diastereoselectivity issue, the epoxide opening had to be demonstrated. Unfortunately, several attempts to accomplish this transformation failed to yield **59**, instead cleaving the *p*-methoxybenzyl group. While this general strategy still might have provided the completed aliphatic fragment,³⁹ a more direct strategy was already showing promise (*vide infra*).

One potential method to prepare the aliphatic fragment would be to couple a fully saturated aldehyde and ketone through the direct aldol methodology. Utilization of such an unstabilized ketone enolate with nearly 1:1 stoichiometry had not previously been demonstrated in this reaction (acetone and acetophenone are commonly employed as solvent or in excess [5-10 equivalents]). The requisite methyl ketone **61** was prepared easily via Wacker oxidation⁴⁰ of the known⁴¹ homoallylic alcohol **60** (Scheme 10). By using a preformed lithium enolate of ketone **61** with aldehyde **53** the expected aldol product could be generated, but low

stereoselectivity was observed as expected (~1:1 d.r., Entry 1). The standard zinc aldol conditions failed entirely to give the desired product and only the reduced primary alcohol (**63**) and recovered ketone (**61**) were observed (Entry 2). Presumably this reduction must proceed *via* a Meerwein/Ponndorf/Verley pathway. Employing *t*-BuOH instead of *i*-PrOH resulted in some yield of the desired product (**62**, Entry 3), while switching the solvent to 1,4-dioxane and increasing the catalyst loading gave an improved yield of aldol **62** in excellent stereoselectivity (Entry 4). Stoichiometric ProPhenol and diethylzinc failed to increase this yield and instead led to contamination of the product with various side products (Entry 5). Fortunately, the aldehyde and ketone could be recovered and resubjected to the reaction conditions, leading to an acceptable yield (65%) after three total runs. With ketone **62** in hand *syn*-reduction and acetonide formation gave **64**, which was converted to aliphatic fragment **65** in three additional steps including a Horner/Wadsworth/Emmons olefination.

Further attempts to optimize the aldol reaction were fruitless and short of modifying the ligand there seemed little hope for solving the problem of catalyst turnover that required several recycling iterations. To this end, we considered another aldol disconnection, wherein the C₇-C₈ bond instead of the C₈-C₉ would be formed in the aldol reaction. To attempt this, preparation of methyl ketone **67** was required rather than aldehyde **53**. The ideal method to accomplish this would be a stereoselective allylation of 2,3-butanedione (Equation 6). This reaction was unknown in the literature and a number of attempts were made to accomplish it, which are outlined below. Unfortunately, the best enantioselectivity obtained was 45% *ee*, (Entry 5) too low to carry forward with the desired route.

Entry	M =	Conditions	Yield (<i>ee</i>)*
1	(+)-Ipc ₂ B	AllylMgBr, (+)-Ipc ₂ OMe, Et ₂ O	>10% (9%)
2	Br	(+)-DIPT, CuI, DBU, Sn ^{II} (catecholate), CH ₂ Cl ₂	20% (7%)
3	PinB	20 mol% In(OTf) ₃ , 10 mol% (<i>S,S</i>)-ProPhenol, THF, 4 °C	60% (0%)
4	PinB	20 mol% Et ₂ Zn, 10 mol% (<i>S,S</i>)-ProPhenol, THF, 4 °C	25% (45%)
5	PinB	20 mol% Et ₂ Zn, 10 mol% (<i>S,S</i>)-ProPhenol, THF, <i>i</i> -PrOH, 4 °C	52% (44%)

* Yield and *ee* were determined by achiral and chiral GC respectively. For achiral GC naphthalene was employed as an internal standard.

(6)

Alternatively aldehyde **53** was transformed to silyl ether **68** in three steps. Addition of methyl magnesium bromide gave a mixture of secondary alcohols, which were oxidized without further purification. Finally, formation of the lithium enolate and silylation afforded the desired silyl enol ether.

(7)

Aldehyde partner **69** was prepared *via* oxidative cleavage of **60**. Moderate yield and diastereoselectivity were obtained in the Mukaiyama, aldol reaction⁴² using a slight excess of

68 in toluene with boron trifluoride diethyl etherate to yield the Felkin-Anh aldol product (**70**).⁴³ Reduction of this hydroxy ketone afforded the *syn*-diol, which was converted to an acetonide (**64**) identical to that prepared in our previous strategy. Alternatively, the PMB group was cyclized to the *p*-methoxyphenyl acetal (**71**). Upon irradiation of the benzylic proton strong NOE correlations were observed to H₅ and H₇. Coupling constant analysis ($J_{H5-H6} = 2.0\text{Hz}$ [axial-equatorial] and $J_{H6-H7} = 2.0\text{Hz}$ [axial-equatorial]) was also consistent with the chair form indicated, further supporting that the expected stereochemistry had been observed in both aldol reactions.

Completion of the Synthesis

Completion of the synthesis rested on two key bond-forming reactions – an $\text{sp}^3\text{-sp}^2$ Suzuki coupling⁴⁴ and an esterification. While a great many syntheses have relied on macrolactonization,⁴⁵ only a few have utilized Suzuki coupling (particularly an $\text{sp}^3\text{-sp}^2$ coupling) for the formation of macrocycles.⁴⁶ We therefore elected to attempt the Suzuki macrocyclization first in order to explore its potential in this arena (Scheme 12). Yamaguchi esterification⁴⁷ allowed for the direct coupling of **47a** and **65** to yield ester **72**. Hydroboration of **72** (9-BBN, THF) gave the expected preference for reaction of the terminal olefin as observed by ¹H NMR analysis. However, under a variety of conditions, which were later shown to be viable in an intermolecular reaction, macrocycle **73** could not be detected or isolated. The borate described above was injected *via* syringe pump to a solution containing palladium catalyst, base and solvent (palladium sources: Pd[PPh₃]₄, PdCl₂•(dppf) and Pd₂(dba)₃•CHCl₃; bases: Cs₂CO₃, TIOEt; solvents: dioxane, THF, DMF each with co-solvent of water). In many of these attempts the vinyl iodide was clearly consumed, unfortunately a number of products were formed, none of which corresponded to the macrocycle **73**, which was successfully isolated later on.

The alternative pathway (Suzuki coupling then macrolactonization) was attempted. From the unsaturated acid (**65**) hydroboration was feasible with 2.5 equivalents of 9-BBN. Immediate gas evolution gave evidence that the carboxylic acid was reacting with one equivalent of 9-BBN, however, this fast reaction was followed by consumption of the terminal olefin. The crude borate (**74**) was added along with vinyl iodide **47a** to a flask containing PdCl₂•(dppf), triphenylarsine and cesium carbonate.⁴⁸ At room temperature this gave *seco*-acid **75** directly.

Several conditions were applied to the *seco*-acid (**75**) to effect macrolactonization. Yonemitsu's modification of the Yamaguchi esterification protocol^{8b} failed to give any product (2,4,6-trichlorobenzoyl chloride, Et₃N, 4-DMAP, benzene). Applying the Wasserman-Kita esterification to macrolactonization⁴⁹ gave the first successful isolation of macrocycle **73** in ~30% yield, but attempts to improve the yield were unsuccessful (ethoxy acetylene, 2 mol% [RuCl₂(*p*-cymene)]₂, toluene, then 10 mol% CSA). The conditions of Shiina,⁵⁰ gave the best result (65%) for preparation of macrolactone **73**. Following cyclization, the secondary *tert*-butyldimethylsilyl ether was cleaved employing buffered HF•pyridine and the resulting alcohol was oxidized with Dess-Martin periodinane. The final deprotection required DDQ to oxidatively cleave the *p*-methoxybenzyl ethers and then acid to hydrolyze the cyclic acetonide. It proved to be essential to perform the oxidative deprotection first, as prior deprotection of the acetonide led to undesired cyclization during oxidative removal of the PMB group (PMP acetal or *p*-methoxybenzoate formation). When the PMB groups were removed first, the resulting triol could be cleanly isolated without interference from neighboring groups. In the final acid catalyzed deprotection the starting acetonide triol quickly interconverted to two other compounds of similar polarity as evidenced by TLC analysis (presumably these were migration of the acetonide to the C₅ or C₁₀ oxygens respectively), however, on prolonged stirring at room temperature (~4 hrs) these three compounds converged to a single spot of indistinguishable *R_f* when compared to an authentic sample of ushikulide A. Upon isolation, this compound

exhibited identical ^1H , ^{13}C , IR, HRMS, and HPLC properties when compared to the same authentic sample. Measurement of the optical rotation $\{[\alpha]_{\text{D}} (24\text{ }^\circ\text{C}): -12^\circ (c\ 0.28, \text{MeOH})\}$ confirmed that the absolute stereochemistry was the same as that of natural ushikulide A $\{[\alpha]_{\text{D}} (24\text{ }^\circ\text{C}): -13^\circ (c\ 0.50, \text{MeOH})\}$. In total, these data confirmed our tentative assignment and led us to assign the complete structure of (–)-ushikulide A as depicted in Scheme 12.

Summary

We have developed a flexible synthetic route, which establishes the full structure of (–)-ushikulide A for the first time. This route, which requires 21 linear and 40 total steps, provides the natural product in 2.2% yield over the longest linear sequence a significant accomplishment given the complexity of **1**. The key gold-catalyzed spiroketalization to yield **41** is the first example of a metal-catalyzed spiroketalization in which regioselectivity is controlled by the electronic bias of the alkyne. This implementation extends the methodology to 1,7-dioxaspiro [5.5]undecane ring systems, a motif found in many natural products. Application of our zinc aldol methodology to aldehyde **53** and ketone **61** gave excellent control of diastereoselectivity, although catalyst turnover still needs to be improved. Control of diastereoselectivity in this manner allowed for an otherwise unavailable synthetic disconnection resulting in the preparation of aliphatic fragment **65** in only 11 linear steps. Finally, effective use of Suzuki coupling, Shiina macrolactonization and the Marshall/Tamaru propargylation all contributed to the overall success of this synthetic endeavor.

This synthesis confirmed the ‘educated guess’ that was made when we began our synthesis, however, it is likely that almost any diastereomer of ushikulide might have been prepared *via* this general strategy. Furthermore, a large number of unassigned natural products could be targeted by this method. In total, there are at least 12 natural products as close in similarity to ushikulide as cytovaricin,⁵¹ and many other unknown natural products could be approached in such a manner. Each specific case would lead to a slightly different implementation of this method. For example, access to greater amounts of material would allow for degradation studies, whereas a less conformational flexible target would allow for more computational analysis than was possible here. The success of this project is a testament to the power of organic synthesis in the structural elucidation of complex natural products. Further studies to apply and improve upon this strategy in the synthesis of other members of the oligomycin-rutamycin family are underway and will be reported in due course.

Experimental Section

(2S,3S,4S,6R,8S)-2-((1,3-dithian-2-yl)methyl)-8-((R)-2-(4-methoxybenzyloxy)butyl)-3,9-dimethyl-1,7-dioxaspiro[5.5]undecan-4-yl benzoate (**41**)

A solution of diol **29c** (1.1492g, 1.82mmoles) in dry THF (0.05M, 36mL) is degassed (2x freeze/pump/thaw) and allowed to warm under argon. Gold (I) chloride (42.3mg, 0.18mmoles) and PPTS (45.7mg, 0.18mmoles) are added and degassing is repeated for 2 additional cycles. The reaction is then warmed to 50 $^\circ\text{C}$ for a period of 22 hours. After this time the solvent is removed under reduced pressure, a drop of triethyl amine is added, and the residue is purified by flash chromatography (9:1 hexane/THF) yielding unsaturated spiroketal **41** (325.6mg, 28%) followed by the desired spiroketal product **42** (729.6mg, 63%).

Data for **41**

R_f = 0.23 (9:1 Hexane/THF); $[\alpha]_{\text{D}} (23\text{ }^\circ\text{C}): -5.2^\circ (c=0.71, \text{CHCl}_3)$; IR (neat): 2936, 1719 (sharp) cm^{-1} ; ^1H NMR (500 MHz, C_6D_6): δ 8.17-8.16 (m, 2H), 7.39 (d, $J=8.5\text{Hz}$, 2H), 7.16-7.07 (m, 3H), 6.85 (d, $J=8.5\text{Hz}$, 2H), 5.83 (dt, 1H, $J=4.7, 12.1\text{Hz}$), 4.56 (d, $J=11.4\text{Hz}$, 1H), 4.50 (d, $J=11.4\text{Hz}$, 1H), 4.34 (ddd, 1H, $J=3.4, 3.6, 11.4\text{Hz}$), 4.26 (dd, 1H, $J=5.9, 8.5\text{Hz}$), 4.19 (dt,

$J=2.4, 6.8\text{Hz}, 1\text{H}$), 3.71-3.68 (m, 1H), 3.31 (s, 3H), 2.43-2.24 (m, 4H), 2.09 (t, $J=7.0\text{Hz}, 1\text{H}$), 1.92-1.84 (m, 3H), 1.73-1.58 (m, 4H), 1.50-1.20 (m, 8H), 1.14 (t, $J=7.4\text{Hz}, 3\text{H}$), 0.98 (d, $J=6.9\text{Hz}, 3\text{H}$), 0.87 (d, $J=6.9\text{Hz}, 3\text{H}$); ^{13}C NMR (125 MHz, C_6D_6): δ 165.3, 159.4, 132.8, 132.1, 129.8, 129.1, 113.9, 97.7, 78.3, 71.0, 69.9, 67.4, 54.7, 44.4, 39.3, 37.5, 36.0, 35.8, 30.5, 30.3, 30.2, 29.9, 29.9, 27.4, 26.8, 25.9, 11.4, 9.6, 5.4 ppm; HRMS (ESI+) calc'd for $\text{C}_{35}\text{H}_{48}\text{NaO}_6\text{S}_2$ ($\text{M}+\text{Na}$) $^+$ 651.2790, found 651.3694.

(*R*)-methyl 4-hydroxy-6-methylhept-2-ynoate (43)

A flask containing (*S,S*) ProPhenol (479.1mg, 0.75mmoles) dissolved in dry toluene (50mL) is sequentially treated with methyl propiolate (1.87mL, 21.0mmoles) followed by dimethyl zinc (1.2M in hexane, 17.5mL, 21.0mmoles). After 30 minutes isovaleraldehyde is injected (804 μL , 7.5mmoles). The reaction is transferred to the cold room (ca 4 °C) and stirred for a period of 60 hours. Concentrated aqueous ammonium chloride (50mL) and diethyl ether (50mL) are added and the reaction is transferred to a separatory funnel. The organic phase is dried over magnesium sulfate, filtered and concentrated under reduced pressure (ca 100 torr). The resulting solution is purified by flash chromatography (4:1 PE/Et₂O) and concentrated under reduced pressure (~100 torr) to yield propargyl alcohol **44a** as a clear oil (1.1193g, 88%).

Data for 43⁵²

Enantiomeric excess is determined to be 95% on Chiracel OD column; 98:2 heptane/isopropanol; Flowrate=0.8mL/min; Detector set to 220nm; R_t (major *S*): 21.6min, R_t (minor *R*): 23.67min; $R_f=0.30$ (2:1 Hexanes/Et₂O); $[\alpha]_D$ (23 °C): 13.8° ($c=0.93$, CHCl_3); IR (neat): 3419, 2959, 2237, 1719, 1436, 1254, 1070 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3): δ 4.52 (t, 1H, $J=7.0\text{Hz}$), 3.77 (s, 3H), 2.70 (s, broad, 1H), 1.89-1.81 (m, 1H), 1.70 (ddd, $J=7.0, 7.5, 13.5\text{Hz}$, 1H), 1.60 (ddd, $J=7.0, 7.0, 14.0\text{Hz}$, 1H), 0.94 (d, $J=7.0\text{Hz}$, 3H), 0.94 (d, $J=7.0\text{Hz}$, 3H) ppm; ^{13}C NMR (125 MHz, CDCl_3) δ 153.9, 88.8, 75.9, 60.5, 52.8, 45.6, 24.4, 22.4, 22.1 ppm.

(2*S*,3*S*,4*R*,6*R*,8*S*)-2-((2*R*,3*S*)-3-ethynyl-2-hydroxy-5-methylhexyl)-8-((*R*)-2-(4-methoxybenzyloxy)butyl)-3,9-dimethyl-1,7-dioxaspiro[5.5]undecan-4-yl benzoate (46)

Aldehyde **45** (132.9mg, 0.25mmoles) and propargyl mesylate **44** (71.5mg, 0.38mmoles) are dissolved in dry THF (1.7mL) and cooled to -78 °C. A solution of palladium acetate (5.6mg, 25.1 μmoles) and triphenylphosphine (6.6mg, 25.1 μmoles) in THF (1mL) is transferred into the reaction flask *via* syringe, followed by a solution of diethyl zinc in hexanes (0.75mL, 1M, 0.75mmoles). The reaction is allowed to warm to -20 °C for a period of 14hrs, after which time it is poured into a saturated solution of ammonium chloride. The organic phase is diluted with ethyl acetate, separated and dried over sodium sulfate. Following filtration and concentration the crude residue is purified by flash chromatography (19:1 toluene/EtOAc) yielding the major diastereomer **46a** (113.2 mg, 71%) followed by its C-17 epimer **46b** (30.0mg, 19%).

Data for 46a

$R_f=0.20$ (19:1 Toluene/EtOAc); $[\alpha]_D$ (23 °C): -69° ($c=0.93$, CHCl_3); IR (neat): 3505 (broad), 3305, 2955, 1718 (sharp), 1514 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3): δ 8.04 (d, $J=7.2\text{Hz}$, 2H), 7.57 (t, $J=7.4\text{Hz}$, 1H), 7.45 (t, $J=7.6\text{Hz}$, 2H), 7.29 (d, $J=8.5\text{Hz}$, 2H), 6.85 (d, $J=8.6\text{Hz}$, 2H), 5.54 (td, $J=12.0, 4.8\text{Hz}$, 1H), 4.50 (d, $J=11.4\text{Hz}$, 1H), 4.45 (d, $J=11.2\text{Hz}$, 1H), 4.17 (m, 1H), 4.02-3.95 (m, 1H), 3.87 (s, 1H), 3.78 (s, 3H), 3.73 (dd, $J=9.7, 3.0\text{Hz}$, 1H), 3.64 (dt, $J=10.0, 5.4\text{Hz}$, 1H), 2.34-2.22 (m, 1H), 2.17-2.10 (m, 1H), 2.04 (d, $J=2.3\text{Hz}$, 1H), 1.99-1.90 (m, 3H), 1.84-1.73 (m, 2H), 1.73-1.41 (m, 9H), 1.38 (dd, $J=14.36, 1.78\text{Hz}$, 1H), 1.17-1.05 (m, 1H), 0.97 (d, $J=6.9\text{Hz}$, 3H), 0.96 (d, $J=7.3\text{Hz}$, 3H), 0.95-0.90 (m, 6H), 0.84 (d, $J=6.6\text{Hz}$, 3H) ppm; ^{13}C NMR (125 MHz, CDCl_3): δ 165.7, 158.8, 132.9, 131.3, 130.5, 129.5, 128.7, 128.4, 113.7, 98.2, 84.6, 76.9, 73.0, 71.6, 70.8, 70.6, 69.3, 68.2, 55.2, 39.0, 37.0, 36.6, 36.2, 35.7,

30.1, 29.1, 26.6, 26.2, 25.8, 23.6, 21.1, 11.2, 9.2, 5.6 ppm; HRMS (ESI+) calculated for $C_{39}H_{54}O_7Na$ ($M+Na$)⁺ 657.3766 found 657.3767.

(2*S*,3*R*,4*S*,7*R*,8*R*)-1-(*tert*-butyldimethylsilyloxy)-7-hydroxy-3,8-bis(4-methoxybenzyloxy)-2,4,8-trimethylundec-10-en-5-one (62)

A solution of (*S,S*) ProPhenol (590.2mg, 0.92mmoles) in dioxane (8mL) is treated with diethyl zinc (1M in hexanes, 1.84mL) at ambient temperature. A separate flask is charged with ketone **61** (913.6mg, 2.31mmoles), aldehyde **53** (759.5mg, 3.24mmoles) and activated 4Å powdered molecular sieves (500mg). After 30 minutes the catalyst solution is transferred into the reaction flask at ambient temperature. The reaction is stirred for 45 hours and quenched with 0.1N HCl (50mL) and diluted with ethyl acetate (100mL). The solution is filtered and the organic phase is washed with an addition portion of 0.1N HCl followed by sodium bicarbonate (25mL portions). The organic phase is dried over sodium sulfate, filtered and concentrated. Flash chromatography (85:15 Hexane/EtOAc) gives aldol adduct **62** (518.5mg, 35%) as a single diastereomer along with a 1.4:1 mixture of the aldehyde and ketone (1.026g, 60%). Repetition of the procedure above 2 additional times (identical relative stoichiometry) with recovered starting material gives a total of 65% yield based on recovered starting material.

Data for 62

R_f = 0.33 (4:1 Hexane/EtOAc); $[\alpha]_D$ (23 °C): +28° (c =0.28, CH_2Cl_2); IR (neat): 3525 (broad), 2931, 1703 (sharp), 1514 cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$): δ 7.21 (d, J =9.3Hz, 4H), 6.84 (d, J =8.7Hz, 4H), 5.94-5.82 (m, 1H), 5.20-5.05 (m, 2H), 4.43-4.39 (m, 4H), 4.19-4.06 (m, 1H), 3.88-3.71 (m, 7H), 3.54-3.40 (m, 2H), 3.04-2.82 (m, 2H), 2.63-2.48 (m, 2H), 2.45-2.32 (m, 1H), 1.73-1.69 (m, 1H), 1.21 (s, 3H), 1.14 (d, J =6.7Hz, 3H), 0.85 (d, J =6.8Hz, 3H), 0.90 (s, 9H), 0.03 (s, 3H), 0.03 (s, 3H) ppm; ^{13}C NMR (125 MHz, $CDCl_3$): δ 213.9, 159.1, 133.7, 131.2, 130.8, 129.3, 128.8, 117.9, 113.8, 113.7, 79.7, 78.5, 73.9, 71.7, 65.5, 63.8, 55.3, 50.0, 43.5, 39.0, 38.8, 29.7, 25.9, 19.6, 18.2, 12.6, 11.5, -5.4, -5.5; HRMS (ESI+) calculated for $C_{36}H_{56}O_7NaSi$ ($M+Na$)⁺ 651.3693 found 651.3694.

Seco Acid (75)

Carboxylic acid **65** (64.8mg, 0.109mmoles) and 9-BBN (33.1mg, 0.271mmoles) are dissolved in THF (1mL) and stirred under a nitrogen atmosphere for a period of 14 hours. The solvent is removed under reduced pressure and water is injected (60 μ L). In a separate flask vinyl iodide **47** (109.5mg, 0.141mmoles) is dissolved in DMF (2mL) and the solvent is degassed (2 $\times$ freeze/pump/thaw). $PdCl_2(dppf)$ (7.9mg, 10.9 μ moles), cesium carbonate (106.5mg, 0.327mmoles) and triphenyl arsine (3.3mg, 11.0 μ moles) are added to the flask containing **74**. The DMF solution of **47** is transferred into the reaction flask and the reaction is stirred at ambient temperature under argon for 21 hours. The reaction is then cooled in an ice bath, 30% aqueous peroxide (2mL) is injected dropwise. After warming to ambient temperature for 30 minutes, the reaction is poured into sodium thiosulfate (5mL) acidified with 1M sodium bisulfate (pH=2, ~15mL) and further diluted with water (30mL). The aqueous solution is washed with ethyl acetate (3 $\times$ 25mL) and the combined organic extracts are dried over sodium sulfate, filtered and concentrated. Flash chromatography (gradient 10% to 20% to 35% chloroform/EtOAc with 1% methanol) gives *seco*-acid **75** as a white foam (89.1mg, 67%).

Data for 75

R_f =0.35 (2:1 chloroform/EtOAc with 1% methanol); $[\alpha]_D$ (23 °C): -16° (c =0.26, benzene); IR (neat): 3419 (broad), 2956, 1697 cm^{-1} ; 1H NMR (500 MHz, $CDCl_3$): δ 7.37-7.18 (m, 6H), 7.10 (dd, J =15.7, 7.9Hz, 1H), 6.91-6.85 (m, 6H), 5.83 (d, J =15.8Hz, 1H), 5.40 (dd, J =14.3, 7.5Hz, 1H), 5.19 (dd, J =15.2, 8.9Hz, 1H), 4.63-4.34 (m, 7H), 4.25-4.13 (m, 1H), 3.88-3.76 (m, 12H), 3.71-3.63 (m, 1H), 3.51-3.38 (m, 2H), 2.82-2.68 (m, 1H), 2.27-2.15 (m, 1H), 2.12-2.00

(m, 2H), 1.95-1.81 (m, 2H), 1.82-1.44 (m, 17H), 1.39 (s, 1H), 1.36 (s, 3H), 1.20-1.30 (m, 2H), 1.14 (s, 3H), 1.12 (d, $J=6.8\text{Hz}$, 3H), 0.98 (d, $J=6.9\text{Hz}$, 3H), 0.98-0.93 (m, 9H), 0.88 (s, 9H), 0.82 (d, $J=6.6\text{Hz}$, 3H), 0.81-0.77 (m, 6H), 0.07 (s, 6H) ppm; ^{13}C NMR (125 MHz, CDCl_3): δ 170.2, 159.1, 159.0, 158.7, 154.4, 132.4, 132.0, 131.1, 130.8, 129.2, 128.8, 128.3, 119.9, 113.7, 113.7, 113.7, 98.4, 97.5, 82.5, 77.9, 74.1, 73.7, 73.7, 73.5, 70.7, 69.6, 68.2, 67.3, 64.3, 55.2 (3C), 47.1, 41.1, 40.0, 39.4, 37.4, 36.4, 34.6, 30.3, 30.1, 29.8, 28.3, 27.8, 27.4, 27.0, 26.8, 26.5, 26.0, 25.6, 24.3, 22.6, 20.6, 19.3, 18.1, 17.5, 14.9, 13.6, 10.9, 9.8, 9.3, -4.1, -4.3 ppm; HRMS (ESI+) calculated for $\text{C}_{73}\text{H}_{114}\text{O}_{14}\text{NaSi}$ ($\text{M}+\text{Na}$) $^{+}$ 1265.7876 found 1265.7885.

Macrocycle (73)

A solution of *seco*-acid **75** (84.3mg, 68.8 μmoles) in dry DCE (5mL) is transferred to a gastight syringe and injected into a flask containing 2-nitrobenzoic anhydride (freshly recrystallized, 38.6mg, 0.105mmoles) and 4-DMAP (18.0mg, 0.126mmoles) in additional DCE (55mL) over a period of 18 hours. After 2 additional hours the reaction is poured into a separatory funnel containing saturated sodium bicarbonate (30mL) and ethyl acetate (90mL). The aqueous phase is separated and the organic phase is washed with ammonium chloride and sodium bicarbonate (30mL portions). The organic phase is then dried over sodium sulfate, filtered and concentrated. Flash chromatography (95:5 toluene/ethyl acetate) gives the macrocycle as a white foam (55.4mg, 65%).

Data for 73

$R_f=0.54$ (9:1 toluene/EtOAc); $[\alpha]_D$ (23 °C): -20° ($c=0.10$, chloroform); IR (neat): 2926, 1718, 1514 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3): δ 7.29-7.21 (m, 6H), 6.88-6.82 (m, 6H), 6.83 (dd, $J=15.6$, 7.9Hz, 1H), 5.83 (d, $J=15.8\text{Hz}$, 1H), 5.54-5.28 (m, 2H), 5.19 (dd, $J=15.2$, 9.3Hz, 1H), 4.58-4.35 (m, 6H), 4.00-3.95 (m, 1H), 3.85-3.83 (m, 1H), 3.82-3.75 (m, 10H), 3.67-3.61 (m, 1H), 3.48-3.34 (m, 2H), 3.18 (d, $J=7.0\text{Hz}$, 1H), 2.78-2.61 (m, 1H), 2.12-1.91 (m, 4H), 1.90-1.77 (m, 2H), 1.79-1.42 (m, 19H), 1.39 (s, 3H), 1.36 (s, 3H), 1.28-1.22 (m, 2H), 1.14 (s, 3H), 1.12 (d, $J=6.8\text{Hz}$, 3H), 0.98 (d, $J=6.9\text{Hz}$, 3H), 0.96-0.90 (m, 6H), 0.88 (s, 9H), 0.85-0.79 (m, 3H), 0.78-0.74 (m, 6H), 0.07 (s, 3H), 0.03 (s, 3H) ppm; ^{13}C NMR (125 MHz, CDCl_3): δ 164.5, 159.1, 158.9, 158.7, 150.6, 132.5, 131.2, 130.9, 129.4, 129.2, 128.6, 128.2, 122.2, 113.7, 113.7, 113.6, 98.4, 97.4, 84.4, 78.2, 77.9, 74.2, 73.9, 73.4, 70.7, 69.7 (3C), 67.4, 63.7, 55.2 (3C), 47.1, 42.3, 42.1, 37.5, 36.0, 34.3, 33.7, 33.0, 30.2, 30.1, 29.9, 29.7, 26.9, 26.4, 26.0, 25.5, 24.3, 22.6, 21.3, 20.3, 19.5, 18.1, 17.7, 10.9, 9.1, 7.8, 4.7, -4.2, -4.4 ppm; HRMS (ESI+) calculated for $\text{C}_{73}\text{H}_{112}\text{O}_{13}\text{NaSi}$ ($\text{M}+\text{Na}$) $^{+}$ 1247.7770 found 1247.7766.

(-)-Ushikulide A (1)

A plastic vial is charged with THF (10mL), pyridine (5.7mL) and HF-pyridine (Aldrich, 3.0mL). To a second plastic vial containing macrocycle **73** (10.6mg, 8.7 μmoles) is added the stock solution prepared above (5mL). The reaction is stirred under nitrogen for 26 hours and diluted with ethyl acetate (100mL) and diluted with water (25mL). Finally, the biphasic mixture is neutralized with saturated sodium bicarbonate (~20mL). The organic phase is then washed with copper sulfate, water, and brine (30mL portions). The organic phase is then dried over sodium sulfate, filtered and concentrated. Flash chromatography (9:1 toluene/EtOAc) gives the alcohol as a white foam.

The alcohol prepared above is immediately dissolved in dichloromethane (1mL) and Dess-Martin reagent is added (15.0mg). The milky white solution is stirred for 4 hours and quenched with sodium thiosulfate (5mL) and diluted with diethyl ether (40mL). The organic phase is washed with sodium bicarbonate (2 $\times$ 15mL) dried over sodium sulfate, concentrated and used directly in the next step without further purification.

The crude ketone is dissolved in dichloromethane (3mL) and water (0.5mL). DDQ (10.2mg) is added at ambient temperature and the red solution is stirred vigorously for 45 minutes. Silica gel (300mg) is added along with additional dichloromethane (10mL). The solution is concentrated to dryness and the residual solid is loaded onto a short column of silica gel. The product elutes (single fraction collected, 9:1 DCM/EtOAc) separating easily from the red catechol byproduct. After concentration the crude triol is suspended in a 3:2 mixture of glacial acetic acid and water (3mL) and the reaction is stirred vigorously under nitrogen for 5 hours. The solvent is removed *in vacuo*, and the residue is dissolved three times in toluene and concentrated to dryness (for the first cycle a minimum amount of methanol is added to give a homogeneous solution). The resulting oil is purified by flash chromatography (10:1 chloroform/methanol) yielding **1** as a white powder (3.3mg, 52% over 3 steps).

Data for Synthetic (–)-Ushikulide A⁵³

R_f =0.30 (9:1 chloroform/methanol); $[\alpha]_D$ (23 °C): -12° (c =0.28, CH₃OH); IR (neat): 3440 (broad), 1711 (sharp), 1463, 1276, 985 cm⁻¹; ¹H NMR (600 MHz, CD₃OD): δ 6.76 (dd, J =15.7, 10.1Hz, 1H), 5.81 (d, J =15.7Hz, 1H), 5.60 (ddd, J =15.0, 9.4, 4.2Hz, 1H), 5.44 (dd, J =15.4, 8.8Hz, 1H), 5.30 (ddd, J =11.6, 5.0, 5.0Hz, 1H), 4.33 (t, broad, J =8.2Hz, 1H), 4.09 (m, 1H), 4.02 (m, 1H), 3.78 (m, 1H), 3.62 (J =dd, 9.5, 1.5Hz, 1H), 3.18 (J =dd, 11.0, 2.0Hz, 1H), 3.07 (ddd, J =15.1, 9.0, 8.0Hz, 1H), 2.78 (dd, J =17.6, 6.6Hz, 1H), 2.73 (J =dd, 17.4, 8.1Hz, 1H), 2.46 (m, 1H), 2.23 (m, 1H), 2.19 (m, 1H), 2.15-2.00 (m, 3H), 1.80-1.40 (m, 18H), 1.28 (m, 1H), 1.14 (d, J =6.6Hz, 3H), 1.06 (s, 3H), 0.98 (t, J =7.3Hz, 3H), 0.94 (d, J =7.1Hz, 3H), 0.91 (d, J =6.8Hz, 3H), 0.87 (d, J =7.1Hz, 3H), 0.85 (d, J =7.0Hz, 3H), 0.79 (d, J =7.0Hz, 3H) ppm; ¹³C NMR (125 MHz): δ 212.5, 166.5, 152.1, 136.6, 129.5, 122.7, 98.8, 81.3, 75.9, 75.4, 74.9, 71.2, 70.5, 69.0, 67.1, 57.8, 42.9, 41.6, 41.9, 41.3, 39.7, 38.4, 36.3, 36.3, 34.6, 34.5, 33.0, 32.0, 30.6, 27.5, 26.6, 24.1, 23.2, 22.7, 22.3, 17.9, 11.4, 10.2, 6.1, 4.2 ppm; HRMS (ESI+) calculated for C₄₀H₆₈NaO₁₀ (M+Na)⁺ 731.4710, found 731.4700.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We gratefully acknowledge K. Takahashi for providing an authentic sample of ushikulide A and D. A. Evans for providing spectra of cytovaricin. We also acknowledge S. Lynch for his invaluable help in the acquisition of 2-D NMR data. We thank the Stanford Graduate Fellowship Program and NIH (GM 13598) for their financial support of our program, Johnson Matthey for their supply of precious metal salts and Aldrich for ProPhenol.

References and Footnotes

- (1). Nicolaou KC, Snyder SA. Angew. Chem. Int. Ed 2005;44:1012.
- (2)(a). Trost BM, Harrington PE. J. Am. Chem. Soc 2004;126:5028. [PubMed: 15099060] (b) Trost BM, Wroblewski ST, Chisholm JD, Harrington PE, Jung M. J. Am. Chem. Soc 2005;127:13589. [PubMed: 16190724] (c) Trost BM, Harrington PE, Chisholm JD, Wroblewski ST. J. Am. Chem. Soc 2005;127:13598. [PubMed: 16190725]
- (3). Takahashi K, Yoshihara T, Kurosawa K. J. Antibiotics 2005;58:420. [PubMed: 16156521]
- (4). Sakurai T, Kihara T, Isono K. Acta. Cryst 1983;C39:295.
- (5). Evans DA, Kaldor SW, Jones TK, Clardy J, Stout TJ. J. Am. Chem. Soc 1990;112:7001.
- (6). Takahashi, K. Attempts to prepare crystals of ushikulide A suitable for X-ray diffraction were unsuccessful in this case. Jul 22. 2008 personal communication
- (7). Trost BM, O'Boyle BM. J. Am. Chem. Soc 2008;130:16190. [PubMed: 18989964]
- (8)(a). References 7, ^{9a} and ^b use a similar Suzuki/macrolactonization strategy, whereas refs ^{9c} and ^{9d} employ a related Stille/macrolactonization strategy: White JD, Hanselmann R, Jackson RW, Porter

- WJ, Ohba Y, Tiller T, Wang S. *J. Org. Chem* 2001;66:5217. [PubMed: 11463276] (b) Evans DA, Ng HP, Rieger DL. *J. Am. Chem. Soc* 1993;115:11446. (c) Panek JS, Jain NF. *J. Org. Chem* 1998;63:4572. (d) Panek JS, Jain NF. *J. Org. Chem* 2001;66:2747. [PubMed: 11304197]
- (9). Chemler SR, Trauner D, Danishefsky SJ. *Angew. Chem. Int. Ed* 2001;40:4544.
- (10). Utimoto K. *Pure Appl. Chem* 1983;55:1845.
- (11). Burns NZ, Baran PS, Hoffmann RW. *Angew. Chem. Int. Ed* 2009;48:2854.
- (12)(a). For previous examples which employ the alkyne as a spiroketal surrogate see: Li Y, Zhou F, Forsyth CJ. *Angew. Chem. Int. Ed* 2007;46:279. (b) Trost BM, Weiss AH. *Angew. Chem. Int. Ed* 2007;46:7664.
- (13)(a). Methyl Ketones: Trost BM, Ito H. *J. Am. Chem. Soc* 2000;122:12003. (b) Trost BM, Ito H, Silcoff ER. *J. Am. Chem. Soc* 2001;123:3367. [PubMed: 11457073] (c) Trost BM, Fettes A, Shireman BT. *J. Am. Chem. Soc* 2004;126:2660. [PubMed: 14995157] (d) Trost BM, Shin S, Sclafani JA. *J. Am. Chem. Soc* 2005;127:8602. [PubMed: 15954759] (e) Trost BM, Silcoff ER, Ito H. *Org. Lett* 2001;3:2497. [PubMed: 11483044] (f) Trost BM, Malhotra S, Fried BA. *J. Am. Chem. Soc* 2009;131:1674. [PubMed: 19191696]
- (14)(a). Crimmins MT, She J. *Synlett* 2004;8:1371. (b) Crimmins MT, King BW, Tabet EA, Chaudhary K. *J. Org. Chem* 2001;66:894. [PubMed: 11430110]
- (15)(a). Barry CS, Bushby N, Harding JR, Willis CL. *Org. Lett* 2005;7:2683. [PubMed: 15957921] (b) Dickschat JS, Bode HB, Mahmud T, Müller R, Schulz S. *J. Org. Chem* 2005;70:5174. [PubMed: 15960521] (c) Hatakeyama S, Kawamura M, Shimanuki E, Saijo K, Takano S. *Synlett* 1992;2:114.
- (16). Kitamura M, Tokunaga M, Ohkuma T, Noyori R. *Org. Synth* 1998;71:589.
- (17)(a). Keck GE, Abbott DE, Wiley MR. *Tetrahedron Lett* 1987;28:139. (b) Keck GE, Boden EP. *Tetrahedron Lett* 1984;25:265. (c) Keck GE, Abbott DE. *Tetrahedron Lett* 1984;25:1883. (d) Keck GE, Abbott DE, Boden EP, Enholm EJ. *Tetrahedron Lett* 1984;25:3927.
- (18)(a). Nicolaou KC, Li Y, Fylaktakidou KC, Mitchell HJ, Wei H-X, Weyershausen B. *Angew. Chem. Int. Ed* 2001;40:3849. (b) Chen S-H, Horvath RF, Joglar J, Fisher MJ, Danishefsky SJ. *J. Org. Chem* 1991;56:5834.
- (19). Brown HC, Bhat KS. *J. Am. Chem. Soc* 1986;108:5919.
- (20). Nakatsuka M, Ragan JA, Sammakia T, Smith DB, Uehling DE, Schreiber SL. *J. Am. Chem. Soc* 1990;112:5583.
- (21). Rychnovsky SD, Rogers B, Yang G. *J. Org. Chem* 1993;58:3511.
- (22). Matsumura K, Hashiguchi S, Ikariya T, Noyori R. *J. Am. Chem. Soc* 1997;119:8738.
- (23)(a). Dieguez-Vazquez A, Tzschucke CC, Lam WY, Ley SV. *Angew. Chem. Int. Ed* 2008;47:209. (b) Liu B, De Brabander JK. *Org. Lett* 2006;8:4907. [PubMed: 17020333]
- (24)(a). Antoniotti S, Genin E, Michelet V, Genêt JP. *J. Am. Chem. Soc* 2005;127:9976. [PubMed: 16011342]
- (25). Aponick A, Li C-Y, Palmes JA. *Org. Lett* 2009;11:121. [PubMed: 19049406]
- (26)(a). Bordwell FG. *Acc. Chem. Res* 1988;21:456. (b) Harned HS, Robinson RA. *Trans. Faraday Soc* 1940;36:973.
- (27)(a). This electronic bias has been observed in the mechanistically related Wacker oxidation of terminal alkenes with allylic alkoxy substituents: Kang S-K, Jung K-Y, Chung J-U, Namkoong E-Y, Kim T-H. *J. Org. Chem* 1995;60:4678.
- (28)(a). Evidence for *syn* oxymetalation in gold-catalyzed hydration: Teles JH, Brode S, Chabanas M. *Angew. Chem. Int. Ed* 1998;37:1415. And in a related palladium-catalyzed Wacker-type cyclization: (b) Hayashi T, Yamasaki K, Mimura M, Uozumi Y. *J. Am. Chem. Soc* 2004;126:3036. [PubMed: 15012118]
- (29). Crimmins MT, McDougall PJ, Ellis JM. *Org. Lett* 2006;8:4079. [PubMed: 16928078]
- (30)(a). The method of Carreira is the most general, however, its catalytic implementation, particularly for α -mono-substituted or acidic aldehydes is somewhat unreliable: Anand NK, Carreria EM. *J. Am. Chem. Soc* 2001;123:9687. [PubMed: 11572696] (b) Tullington M, DeBerardinis AM, Pu L. *Org. Lett* 2009;11:2441. [PubMed: 19422198]
- (31). Trost BM, Weiss AH, von Wangelin AJ. *J. Am. Chem. Soc* 2006;128:8. [PubMed: 16390095]
- (32). Trost BM, Weiss AH. *Org. Lett* 2006;8:4461. [PubMed: 16986925]

- (33). Marshall JA, Schaaf GM. *J. Org. Chem* 2001;66:7825. [PubMed: 11701042]
- (34). Mulzer J, Berger M. *J. Org. Chem* 2004;69:891. [PubMed: 14750819]
- (35)(a). Corey EJ, Fuchs PL. *Tetrahedron Lett* 1972;36:3769. (b) Orita A, Otera J. *Chem. Rev* 2006;106:5387. [PubMed: 17165692]
- (36). Trost BM, Ball ZT. *J. Am. Chem. Soc* 2005;127:17644. [PubMed: 16351094]
- (37). Yamada K, Tozawa T, Nishida M, Mukaiyama T. *Bull. Chem. Soc. Jpn* 1997;70:2301.
- (38). Trost BM, Braslau R. *Tetrahedron Lett* 1989;30:4657.
- (39). Torres W, Rodriguez RR, Prieto JA. *J. Org. Chem* 2009;74:2447. [PubMed: 19236033]
- (40). Tsuji J. *Synthesis* 1984:369.
- (41)(a). Keck GC, Abbott DE, Boden EP, Enhom EJ. *Tetrahedron Lett* 1984;25:3927. For detailed experimental conditions see Reference 10a.
- (42). Mahrwald R. *Chem. Rev* 1999;99:1095. [PubMed: 11749441]
- (43). Mitchell IS, Pattenden G, Stonehouse J. *Org. Biomol. Chem* 2005;3:4412. [PubMed: 16327903]
- (44). Chemler SR, Trauner D, Danishefsky SJ. *Angew. Chem., Int. Ed* 2001;40:4544.
- (45). Parenty A, Moreau X, Campagne JM. *Chem. Rev* 2006;106:911. [PubMed: 16522013]
- (46)(a). White JD, Tiller T, Ohba Y, Porter WJ, Jackson RW, Wang S, Hanselmann R. *Chem. Comm* 1998;79 (b) Chemler SR, Danishefsky SJ. *Org. Lett* 2000;2:2695. [PubMed: 10990430] (c) Kallan NC, Halcomb RL. *Org. Lett* 2000;2:2687. [PubMed: 10990428] (e) Mohr PJ, Halcomb RL. *J. Am. Chem. Soc* 2003;125:1712. [PubMed: 12580592] (c) Williams DR, Walsh MJ, Miller NA. *J. Am. Chem. Soc* 2009;131:9038. [PubMed: 19485326]
- (47). Füxsrstner A, Kattnig E, Lepage O. *J. Am. Chem. Soc* 2006;128:9194. [PubMed: 16834393]
- (48)(a). Miyaoura N, Suzuki A. *Chem. Rev* 1995;95:2457. (b) Kapur M, Khartulyari A, Maier ME. *Org. Lett* 2006;8:1629. [PubMed: 16597127]
- (49). Trost BM, Chisholm JD. *Org. Lett* 2002;4:3743. [PubMed: 12375933]
- (50)(a). Shiina I, Kubota M, Oshiumi H, Hashizume M. *J. Org. Chem* 2004;69:1822. [PubMed: 15058924] (b) Shiina I. *Chem. Rev* 2007;107:239. [PubMed: 17212476]
- (51)(a). The Dunaimycins are the largest class of related natural products: Karwowski JP, Jackson M, Maus ML, Kohl WL, Humphrey PE, Tillis PM. *J. Antibiotics* 1991;44:1312. [PubMed: 1778784] (b) Hochlowski JE, Mullally MM, Brill GM, Whittern DN, Buko AM, Hill P, McAlpine JB. *J. Antibiotics* 1991;44:1318. [PubMed: 1778785] (c) Burres NS, Premachandran U, Frigo A, Swanson SJ, Mollison KW, Fey TA, Krause RA, Thomas VA, Lane B, Miller LN, McAlpine JB. *J. Antibiotics* 1991;44:1331. [PubMed: 1723403] (d) Yamazaki M, Yamashita T, Harada T, Nishikiori T, Saito S, Shimada N, Fujji A. *J. Antibiotics* 1992;45:171. [PubMed: 1556008] (e) Kirst HA, Mynderse JS, Martin JW, Patrick JB, Paschal JW, Steiner JLR, Lobkovsky E, Clardy J. *J. Antibiotics* 1996;49:162. [PubMed: 8621357] (f) Hosotani N, Kumagai K, Nakagawa H, Shimatani T, Saji I. *J. Antibiotics* 2005;58:409. [PubMed: 16156518]
- (52). **43** has been synthesized previously: Rajaram AR, Pu L. *Org. Lett* 2006;8:2019. [PubMed: 16671771]
- (53). The

Figure 1.
Tentative Stereochemical Assignment of (-)-Ushikulide A

Figure 2.
Comparison of Cytovaricin and Ushikulide A

Figure 3.
NOE and Coupling Constant Analysis of the Spiroketal

Scheme 1.
Retrosynthetic Analysis of Ushikulide A

Scheme 2.

Preparation of Several Aldehyde Partners through a Crimmins Aldol Reaction

Entry	M =	Conditions	Yield (19a:19b)*
1	Bu ₃ Sn	CH ₂ Cl ₂ , BF ₃ •OEt ₂	64% (2:1)
2	Bu ₃ Sn	CH ₂ Cl ₂ , SnCl ₄	57% (7:3)
3	(+)-Ipc ₂ B	i. <i>cis</i> -2-butene, <i>t</i> -BuOK, <i>n</i> -BuLi, THF ii. (+)-Ipc ₂ BOMe iii. 17 , BF ₃ •OEt ₂	80% (20:1)
4	(-)-Ipc ₂ B	i. <i>cis</i> -2-butene, <i>t</i> -BuOK, <i>n</i> -BuLi, THF ii. (-)-Ipc ₂ BOMe iii. 17 , BF ₃ •OEt ₂	75% (1:20)

*Isolated yields after purification by flash column chromatography. Diastereomeric ratios were determined by ¹H NMR analysis of the crude reaction mixture prior to purification.

Scheme 3.
Preparation of the Alkyne Fragment

Entry	R =	Conditions	Yield (<i>anti:syn</i>)*
1	TBS	<i>n</i> -BuLi, THF	90% (2:3)
2	TBS	<i>n</i> -BuLi, LiBr, 4 Å M.S., THF	77% (1:6)
3	TBS	<i>n</i> -BuLi, Et ₂ AlCl, toluene	38% (1:2.5)
4	MOM	LiN(<i>i</i> -Pr) ₂ , THF	43% (4:1)

*Isolated yields after purification by flash column chromatography. Diastereomeric ratios were determined by ¹H NMR analysis of the crude reaction mixture prior to purification.

Scheme 4.
Attempted Diastereoselective Alkyne Addition

Scheme 5.
Modification of the Spiroketalization Substrate

Entry	R =	Conditions	Yield (31:32)*
1	H	10 mol% $\text{PdCl}_2(\text{CH}_3\text{CN})_2$, PPTS, MeOH, 50 °C	No Pdt. observed
2	H	10 mol% $\text{Pt}(\text{COD})\text{Cl}_2$, <i>i</i> -Pr ₂ O, PPTS, 60 °C	No Pdt. observed
3	H	5 mol% AuCl, MeOH, PPTS, r.t.	Trace : 61%
4	Bz	5 mol% AuCl, MTBE, r.t.	17% : 47%
5	Bz	5 mol% AuCl, PPTS, MTBE, r.t.	41% : 41%
6	Bz	5 mol% AuCl, CSA, MTBE, r.t.	24% : 34%
7	Bz	5 mol% AuCl, PPTS, THF, r.t.	69% : 15%

* Yields for Entries 1-6 were determined through ¹H NMR analysis of the crude reaction mixture using benzyl benzoate as an internal standard. Entry 7 is the average isolated yield of two reactions, after separation of **31b** and **32** by column chromatography. Both reactions were performed on greater than 150mg scale.

Scheme 6.
Optimization of the Spiroketalization Reaction

Scheme 7.
Mechanism of the Spiroketalization Reaction

Scheme 8.
Completion of the Spiroketal Fragment

Scheme 9.
First Approach Toward the Aliphatic Fragment

Entry	Conditions	Yield (d.r.)*
1	LiN(<i>i</i> -Pr) ₂ , THF, then add 53	60% (~1:1)
2	10 mol% (<i>S,S</i>)-ProPhenol, 20 mol% Et ₂ Zn, <i>i</i> -PrOH, THF	no pdt; 63 isolated
3	10 mol% (<i>S,S</i>)-ProPhenol, 20 mol% Et ₂ Zn, <i>t</i> -BuOH, THF	~10% (>20:1)
4	30 mol% (<i>S,S</i>)-ProPhenol, 60 mol% Et ₂ Zn, <i>t</i> -BuOH, 1,4-dioxane	35% (>20:1)
5	1.1 eq. (<i>S,S</i>)-ProPhenol, 2.2 eq. Et ₂ Zn, <i>t</i> -BuOH, 1,4-dioxane	not clean
6	Same as Entry 4, three total runs with recovered starting material	65% (>20:1)

*Isolated yields after purification by flash column chromatography. Diastereomeric ratios were determined by ¹H NMR analysis of the crude reaction mixture prior to purification. Entry 6 is based on recovered starting material (b.o.r.s.m.) after three total reactions.

Scheme 10.
Second Generation and Completion of the Aliphatic Fragment

Scheme 11.

An Alternative Aldol Disconnection in the Synthesis of the Aliphatic Fragment

Scheme 12.
Completion of the Synthesis