Undertreatment of Cancer Patients With Chemotherapy Is a Global Concern

By Gary H. Lyman, MD, MPH

Gary H. Lyman,
MD, MPH

The efficacy of cancer chemotherapy is generally established on the basis of randomized controlled clinical trials evaluating a particular drug or combination using a specific dose and schedule. In adult patients with cancer, drug dosing has traditionally been based on a patient's estimated body-surface area (BSA), although there are little data supporting such a strategy. Despite continuing

controversy concerning the value of dose escalation and intensification schedules, there exists compelling preclinical and clinical evidence indicating that reductions in standard dose intensity may compromise disease-free and overall survival in the curative setting.¹⁻⁴ In practice, however, the delivery of full standard dose intensity is often not achieved for a variety of reasons including fear of excessive toxicity in elderly and obese patients.^{5,6}

Cancer is largely a disease of aging, and the treatment of cancer in elderly patients should be cognizant of both limited bone marrow reserves and any associated comorbidity.⁷ With modern supportive care, however, it has been demonstrated that most elderly patients with cancer can both tolerate and benefit from standard chemotherapy regimens.⁷⁻¹⁰ Likewise, concerns about overdosing the obese patient with cancer based on the use of actual body weight are largely unfounded.^{5,11-13} In fact, an obese patient with cancer often experiences less toxicity while deriving less benefit from such undertreatment, perhaps explaining the poorer outcomes often observed in such patients.^{11,14} Recent pharmacokinetic studies have clearly demonstrated that actual rather than ideal body weight should be used in dose calculations.¹⁵

Nevertheless, recent investigations, including the study from Australia reported in this issue of the *Journal of Oncology Practice*, confirm that many oncologists continue to either use "ideal" rather than actual body weight to calculate BSA or cap the BSA at 2.0 m², resulting in a systematic underdosing of

chemotherapy among overweight and obese patients.¹⁶ A large proportion of practicing oncologists and trainees in Australia responded to a survey soliciting their approach to dosing and dose reduction among the fit, the elderly, the obese, and the thin patient with cancer. While the majority stated that they dose based on actual body weight, half of all respondents indicated that they cap at a BSA 2.0 m² when treating the obese patient with cancer. The majority of the remaining respondents dose based on ideal body weight, leaving only 6% who dose on the basis of actual weight. Only one fourth of oncologists indicated that they rarely dose reduce in this setting while another fourth do so routinely and half reduce in some cases. The investigators in the current study did not survey chemotherapy dosing in actual practice. However, the responses obtained are consistent with those found among oncology practices in the United States and elsewhere, demonstrating considerable variation in chemotherapy dosing of the elderly and the obese.^{5,8,16}

Unfortunately, the problem identified here may be but the tip of the iceberg as it does not address those patients with limited access to oncology care, those who may not see or are not treated by an oncologist, and those who do not comply with treatment. Some authors have argued that chemotherapy dose intensity delivered in the curative cancer setting should be considered a quality-of-care measure.^{1,17} There does appear to be sufficient uncertainty on the part of oncologists on how best to dose elderly and obese patients with cancer that the American Society of Clinical Oncology might consider either a policy statement or the development of a clinical practice guideline providing appropriate recommendations for chemotherapy dosing based on a comprehensive systematic review of the available evidence. While more research is clearly needed in this important area, it is essential that the science that we do have is applied appropriately and effectively in the optimal treatment of patients with cancer.

Gary H. Lyman, MD, MPH, is Professor of Medicine and Director, Health Services and Outcomes Research in Oncology and Senior Fellow, Center for Clinical Health Policy Research at Duke University in Durham, North Carolina. Dr Lyman also serves as Associate Editor on the JOP Editorial Board.

DOI: 10.1200/JOP.0831502

References

1. Lyman GH: Chemotherapy dose intensity and quality cancer care. *Oncology* 20:16-25, 2006 (suppl 9)
2. Bonnetterre J, Roche H, Kerbrat P, et al: Epirubicin increases long-term survival in adjuvant chemotherapy of patients with poor-prognosis, node-positive, early breast cancer: 10-year follow-up results of the French Adjuvant Study Group 05 randomized trial. *J Clin Oncol* 20 23:2686-2693, 2005

3. Budman DR, Berry DA, Cirincione CT, et al: Dose and dose intensity as determinants of outcome in the adjuvant treatment of breast cancer: The Cancer and Leukemia Group B. *J Natl Cancer Inst* 90:1205-1211, 1998
4. Lepage E, Gisselbrecht C, Haioun C, et al: Prognostic significance of received relative dose intensity in non-Hodgkin's lymphoma patients: Application to

LNH-87 protocol—The GELA (Groupe d'Etude des Lymphomes de l'Adulte). *Ann Oncol* 4:651-656, 1993

5. Lyman GH, Dale DC, Crawford J: Incidence and predictors of low dose-intensity in adjuvant breast cancer chemotherapy: A nationwide study of community practices. *J Clin Oncol* 21:4524-4531, 2003

6. Lyman GH, Dale DC, Friedberg J, et al: Incidence and predictors of low chemotherapy dose-intensity in aggressive non-Hodgkin's lymphoma: A nationwide study. *J Clin Oncol* 22:4302-4311, 2004

7. Balducci L, Hardy CL, Lyman GH: Hemopoiesis and aging. *Cancer Treat Res* 124:109-134, 2005

8. Shayne M, Culakova E, Poniewierski MS, et al: Dose intensity and hematologic toxicity in older cancer patients receiving systemic chemotherapy. *Cancer* 110:1611-1620, 2007

9. Smith TJ, Khatcheressian J, Lyman GH, et al: 2006 update of recommendations for the use of white blood cell growth factors: An evidence-based clinical practice guideline. *J Clin Oncol* 24:3187-3205, 2006

10. Kuderer NM, Dale DC, Crawford J, et al: Impact of primary prophylaxis with granulocyte colony-stimulating factor on febrile neutropenia and mortality in adult cancer patients receiving chemotherapy: A systematic review. *J Clin Oncol* 25:3158-3167, 2007

11. Rosner GL, Hargis JB, Hollis DR, et al: Relationship between toxicity and obesity in women receiving adjuvant chemotherapy for breast cancer: Results from cancer and leukemia group B study 8541. *J Clin Oncol* 14:3000-3008, 1996

12. Griggs JJ, Culakova E, Sorbero ME, et al: Effect of patient socioeconomic status and body mass index on the quality of breast cancer adjuvant chemotherapy. *J Clin Oncol* 25:277-284, 2007

13. Griggs JJ, Sorbero ME, Lyman GH: Undertreatment of obese women receiving breast cancer chemotherapy. *Arch Intern Med* 165:1267-1273, 2005

14. Colleoni M, Li S, Gelber RD, et al: Relation between chemotherapy dose, oestrogen receptor expression, and body-mass index. *Lancet* 366:1108-1110, 2005

15. Sparreboom A, Wolff AC, Mathijssen RH, et al: Evaluation of alternate size descriptors for dose calculation of anticancer drugs in the obese. *J Clin Oncol* 25:4707-4713, 2007

16. Field KM, Kosmider S, Jefford M, et al: Chemotherapy dosing strategies in the obese, elderly, and thin patients: Results of a nationwide survey. *J Oncol Pract* 4:111-116, 2008

17. Malin JL, Schneider EC, Epstein AM, et al: Results of the National Initiative for Cancer Care Quality: How can we improve the quality of cancer care in the United States? *J Clin Oncol* 24:626-634, 2006

HOUSING AND REGISTRATION OPEN FOR THE 2008 ASCO ANNUAL MEETING

The ASCO Annual Meeting brings together oncologists, researchers, nurses, and patient advocates in the fight against cancer. Register by April 25, 2008, to take advantage of reduced rates. When registering, remember to save time by adding Virtual Meeting and podcast subscriptions to your package. For more details, visit www.asco.org/annualmeeting.