Is weight loss the optimal target for obesity-related cardiovascular disease risk reduction?

Robert Ross PhD^{1,2}, Peter M Janiszewski MSc¹

R Ross, PM Janiszewski. Is weight loss the optimal target for obesity-related cardiovascular disease risk reduction? *Can J Cardiol* 2008;24(Suppl D):25D-31D.

In the present review, it is argued that while weight loss is associated with substantial reduction in obesity-related cardiovascular disease risk and remains a desired outcome of relevant treatment strategies, increasing physical activity is associated with marked reduction in waist circumference, visceral fat and cardiometabolic risk factors, concurrent with an increase in cardiorespiratory fitness despite minimal or no change in body weight. Failure to recognize the benefits of exercise independent of weight loss masks opportunities to counsel and educate patients whose sole criteria for gauging obesity reduction success is the bathroom scale.

Key Words: *Cardiorespiratory fitness; Exercise; Physical activity; Visceral fat; Waist circumference*

Obesity and its attendant diseases represent a threat to public health and a major challenge to the allied health care profession. Clinicians remain a cornerstone of health care delivery in North America and, given that one in two Canadians are overweight (1), are often faced with the burden of educating overweight and obese persons on the associated health risks of obesity and appropriate strategies for its reduction. Counselling begins with accurate identification of those at risk, which is traditionally determined using the body mass index (BMI). BMI remains a well-established marker of health risk, and race-specific cutpoints for determination of overweight and obese individuals have been established (2). Given the strong association between BMI and cardiovascular disease (CVD), it follows that strategies designed to reduce obesity and CVD would identify weight loss as a primary end point for treatment success. It is now apparent, however, that obesity is a heterogeneous condition and that excess accumulation of abdominal fat presents the greatest health risk (3). Waist circumference (WC) is a simple marker of abdominal obesity, and is a strong predictor of morbidity and mortality independent of BMI (4-6). Although the causal link between WC and CVD is not firmly established, mounting evidence suggests that intra-abdominal or visceral fat is the depot that best explains the association between abdominal obesity and CVD (7). These observations suggest that both WC and visceral fat should be primary targets for strategies designed to reduce obesity-related CVD.

Consistent with the argument that weight loss is the primary end point for obesity reduction, leading health authorities suggest that the utility of exercise as a strategy for obesity reduction depends in large measure on body weight reduction (8). Although exercise-induced weight loss is associated with attenuated CVD risk, several lines of evidence suggest that weight loss is not absolutely necessary to observe substantial benefit. First, it is well established that increasing physical

La perte de poids est-elle l'objectif optimal de réduction du risque de maladie cardiovasculaire reliée à l'obésité ?

Dans la présente analyse, on postule que même si la perte de poids s'associe à une importante réduction du risque de maladie cardiovasculaire reliée à l'obésité et qu'elle demeure une issue souhaitée des stratégies de traitement pertinentes, l'accroissement de l'activité physique s'associe à une diminution marquée du tour de taille, de la graisse viscérale et des facteurs de risque cardiométaboliques, conjointement avec une augmentation de la capacité aérobie, malgré une perte de poids minime ou même l'absence de perte de poids. Le fait de ne pas reconnaître les bienfaits de l'exercice qui n'entraîne pas de perte de poids enlève la possibilité de conseiller et d'éduquer les patients qui se fient seulement au pèse-personne pour évaluer la réussite de leur réduction d'obésité.

activity and associated improvements in cardiorespiratory fitness (CRF) are associated with profound reductions in coronary artery disease and CVD mortality, independent of weight or BMI (9,10). Second, acute and chronic exercise is associated with substantial reductions in several cardiometabolic risk factors despite minimal or no change in body weight (11-13). Third, while WC and visceral fat are reduced consequent to weight loss (14-16), it is also true that both variables are substantively reduced in response to exercise with minimal or no weight loss (14,15). These observations bring one to question whether weight loss is the optimal outcome for determining the efficacy of treatment strategies designed to reduce obesity and related comorbidities.

WC, VISCERAL OBESITY AND CARDIOMETABOLIC RISK

Since the seminal work of Jean Vague (17) more than 50 years ago, numerous investigators have confirmed that abdominal obesity, as measured by WC, conveys a substantial health risk, and that WC explains morbidity and mortality risk beyond that explained by BMI (18). The recently published findings of the International Day for Evaluation of Abdominal Obesity (IDEA) (19) provide compelling evidence that WC explains both diabetes and CVD risk beyond that explained by BMI alone. The IDEA study involved 6407 randomly chosen primary care physicians in 63 countries, who evaluated 168,159 patients aged 18 to 80 years. The authors observed a graded increase in the frequency of CVD and diabetes mellitus with both BMI and WC; there was a stronger relationship for WC than for BMI across regions for both sexes. Extending these observations is the recent report from Janiszewski et al (5) demonstrating that WC predicts diabetes risk beyond BMI and other commonly measured

¹School of Kinesiology and Health Studies, ²Department of Medicine, Division of Endocrinology and Metabolism, Queen's University, Kingston, Ontario
Correspondence: Dr Robert Ross, School of Kinesiology and Health Studies, Queen's University, 155 North Street, Kingston, Ontario K7L 3N6.

Telephone 613-533-6583, fax 613-533-2009, e-mail rossr@queensu.ca
Received for publication March 24, 2008. Accepted March 26, 2008

TABLE 1

Changes in body composition, fitness and metabolic parameters in exercise groups with nonsignificant change in weight from randomized and nonrandomized trials (1997–2007)

Reference	Experimental design	Subjects, description	Exercise time, type	Study length (weeks)	Weight Δ (kg)	Waist circumference Δ (cm)	ASAT Δ (%)	VAT Δ (%)	VO ₂ max Δ (mL/kg/min)	Metabolic improvement
Mourier et al, 1997 (31)	RCT	11 T2D (45 y)	112 min/wk, AE	8	−1.5	−1.0	−17.9*	−48.5*	+9.3*	✓ IS
Owens et al, 1999 (73)	RCT	35 M/F (10 y)	160 min/wk, AE	16	+1.1		−1.0	+0.5*		✓ TG†
Wilmore et al, 1999 (74)	Non-RCT	250 M (17–65 y)	150 min/wk, AE	20	−2.4	−1.0	−4.8	−7.1*		✓ HDL, TG†
Wilmore et al, 1999 (74)	Non-RCT	252 F (17–65 y)	150 min/wk, AE	20	−0.1	−0.9	−2.9	−4.6*		✓ HDL
Boudou et al, 2000 (30)	RCT	8 T2D M (45 y)	115 min/wk, AE	8	−1.9		−18.1*	−44.7*	+9.4*	✓ TG, IS
Eliakim et al, 2000‡ (75)	RCT	22 M (16 y)	540 min/wk, AE	5	+0.8		−7.6*	−0.2		✓ HDL
Poehlman et al, 2000‡ (76)	RCT	14 F (29 y)	99 min/wk, AE	24	0.0		−0.5	+2.5	+10.2*	✓ IS
Poehlman et al, 2000‡ (76)	RCT	17 F (28 y)	81 set/wk, RE	24	+2.0		0	0	+1.1	
Ross et al, 2000 (14)	RCT	14 M (45 y)	443 min/wk, AE	12	−0.5	−1.8		−11.8*	+7.4*	
Thomas et al, 2000 (32)	Non-RCT	17 F (25–45 y)	120 min/wk, AE	24	−0.6		−6.6	−23.5*	+3.9*	✓ HDL
Gan et al, 2003 (53)	Non-RCT	18 M (37 y)	164 min/wk, AE	9.7	−1.3	−2.0*	−4.6*	−5.4	+5.2*	✓ Chol, TG, IS
Short et al, 2003‡ (77)	RCT	65 M/F (22–87 y)	160 min/wk, AE	16	−0.8	−4.1	−6.5	−6.0	+2.7	✓ TG, IS
Green et al, 2004 (78)	Non-RCT	30 F (56 y)	150 min/wk, AE	20	+0.1	−0.9	−1.2	+3.1	+4.3*	
Green et al, 2004 (78)	Non-RCT	18 F (52 y)	150 min/wk, AE	20	+0.5	−0.1	+1.6	−6.7	+4.8*	✓ LDL
Ross et al, 2004 (15)	RCT	12 F (41 y)	441 min/wk, AE	14	−0.5	−3.1*	−4.9*	−18.2*	+5.9*	
Giannopoulou et al, 2005 (29)	Non-RCT	11 T2D F (56 y)	150 min/wk, AE	14	−1.7	−3.0*	−5.3*	−10.2*	+3.2*	✓ IS
Ibanez et al, 2005 (79)	Non-RCT	9 T2D M (67 y)	105 min/wk, RE	16	−0.5		−13.3*	−9.6*		✓ F Glu, IS
Lee et al, 2005 (12)	Non-RCT	8 M (48 y)	300 min/wk, AE	13	−0.4	−2.7*		−10.0*	+10.5*	
Lee et al, 2005 (12)	Non-RCT	8 M (47 y)	300 min/wk, AE	13	−0.4	−2.6*		−15.0*	+6.6*	
Lee et al, 2005 (12)	Non-RCT	7 T2D M (51 y)	300 min/wk, AE	13	−0.8	−2.7*		−23.7*	+11.3*	
O'Donovan et al, 2005 (80)	RCT	14 M (41 y)		24	−1.1	−1.0			+4.9*	
O'Donovan et al, 2005 (80)	RCT	13 M (41 y)		24	−0.5	−2.0*			+7.1*	✓ Chol, LDL
Dolan et al, 2006 (81)	Non-RCT	20 HIV F (43 y)	90 min/wk, AE	16	+0.4§	−1.0	+0.8	−1.4	+1.5*	
Church et al, 2007 (82)	RCT	155 F (58 y)	72 min/wk, AE	24	−0.4	−1.9*			+0.7*¶	
Church et al, 2007 (82)	RCT	104 F (57 y)	136 min/wk, AE	24	−2.2	−2.9*			+1.5*¶	
Church et al, 2007 (82)	RCT	103 F (57 y)	191 min/wk, AE	24	−0.6	−1.4*			+0.8*¶	
Ramvalho et al, 2006 (83)	Non-RCT	7 T1D M/F (13–30 y)	120 min/wk, AE	12	+0.05§	−1.9*			+2.0	✓ HbA1c
Ramvalho et al, 2006 (83)	Non-RCT	6 T1D M/F (13–30 y)	120 min/wk, RE	12	+0.0§	+0.9			+5.1	
Shojaee-Moradie et al, 2007 (33)	RCT	10 M (50 y)	60 min/wk, AE	6	+0.2		−3.0	−18.0	+7.0*	

Age is presented as mean or range. *Significant improvement ($P < 0.05$); †Metabolic variables from the Wilmore et al (74) are reported in reference 52; ‡Only within-group difference reported; §Change in body mass index (kg/m^2); ¶Peak maximal oxygen consumption ($\text{VO}_2 \text{ max}$). Δ Change in; AE Aerobic exercise; ASAT Abdominal subcutaneous adipose tissue; Chol Total cholesterol; F Glu Fasting glucose; F Female; HbA1c Glycosylated hemoglobin; HDL High-density lipoprotein; IS Insulin sensitivity; LDL Low-density lipoprotein; M Male; RCT Randomized controlled trial; RE Resistance exercise; T1D Type 1 diabetes; T2D Type 2 diabetes; TG Triglycerides; VAT Visceral adipose tissue; wk Week; y Year

cardiometabolic risk factors, such as smoking, dyslipidemia and blood pressure. Together, these observations underscore the importance of WC as a routine measure in clinical practice to help to identify the high-risk, abdominally obese patient. Accordingly, these observations support the notion that WC should be a primary treatment target for strategies designed to reduce obesity-related CVD risk.

Overwhelming evidence suggests that, of the two abdominal fat depots (subcutaneous and visceral), visceral fat is the depot that best explains the association between abdominal obesity and CVD risk (please see Després et al in the present issue, pages 7D-12D). Indeed, independent of subcutaneous fat, visceral fat is a strong predictor of dyslipidemia (20,21), glucose tolerance (22,23), insulin resistance (24) and systemic inflammation (25), as well as incidence of hypertension (26),

CVD (7), type 2 diabetes (27) and all-cause mortality (28). These observations underscore the notion that both WC and visceral fat should be primary targets for strategies designed to reduce obesity and related CVD risk. While it is clear that diet- or exercise-induced weight loss is associated with substantial reductions in both WC and visceral fat (14-16), increasing evidence supports the view that both WC and visceral fat are substantially reduced in response to exercise with minimal or no weight loss (14,15,29).

Exercise-induced reduction in WC with minimal or no weight loss

Table 1 reveals that short-term, regular, physical activity is associated with a marked reduction in WC, independent of sex or age, despite no statistically significant change in body mass. In these

Figure 1 Relationship between changes in body weight and waist circumference (WC). **A** Data are taken from prior publications (14,15,71,72) and illustrate the relationship between change in weight (kg) and change in WC (cm) in a sample of 105 men and women combined. **B** Based on the same subjects in Panel A, the data were separated into tertiles of weight reduction (<6.1 kg, 6.2 kg to 9.8 kg, and 9.9 kg to 20.7 kg). Each tertile of weight and WC reduction was significantly different from the other two tertiles ($P < 0.05$)

studies, the negative energy balance was induced by the increase in physical activity, because all participants were instructed not to reduce caloric intake. It is also noteworthy that in the majority of studies, the reduction in WC without significant weight loss was associated with significant reductions in cardiometabolic risk factors. These are encouraging findings and suggest that reliance on BMI alone to determine the utility of strategies to reduce obesity and related cardiometabolic risk factors are misguided.

The lack of association between changes in body weight and WC in response to increases in physical activity are underscored in Figure 1. The data presented clearly show that change in body weight is a poor indicator of corresponding changes in WC on an individual basis, and perhaps more importantly, that the lack of association is greatest for the lowest tertile of change in weight and WC. This is not a trivial observation, because resistance to apparent weight loss often occurs during the early stage of initiating a program of obesity reduction. In other words, during the early phase of attempts to change behaviour (increase physical activity combined with a healthful diet) to reduce obesity, negligible changes in body weight may be discouraging to the participant to the point that they discontinue attempts to change behaviour. On the other hand, recognition that changes in WC are occurring in the absence of weight change would be extremely positive and empowering for the participant, and this presents the clinician with an excellent opportunity to acknowledge benefit.

Exercise-induced reduction in visceral fat with minimal or no weight loss

Consistent with the observation that exercise is associated with WC reduction despite minimal weight loss, it is also apparent that regular physical activity is associated with a marked reduction in visceral fat despite little or no change in weight (Table 1). Several studies have examined the effect of physical activity on visceral fat independent of weight loss by having study participants consume compensatory kilocalories equivalent to the amount expended during physical activity (12,14,15). The results suggest that in Caucasian lean and obese men and obese women, significant reductions in visceral fat (–10% to –19%) occur consequent to three months of regular physical activity (40 min/day to 60 min/day) despite no significant change in body

weight. These observations are consistent with findings from studies in type 2 diabetic subjects suggesting that two to three months of moderate intensity aerobic exercise is associated with substantive reductions in visceral fat (–27% to –45%) despite little or no change in weight (12,30,31). Similar results have been documented in healthy, nonobese, premenopausal women (32) and middle-aged men (33).

These observations reinforce the long-standing notion that changes in body composition, rather than body weight, are more meaningful outcomes in response to physical activity (34). In fact, significant reductions in fat mass, especially visceral fat, often occur concurrent with equal increases in lean body mass in response to physical activity (15). These equal but opposite changes in fat mass and lean mass are not detected by alterations in body weight (15) but may be interpreted from changes in WC. Indeed, as illustrated by the data presented in Figure 2, while increases in lean mass offset corresponding changes in total fat or body weight, substantial reductions in WC are observed. Combined with the increase in CRF and insulin sensitivity (please see below), the reductions in WC offer the clinician a valuable opportunity to counsel the patient with respect to the benefits of exercise-induced reductions in abdominal obesity despite no change in scale weight.

It is important to emphasize, however, that the reductions in WC and visceral fat observed in exercisers who maintain body weight are substantially less than those observed in exercisers who lose weight (14,15). Thus, from a clinical perspective, those at obesity-related CVD risk should be advised that exercise-induced weight loss is likely associated with the greatest benefit. On the other hand, given the challenges associated with attaining substantial weight loss for many obese individuals, in particular, older adults – because of ingrained, lifelong habits – it is equally important to recognize that abdominal obesity and related health risk can be markedly reduced with minimal weight loss.

Exercise-induced reduction in cardiometabolic risk factors without weight loss

It is important to recall that significant improvements in a number of cardiometabolic risk factors are known to occur after only a single session of moderate-intensity aerobic exercise (11). For instance, insulin

Figure 2) Disassociation between changes in body weight, body composition and health risk. Individual body composition change data are shown for a sample of 20 obese men who participated in studies designed to reduce obesity by increasing exercise without changing caloric intake (14). AT Adipose tissue; CRF Cardiorespiratory fitness; LT Lean tissue; WC Waist circumference; WL Weight loss

resistance is known to improve by approximately 20% after 1 h of aerobic exercise in healthy (35,36), insulin-resistant (37) and diabetic subjects (36,38) – improvements that are equivalent in magnitude to those achieved through chronic pharmacological intervention (39,40). The augmented insulin sensitivity is apparent immediately after exercise (36) and persists for 20 h to 48 h thereafter (35,37,38). It is also well established that a single bout of aerobic exercise results in 10% to 25% reductions in triglycerides (TGs) and 7% to 15% increases in high-density lipoprotein (HDL) cholesterol levels (11,41-43). These lipid changes become apparent between 24 h and 48 h after exercise in both untrained (41,42) and trained subjects (43) in response to caloric expenditures of 350 kcal to 500 kcal and 1000 kcal, respectively. Additionally, the blood pressure-lowering effect of acute aerobic exercise has been long established (44). This effect, termed postexercise hypotension, is especially pronounced among hypertensive individuals with maximal reductions in systolic and diastolic blood pressure of -11 mmHg and -6 mmHg, respectively (45,46). The onset of postexercise hypotension is immediate and can persist for up to 22 h after an acute bout of aerobic physical activity (47). Indeed, as little as 15 min of low-intensity (40% maximal oxygen consumption) aerobic exercise can induce marked postexercise hypotension that persists throughout the day (48). In unison, these encouraging results suggest that independent of any training effects, repeated bouts of aerobic exercise may be a viable treatment strategy for the treatment of insulin resistance, as well as dyslipidemia and hypertension.

Chronic exercise training, or the summation of acute bouts of exercise, can lead to further improvements in cardiometabolic risk factors, even in the absence of clinically significant weight loss. Indeed, numerous studies have shown that improvements in insulin sensitivity through chronic exercise can occur in the absence of significant weight reduction (13,14,29,30). For example, three months of daily aerobic training in obese men, who consumed compensatory kilocalories equivalent to the amount expended during exercise, resulted in a 30% improvement in insulin sensitivity despite no change in weight (14). Furthermore, while improvements in HDL cholesterol and TGs are generally greater in those who lose weight, these improvements

can be seen even when weight remains virtually unchanged (49-51). For example, a number of physical activity interventions that do not significantly alter body weight have documented 5% to 35% reductions in TG levels (52,53) and/or 3% to 5% increases in HDL values (32,52). Lastly, the blood pressure reductions induced through chronic exercise do not appear to be related to alterations in body weight or abdominal obesity (54). In fact, a large meta-analysis (54), which reported an overall beneficial effect of chronic exercise on blood pressure, found that the average weight loss across the reviewed trials was largely inconsequential (-1.2 kg). Nevertheless, some studies report that individuals who become active and also lose a significant amount of weight (-7.9 kg), in comparison with those who lose minimal weight (-1.8 kg), show greater reductions in systolic and diastolic blood pressure (-7 mmHg and -5 mmHg versus -4 mmHg and -4 mmHg, respectively) (55).

The improvement in insulin resistance through chronic physical activity independent of weight change likely involves changes in body composition – in particular, reductions in visceral fat (14), as well as augmented expression of GLUT-4 protein in skeletal muscle (56,57), and enhanced metabolic efficiency of muscle (58,59). The lipid improvements independent of significant weight reduction may also be mediated by improvements in body composition, such as reductions in visceral fat (60), but apparently are not due to improvements in CRF (51). Lastly, it appears that physical activity and weight loss may have independent effects on blood pressure, because reductions in blood pressure are significantly related to training-induced improvements in CRF (54), as well as the magnitude of weight loss (55).

Consistent with the effects of chronic exercise described above, prospective studies with long-term follow-up have demonstrated an inverse association between physical activity and fitness on the one hand, and morbidity and mortality on the other, independent of obesity status. For example, Helmrigh et al (61) have previously shown, in a 14-year prospective study of approximately 6000 men, that a high level of physical activity is associated with the lowest risk of diabetes irrespective of the level of obesity. Specifically, it was reported that for every 500 kcal/week increase in physical activity, the risk of diabetes

decreased by 6% (61). Results of other prospective studies in men and women further suggest that high levels of physical activity protect against the incidence of coronary artery disease, independent of BMI, as well as other factors such as age, smoking, diabetes and hypertension (62). Furthermore, prospective analyses from the Aerobics Center Longitudinal Study have consistently reported that CRF is a significant predictor of mortality from all causes (9,10,63,64), CVD (9,10) and type 2 diabetes (65) in men and women, independent of BMI or WC. These relationships have been confirmed in older adults (66) and Canadian women (67).

Beyond weight loss as the only indicator of obesity treatment success

From the observations described in the present overview, it is apparent that the time has come to look beyond weight loss as the only criteria for successful obesity treatment. Indeed, decreases in WC combined with increases in skeletal muscle mass, improvements in cardiometabolic risk factors and CRF consequent to exercise, with little or no weight loss, underscore the recommendation that clinicians include WC as a routine measure in clinical practice and encourage their patients to monitor WC in addition to scale weight to gauge treatment success (Figure 3). However, as reviewed here, it is also clear that increasing physical activity is associated with marked attenuation in CVD risk factors, at least partly mediated by increased CRF independent of BMI or WC. Furthermore, it is reported that CRF is associated with a substantial reduction in cardiometabolic risk factors independent of visceral fat (68). These findings support the recommendation that CRF be assessed by clinicians as an objective measure of changes in physical activity level and associated reductions in CVD risk. Although the measurement of CRF in clinical practice may be difficult for pragmatic reasons, referral of patients to properly trained Certified Exercise Physiologists (69) working at commercial and/or public health facilities offers a reasonable alternative to practitioners.

SUMMARY AND RECOMMENDATIONS

In the present brief report, we propose the notion that, while weight loss is associated with marked reduction in obesity-related CVD risk and remains a desired outcome of strategies designed to reduce obesity-related CVD, increasing physical activity is often associated with a decrease in WC and cardiometabolic risk, as well as an increase in CRF, despite little or no change in body weight or BMI. These observations support the recommendation that clinicians look beyond weight loss as the only indicator of treatment success when targeting obesity-related CVD risk. Indeed, as illustrated in Figure 3, we propose three distinct scenarios for which an increase in physical activity is associated with substantial reduction in obesity-related CVD risk with or without weight loss. That all three scenarios described are associated with attenuated CVD risk is encouraging, and offers both the clinician and the patient treatment options. Indeed, recognition of the distinct benefits of exercise, combined with the consumption of a

Figure 3) Favourable effects of physical activity on obesity-related health risk: three possible scenarios. The scenarios presented represent three possible outcomes when increasing physical activity combined with a balanced (healthful) diet for the purpose of reducing obesity and related comorbid conditions. BMI Body mass index; CRF Cardiorespiratory fitness; WC Waist circumference

balanced diet, on CVD risk factors offers the clinician several unique opportunities to counsel and educate patients on the benefits of adopting healthy behaviours for the purpose of reducing obesity and related health risks despite possible resistance to weight loss. In this way, it is recommended that WC should be a routine measure in clinical practice for identifying the high-risk, abdominally obese patient, and that patients should be educated on the proper methods for self-measurement of WC (70). Finally, overwhelming evidence supports the recommendation that CRF be a routine measure in clinical practice. Although this may be impractical for many, if not most, clinicians, referral to appropriately trained personnel working at commercial or public health centres is a realistic option for both the clinician and the participant. Obesity is a multidimensional problem that requires a multidimensional response. Close collaboration among practitioners at all levels of health care is a small step toward the resolve of a major public health problem

DISCLOSURE: Dr Robert Ross is a speaker for GlaxoSmithKline and sanofi-aventis. He has received research funding from MARS Inc, Roche Pharmaceutical and sanofi-aventis; is on the advisory board for sanofi-aventis; has received consulting fees from GlaxoSmithKline, Roche Pharmaceutical, sanofi-aventis and Theratechnologies Inc; and receives royalties from Human Kinetics. Peter M Janiszewski is supported by a Doctoral Research Award from the Institutes of Health Research.

REFERENCES

- Katzmarzyk PT. The Canadian obesity epidemic: An historical perspective. *Obes Res* 2002;10:666-74.
- Clinical Guidelines on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults – The Evidence Report. National Institutes of Health. *Obes Res* 1998;6(Suppl 2):51S-209S. (Erratum in 1998;6:464).
- Despres JP, Lemieux I, Prud'homme D. Treatment of obesity: Need to focus on high risk abdominally obese patients. *BMJ* 2001;322:716-20.
- Yusuf S, Hawken S, Ounpuu S, et al. Obesity and the risk of myocardial infarction in 27,000 participants from 52 countries: A case-control study. *Lancet* 2005;366:1640-9.
- Janiszewski PM, Janssen I, Ross R. Does waist circumference predict diabetes and cardiovascular disease beyond commonly evaluated cardiometabolic risk factors? *Diabetes Care* 2007;30:3105-9.
- Bigaard J, Tjonneland A, Thomsen BL, Overvad K, Heitmann BL, Sorensen TI. Waist circumference, BMI, smoking, and mortality in middle-aged men and women. *Obes Res* 2003;11:895-903.
- Fujimoto WY, Bergstrom RW, Boyko EJ, et al. Visceral adiposity and incident coronary heart disease in Japanese-American men. The 10-year follow-up results of the Seattle Japanese-American Community Diabetes Study. *Diabetes Care* 1999;22:1808-12.
- Lau DC, Douketis JD, Morrison KM, Hramiak IM, Sharma AM, Ur E; Obesity Canada Clinical Practice Guidelines Expert Panel. 2006 Canadian clinical practice guidelines on the management and prevention of obesity in adults and children [summary]. *CMAJ* 2007;176:S1-13.
- Wei M, Kampert JB, Barlow CE, et al. Relationship between low cardiorespiratory fitness and mortality in normal-weight, overweight, and obese men. *JAMA* 1999;282:1547-53.

10. Lee CD, Blair SN, Jackson AS. Cardiorespiratory fitness, body composition, and all-cause and cardiovascular disease mortality in men. *Am J Clin Nutr* 1999;69:373-80.
11. Thompson PD, Crouse SF, Goodpaster B, Kelley D, Moyna N, Pescatello L. The acute versus the chronic response to exercise. *Med Sci Sports Exerc* 2001;33:S438-45.
12. Lee S, Kuk JL, Davidson LE, et al. Exercise without weight loss is an effective strategy for obesity reduction in obese individuals with and without Type 2 diabetes. *J Appl Physiol* 2005;99:1220-5.
13. Dengel DR, Pratley RE, Hagberg JM, Rogus EM, Goldberg AP. Distinct effects of aerobic exercise training and weight loss on glucose homeostasis in obese sedentary men. *J Appl Physiol* 1996;81:318-25.
14. Ross R, Dagnone D, Jones PJ, et al. Reduction in obesity and related comorbid conditions after diet-induced weight loss or exercise-induced weight loss in men. A randomized, controlled trial. *Ann Intern Med* 2000;133:92-103.
15. Ross R, Janssen I, Dawson J, et al. Exercise-induced reduction in obesity and insulin resistance in women: A randomized controlled trial. *Obes Res* 2004;12:789-98.
16. Slentz CA, Duscha BD, Johnson JL, et al. Effects of the amount of exercise on body weight, body composition, and measures of central obesity: STRRIDE – a randomized controlled study. *Arch Intern Med* 2004;164:31-9.
17. Vague P. The degree of masculine differentiation of obesity: A factor determining predisposition to diabetes, atherosclerosis, gout, and uric calculous disease. *Am J Clin Nutr* 1956;4:20-34.
18. Despres J-P. Abdominal obesity: The most prevalent cause of the metabolic syndrome and related cardiometabolic risk. *Eur Heart J Suppl* 2006;8:B4-12.
19. Balkau B, Deanfield JE, Despres JP, et al. International Day for the Evaluation of Abdominal Obesity (IDEA): A study of waist circumference, cardiovascular disease, and diabetes mellitus in 168,000 primary care patients in 63 countries. *Circulation* 2007;116:1942-51.
20. Nguyen-Duy TB, Nichaman MZ, Church TS, Blair SN, Ross R. Visceral fat and liver fat are independent predictors of metabolic risk factors in men. *Am J Physiol Endocrinol Metab* 2003;284:E1065-71.
21. Nieves DJ, Cnop M, Retzlaff B, et al. The atherogenic lipoprotein profile associated with obesity and insulin resistance is largely attributable to intra-abdominal fat. *Diabetes* 2003;52:172-9.
22. Brochu M, Starling RD, Tchernof A, Matthews DE, Garcia-Rubi E, Poehlman ET. Visceral adipose tissue is an independent correlate of glucose disposal in older obese postmenopausal women. *J Clin Endocrinol Metab* 2000;85:2378-84.
23. Janssen I, Fortier A, Hudson R, Ross R. Effects of an energy-restrictive diet with or without exercise on abdominal fat, intermuscular fat, and metabolic risk factors in obese women. *Diabetes Care* 2002;25:431-8.
24. Ross R, Aru J, Freeman J, Hudson R, Janssen I. Abdominal adiposity and insulin resistance in obese men. *Am J Physiol Endocrinol Metab* 2002;282:E657-63.
25. Forouhi NG, Sattar N, McKeigue PM. Relation of C-reactive protein to body fat distribution and features of the metabolic syndrome in Europeans and South Asians. *Int J Obes Relat Metab Disord* 2001;25:1327-31.
26. Hayashi T, Boyko EJ, Leonetti DL, et al. Visceral adiposity is an independent predictor of incident hypertension in Japanese Americans. *Ann Intern Med* 2004;140:992-1000.
27. Boyko EJ, Fujimoto WY, Leonetti DL, Newell-Morris L. Visceral adiposity and risk of type 2 diabetes: A prospective study among Japanese Americans. *Diabetes Care* 2000;23:465-71.
28. Kuk JL, Katzmarzyk PT, Nichaman MZ, Church TS, Blair SN, Ross R. Visceral fat is an independent predictor of all-cause mortality in men. *Obesity (Silver Spring)*. 2006;14:336-41.
29. Giannopoulou I, Ploutz-Snyder LL, Carhart R, et al. Exercise is required for visceral fat loss in postmenopausal women with type 2 diabetes. *J Clin Endocrinol Metab* 2005;90:1511-8.
30. Boudou P, De Kerviler E, Vexiau P, Fiet J, Cathelineau G, Gautier J. Effects of a single bout of exercise and exercise training on steroid levels in middle-aged type 2 diabetic men: Relationship to abdominal adipose tissue distribution and metabolic status. *Diabetes Metab* 2000;26:450-7.
31. Mourier A, Gautier JF, De Kerviler E, et al. Mobilization of visceral adipose tissue related to the improvement in insulin sensitivity in response to physical training in NIDDM. Effects of branched-chain amino acid supplements. *Diabetes Care* 1997;20:385-91.
32. Thomas EL, Brynes AE, McCarthy J, et al. Preferential loss of visceral fat following aerobic exercise, measured by magnetic resonance imaging. *Lipids* 2000;35:769-76.
33. Shojaae-Moradie F, Baynes KC, Pentecost C, et al. Exercise training reduces fatty acid availability and improves the insulin sensitivity of glucose metabolism. *Diabetologia* 2007;50:404-13.
34. Janiszewski PM, Ross R. Physical activity in the treatment of obesity: Beyond body weight reduction. *Appl Physiol Nutr Metab* 2007;32:512-22.
35. Mikines KJ, Sonne B, Farrell PA, Tronier B, Galbo H. Effect of physical exercise on sensitivity and responsiveness to insulin in humans. *Am J Physiol* 1988;254:E248-59.
36. Burstein R, Epstein Y, Shapiro Y, Charuzi I, Karnieli E. Effect of an acute bout of exercise on glucose disposal in human obesity. *J Appl Physiol* 1990;69:299-304.
37. Perseghin G, Price TB, Petersen KF, et al. Increased glucose transport-phosphorylation and muscle glycogen synthesis after exercise training in insulin-resistant subjects. *N Engl J Med* 1996;335:1357-62.
38. Devlin JT, Hirshman M, Horton ED, Horton ES. Enhanced peripheral and splanchnic insulin sensitivity in NIDDM men after single bout of exercise. *Diabetes* 1987;36:434-9.
39. Widén EL, Eriksson JG, Groop LC. Metformin normalizes nonoxidative glucose metabolism in insulin-resistant normoglycemic first-degree relatives of patients with NIDDM. *Diabetes* 1992;41:354-8.
40. Nolan JJ, Ludvik B, Beersden P, Joyce M, Olefsky J. Improvement in glucose tolerance and insulin resistance in obese subjects treated with troglitazone. *N Engl J Med* 1994;331:1188-93.
41. Crouse SF, O'Brien BC, Rohack JJ, et al. Changes in serum lipids and apolipoproteins after exercise in men with high cholesterol: Influence of intensity. *J Appl Physiol* 1995;79:279-86.
42. Grandjean PW, Crouse SF, Rohack JJ. Influence of cholesterol status on blood lipid and lipoprotein enzyme responses to aerobic exercise. *J Appl Physiol* 2000;89:472-80.
43. Bounds RG, Grandjean PW, O'Brien BC, Inman C, Crouse SF. Diet and short term plasma lipoprotein-lipid changes after exercise in trained men. *Int J Sport Nutr Exerc Metab* 2000;10:114-27.
44. Kenney MJ, Seals DR. Postexercise hypotension. Key features, mechanisms, and clinical significance. *Hypertension* 1993;22:653-64.
45. Rueckert PA, Slane PR, Lillis DL, Hanson P. Hemodynamic patterns and duration of post-dynamic exercise hypotension in hypertensive humans. *Med Sci Sports Exerc* 1996;28:24-32.
46. Pescatello LS, Miller B, Danias PG, et al. Dynamic exercise normalizes resting blood pressure in mildly hypertensive premenopausal women. *Am Heart J* 1999;138:916-21.
47. Brandao Rondon MU, Alves MJ, Braga AM, et al. Postexercise blood pressure reduction in elderly hypertensive patients. *J Am Coll Cardiol* 2002;39:676-82.
48. Guidry MA, Blanchard BE, Thompson PD, et al. The influence of short and long duration on the blood pressure response to an acute bout of dynamic exercise. *Am Heart J* 2006;151:1322, e5-12.
49. Carroll S, Dudfield M. What is the relationship between exercise and metabolic abnormalities? A review of the metabolic syndrome. *Sports Med* 2004;34:371-418.
50. Durstine JL, Grandjean PW, Davis PG, Ferguson MA, Alderson NL, DuBose KD. Blood lipid and lipoprotein adaptations to exercise: A quantitative analysis. *Sports Med* 2001;31:1033-62.
51. Kraus WE, Houmard JA, Duscha BD, et al. Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med* 2002;347:1483-92.
52. Leon AS, Rice T, Mandel S, et al. Blood lipid response to 20 weeks of supervised exercise in a large biracial population: The HERITAGE Family Study. *Metabolism* 2000;49:513-20.
53. Gan SK, Kriketos AD, Ellis BA, Thompson CH, Kraegen EW, Chisholm DJ. Changes in aerobic capacity and visceral fat but not myocyte lipid levels predict increased insulin action after exercise in overweight and obese men. *Diabetes Care* 2003;26:1706-13.
54. Cornelissen VA, Fagard RH. Effects of endurance training on blood pressure, blood pressure-regulating mechanisms, and cardiovascular risk factors. *Hypertension* 2005;46:667-75.
55. Blumenthal JA, Sherwood A, Gullette EC, et al. Exercise and weight loss reduce blood pressure in men and women with mild

- hypertension: Effects on cardiovascular, metabolic, and hemodynamic functioning. *Arch Intern Med* 2000;160:1947-58.
56. Houmard JA, Shinebarger MH, Dolan PL, et al. Exercise training increases GLUT-4 protein concentration in previously sedentary middle-aged men. *Am J Physiol* 1993;264:E896-901.
 57. Hughes VA, Fiatarone MA, Fielding RA, et al. Exercise increases muscle GLUT-4 levels and insulin action in subjects with impaired glucose tolerance. *Am J Physiol* 1993;264:E855-62.
 58. Menshikova EV, Ritov VB, Toledo FG, Ferrell RE, Goodpaster BH, Kelley DE. Effects of weight loss and physical activity on skeletal muscle mitochondrial function in obesity. *Am J Physiol Endocrinol Metab* 2005;288:E818-25.
 59. Toledo FG, Menshikova EV, Ritov VB, et al. Effects of physical activity and weight loss on skeletal muscle mitochondria and relationship with glucose control in type 2 diabetes. *Diabetes* 2007;56:2142-7.
 60. Fujioka S, Matsuzawa Y, Tokunaga K, et al. Improvement of glucose and lipid metabolism associated with selective reduction of intra-abdominal visceral fat in premenopausal women with visceral fat obesity. *Int J Obes* 1991;15:853-9.
 61. Helmrich SP, Ragland DR, Leung RW, Paffenbarger RS Jr. Physical activity and reduced occurrence of non-insulin-dependent diabetes mellitus. *N Engl J Med* 1991;325:147-52.
 62. Haapanen N, Miilunpalo S, Vuori I, Oja P, Pasanen M. Association of leisure time physical activity with the risk of coronary heart disease, hypertension and diabetes in middle-aged men and women. *Int J Epidemiol* 1997;26:739-47.
 63. Farrell SW, Braun L, Barlow CE, Cheng YJ, Blair SN. The relation of body mass index, cardiorespiratory fitness, and all-cause mortality in women. *Obes Res* 2002;10:417-23.
 64. Church TS, Cheng YJ, Earnest CP, et al. Exercise capacity and body composition as predictors of mortality among men with diabetes. *Diabetes Care* 2004;27:83-8.
 65. Sui X, Hooker SP, Lee IM, et al. A prospective study of cardiorespiratory fitness and risk of type 2 diabetes in women. *Diabetes Care* 2008;31:550-5.
 66. Sui X, LaMonte MJ, Laditka JN, et al. Cardiorespiratory fitness and adiposity as mortality predictors in older adults. *JAMA* 2007;298:2507-16.
 67. Katzmarzyk PT, Craig CL. Independent effects of waist circumference and physical activity on all-cause mortality in Canadian women. *Appl Physiol Nutr Metab* 2006;31:271-6.
 68. Lee S, Kuk JL, Katzmarzyk PT, Blair SN, Church TS, Ross R. Cardiorespiratory fitness attenuates metabolic risk independent of abdominal subcutaneous and visceral fat in men. *Diabetes Care* 2005;28:895-901.
 69. Canadian Society for Exercise Physiology. <www.csepmembers.ca/english/search.asp> (Version current at April 15, 2008).
 70. Centre for Obesity Research and Education. Instructional material for professional and self-measurement of waist circumference. <www.corecanada.net> (Version current at April 15, 2008).
 71. Ross R, Pedwell H, Rissanen J. Effects of energy restriction and exercise on skeletal muscle and adipose tissue in women as measured by magnetic resonance imaging. *Am J Clin Nutr* 1995;61:1179-85.
 72. Ross R, Rissanen J, Pedwell H, Clifford J, Shragge P. Influence of diet and exercise on skeletal muscle and visceral adipose tissue in men. *J Appl Physiol* 1996;81:2445-55.
 73. Owens S, Gutin B, Allison J, et al. Effect of physical training on total and visceral fat in obese children. *Med Sci Sports Exerc* 1999;31:143-8.
 74. Wilmore JH, Despres JP, Stanforth PR, et al. Alterations in body weight and composition consequent to 20 wk of endurance training: The HERITAGE Family Study. *Am J Clin Nutr* 1999;70:346-52.
 75. Eliakim A, Makowski GS, Brasel JA, Cooper DM. Adiposity, lipid levels, and brief endurance training in nonobese adolescent males. *Int J Sports Med* 2000;21:332-7.
 76. Poehlman ET, Dvorak RV, DeNino WF, Brochu M, Ades PA. Effects of resistance training and endurance training on insulin sensitivity in nonobese, young women: A controlled randomized trial. *J Clin Endocrinol Metab* 2000;85:2463-8.
 77. Short KR, Vittone JL, Bigelow ML, et al. Impact of aerobic exercise training on age-related changes in insulin sensitivity and muscle oxidative capacity. *Diabetes* 2003;52:1888-96.
 78. Green JS, Stanforth PR, Rankinen T, et al. The effects of exercise training on abdominal visceral fat, body composition, and indicators of the metabolic syndrome in postmenopausal women with and without estrogen replacement therapy: The HERITAGE family study. *Metabolism* 2004;53:1192-6.
 79. Ibanez J, Izquierdo M, Arguelles I, et al. Twice-weekly progressive resistance training decreases abdominal fat and improves insulin sensitivity in older men with type 2 diabetes. *Diabetes Care* 2005;28:662-7.
 80. O'Donovan G, Owen A, Bird SR, et al. Changes in cardiorespiratory fitness and coronary heart disease risk factors following 24 wk of moderate- or high-intensity exercise of equal energy cost. *J Appl Physiol* 2005;98:1619-25.
 81. Dolan SE, Frontera W, Librizzi J, et al. Effects of a supervised home-based aerobic and progressive resistance training regimen in women infected with human immunodeficiency virus: A randomized trial. *Arch Intern Med* 2006;166:1225-31.
 82. Church TS, Earnest CP, Skinner JS, Blair SN. Effects of different doses of physical activity on cardiorespiratory fitness among sedentary, overweight or obese postmenopausal women with elevated blood pressure: A randomized controlled trial. *JAMA* 2007;297:2081-91.
 83. Ramalho AC, de Lourdes Lima M, Nunes F, et al. The effect of resistance versus aerobic training on metabolic control in patients with type-1 diabetes mellitus. *Diabetes Res Clin Pract* 2006;72:271-6.