Published in final edited form as:

Neuroscience. 2010 January 20; 165(2): 500–514. doi:10.1016/j.neuroscience.2009.10.037.

Cocaine action on peripheral, non-monoamine neural substrates as a trigger of EEG desynchronization and EMG activation following intravenous administration in freely moving rats

Michael S. Smirnov, Eugene A. Kiyatkin^{*}, and Y. Smith

Behavioral Neuroscience Branch, National Institute on Drug Abuse – Intramural Research Program, National Institutes of Health, DHHS, 333 Cassell Drive, Baltimore, Maryland 21224

Abstract

Many important physiological, behavioral and subjective effects of intravenous (iv) cocaine (COC) are exceptionally rapid and transient, suggesting a possible involvement of peripheral neural substrates in their triggering. In the present study, we used high-speed EEG and EMG recordings (4-s resolution) in freely moving rats to characterize the central electrophysiological effects of iv COC at low doses within a self-administration range (0.25–1.0 mg/kg). We found that COC induces rapid, strong, and prolonged desynchronization of cortical EEG (decrease in alpha and increase in beta and gamma activity) and activation of the neck EMG that begin within 2–6 s following the start of a 10-s injection; immediate components of both effects were dose-independent. The rapid effects of COC were mimicked by iv COC methiodide, a derivative that cannot cross the blood-brain barrier. At equipotential doses (0.33–1.33 mg/kg), COC methiodide had equally fast and strong effects on EEG and EMG total powers, decreasing alpha and increasing beta and gamma activities. Rapid EEG desynchronization and EMG activation was also induced by iv procaine, a structurally similar, short-acting local anesthetic with virtually no effects on monoamine uptake; at equipotential doses (1.25–5.0 mg/kg), these effects were weaker and shorter in duration than those of COC. Surprisingly, iv saline injection delivered during slow-wave sleep (but not during quiet wakefulness) also induced a transient EEG desynchronization but without changes in EMG and motor activity; these effects were significantly weaker and much shorter than those induced by all tested drugs. These data suggest that in awake animals, iv COC induces rapid cortical activation and a subsequent motor response via its action on peripheral non-monoamine neural elements, involving neural transmission via visceral sensory pathways. By providing a rapid neural signal and triggering neural activation, such an action might play a crucial role in the sensory effects of COC, thus contributing to the learning and development of drug-taking behavior.

Keywords

electroencephalography; electromyography; arousal; neural activation; sensory effects; cocaine methiodide; procaine; reinforcement

^{*} Correspondence should be addressed to Eugene A. Kiyatkin at the above address. Tel: (443) 740-2844; Fax: (443) 740-2155; ekiyatki@intra.nida.nih.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

It is generally believed that intravenously (iv) delivered cocaine (COC) needs to interact with specific brain substrates to induce its psycho-emotional, physiological and behavioral effects. Although COC has high affinity to a number of ionic channels (i.e., Na^+ , K^+ , Ca^{++} , possibly TRP) on central and peripheral neural elements, a large body of evidence suggests that brain monoamine transporters, specifically the dopamine (DA) transporter, are the major interaction sites underlying its psychoactive and reinforcing properties (Ritz et al., 1987; Rothman and Baumann, 2003; Wise and Bozarth, 1987). However, many important centrally mediated effects of COC are extremely fast and short-lived to be a reflection of the drug's direct interaction with brain substrates. For example, COC-induced euphoria (Fischman and Shuster, 1983; Zernig et al., 2002), increase in arterial blood pressure (Poon and van den Buuse, 1998; Tella, 1996), and peripheral vasoconstriction (Brown and Kiyatkin, 2006) each occur either during or within seconds after iv injection. Although COC easily penetrates the blood-brain barrier (BBB), promoting its rapid entry to the brain tissue, this is a timely process that depends on circulation, crossing of the BBB, and intra-brain diffusion to centrally located neural substrates. In addition to pharmacokinetic considerations, COC-induced euphoria, vasoconstriction, and the acute pressor effect remain generally intact during a DA receptor blockade (Gawin, 1986; Kiritsy-Roy et al., 1990; Poon and van den Buuse, 1998; Sherer et al., 1989; Tella, 1996), suggesting the role of non-DA mechanisms in their mediation. Since ionic channels are abundantly expressed on the terminals of sensory nerves (Lee et al., 2005) that densely innervate blood vessels (Cliff, 1976), COC could directly interact with these peripheral neural substrates, thus triggering some of its rapid physiological effects that could occur before the drug is able to physically interact with brain receptor sites.

Direct support for this mechanism was first obtained in our thermorecording studies, which showed that COC-methiodide (COC-MET), a COC derivative that cannot cross the BBB (Shriver and Long, 1971), mimics regular COC in its ability to increase brain and body temperatures and transiently decrease skin temperature in awake, freely moving rats (Kiyatkin and Brown, 2006). Since a similar increase in brain and body temperatures is also induced by various somato-sensory stimuli, mainly reflecting metabolic neural activation (Kiyatkin et al., 2002), it appears that the peripheral action of COC is sufficient for triggering some of its rapid central excitatory effects. A similar decrease in skin temperature induced by both COC analogues points at the peripheral triggering of COC-induced vasoconstriction, another centrally mediated effect (Knuepfer and Branch, 1992; Crandall et al., 2002) also induced by various natural arousing stimuli (Altschule, 1951; Baker et al., 1976; Solomon et al., 1964). Despite a similarity in central and peripheral temperature responses as well as acute pressor effects (Dickerson et al., 1999), COC-MET did not mimic regular COC in its ability to stimulate locomotion; this effect of COC was fully blocked by DA antagonists (Brown and Kiyatkin, 2006), pointing at the DA system as its critical underlying substrate.

Central and peripheral temperature responses induced by COC were also mimicked by procaine (PRO), a structurally similar local anesthetic drug with negligible effects on monoamine transporters (100-2000-fold lower affinity to DA, norepinephrine and serotonin transporters; Catterall and Mackie, 1996; Ritz et al., 1987). At low doses equipotential to COC for a Na^+ channel blockade, iv PRO, similar to COC, induces increases in brain and body temperatures and transient decreases in skin temperature (Brown and Kiyatkin, 2006). Although PRO is not an addictive drug and has no locomotor stimulatory effects of COC, with rapid iv administration it induces strong sensory and subjective effects in drug-naïve individuals (Servan-Shreiber et al., 1998) and transient sympathoexcitation in animals (Pitts et al., 1988). In COC-experienced individuals, iv PRO induces COC-like acute euphoria (Adiniff et al., 1998) and is self-administered by animals (Johanson, 1980; Kiyatkin and Stein, 1995; Woolverton, 1995). Therefore, it appears that COC interaction with peripherally located non-monoamine substrates could trigger some of its central excitatory effects.

This study was designed to test this hypothesis by comparing acute electroencephalographic (EEG) and electromyographic (EMG) effects of iv COC and related drugs in freely moving rats. First, we examined alterations in cortical EEG and neck EMG induced by iv COC administered at low doses within the reinforcing range (0.25, 0.5 and 1.0 mg/kg or 0.7-2.9 μ M). Second, to verify the peripheral source of central excitatory effects of iv COC, we evaluated EEG and EMG effects of peripherally acting COC-MET iv administered at equimolar doses (0.33, 0.67 and 1.33 mg/kg). Third, to evaluate the contribution of non-monoamine, ion channel mechanisms in the triggering of COC's central effects, we examined EEG and EMG responses induced by PRO at doses (1.25, 2.50 and 5.0 mg/kg) equipotential to COC in its ability to block Na⁺ transport. For accurate capture of the time-course of drug-induced alterations in electrophysiological parameters, analyses were done with a high temporal resolution (4-s bins).

Materials and Methods

Animals and Surgery

Data were obtained from 9 male Long-Evans rats (400 $\pm$ 50 g) supplied by Charles River Laboratories (Greensboro, NC). All animals were housed individually under standard laboratory conditions (12-hr light cycle beginning at 07:00) with free access to food and water. Protocols were performed in compliance with the Guide for the Care and Use of Laboratory Animals (NIH, Publication 865-23) and were approved by the NIDA-IRP Animal Care and Use Committee.

Each rat was surgically prepared for chronic EEG and EMG recording and had a jugular iv catheter. Under general anesthesia (Equithesin 0.33 ml/100 g i.p.; dose of sodium pentobarbital 32.5 mg/kg and chloral hydrate 145 mg/kg), each rat was implanted with 3 stainless steel screws threaded into the skull and two stainless steel wire electrodes that were implanted bilaterally in deep neck muscle. The EMG electrodes were prepared in-house from four insulated 50 μ m wires, which were twisted together and covered (except the last ~4 mm at the tip) by a plastic catheter. All these wires were soldered to a gold-pin connector, similar to that for the EEG electrode. This electrode design provided the necessary strength and flexibility for stable EMG recordings for at least two weeks. After implantation, all connectors were inserted into a plastic socket and fixed with dental acrylic as a head assembly. During the same surgical session, each rat was implanted with a chronic right jugular catheter, which was run subcutaneously to the head mount and secured with dental acrylic. After a 3-4 day period of recovery and habituation to the experimental chamber, recording sessions were held once daily over the next 5-10 days. The top of the plastic socket was protected between sessions by a plastic cap.

EEG/EMG recording

EEG signals were recorded differentially, using two active (2 mm anterior from bregma and 2 mm lateral from midline; 4 mm posterior from bregma and 1.5 mm lateral from midline) and one reference (5 mm anterior from bregma and 1.0 mm lateral from midline; other side of skull) screws implanted into the skull surface according to stereotaxic coordinates of Paxinos and Watson (1998). The latter screw was also used as a reference electrode for differential EMG recording. Electrical activity from EEG and EMG electrodes passed through the pre-amplifier (Pinnacle Technology Inc., Lawrence, Kansas) incorporated inside of an extension cord and electrical swivel to main amplifiers (P15D and P55; Grass Electronics, USA), which were used for additional signal amplification and filtering. EEG signals were filtered from 1 to 30 Hz and EMG signals were filtered between 10 and 1000Hz. The filtered signals then enter a Micro 1401 MK2 interface (Cambridge Electronic Design, Cambridge, UK), allowing its acquisition, recording and analysis using a Spike2 interface (Cambridge Electronic Design).

Experimental protocol

Experiments were conducted in an electrically insulated cage (38 × 47 × 47 cm) placed inside a sound- and light-attenuated plastic box (60 × 566 × 70 cm) under continuous low-intensity red light insulation (20 W) and in view of a small USB camera mounted above the cage. After placement in the cage, the rat was connected to the recording cable and a plastic catheter extension. This extension passed through a liquid swivel outside of the cage, allowing for drug administration without detection of the injection procedure by an animal. Each rat was habituated to the future recording environment both before (2-3 daily sessions ~5-7 hours each) and after (2-3 sessions) surgery without recordings. The first recording session was always an additional habituation session, during which we examined natural fluctuations in EEG and EMG in relation to animal behavior as well as EEG and EMG changes induced by natural sensory stimuli.

Each rat received one or several drugs (COC, COC-MET, PRO), each drug at several injections of three distinct doses. The effects of each drug were tested during two consecutive daily sessions, with two injections at low and moderate doses on Day 1 and two injections at moderate and high doses on Day 2. One drug-free session was always between drug sessions. To exclude possible influences of one drug on the effects of other drugs, the order of drug testing was stochastic and equally balanced in the animal group (i.e., COC was the first testing drug in three rats, COC-MET was the first drug in other three rats, and PRO was first tested in the last three rats); the duration of the testing program was determined by the patency of the iv catheter.

COC HCl was tested in three doses (0.25, 0.50 and 1.0 mg/kg or ~0.75-3 μ M). While the latter two doses are traditionally used in self-administration experiments, COC at a 0.25 mg/kg dose is detected by rat, self-injected, and induces significant locomotor activation (Wellman et al., 2007; Wee et al., 2007). The intervals between individual injections were at least 60 min for low and moderate doses and at least two hours for a high dose. A similar timing was used for the two other drugs, which were injected at doses either equimolar to COC (COC-MET; 0.33, 0.67 and 1.33 mg/kg) or equipotential to COC for a Na⁺ channel blockade (PRO 1.25, 2.50 and 5.0 mg/kg). Each drug was dissolved in saline to provide a small dose in 0.1 ml (over 10 s) and a moderate dose in 0.2 ml (over 20 s). On the next testing day, drugs were dissolved at doubled concentrations. Therefore, a small dose of each drug was always delivered over 10 s, a high dose over 20 s, and a moderate dose over 10 or 20 s (mean=15 s). This bivolumetric injection protocol was used to eliminate refilling of catheters and extension tubing in the middle of a recording session, thus allowing us to bypass the strong stressful influences associated with this procedure. When several drugs were used in the same animals, catheters were flushed twice with saline after completion of a daily session and saline flushing was repeated during the next, drug-free session. In separate sessions (before drug treatments) or in drug-free rats, we also tested the effects of iv saline administration (0.1-0.2 ml over 10-20 s).

Drug injections were made when the rat was in quiet wakefulness or drowsiness/sleep with no motor activity during for 60 s prior to injection (see Results section). Cases, where we observed movements or sudden changes in EMG either spontaneously or due to sensory influences within the 60 s pre-injection time were removed from analysis.

Data analysis

EEG and EMG signals were analyzed based on 6-min durations of recording, with one min preceding and 5 min following drug or saline administration. Although the physiological effects of COC and related drugs are much longer, the 5-min post-injection interval is sufficient for observing pharmacological effects of COC mediated via brain substrates; PET evaluations of drug entry to the brain suggest its peak at 100-120 s after iv injection (Fowler et al., 1998). The following parameters were analyzed for each response: change in total power of EEG (an

integral measure of signal dependent upon both amplitude and frequencies), change in total power of EMG, change in delta (2-4 Hz), theta (4-8 Hz), alpha (8-15 Hz), beta (15-29 Hz), and gamma (29-58 Hz) bands.

Since EEG and EMG signals in individual rats differ in their magnitude and were often saved at different amplification, absolute values of total power were transferred into relative changes, taking a basal value (mean for the 60 s pre-injection interval) as 100%. Therefore, our data represent fluctuations in EEG and EMG total powers relative to their pre-injection state assessed with a 4-s averaging bin. Changes in each individual EEG band (frequency analysis) were analyzed in percents with respect to the pre-event baseline (the 60 s pre-injection interval). A sum of all five bands is equal to EEG total power (=100%).

One-way ANOVA with repeated measures (followed by Fisher post-hoc test) was used as a primary tool for evaluating statistical significance of EEG and EMG changes induced by the drugs. To compare the strength of the effects induced by different drugs and by the same drug at different doses, we calculated mean changes in EEG and EMG total powers at two time scales, representing an immediate (for 20 s from the injection start) and long-term (three minutes from the injection start) effects. Student's t-test was used for between-group comparisons. The use of the words “increase”, “decrease” and “significant” means the presence of a statistically significant change in the parameter or differences between the compared groups or conditions (with at least $p < 0.05$) revealed by either ANOVA or Student's t-test. To simplify the text, most quantitative results of statistical evaluations and comparisons are presented in figure captions and the Table.

Results

1. EEG and EMG response to COC HCl

As shown in Fig. 1 (top row), iv COC administration induced a rapid, robust, and prolonged decrease in EEG total power at each dose tested. While the EEG signal was relatively stable in baseline (fluctuations around 100% values), it dropped within the first analysis bin (0-4 s), reached its minimum at 6-10 s, and remained at these low levels for the entire duration of analysis (5 min). The change was similar at each dose tested, but the response following the lowest dose began to restore from 3-4 min (mean EEG total power $47.69 \pm 1.97\%$ for the last vs. $31.36 \pm 0.92\%$ for the first post-injection minutes, $p < 0.01$), while the decreases at moderate and high doses were stronger and had lower variability (see standard errors of mean). COC injections also resulted in rapid, strong, and prolonged increases in EMG total power (Fig. 1, bottom row), which was similar with each dose tested (see Table). While the increase was evident within the first data point, it peaked (10-18-fold vs. baseline) within 16-24 s (i.e. later than EEG peak) and remained at high levels during the entire analysis interval. An original example of EEG and EMG changes following COC administration at the 0.5 mg/kg dose is shown in Fig. 2 (S76-d-2). Statistical analysis (see Table) revealed no differences in the strength of the COC effect on EEG and EMG total powers at each dose tested in both immediate (0-20 s) and more prolonged (0-180 s) time scales.

EEG frequency analysis (Fig. 3) revealed three prominent and consistent changes induced by COC administration at each dose. Low-frequency alpha activity decreased and high-frequency beta and gamma activities increased. Increases in both beta and gamma frequencies were strong and rapid, peaked at 2-6 s after the injection start, but then beta activity decreased below baseline (~30 s), while gamma activity remained elevated for the entire duration of analysis. Decreases in alpha activity induced by COC had longer onset latencies and were more prolonged. Phasic increases in high-frequency beta and gamma activities were not dose-dependent, but a subsequent tonic decrease appeared to be stronger with increases in dose. While only a tendency to decrease in beta frequencies was seen with the lowest COC dose, the

decrease was stronger and more prolonged at moderate and especially high doses (see Fig. 3; most individual values from ~45 s post-injection were significantly lower than baseline). COC also induced a tonic increase in theta activity, which developed with ~30-s latency and was maintained for about 3-5 min; this effect was the strongest at the 1 mg/kg dose. Delta activity did not show change at low and moderate COC doses, but a weak tonic decrease was seen at the highest dose used.

2. EEG and EMG response to COC-MET

COC-MET used at equimolar doses mimics regular COC HCl in its ability to induce rapid, strong, and prolonged EEG desynchronization as well as EMG activation (Fig. 4 and Table). Similar to regular COC, EEG total power decreased within the injection duration and slowly returned toward baseline, being significantly lower than baseline within the entire 5-min post-injection interval. The effect was significant and strong at each dose, but was maximal at the largest drug dose (see lower standard errors and deeper decrease). The elevation of EMG total power also followed this pattern, but the response was equally strong at each dose tested. An original example of EEG and EMG changes following COC-MET administration at 0.67 mg/kg dose is shown in Fig. 2 (S72-e-3). Although the decrease in EEG total power induced by COC-MET was lower than that for regular COC, differences were minimal and never reached statistical significance (see Table).

COC-MET also mimicked regular COC by the pattern of changes of individual EEG frequencies (Fig. 5). All three basic desynchronization features, decrease in alpha and increases in beta and gamma activities revealed with regular COC, were also seen with COC-MET. However, the beta response was transient and monophasic without the subsequent decrease typical of COC. Similar to COC HCl, COC-MET tonically increased theta frequencies and had minimal effects on delta activity.

3. EEG and EMG response to PRO

PRO used at doses equipotential to COC HCl in blocking Na⁺ transport also induced rapid and strong decrease in EEG total power and EMG activation (Fig. 6). While mimicking the pattern and time-course seen with two COC derivatives, the effects of PRO were weaker, with the EEG total power returning to baseline within 1-3 min after the injection. Only at the largest dose did EEG and EMG changes remain significantly different from baseline for the entire 5-min analysis interval. An original example of EEG and EMG changes following procaine administration at 2.5 mg/kg dose is shown in Fig. 2 (S76-g-2). While the immediate change induced by PRO was similar to that induced by COC, the PRO-induced decrease in EEG total power was significantly weaker over the long term ($p < 0.001$ and $p < 0.05$ for small and moderate doses, respectively; see Table). Similarly, the EMG response induced by PRO at the lowest dose was significantly lower than that produced by COC at the lowest dose.

PRO also mimicked COC in the pattern of changes in individual EEG frequencies. As shown in Fig. 7, beta and gamma frequencies rapidly increased and alpha frequencies rapidly decreased after an iv PRO injection at each dose tested. Similar to COC, beta activation was transient (~20 s) but in contrast to COC it was not followed by a tonic decrease. Similar to COC, gamma activation slowly decreased after the initial peak seen during the injection, but remained at higher levels for the entire analysis interval. The decrease in alpha frequencies was also prolonged and stronger with an increase in drug dose. Similar to COC, PRO also increased theta activity. While COC has a tendency to decrease delta activity, PRO had the opposite pattern, slowly increasing this low-frequency activity in a dose-dependent manner.

4. Dependence of EEG desynchronization upon baseline EEG activity

A rapid, transient increase in beta activity is a typical feature of EEG desynchronization seen with iv injections of all three drugs and at each dose tested. Although the mean magnitude of the phasic beta increase was about the same for each drug and dose (15-20%), certain differences in its changes were seen depending upon the preceding baseline. To assess these relationships, we evaluated how acute beta desynchronization (value of beta power within the first 12 s post-injection) depends upon basal beta activity (mean values of beta power for 60 s preceding injections). Since this change was dose-independent (see Fig. 1, 3 and 5), data were grouped for each drug, combining all three doses tested (Fig. 8).

As can be seen in Fig. 8A, in most cases (135/145) beta power increased after the injection of each tested drug; in a few cases it did not change (5/145) or weakly decreased (5/145). The absolute values of beta increase were approximately the same at different basal values of beta activity (no significant correlations for each drug). However, regression lines crossed the line of equality (response=basal value) at about 20-24%, suggesting that the increase in beta power disappears at a high basal beta activity. These relationships are evident in Fig. 8B, which shows the change in beta power (response minus basal) depending upon basal beta activity. In each case, the change was significantly dependent upon the baseline, with the largest increases at low basal activity and the smallest increases at higher basal activity. Correlation was maximal for PRO ($r=-0.54$, $p<0.001$), slightly lower for COC-HCl ($r=-0.45$, $p<0.01$), and minimal for COC-MET ($r=0.34$, $p<0.05$). Again, the regression lines for each group crossed the line of no effect ($=0\%$) at about the same values of basal beta activity (20-24%). Therefore, the EEG desynchronization is strong when the basal EEG signal is minimally desynchronized (or synchronized) and weak when the basal EEG is already desynchronized. The point of a hypothetical disappearance of beta response (intersection of regression lines with line of no effect) is close to the mean values of beta power increase actually induced by each drug ($18.97 \pm 0.73\%$, $20.18 \pm 1.08\%$, and $22.08 \pm 1.00\%$ for COC HCl, COC-MET, and PRO, respectively). While absolute differences were small, beta activation was maximal with PRO, smaller with COC-MET, and minimal with regular COC ($p<0.05$ vs. PRO).

5. Control iv saline injections

In many cases, saline injections ($n=30$ in 5 rats) did not induce evident changes in EEG and EMG signals. However, when stable high-amplitude activity preceded saline administration, we also observed phasic decreases in amplitude of the EEG signal. An example of transient EEG desynchronization following iv saline injection is shown in Fig. 2 (S79-c-4). When quantified as a group mean (see Table), the effect of saline on EEG total power was significant as an immediate change, but absent as a long-term change. Interestingly, there were no changes in EMG activity following saline injection at both time intervals. Despite a significant decrease in EEG total power as an immediate change, this decrease was significantly weaker than that induced by any drug and in any dose. The difference between the effects of any drug and saline was also highly significant at a prolonged time scale when mean values of EEG total power for saline did not differ from baseline.

Discussion

Hans Berger was the first who reported rapid EEG desynchronization in human subjects following COC injection (Berger, 1937). While confirming this and subsequent results (Chang et al., 1994; Herning et al., 1985; Liu et al., 1998; Luoh et al., 1994; Matsuzaki et al., 1978), the present study demonstrates that EEG desynchronization induced by iv COC in freely moving rats has very short onset latency, appearing within the duration of injection (2-6 s). This rapid effect of COC is dose-independent, being equally strong at all doses effective in maintaining drug self-administration. COC-induced EEG desynchronization manifests as a

profound, three-fold decrease in signal total power associated with phasic relative increases in high-frequency beta and gamma activities and a more prolonged relative decrease in alpha activity. While the increase in beta activity was transient, followed by its decrease below baseline, gamma activation continued for the entire analysis interval (>5 min). COC administration also resulted in weak decreases in delta and increases in theta EEG frequencies. EMG activity also followed a similar but inverted pattern, showing rapid and strong increases that began within the injection interval and maintained for the entire 5-min interval following the injection. Since EEG desynchronization is a typical manifestation of neural activation induced by various somato-sensory stimuli (Pfurtscheller and Lopez de Silva, 1999; Sasaki et al., 1996), these data indicate that iv COC induces rapid neural activation and strong EMG response that are time-locked to the moment of its injection. Therefore, this change should be considered as an essential contributor in mediating the acute physiological effects of iv COC.

The second major finding of this study, the ability of COC-MET to mimic the rapid effects of regular COC, points at the peripheral mechanism involved in triggering the rapid excitatory effects of iv COC. Although COC-MET is able to reach all peripheral substrates of COC interaction, it cannot cross the BBB, acting entirely on the periphery. Therefore, peripherally located monoamine transporters and ionic channels could be viewed as possible substrates that trigger rapid electrophysiological effects of iv COC. In contrast to regular COC, its methiodide derivative has much lower affinity (30 to 60-fold) to DA, norepinephrine and serotonin transporters (Abraham et al., 1992; Ritz et al., 1990). In contrast to the CNS, where all three monoamine transporters are expressed abundantly, only norepinephrine transporters are present on peripheral neural elements. While this transporter could be viewed as a possible substrate, affected by both COC salts, its involvement seems unlikely in light of the very small drug doses (0.7-3.0 μ M) and exceptionally short onsets of EEG and EMG responses. COC from a peripheral vein cannot reach the norepinephrine transporters located on the presynaptic terminals of postganglionic neurons of the sympathetic nervous system with this kind of speed.

To further clarify the possible substrates that could be involved in triggering rapid central effects of COC we used PRO, a structurally similar short-acting anesthetic drug, which shares COC's action on ionic channels (with a ~5-fold lower potency; Reith et al., 1986; Wilcox et al., 1998). The affinity of PRO to DA, norepinephrine, and serotonin transporters is 164-, 136-, and 1970-fold lower than that for COC (K_i within 100-300 μ M range; Ritz et al., 1987; Wilcox et al., 2001), making its interaction with these substrates highly unlikely at the relatively low doses used in this study (4.5-18 μ M). This drug used at doses equipotential to COC in inhibiting Na^+ transport also induced rapid and strong EEG desynchronization and an EMG response that were similar to those induced by regular COC. These effects, however, were weaker in terms of response magnitude (especially for the lowest dose of 1.25 mg/kg) and much less prolonged. Therefore, this finding points at the critical role of non-monoamine neural substrates, particularly ionic channels (Na^+ , K^+ , Ca^{++} , possibly TRP) in triggering rapid excitatory effects of iv COC.

1. Peripheral triggering of central responses via neural sensory mechanism

Cortical EEG desynchronization coupled with EMG activation is a classic neurophysiological response elicited by various somato-sensory stimuli and suggesting neural activation. While the stimulation of somatic or visceral afferents of sensory nerves is an obvious cause for the rapid central responses elicited by somato-sensory stimuli, the same neural mechanism could be invoked by iv COC acting locally at the injection site (blood vessels) before the drug is able to reach the brain and interact with brain receptor sites. Peripheral vessels, especially the veins, are densely innervated by terminals of sensory nerves (Cliff, 1976; Michaelis et al., 1994), and these terminals express various ionic channels, which could be affected by mechanical, thermal and chemical stimuli (Lee et al., 2005). Activation of these vascular receptors results in rapid

excitatory cortical responses that occur via the thalamocortical pathways similar to those involved in transmission of cutaneous sensations (Ploner et al., 2002). Although Na⁺ channels are traditionally viewed as the primary substrate underlying a potent local anesthetic action of COC, this drug has a high affinity to other ionic channels (i.e., K⁺, Ca⁺⁺ and possibly TRP). In contrast to the slower, concentration-dependent action of COC on Na⁺ channels which eventually results in inhibition of neuronal activity (Kiyatkin and Rebec, 2000) and nerve conductance (Caterall and Mackie, 1996), COC's binding to the K⁺ channels has higher affinity (ED₅₀=5-7 μM), ultra-fast dynamics (Ferreira et al., 2001; Premkumar et al., 2005; Wu et al., 2006), and could increase neural activity (Chen et al., 2006), thus producing an excitatory afferent signal, which is transmitted to the CNS via visceral sensory pathways. The issue of primary neural substrates that could be affected by cocaine in the periphery is a complex one and could be clarified in special experiments, using other techniques and experimental paradigms.

While the effective concentrations achieved in brain tissue after iv COC injections at low, self-administering doses could be within a 1-10 μM range (see Fowler et al., 1998) and therefore high enough to interact strongly with centrally located monoamine transporters (K_i between 0.15-1.60 μM for different transporters; Ritz et al., 1987, 1990), much higher drug concentrations will transiently occur in peripheral plasma immediately after iv administration. For example, if iv administered COC is equally distributed in blood plasma (15 ml for a 400 g rat; Lee and Blaurox, 1984), its levels could reach 54-218 μM with 0.25-1.0 mg/kg doses. Since the drug is injected as a high-concentration bolus, actual local levels inside blood vessels could transiently reach even higher concentrations. Such levels greatly exceed COC's affinity to most types of ionic channels, which fluctuate within 5-100 μM range (Woodward et al., 1995; Reith et al., 1986, Wilcox et al., 1998). These phasic peaks of PRO blood plasma levels could be estimated to be within a 0.27-1.1 mM range (for 1.25-5.0 mg/kg doses). Although special work is necessary to clarify the nature of the primary neural elements activated by COC, this activation appears to be the core of a transient arousal and unusual sensory feelings induced in humans induced by iv administrations of COC and other local anesthetic drugs. Both humans and animals always detect COC as a sensory stimulus with sc and iv injection and, although the subjective effects of iv PRO in drug-naïve individuals differ from those of COC, they are equally rapid and powerful, combining euphoria, anxiety, depression, fear, as well as strong sensory and somatic sensations (Servan-Shreiber et al., 1998).

2. Cocaine-induced EEG desynchronization as a reflection of generalized neural activation

While the cellular mechanisms underlying cortical EEG desynchronization are currently a matter of debate (see Buzsaki, 2006; Hobson, 1999 for review), it is clear that this electrophysiological phenomenon reflects CNS activation, which could be triggered by various salient environmental stimuli, ranging from simple sensory to noxious somatic or visceral challenges. While the neural pathways conveying sensory afferent signals differ depending upon the stimulus' modality and strength, these signals converge to a number of subcortical structures, finally resulting in generalized neural activation (Steriade and McCarley, 2005). Since iv COC injection mimics sensory stimuli in its ability to induce rapid EEG desynchronization and EMG activation, it appears that it has a similar central excitatory action, inducing a generalized neural activation.

Surprisingly, the cortical EEG signal was affected by iv saline injection. When the rat was fully inactive and showed large-amplitude EEG fluctuations (slow-wave sleep), saline injection typically induced a transient decrease in EEG total power, with a drop in alpha activity and a rise in beta and gamma activities, the pattern typical to desynchronization induced by both sensory stimuli and all three drugs tested in this study. However, the EMG did not show any changes in this case. In contrast, when the rat was in quiet wakefulness and the basal EEG

signal was more desynchronized, the EEG signal did not change at all following saline injection. Therefore, this finding suggests that the iv saline administration is detected by animals and results in transient EEG desynchronization, but only in freely moving conditions following slow-waves sleep. Since our protocol excluded the possibility of additional sensory stimuli associated with the procedure of iv injection, it appears that changes in pressure or/and temperature are likely factors that trigger this transient EEG response. While this finding was unexpected, it suggests that EEG is sensitive to detect both spontaneous fluctuations in the CNS activity state and transient desynchronization episodes induced by weak visceral sensory stimuli. On the other hand, this finding also suggests that interoceptive sensory stimuli associated with the injection procedure (i.e., its speed, temperature, and volume) could modulate to some extent the effects induced by the drug *per se*.

While ultra-fast EEG desynchronization and EMG activation induced by iv COC is a manifestation of central excitatory action, this action appears to be non-specific (i.e. occurring with other visceral and somatic sensory challenges), manifesting in rapid changes of other physiological parameters. For example, an acute increase in arterial blood pressure, a known centrally mediated effect of COC (Knuepfer and Branch, 1992), peaks at 10-15 s after the start of iv injection with a subsequent significant decrease toward baseline within the first post-injection minute (Poon and van den Buuse, 1998; Tella, 1996). Similar to an apparent lack of dose dependency in the immediate effects of COC on EEG and EMG, peak increases of arterial blood pressure induced by COC were equally strong within 0.3-3.0 mg/kg doses (Poon and van den Buuse, 1998), covering the entire range of drug self-administration, and were resistant to a DA receptor blockade (Kiritsy-Roy et al., 1990; Poon and van den Buuse, 1998; Tella, 1996). In contrast to the inability of DA antagonists to block the immediate pressor effect of iv COC, subsequent tonic blood pressure elevations were dose-dependent and were attenuated by a DA receptor blockade. Despite different triggering mechanisms, the acute pressor effect of COC was similar to that elicited by arousing somato-sensory stimuli such as sound, light, and tactile stimulation (Stojicić et al., 2008). Iv COC also induced a surprisingly rapid drop in skin temperature (~20 s from the injection start; Kiyatkin and Brown, 2005), suggesting acute peripheral vasoconstriction (Baker et al., 1976)—another centrally mediated effect of COC (Gillis et al., 1995; Vongpatanasin et al., 1999) that is mimicked by various arousing somato-sensory stimuli (Altschule, 1951; Baker et al., 1976; Solomon et al., 1964). Rapid, transient drop in skin temperature induced by iv COC (1 mg/kg) was both quantitatively and qualitatively similar to that induced by such arousing stimuli as tail-pinch and social interaction with another animal; this effect was also fully resistant to a DA receptor blockade (Kiyatkin and Brown, 2005). Finally, acute COC-induced euphoria has exceptionally short latencies and a transient nature (Zernig et al., 2003), being generally resistant to a DA receptor blockade (Gawin, 1986; Sherer et al., 1989) and mimicked by PRO (Fischman and Schuster, 1983; Adinoff et al., 1998), which has incomparably lower affinity to monoamine transporters.

It is well known that EEG desynchronization is a state-dependent phenomenon, which manifests following highly synchronized activity (Steriade and McCarley, 2005). Our data revealed that all three tested drugs also shared this property (Fig. 8), inducing EEG desynchronization strongly dependent upon basal EEG activity. Being quantified as a change in beta power, the response was strong when basal beta activity was at its low levels (4-8%), progressively decreased with increases in basal beta activity (8-14%), and fully disappeared when basal activity was highly desynchronized (18-22%). Importantly, this level of beta activation was the phasic response that in fact occurred following administration of these drugs.

3. Relationships between rapid sensory and slow pharmacological effects of COC

In addition to COC's ability to induce rapid EEG desynchronization and an acute EMG response, the effects shared with other tested drugs, COC also had unique effects that were absent with

other drugs. While a common mechanism invoking afferents of sensory nerves, rapid neural transmission, and generalized neural activation could explain the ultra-fast nature of many important physiological, behavioral and psycho-emotional effects induced by iv COC and the unusual parallels in these effects observed with peripherally acting COC-MET and a “pure” local anesthetic PRO, COC is a powerful addictive drug, which has specific, dose-dependent effects, which depend on its direct interaction with multiple brain substrates, particularly monoamine transporters (Ritz et al., 1987; Rothman and Baumann, 2003; Wise and Bozarth, 1987). However, at the level of EEG it is difficult to distinguish these COC-specific effects from its non-specific effects shared with other drugs and arousing stimuli. Moreover, these effects are superimposed on the non-specific effects common to all three drugs, making their identification even more difficult. This task is especially difficult since it is impossible to selectively block COC's action on ionic channels using pharmacological tools in freely moving animal. Other strategies should be developed to find a way of blocking the rapid sensory actions of COC and examine its physiological and behavioral effects, which are mediated via its action on monoamine transporters. Since multiple peripheral and central mechanisms are simultaneously engaged by COC under natural conditions, it is also unclear how the behavioral and physiological effects of this drug will be modulated if its rapid arousing effects will be somehow eliminated. Converging to the common CNS substrates but having quite a different time-course, this rapid action could strongly modulate more slow and prolonged effects mediated via a direct drug action on brain substrates.

4. Functional implications

Our present data suggest that iv COC induces neural activation and point at the critical role of peripheral non-monoamine neural substrates in the triggering of this rapid and generalized effect. Although this peripherally triggered action of COC, which it shares with other related drugs and a variety of somato-sensory stimuli, is not sufficient for making COC a unique drug of abuse, it could play a crucial role in mediating the physiological and subjective effects of COC, thus contributing to mechanisms of COC reinforcement.

By providing a rapid sensory signal immediately following drug administration and by triggering neural activation, this action appears to play a critical role in the sensory effects of COC. Therefore, the acute somato-sensory (= subjective or psychoemotional) experience associated with iv COC administration could occur as a consequence of the drug's direct interaction with neural substrates of visceral sensory systems, which is detected by the CNS, inducing a wide-spreading neural activation. This rapid action, however, could be dramatically modulated (“conditioned”) due to its repeated preceding (pairing with) of slower and prolonged direct actions on brain neural substrates, particularly monoamine transporters that are essential for COC reinforcement (Ritz et al., 1987; Wise and Bozarth, 1987). Therefore, acute COC-induced euphoria, a rapid, powerful but transient psycho-emotional experience occurring immediately after iv or smoked COC, could be the result of COC's direct interaction with peripheral sensory substrates being conditioned due to drug experience.

By affecting various central neurons that are involved in the processing of somato-sensory information and organization of brain activational processes, rapid sensory action of COC may interact with other pharmacological actions, particularly on monoamine transporters and monoamine uptake. It is known that central neurons containing DA, norepinephrine and serotonin are affected by sensory stimuli, producing in most cases phasic neuronal excitations (Aston-Jones and Bloom, 1981; Heim et al., 1982; Kiyatkin, 1988; Steinfels et al., 1983). If, similar to sensory stimuli, iv COC will induce phasic excitation of DA neurons, it will provide impulse-dependent DA release, which will be potentiated by its later action on DA uptake, altering synaptic DA levels. Direct evidence for such an action was recently obtained independently with electrochemistry (Aragona et al., 2008) and single-unit recording (Brown

and Kiyatkin, 2008). As shown with fast-cyclic voltammetry, extrasynaptic DA levels rapidly grow following iv COC injection in awake rats administered with a D2 antagonist (Aragona et al., 2008). Similarly, in awake rats treated with DA antagonists, most presumed DA VTA neurons show phasic, transient excitations that begin during iv COC injection (Brown and Kiyatkin, 2008). Therefore, COC could affect the DA system indirectly via rapid neural mechanism before it physically reaches the DA transporters. Thus, COC's effects on DA transmission could be determined by at least two independent pharmacological actions, which may interact with and influence each other.

This interaction of two independent actions could play a crucial role in COC reinforcement (conditioning, learning), which always includes interaction between two initially independent forces (actions). While COC learning (conditioning) is usually viewed as an interaction between the drug's pharmacological actions and sensory stimuli or voluntary movement acts that precede these actions, our present data suggest that COC administration provides its own pharmacologically-mediated sensory signal, which may interact with other pharmacological actions. At neuronal and neurochemical levels, this is an interaction between a rapid neural effect which is triggered by drug administration, signals drug delivery, and induces acute psychophysiological activation, and subsequent slow and more prolonged direct pharmacological actions on monoamine uptake. While cellular mechanisms underlying these interactions following repeated drug use (experience) remain unclear, they may play an important role in the development of drug-seeking behavior. This does not mean that external stimuli associated with COC intake are not important in development of drug-taking behavior. However, it appears that, similar to natural reinforcing stimuli, COC has its acute sensory effects that are involved in mechanisms of COC reinforcement, thus playing a crucial role in the development and maintenance of drug-seeking and drug-taking behavior.

Acknowledgments

The research described in this manuscript was supported by the Intramural Research Program of the NIH, NIDA.

References

- Abraham P, Pitner JB, Lewin AH, Boja JW, Kuhar MJ, Carroll FI. N-modified analogues of cocaine: synthesis and inhibition of binding to the cocaine receptor. *J Med Chem* 1992;35:141–144. [PubMed: 1732521]
- Adinoff B, Brady K, Sonne S, Mirabella RF, Kellner CH. Cocaine-like effects of intravenous procaine in cocaine addicts. *Addiction Biol* 1998;3:189–196.
- Altschule MD. Emotion and circulation. *Circulation* 1951;3:444–454. [PubMed: 14812675]
- Aragona BJ, Cleaveland NA, Stuber GD, Day JJ, Carelli RM, Wightman RM. Preferential enhancement of dopamine transmission within the nucleus accumbens shell by cocaine is attributable to a direct increase in phasic dopamine release events. *J Neurosci* 2008;28:8821–8831. [PubMed: 18753384]
- Aston-Jones G, Bloom FE. Norepinephrine-containing locus coeruleus neurons in behaving rats exhibit pronounced responses to non-noxious environmental stimuli. *J Neurosci* 1981;1:887–900. [PubMed: 7346593]
- Baker M, Cronin M, Mountjoy D. Variability of skin temperature in the waking monkey. *Am J Physiol* 1976;230:449–455. [PubMed: 816208]
- Berger H. *Arch Psychiatr Nervenkr* 1937;106:577–584. Hans Berger on the electroencephalography of man. *EEG Clin Neurophysiol* 1969;28:95–132.
- Brown PL, Kiyatkin EA. The role of peripheral Na⁺ channels in triggering the central excitatory effects of intravenous cocaine. *Eur J Neurosci* 2006;24:1182–1192. [PubMed: 16930444]
- Brown PL, Kiyatkin EA. Sensory effects of intravenous cocaine on dopamine and non-dopamine ventral tegmental area neurons. *Brain Res* 2008;1218:230–249. [PubMed: 18514638]
- Buzsaki, G. *Rhythms of the brain*. Oxford University Press; Oxford: 2006.

- Catterall, W.; Mackie, K. Local anesthetics.. In: Hardman, JC.; Limbird, LE., editors. Goodman and Gilman's The Pharmacological Basis of Therapeutics. Vol. 9th Ed.. McGraw; NY: 1996. p. 331-347.
- Chang AY, Kuo TB, Chan SH. Power spectral analysis of electroencephalographic desynchronization induced by cocaine in the rat. *Neurosci Lett* 1994;170:175–178. [PubMed: 8041501]
- Chen YH, Lin CH, Lin PL, Tsai MC. Cocaine elicits action potential burst in a central snail neuron: the role of delayed rectifying K⁺ channels. *Neuroscience* 2006;138:257–280. [PubMed: 16377093]
- Cliff, WJ. Biological Structure and Functions. Cambridge University Press; Cambridge and NY: 1976. Blood vessels..
- Crandall CG, Vongpatanasin W, Victor RG. Mechanism of cocaine-induced hyperthermia in humans. *Ann Intern Med* 2002;136:785–791. [PubMed: 12044126]
- Dickerson LW, Rodak DJ, Kuhn FE, Wahlstrom SK, Tessel RE, Visner MS, Schaer GL, Gillis RA. Cocaine-induced cardiovascular effects: Lack of evidence for a central nervous system site of action based on hemodynamic studies with cocaine methiodide. *J Cardiovasc Pharmacol* 1999;33:36–42. [PubMed: 9890394]
- Ferreira S, Crumb WJ, Carlton CG, Clarkson CW. Effects of cocaine and its major metabolites on the HERG-encoded potassium channel. *J Pharmacol Exp Ther* 2001;299:220–226. [PubMed: 11561083]
- Fischman MW, Schuster CR. A comparison of the subjective and cardiovascular effects of cocaine and procaine in humans. *Pharmacol Biochem Behav* 1983;18:711–716. [PubMed: 6856646]
- Fowler JS, Volkow ND, Logan J, Gatley SJ, Pappas N, King P, Ding Y-S, Wang G-J. Measuring dopamine transporter occupancy by cocaine in vitro: radiotracer considerations. *Synapse* 1998;28:111–116. [PubMed: 9450511]
- Gawin F. Neuroleptic reduction of cocaine-induced paranoia but not euphoria? *Psychopharmacology* 1986;90:142–143. [PubMed: 2876452]
- Gillis RA, Hernandez YM, Erzuoki HK, Raczkowski VF, Mandal AK, Kuhn FE, Dretchen KL. Sympathetic nervous system mediated cardiovascular effects of cocaine are primary due to a peripheral site of action of the drug. *Drug Alcohol Depend* 1995;37:217–230. [PubMed: 7796716]
- Heym J, Steinfels GF, Jacobs BL. Activity of serotonin-containing neurons in the nucleus raphe pallidus of freely moving rats. *Brain Res* 1982;251:259–276. [PubMed: 7139326]
- Herning RI, Jones RT, Hooker WD, Mendelson J, Blackwell L. Cocaine increases EEG beta: a replication and extension of Hans Berger's historic experiments. *EEG Clin Neurophysiol* 1985;60:470–477. 9.
- Hobson, JA. Sleep and dreaming.. In: Zigmond, MJ.; Bloom, FE.; Landis, SC.; Roberts, JL.; Squire, LR., editors. *Fundamental Neuroscience*. Academic Press; San Diego: 1999. p. 1207-1227.
- Johanson CE. The reinforcing properties of procaine, chlorprocaine and proparacaine in rhesus monkeys. *Psychopharmacology* 1980;67:189–194. [PubMed: 6768094]
- Kiritsy-Roy JA, Halter JB, Gordon SM, Smith MJ, Terry LC. Role of the central nervous system in hemodynamic and sympathoadrenal responses to cocaine in rats. *J Pharmacol Exp Ther* 1990;256:154–160. [PubMed: 2213551]
- Kiyatkin EA. Functional properties of presumed dopamine-containing and other ventral tegmental area neurons in conscious rats. *Int J Neurosci* 1988;42:21–43. [PubMed: 3209370]
- Kiyatkin EA, Brown PL, Wise R. Brain temperature fluctuation: a reflection of functional neural activation. *Eur J Neurosci* 2002;16:164–168. [PubMed: 12153543]
- Kiyatkin EA, Rebec GV. Dopamine-independent action of cocaine on striatal and accumbal neurons. *Eur J Neurosci* 2000;12:1–13. [PubMed: 10651854]
- Kiyatkin EA, Stein EA. Fluctuations in nucleus accumbens dopamine during cocaine self-administration behavior: an in vivo electrochemical study. *Neuroscience* 1995;64:599–617. [PubMed: 7715774]
- Knuepfer MM, Branch CA. Cardiovascular responses to cocaine are initially mediated by the central nervous system in rats. *J Pharmacol Exp Ther* 1992;263:734–741. [PubMed: 1432699]
- Lee HB, Blafox MD. Blood volume in the rat. *J Nucl Med* 1984;25:72–76. [PubMed: 6233402]
- Lee Y, Lee C-H, Oh U. Painful channels in sensory neurons. *Mol Cells* 2005;20:315–324. [PubMed: 16404144]
- Liu X, Vaupel DB, Grant S, London ED. Effect of cocaine-related environmental stimuli on the spontaneous electroencephalogram in polidrug abusers. *Neuropsychopharmacology* 1998;19:10–17. [PubMed: 9608572]

- Luoh HF, Kuo TB, Chan SH, Pan WH. Power spectral analysis of electroencephalographic desynchronization induced by cocaine in rats: correlation with microdialysis evaluation of dopaminergic neurotransmission at the medial prefrontal cortex. *Synapse* 1994;16:29–35. [PubMed: 8134898]
- Matsuzaki M, Spingler PJ, Whitlock EG, Misra AL, Mule SJ. Comparative effects of cocaine and pseudococaine on EEG activities, cardiorespiratory functions, and self-administration behavior in the rhesus monkey. *Psychopharmacology* (Berlin) 1978;57:12–20.
- Michaelis M, Goder R, Habler HJ, Janig W. Properties of afferent nerve fibers supplying the saphenous vein in the cat. *J Physiol* (London) 1994;474:233–243. [PubMed: 8006810]
- Paxinos, J.; Watson, C. *The Rat Brain in Stereotaxic Coordinates*. Academic Press; Sydney: 1998.
- Pfurtscheller G, Lopes de Silva FH. Event-related EEG/EMG synchronization and desynchronization: basic principles. *Clin Neurophysiol* 1999;110:1842–1857. [PubMed: 10576479]
- Pitts DK, Udom CE, Marwah J. Cardiovascular effects of cocaine in anesthetized and conscious rats. *Life Sci* 1987;40:1099–1111. [PubMed: 3821374]
- Ploner M, Holthusen H, Noetges P, Schnitzler A. Cortical representation of venous nociception in humans. *J Neurophysiol* 2002;88:300–305. [PubMed: 12091555]
- Poon J, van den Buuse M. Autonomic mechanisms in the acute cardiovascular effects of cocaine in conscious rats. *Eur J Pharmacol* 1998;363:147–152. [PubMed: 9881582]
- Premkumar LS. Block of a Ca^{++} -activated potassium channel by cocaine. *J Membr Biol* 2005;204:129–136. [PubMed: 16245035]
- Reith MEA, Kim SS, Lajtha A. Structural requirements for cocaine congeners to interact with [^3H] batrachotoxin A20- α -benzoate binding sites on sodium channels in mouse brain synaptosomes. *J Biol Chem* 1986;261:7300–7305. [PubMed: 2423518]
- Ritz MC, Lamb RJ, Goldberg SR, Kuhar MJ. Cocaine receptors of dopamine transporters are related to self-administration of cocaine. *Science* 1987;237:1219–1223. [PubMed: 2820058]
- Ritz MC, Cone EJ, Kuhar MJ. Cocaine inhibition of ligand binding at dopamine, norepinephrine and serotonin transporters: a structure-activity study. *Life Sci* 1990;46:635–645. [PubMed: 2308472]
- Rothman RB, Baumann MH. Monoamine transporters and psychostimulant drugs. *Eur J Pharmacol* 2003;479:23–40. [PubMed: 14612135]
- Sasaki H, Coffey P, Villegas-Perez MP, Vidal-Sanz M, Young MJ, Lund RD, Fukuda Y. Light induced EEG desynchronization and behavioral arousal in rats with restored retinocollicular projection by peripheral nerve graft. *Neurosci Lett* 1996;218:45–48. [PubMed: 8939477]
- Servan-Shreiber D, Perlstein WM, Cohen JD, Mintun M. Selective pharmacological activation of limbic structures in human volunteers: A positron emission tomography study. *J Neuropsychiatry* 1998;10:148–159.
- Sherer MA, Kumor KM, Jaffe JH. Effects of intravenous cocaine are partially attenuated by haloperidol. *Psychiatry Res* 1989;27:117–125. [PubMed: 2652168]
- Shriver DA, Long IP. A pharmacological comparison of some quaternary derivatives of cocaine. *Arch Int Pharmacodyn* 1971;189:198–208. [PubMed: 5130150]
- Solomon GF, Moos RH, Stone GC, Fessel WJ. Peripheral vasoconstriction induced by emotional stress in rats. *Angiology* 1964;15:362–365. [PubMed: 14198214]
- Steinfels GF, Heym J, Stricker RE, Jacobs BL. Behavioral correlates of dopaminergic unit activity in freely moving cats. *Brain Res* 1983;258:217–228. [PubMed: 6824912]
- Steriade, M.; McCarley, RW. *Brain control of wakefulness and sleep*. Kluwer Academic/Plenum Publishers; NY: 2005.
- Stojčić S, Milutinović-Smiljanić S, Sarenac O, Milosavljević S, Paton JF, Murphy D, Japundžić-Zigon N. Blockade of central vasopressin receptors reduces the cardiovascular response to acute stress in freely moving rats. *Neuropharmacology* 2008;54:824–836. [PubMed: 18339407]
- Tella SR. Possible novel pharmacodynamic action of cocaine: cardiovascular and behavioral evidence. *Pharmacol Biochem Behav* 1996;54:343–354. [PubMed: 8743594]
- Tuncel M, Wang Z, Arbiq D, Fadel PJ, Victor RG, Vongpatanasin W. Mechanism of the blood pressure-raising effect of cocaine in humans. *Circulation* 2002;105:1054–1059. [PubMed: 11877354]

- Vongpatanasin W, Mansour Y, Chavoshan B, Arbique D, Victor RG. Cocaine stimulates the human cardiovascular system via a central mechanism of action. *Circulation* 1999;100:497–502. [PubMed: 10430763]
- Wee S, Specio SE, Koob GF. Effects of dose and session duration on cocaine self-administration in rats. *J Pharmacol Exp Ther* 2007;320:1134–1143. [PubMed: 17202405]
- Wellman PJ, Nation JR, Davis KW. Impairment of acquisition of cocaine self-administration in rats maintained on a high-fat diet. *Pharmacol Biochem Behav* 2007;88:89–93. [PubMed: 17764729]
- Wilcox KM, Paul IA, Ordway GA, Woolverton WL. Role of the dopamine transporter and the sodium channel on the cocaine-like discriminative stimulus effects of local anesthetics in rats. *Psychopharmacology* 2001;157:260–268. [PubMed: 11605081]
- Wise RA, Bozarth MA. A psychomotor stimulant theory of addiction. *Psychol Rev* 1987;94:469–492. [PubMed: 3317472]
- Woodward JJ, Compton DM, Balster RL, Martin BR. In vitro and in vivo effects of cocaine and selected local anesthetics on the dopamine transporter. *Eur J Pharmacol* 1995;13:7–13. [PubMed: 7635175]
- Woolverton WL. Comparison of the reinforcing efficacy of cocaine and procaine in rhesus monkeys responding under a progressive ratio schedule. *Psychopharmacology* 1995;120:296–302. [PubMed: 8524977]
- Wu SN, Chang HD, Sung BJ. Cocaine-induced inhibition of ATP-sensitive K⁺ channels in rat ventricular myocytes and in heart-derived H9c2 cells. *Basic Clin Pharmacol Toxicol* 2006;98:510–517. [PubMed: 16635111]
- Zernig G, Giacomuzzi S, Riemer Y, Wakonig G, Sturm K, Saria A. Intravenous drug injection habits: Drug users' self-reports versus researchers' perception. *Pharmacology* 2003;68:49–56. [PubMed: 12660479]

Fig. 1.

Mean changes in EEG (top) and EMG total powers (bottom, both in %) following iv COC administration at 0.25, 0.50 and 1.0 mg/kg doses. Vertical dashed lines (0, s) show the start of injections and horizontal dashed lines show basal values. Filled symbols show values significantly different (Fisher F-test) from baseline. n=number of tests. ANOVA values for the effect of cocaine on EEG and EMG are: 0.25 mg/kg – $F_{12,402}=20.86, 9.41$; 0.50 mg/kg – $F_{25,805}=26.82, 25.15$; 1.0 mg/kg – $F_{11,371}=37.30, 11.10$; each $p<0.001$).

Fig. 2.

Original examples of EEG and EMG activity (μV at final amplification) recorded in freely moving rats following iv administration of drugs and saline. Vertical hatched lines show injection onsets and offsets. Data were obtained in three different rats (S76-d-2, S76-c-4, S72-e-3, S76-g-2; first three symbols show rat number, the next symbol is session number and the last symbol is test number). While iv saline injection in many cases did not induce evident EEG changes, the example (S79-c-4) shows a transient desynchronization, when the injection was performed during a prolonged episode of slow-wave sleep.

Fig. 3. Mean changes in individual EEG frequencies [total power for delta (A), theta (B), alpha (C), beta (D) and gamma (E) waves, shown in %] following iv COC administration. Similar to Fig. 4, vertical and horizontal hatched lines in each graph show the starts of injections and basal values, respectively. ANOVA values for COC at 0.25 mg/kg (delta: $F_{13,433}=1.08$, NS; theta=3.79, $p<0.001$; alpha=1.90, $p<0.01$; beta=2.85; $p<0.001$; gamma=8.59, $p<0.001$). ANOVA values for COC at 0.50 mg/kg (delta: $F_{21,681}=0.88$, NS; theta=3.40, $p<0.001$; alpha=1.93, $p<0.01$; beta=4.00; $p<0.001$; gamma=8.24, $p<0.001$). ANOVA values for COC at 1.0 mg/kg (delta: $F_{11,371}=1.44$, $p=0.06$; theta=4.61, $p<0.001$; alpha=3.32, $p<0.001$; beta=5.64; $p<0.001$; gamma=8.86, $p<0.001$).

Fig. 4.

Mean changes in EEG (top) and EMG total power (bottom, both in %) induced by iv administration of COC methiodide at 0.33, 0.67 and 1.33 mg/kg doses. Vertical dashed lines (0, s) show the start of injections and horizontal dashed lines show basal values. Filled symbols show values significantly different (Fisher F-test) from baseline. n=number of tests. ANOVA values for the effect of cocaine on EEG and EMG are: 0.33 mg/kg – $F_{12,402}=12.09$, 8.55; 0.67 mg/kg – $F_{18,588}=6.32$, 8.63; 1.33 mg/kg – $F_{11,371}=8.21$, 5.72; each $p<0.001$).

Fig. 5.

Mean changes in individual EEG frequencies [total power for delta (A), theta (B), alpha (C), beta (D) and gamma (E) waves, shown in %] following iv administration of COC methiodide. Similar to Fig. 3, vertical and horizontal hatched lines in each graph show the starts of injections and basal values, respectively. ANOVA values for COC-MET at 0.33 mg/kg (delta: $F_{12,402}=1.08$, NS; theta=2.26, $p<0.01$; alpha=2.64, $p<0.01$; beta=1.62; $p<0.05$; gamma=5.33, $p<0.001$). ANOVA values for COC-METH at 0.67 mg/kg (delta: $F_{18,588}=0.78$, NS; theta=3.25, $p<0.001$; alpha=2.08, $p<0.01$; beta=2.84; $p<0.001$; gamma=3.82, $p<0.001$). ANOVA values for COC-MET at 1.33 mg/kg (delta: $F_{11,371}=1.95$, $p<0.05$; theta=2.59, $p<0.001$; alpha=2.04, $p<0.01$; beta=2.19; $p<0.01$; gamma=3.41, $p<0.001$).

Fig. 6.

Mean changes in EEG (top) and EMG total power (bottom, both in %) induced by iv administration of PRO at 1.25, 2.50 and 5.0 mg/kg doses. Vertical hatched lines (0, s) show the start of injections and horizontal hatched lines show basal values. Filled symbols show values significantly different (Fisher F-test) from baseline. n=number of tests. ANOVA values for the effect of procaine on EEG and EMG are: 1.25 mg/kg – $F_{17,557}=3.58, 4.40$; 2.50 mg/kg – $F_{16,526}=4.29, 6.33$; 5.0 mg/kg – $F_{12,402}=6.79, 11.59$; each $p<0.001$).

Fig. 7.

Mean changes in individual EEG frequencies [total power for delta (A), theta (B), alpha (C), beta (D) and gamma (E) waves, shown in %] following iv administration of PRO. Similar to Fig. 3, vertical and horizontal hatched lines in each graph show the starts of injections and basal values, respectively. ANOVA values for procaine at 1.25 mg/kg (delta: $F_{17,557}=1.86$, $p<0.05$; theta=1.98, $p<0.01$; alpha=4.75, $p<0.001$; beta=2.21; $p<0.01$; gamma=9.73, $p<0.001$). ANOVA values for procaine at 2.50 mg/kg (delta: $F_{16,526}=1.16$, NS; theta=2.01, $p<0.01$; alpha=2.58, $p<0.001$; beta=2.35; $p<0.001$; gamma=5.78, $p<0.001$). ANOVA values for procaine at 5.00 mg/kg (delta: $F_{12,402}=1.27$, NS; theta=1.78, $p<0.01$; alpha=4.45, $p<0.001$; beta=1.89; $p<0.05$; gamma=6.86, $p<0.001$).

Fig. 8.

Relationships between changes in EEG beta frequencies induced by cocaine HCl, cocaine methiodide and procaine depending upon basal beta activity. Upper panel (A) shows relationships between beta activity in baseline and at peak of EEG desynchronization. These values are independent of each other. Lower panel (B) shows relationships between basal beta activity and its change (peak value during desynchronization – basal value). There was significant correlation between these values for each drug (see r , coefficient of correlation, regression equations, and regression lines). Hatched line shows a line of no effect (0% change). Each regression line crosses the line of no effect between 19-24%, suggesting that beta activation is strong at low basal beta activity and disappears at higher basal activity.

Table

Rapid and long-term changes in EEG and EMG total powers induced by iv injections of COC HCl (COC), COC methiodide (COC-MET), procaine (PRO), and saline in freely moving rats

Group	Immediate change (0-20 s)	Long-term change (0-300 s)
EEG, %		
COC 0.25	31.82±2.78***	28.82±2.35***
COC 0.50	35.82±2.70***	32.47±1.76***
COC 1.00	26.34±3.21***	27.50±2.90***
COC-MET 0.33	36.88±4.55***	30.36±2.87***
COC-MET 0.67	43.54±4.80***	40.33±4.95***
COC-MET 1.33	36.41±5.36***	32.61±3.00***
PRO 1.25	34.70±5.59***	55.13±5.36***,ooo
PRO 2.50	40.04±4.91***	45.39±5.72***,o
PRO 5.00	38.61±6.82***	35.05±3.68***
Saline	62.54±5.57*,ooo	99.04±8.81 ^{ooo}
EMG, ln (%)		
COC 0.25	6.67±0.32***	6.76±0.13***
COC 0.50	6.10±0.20***	6.78±0.21***
COC 1.00	6.22±0.20***	6.73±0.12***
COC-MET 0.33	5.99±0.37***	6.08±0.14***
COC-MET 0.67	6.25±0.37***	6.65±0.15***
COC-MET 1.33	6.46±0.28***	6.89±0.14***
PRO 1.25	5.07±0.11*,ooo	5.08±0.12*,oo
PRO 2.50	6.41±0.27***	6.85±0.15***
PRO 5.00	5.66±0.34**	6.61±0.15***
Saline	4.67±0.07 ^{ooo}	4.69±0.06 ^{ooo}

EEG total power data are shown in percents (±sem) relative to baseline (mean of 15 values preceding the injection=100%). The same procedure was used for quantification of EMG total power, but because of high and variable response amplitude, natural logarithmic derivatives of percent values were used for statistical analyses. EMG values shown in the Table are ln (total power, %), i.e. 4.61=100% and 6.67= 788%. *, **, and *** indicate statistically significant decreases differences vs. pre-injection baseline with p<0.05, p<0.01 and p<0.001. o, oo, and ooo indicate statistically significant differences (p<0.05, 0.01 and 0.001, respectively) vs. respective COC groups