CORRESPONDENCE

The Pathophysiology, Diagnosis and Treatment of Constipation

by Prof. Dr. med. Stefan Lissner in volume 25/2009

Obsolescence of X-ray Defecography

Müller-Lissner in his article mentions conventional x-ray defecography as the imaging procedure of choice in the diagnosis of defecation disorders.

This procedure is obsolete, owing to its radiation exposure and its limited validity. Dynamic magnetic resonance imaging (MRI) of the pelvic floor, which is unfortunately not mentioned at all, can be conducted speedily and simply. It does not entail exposure to ionizing radiation and is better tolerated by patients. The reason is that the investigators are outside the room where the patient is lying in the scanner and evacuates into diapers. It was shown years ago that analyzing defecation while the patient is lying down is equivalent to analyzing it while the patient is sitting up (2).

Since defecation disorders are often complex dysfunctions of the pelvic floor, a further advantage lies in the fact that by using MRI, the bladder, urethra, vagina, and uterus can be imaged during one examination, without the need for any further invasive procedures or for contrast medium (1).

MRI also enables the representation of complications such as enterocele or sigmoidocele. High resolution imaging of the pelvic floor structures allows for assessment of atrophies and scarring of the levator sling as well as the fascia and ligaments of the pelvic floor (3).

DOI: 10.3238/arztebl.2009.0765a

REFERENCES

1. Healy JC, et al.: Dynamic MR imaging compared with evacuation proctography when evaluating anorectal configuration and pelvic floor movement. *AJR* 1997; 169: 775–779.
2. Fielding JR, et al.: MR imaging of pelvic floor continence mechanisms in the supine and sitting postitions. *AJR* 1998; 171: 1607–1610.
3. Law YM, et al.: MRI of pelvic floor dysfunction: review. *AJR* 2008; 191: 45–53.
4. Müller-Lissner S: The pathophysiology, diagnosis and treatment of constipation. *Dtsch Arztebl Int* 2009; 106(25): 424–32.

Dr. med. Kirsten Holsteg
Prof. Dr. med. Michael Stehling
 Institut für Bildgebende Diagnostik
 Strahlenberger Str. 110 – AlphaHaus
 63067 Frankfurt am Main/Offenbach, Germany
 holsteg@bilddiagnostik.de

Conflict of interest statement

The authors declare that no conflict of interest exists according to the Guidelines of the International Committee of Medical Journal Editors.

Potential for Dependence

The dependence-forming potential of substances that irritate the colon is, in my opinion, underrated. Someone who has aided his bowel movements with these substances for a few weeks is hardly likely to achieve normal digestion without laxatives. The lowered frequency of bowel movements—notably reduced after laxative intake has ceased—is not something that a person with constipation is likely to accept, given that reduced bowel motility made him or her reach for the laxatives in the first place. If normal bowel actions are achieved at all in people with constipation who have used laxatives for a number of years, this will happen only after months of good practice and high motivation on the patient's part, in spite of concerted application of all the methods described in Müller-Lissner's article.

Hypokalemia, which is described as a rare event if laxatives are used as per the instructions, is nevertheless a common finding when not serum but whole blood is analyzed for potassium. I personally have rarely observed potassium deficiency in serum. However, on analyzing whole blood, such a finding is a lot more common—especially in people with constipation, and then particularly in those taking laxatives. The simplest method to treat constipation was not mentioned: almost all people with constipation are deficient in magnesium—another measurement that becomes obvious in whole-blood analysis.

Because laxatives result not only in potassium deficiency in the smooth intestinal muscle cells but also in reduced magnesium concentrations, potassium substitution (only when deficient in serum and/or whole blood) and administration of magnesium at a high dosage (300–900 mg magnesium ions) is almost always successful. Even if no magnesium deficiency is present, magnesium is effective owing to its saline effects at high dosages. Last, but not least, magnesium salts ingested as a fizzy drink/granulate have a much more agreeable flavor than Glauber's salt or Epsom salts.

DOI: 10.3238/arztebl.2009.0765b

REFERENCES

1. Müller-Lissner S: The pathophysiology, diagnosis and treatment of constipation. *Dtsch Arztebl Int* 2009; 106(25): 424–32.

Dr. med. Volker Schmiedel
 Habichtswald-Klinik Werner Wilhelm Wicker KG
 Wigandstr. 1, 34131 Kassel
 schmiedel@habichtswaldklinik.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the Guidelines of the International Committee of Medical Journal Editors.

A Very Welcome Article

Müller-Lissner's CME article on constipation, a symptom with a high prevalence, is most welcome. The author describes diagnostic and therapeutic options in

an easy to read format, which is sure to improve the current state of knowledge. I wish to make some additions from my own clinical perspective.

- It goes without saying that constipation can be reliably diagnosed with the aid of the Rome III criteria. However, their practical implementation in the real life scenario is difficult because these criteria cannot always be elicited from patients in everyday clinical practice.
- Recommending selective serotonin reuptake inhibitors (SSRIs) as an alternative to tricyclic antidepressants results in constipation less often but is not guaranteed to prevent constipation either. Individual AWMF (Arbeitsgemeinschaft der Wissenschaftlichen Medizinischen Fachgesellschaften e.V.; Association of the Scientific Medical Societies in Germany, reg. assoc.) guidelines list constipation as one possible side effect for SSRIs too. From the pathophysiological point of view, treatment with SSRIs leads to inhibition of gastrointestinal enterochromaffin serotonergic receptors in the enteric nervous system, with simultaneously reduced peristalsis of the colon and lowered secretory capacity.
- From the perspective of pain and palliative medicine, opioid induced constipation should be given more attention as a common reason for patients to stop pain therapy that is otherwise reliable and effective. Constipation prophylaxis is essential when opioid treatment is started.
- The complete compendium of the different therapeutic options and recommendation of certain laxatives with a favorable ratio of effectiveness to side effects is to be welcomed. In the setting of pain and palliative medicine, using different drug mechanisms of action in an escalating stepwise scheme has proved effective. Such a stepwise scheme could be evaluated in other groups of patients too.

DOI: 10.3238/arztebl.2009.0765c

REFERENCES

1. Cash BD, Chey WD: The role of serotonergic agents in the treatment of patients with primary chronic constipation. *Aliment Pharmacol Ther* 2005; 22(11–12): 1047–60.
2. Sittig HB: *Kursbuch Palliative Care*. Unimed 2009: 205–17.
3. Schenk M: *Multimodale Tumorschmerztherapie*. Unimed Verlag 2009: 84–8.
4. Müller-Lissner S: The pathophysiology, diagnosis and treatment of constipation. *Dtsch Arztebl Int* 2009; 106(25): 424–32.

Dr. med. Stefan Wirz

Anästhesie, Interdisziplinäre Intensivmedizin, Schmerztherapie, Palliativmedizin
Katholisches Krankenhaus im Siebengebirge
Schülgenstr. 15, 53604 Bad Honnef, Germany
stefan.wirz@cura.org

Conflict of interest statement

The author has reported associations with the companies Mundipharma, Wyeth, Cephalon, and Norgive.

An Excellent Article

From the perspective of mainstream medicine, the author has summarized the topic in a didactically and thematically excellent way.

I wish to add the following facts, which deserve consideration:

“Abnormal intrinsic motility” is included among the pathophysiological components of chronic constipation. In order to activate or facilitate such abnormal motility, measures from conservative physical therapy or complementary medicine are available that all exert an influence on the autonomous enteric nervous system.

Such measures include colon treatment (5-point treatment according to Vogler, Kraus) and connective tissue massage as reflex zone therapy, in the sense of utilizing cutaneovisceral reflexes, as well as reflex zone therapy of the feet. The existence of reflex zones on the feet has not been neurophysiologically confirmed or proved, but last year, linked reflexes between the plantar surface and the knee were found in cats (1).

The empirical efficacy of plantar reflexology in influencing motility disorders of the gastrointestinal tract has been observed for a long time. We are currently researching this influence in a scientific study. There are data to show that reflex zone therapy of the feet improves the local circulation in the intestine when treating the zone that is associated with the colon (2).

In the context of complementary and extended complementary therapies and naturopathic treatments, moist, hot compresses on the abdominal region and osteopathic (visceral) techniques to stimulate motility and to accelerate delayed motility also deserve mention.

Patients with constipations should have the option of receiving these autoregulatory interventions.

DOI: 10.3238/arztebl.2009.0766a

REFERENCES

1. Rudomin P, et al.: Changes in synaptic effectiveness of myelinated joint afferents during capsaicin-induced inflammation of the footpad in the anesthetized cat. *Exp Brain Res* 2008; 187: 71–84.
2. Mur E, et al.: Influence of reflex zone therapy of the feet on intestinal blood flow measured by color Doppler sonography. *Fokumed* 2001; 8: 86–9.
3. Müller-Lissner S: The pathophysiology, diagnosis and treatment of constipation. *Dtsch Arztebl Int* 2009; 106(25): 424–32.

Prof. Dr. med. Christine Uhlemann

Universitätsklinikum Jena
Klinik Innere Medizin II, Postfach 07740 Jena, Germany
UKJ-KIM2@med.uni-jena.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the Guidelines of the International Committee of Medical Journal Editors.

Confusion

As “appropriate therapeutic management” for opioid induced constipation, the author names a combination of oxycodone with naloxone, or methylnaltrexone, and finally (!) laxatives. I have several comments:

- Every textbook stresses the importance of routine early administration of laxatives when opioid treatment is started, in order to treat any constipation—a common side effect of opioid treatment—consistently from the outset or even to avoid its development completely (1). Laxatives should therefore be named in the first place among the recommendations.
- Changing the way of application or opioid rotation often results in the desired success (2).
- No mention was made of constipation as an important gastrointestinal symptom in patients with advanced tumor disease and the differentiation from chronic subileus/ileus resulting in modified therapeutic consequences. Lubricant laxatives—often used in palliative care—are not mentioned at all.
- Oxycodone/naloxone in the treatment of severe acute (especially postoperative) and chronic pain are justifiably of great importance as they cause little constipation in clinical practice, but the licensed daily dosage is restricted to 80 mg oxycodone/40 mg naloxone. In patients with severe pain, this is not sufficient in many cases. Further, oxycodone is not the suitable opioid for all patients.
- The combination tilidine/naloxone at WHO step II is not mentioned at all, although it has been on the market for years and bears almost no risk of causing constipation (3).
- Methylnaltrexone is expensive and licensed only if conventional laxative therapy fails, so that a more differentiated discourse and weighting is necessary. Within the footnote, the emphasis on methylnaltrexone together with oxycodone/naloxone as the “only measures that have been shown to be effective in randomized, controlled studies” serves to confuse readers rather than offering a rational basis for decisions.

DOI: 10.3238/arztebl.2009.0766b

REFERENCES

1. Bausewein C: Arzneimitteltherapie in der Palliativmedizin. 1. Deutsche Aufl., 2005: 15ff
2. Bausewein C: Arzneimitteltherapie in der Palliativmedizin. 1. Deutsche Aufl., 2005: 173
3. Wörz R: Differenzierte medikamentöse Schmerztherapie. 2. edition., 2001: 190
4. Müller-Lissner S: The pathophysiology, diagnosis and treatment of constipation. Dtsch Arztebl Int 2009; 106(25): 424–32.

Dr. med. Ingo Zerbe
St. Josef-Krankenhaus
Koblenzer Str. 23
54411 Hermeskeil, Germany
i.zerbe@t-online.de

Conflict of interest statement

The author declares that no conflict of interest exists according to the Guidelines of the International Committee of Medical Journal Editors.

In Reply:

I thank my colleagues for their interesting correspondence. Chronic constipation is a very “old” topic that has been the subject of countless hypotheses—many unproved, some proved, and numerous proved wrong (1). A CME article in *Deutsches Ärzteblatt* is restricted—mainly for reasons of space—to points 2 and 3. I have actually never come into contact with untested procedures, and I omitted these for this reason. Uhlemann mentions reflex therapy of the soles of the feet, for which no clinical proof of efficacy exists.

Schmiedel rightly writes that patients with constipation who have enjoyed the advantages of laxative therapy do not develop normal behavior at stool once the medication is stopped. He concludes that laxatives have the potential to cause dependence. Since, however, laxatives represent symptomatic and not causative treatment of constipation, constipation is not likely to disappear after longer term laxative use—in the same way that chronic heart failure is not cured after medical treatment. But even so, to my knowledge nobody has ever claimed that ACE inhibitors have the potential to cause drug dependence. The potassium concentration is always lower in plasma than in serum. This is not associated with constipation or laxatives. Magnesium is a poorly absorbed ion that can be used as a saline laxative (see Table 2 in my article). There is no reason to assume that this depends on the blood concentration of magnesium. Whether magnesium is the easiest method to treat constipation is debatable, or, rather, for the patients to decide.

Wirz comments, correctly, that the Rome criteria are not perfect. Currently, however, I am not aware of a better instrument that is available. I cannot follow the remarks on constipation during treatment with SSRIs, nor can I find these in the cited literature. Serotonin is assumed to have a prokinetic effect via 5-HT₄ receptors, and blockade of the 5-HT₃ receptors can result in severe constipation. Inhibition of serotonin reuptake from the synaptic gap would thus only result in an improvement of the constipation, but such an effect is not clinically relevant. Opiate induced constipation is undoubtedly an important topic within pain therapy, but it affects only a proportion of patients with constipation and thus did not merit any more space in the article.

This is underlined by Zerbe’s letter. The items in my list of possible interventions in opiate-induced constipation caused by opiates were not ranked by order of importance. Comparisons between them are lacking; and as far as I am aware the same is true for data on tilidine/naloxone. The fact that opiate antagonists as a therapeutic principle are not mentioned in the medical textbooks is likely to be due to the fact that the relevant studies were published only in 2008 (2, 3).

Holsteg mentions MRI defecography, for which some literature exists. However, I did not find the cited methodological comparison among the 6 articles by Weishaupt that were listed in PubMed. On comparing conventional defecography and defecography obtained

by open magnet MRI—in a sitting position, not a lying position—the authors conclude that in diagnosing rectal intussusception, defecography is still the method of choice (4).

DOI: 10.3238/arztebl.2009.0767

REFERENCES

1. Müller-Lissner SA, Kamm MA, Scarpignato C, Wald A: Myths and misconceptions about chronic constipation. *Amer J Gastroenterol* 2005; 100: 232–42.
2. Thomas J, Karver S, Cooney GA, Chamberlain BH, Watt CK, Slatkin NE, Stambler N, Kremer AB, Israel RJ: Methylnaltrexone for opioid-induced constipation in advanced illness. *N Engl J Med* 2008; 358: 2332–43.
3. Simpson K, Leyendecker P, Hopp M, et al.: Fixed-ratio combination oxycodone/naloxone compared with oxycodone alone for the relief of opioid-induced constipation in moderate to severe noncancer pain. *Curr Med Res Opin* 2008; 24: 3503–12.

4. Dvorkin LS, Hetzer F, Scott SM, Williams NS, Gedroyc W, Lunniss PJ: Open-magnet MR defaecography compared with evacuation proctography in the diagnosis and management of patients with rectal intussusception. *Colorectal Dis* 2004; 6: 45–53.
5. Müller-Lissner S: The pathophysiology, diagnosis and treatment of constipation. *Dtsch Arztebl Int* 2009; 106(25): 424–32.

Prof. Dr. med. Stefan Müller-Lissner
Park-Klinik Weissensee
Schönstr. 80, 13086 Berlin, Germany
mueli@park-klinik.com

Conflict of interest statement

The author declares that no conflict of interest exists according to the Guidelines of the International Committee of Medical Journal Editors.