Published in final edited form as:

Child Youth Care Forum. 2009 June 1; 38(3): 151–160. doi:10.1007/s10566-009-9074-x.

Parent Depression and Child Anxiety: An Overview of the Literature with Clinical Implications

Christina J.M. Colletti¹, Rex Forehand¹, Emily Garai¹, Aaron Rakow¹, Laura McKee², Jessica M. Fear³, and Bruce E. Compas³

¹ University of Vermont

² University of North Carolina-Chapel Hill

³ Vanderbilt University

Abstract

The association of parental depression with child anxiety has received relatively little attention in the literature. In this paper we initially present several reasons for examining this relationship. We then summarize the empirical support for a link between these two variables. Finally, we discuss directions for future research and clinical implications of an association of parental depression with child anxiety.

Introduction

Major depressive disorder (MDD) represents a significant public health concern. National Comorbidity Study-Replication (NCS-R; Kessler et al., 2003) data indicate that the lifetime and 12-month prevalence rates for depression for 2001 were 16.2% and 6.6%, respectively. The National Epidemiologic Survey on Alcoholism and Related Conditions (NESARC; Hasin, Goodwin, Stinson, & Grant, 2005) provided similar lifetime (13.23%) and 12-month (5.28%) prevalence estimates. Based on these estimates, approximately 32 to 35 million adults are projected to be impacted by depression during their lifetime. Furthermore, the cost of depression in the United States due to lost earnings, treatment, and other factors is approximately \$85 billion annually (Greenberg et al., 2003).

Epidemiological data indicate median and mean recovery times of approximately six and twelve weeks, respectively (Kendler, Walters, & Kessler, 1997). Despite the short duration of episodes, residual depressive symptoms have been shown to persist following remission from a depressive episode in as many as one-third of individuals experiencing depression (Kennedy, Abbott, & Paykel, 2003; Paykel, 1998). Furthermore, longitudinal studies examining the course of MDD indicate that the disorder is highly recurrent. The median number of episodes across all individuals with a history of depression is seven, with approximately 75% to 85% of individuals experiencing more than one episode in their lifetime (Belsher & Costello, 1988; Boland & Keller, 2002).

Impact of Parent Depression on Children

Studies from the last few decades indicate that children and adolescents living with a depressed caregiver are at increased risk for a wide range of emotional and behavioral problems during the preschool, middle childhood, and adolescent years (for a review, see Goodman & Tully,

2006). Substantial research suggests that parental depression is associated with both offspring internalizing and externalizing problems (e.g., Elgar, Mills, McGrath, Waschbusch, & Brownridge, 2007). Although parental depression appears to have non-specific associations across these two types of child problems, and with disorders from within each of these broadband categories (see reviews by Goodman & Tully, 2006; Shanahan, Copeland, Costello, & Angold, 2008), recent investigators (McKee, Colletti, Rakow, Jones, & Forehand, 2008; McMahon, Grant, Compas, Thurm, & Ey, 2003; Shanahan et al., 2008) have emphasized the importance of examining whether particular risk factors, such as parental depression, are associated with specific child outcomes.

When broadband internalizing outcomes have been differentiated into depression or anxiety, the former disorder has received far more attention and support in empirical studies than the latter disorder (see reviews by Goodman, 2007; Goodman & Tully, 2006). This attention to the link between parent and child depression is not surprising, as heritability and environmental factors (e.g., parental modeling) may directly account for the transmission of depression from parent to child (for a review, see Goodman & Tully, 2006). Although the focus on depression outcomes for children of depressed caregivers is logical given our current understanding of the mechanisms of depression risk transmission (for reviews, see Goodman & Gotlib, 1999; Goodman & Tully, 2006), it is equally important that we examine the association of parental depression with child anxiety, the other primary type of internalizing problem. The following three arguments suggest that additional research should more closely examine the association of parental depression and child anxiety.

First, depression and anxiety in both children and adults are frequently comorbid, suggesting that parental depression may be associated with child anxiety in addition to child depression. More specifically, high comorbidity rates, ranging from 30% to 75%, have been reported for children and adolescents (e.g., Angold & Costello, 1993; Clarke, Smith, Neighbors, & Skerlec, 1994; Pilowsky et al., 2006). These comorbidity estimates substantially exceed the rate expected by chance (Williamson, Forbes, Dahle, & Ryan, 2005).

Second, multiple lines of research suggest that anxiety is a precursor for the development of depression in some children: (1) the mean age of children with anxiety disorders is younger than that for children with depressive disorders (e.g., Weissman et al., 2006); (2) most children and adolescents with comorbid depression and anxiety experience anxiety symptoms before they experience depressive symptoms (e.g., Andrade et al., 2003; Kessler et al., 2003; see Moffitt et al., 2008, for an exception); and (3) early anxiety symptoms predict later depressive symptoms in youth (e.g., Cole, Peeke, Martin, Truglio, & Seroczynski, 1998; Feng, Shaw, & Silk, 2008). These lines of evidence suggest that the association of parent and child depression may occur through child anxiety and give import to the task of examining child anxiety outcomes when parents have depression.

Third, our current knowledge of the degree to which anxiety disorders negatively impact youth requires that we examine parental depression as a possible risk factor for child anxiety. Anxiety disorders are among the most common psychiatric problems experienced by children, with prevalence rates ranging from 3% to 15% (Costello & Angold, 1995). Moreover, anxiety disorders appear to remain stable and problematic for youths throughout childhood and adolescence (McLeod, Wood, & Weisz, 2007). Many children with anxiety have difficulty with school performance and making and maintaining friendships, in addition to experiencing significant personal distress (for a review see Silverman & Ginsburg, 1998). Unlike depression, which is often chronic but episodic, anxiety disorders do not tend to remit with time but instead appear to continue into adulthood (Pine, Cohen, Gurley, Brook, & Ma, 1998). Together, these findings suggest that research identifying targets for prevention and treatment of anxiety disorders is necessary (Ginsburg & Schlossberg, 2002). If parental depression is a risk factor

for child anxiety, then increasing our understanding of the relation between these two disorders should lead to more effective prevention and intervention strategies for child anxiety.

The goal of this paper is to provide an overview of current evidence for an association of parental depression with child anxiety. We believe that the relation between these two variables has been understudied, leading to limited support for their association. As a consequence, the clinical implications of parental depression for anxiety in children have not been sufficiently recognized.

Data on the Association between Parent Depression and Child Anxiety

Although the extant literature is limited, both cross-sectional and retrospective studies indicate a link between parental depression and child anxiety. When comparisons have been made between children of parents with current or past clinical depression and children of never-depressed parents, the former group has been approximately two to six times more likely to have an anxiety disorder (e.g., Beidel & Turner, 1997; Fendrich, Warner, & Weissman, 1990; Hammen & Brennan, 2003; Ohannessian et al., 2005). Moreover, in one study, 7- to 12-year-old children of depressed parents were more likely to have an anxiety disorder than a depressive disorder, leading Beidel and Turner (1997) to conclude that “Offspring of depressed parents ... often present with anxious symptomatology” (p. 923).

It is important to note that not all cross-sectional studies find a link between parental depression and child anxiety. In two early studies, Biederman, Rosenbaum, Bolduc, Faraone, and Hirshfeld (1991) and Weissman, Leckman, Merikangas, Gammon, and Prusoff (1984) did not find support for the hypothesized associations between parental depression diagnoses and child anxiety diagnoses. More recently, Biederman and colleagues (Biederman et al., 2001) found support for an association between parental depression and child social phobia but not other anxiety disorders. Even if the data pointed to a consistent relationship between parent depression and child anxiety, reliance on cross-sectional and retrospective data to reach conclusions about these two variables is inherently flawed, as conclusions about the ordering or sequencing of the relationship cannot be reached. Data from longitudinal family studies can begin to provide findings more consistent with causality (Cole & Maxwell, 2003; Loeber & Farmington, 1994) and can address the possible sequencing effect, as a shared genetic diathesis manifests in childhood as anxiety and in early and late adolescence as depression (Silberg, Rutter, & Eaves, 2001).

Most (e.g., Lieb, Isensee, Hofler, Pfister, & Wittchen, 2002; Weissman, Warner, Wickramaratne, Moreau, & Olfson, 1997; Wickramaratne & Weissman, 1998), but not all (Biederman et al., 2006), of the existing longitudinal studies demonstrate that offspring of depressed parents are at increased risk for developing anxiety disorders. For example, findings from a longitudinal examination of psychopathology outcomes among offspring of depressed and non-depressed caregivers (Weissman et al., 2006b) indicated that offspring of depressed caregivers have a threefold increased risk of developing an anxiety disorder that is not otherwise explained by comorbid parental anxiety. In a study using the same sample but including three generations, Weissman and colleagues (2005) found higher rates of anxiety disorders (45%) than depressive disorders (30%) among grandchildren (M age = 12 years) when the previous two generations had experienced major depression. One interpretation of these findings offered by Weissman et al. (2005) is that anxiety during late childhood and early adolescence may be viewed as a precursor for later depression in adolescents and young adults. This is congruent with our earlier argument that anxiety precedes depression for most children.

The most rigorous way to examine the relationship between parent depression and child anxiety is by experimental manipulation of parent depression through medication or psychosocial intervention. There is some *limited* support for the treatment of parental depression alleviating

children's mental health problems (e.g., Weissman et al., 2006a; for a review, see Gunlicks & Weissman, 2008). Unfortunately, anxiety disorders or symptoms have not been examined individually (i.e., constructs such as internalizing symptoms have been examined), random assignment of families to treatment and control conditions has not occurred, or a sufficient sample size to allow statistical analysis of the data has not been included. Thus, current intervention research involving experimental manipulation has not yet been conducted to examine the association of parent depression and child anxiety. In summary, some limited cross-sectional, retrospective, and longitudinal data support an association between parent depression and child anxiety.

Directions for Future Research

Existing data *suggest* there may be a link between parent depression and child anxiety. Future research needs to address a number of areas in order for more definitive conclusions to be reached. These areas include the following: (a) measurement; (b) experimental design; and (c) the study of mediators and moderators.

First, in order to more accurately measure parent depression and child anxiety, it may be necessary to dismantle one or both constructs, as well as address their overlap. Parental depression consists of a number of parameters, including chronicity, severity, and timing of onset in a child's life. With rare exception (e.g., Hammen & Brennan, 2003), depression parameters such as these have been ignored and either a dichotomous measurement (depressed vs. non-depressed) or depressive symptoms have been examined. By refining the measurement of depression, unique relationships between specific parameters of parental depression and child anxiety may emerge. Similarly, with regard to child anxiety, specific types of anxiety disorders (e.g., phobias, social anxiety) should be examined (Shanahan et al., 2008). Biederman et al. (2001) found that parental depression may be related to some, but not all, anxiety disorders. Assessing specific anxiety disorders will increase the sensitivity of the outcome measure, increasing the likelihood of detecting parent depression-child anxiety associations.

With regard to construct overlap, an emerging literature has begun to identify alternatives to the precursor explanation outlined earlier for the comorbidity of anxiety and depression. Briefly, current measures of anxiety and depression may be tapping a broader negative affectivity construct that underlies both disorders (e.g., Mineka et al., 1998). Additional research is needed to clarify whether anxiety and depression are unique constructs or merely overlapping phenotypic expressions of the same underlying disorder. Alternatively, anxiety and depression may not be sufficiently differentiated in children to assess with existing measurement instruments, and a new generation of instruments may be required to more precisely determine the relation between depression and anxiety in both children and adults.

Second, with regard to experimental design, studies should control for parental anxiety and child depression. Without controlling for the symptoms of these disorders, relationships between parental depression and child anxiety may be explained by these comorbid conditions. For example, Biederman et al. (2006) controlled for parent panic disorder and did not find a relationship between parent depression and child anxiety. When comparing their findings to those of other investigators who found a significant association but did not control for comorbid parent anxiety disorders, Biederman et al. (2006) noted that the comorbid anxiety disorders may have accounted for the difference in findings.

Third, longitudinal and experimental designs must be utilized so that conclusions more consistent with causality may be reached. With longitudinal designs, baseline levels of child anxiety can be controlled in data analyses, thereby facilitating assessment of change in anxiety

over time (Cole & Maxwell, 2003). However, experimental manipulation of parental depression will be required to reach the strongest conclusions about causality.

Once a relation between parent depression and child anxiety has been established, attention can turn to mediators and moderators of this relation. For example, an environmental factor, parenting, may be such a mediator, as several parenting behaviors identified as characteristic of depressed parents (e.g., aversiveness, intrusiveness, withdrawal) (Lovejoy, Graczyk, O'Hare, & Neuman, 2000) *also* have been implicated in the development of anxiety problems in children (Ginsburg & Schlossberg, 2002; McLeod, Wood, & Weisz, 2007). Research also suggests the potential role of other environmental factors such as interparental violence (Shanahan et al., 2007). However, attention cannot focus solely on environmental factors, as genetics may well play a role.

Goodman and Tully (2006) noted that the "heritability of depression may not be specific to depression" (p. 253). Congruent with this notion, Dick, Rose, and Kaprio (2006) argued that genetic influences can be "expressed as different phenotypes at different developmental stages" (p. 225). Thus, the genetic transmission of parental depression to offspring *could* be expressed as anxiety in the pre-adolescent years and as depression in the teenage years. Other research suggests that genetic risk factors for depression and anxiety, including generalized anxiety disorder, are highly correlated (Kendler, Gardner, Galz, & Pedersen, 2007). All of these findings suggest that genetic factors may account for a link between parent depression and child anxiety. Furthermore, the interaction of genetic and environmental factors should be considered (Moffitt, Caspi, & Rutter, 2005). For example, it may well be that neither parenting nor genes alone can account for the relation between parent depression and child anxiety whereas the interaction of these two variables does account for this relation.

Beyond mediators, moderators also should be identified. For example, several studies suggest that girls are more sensitive to family stressors like parental depression than boys (Crick & Zahn-Waxler, 2003; Davies & Windle, 1997; Thomas & Forehand, 1991). Examinations of potential moderators, such as child age and gender, family context (e.g., presence of marital conflict), and neighborhood context (e.g., low vs. high risk) will inform us for whom and under what conditions parental depression leads to child anxiety.

Implications for Clinical Practice

Assuming a relation between parent depression and child anxiety, there are several important implications for clinical practice. First, such a relation suggests that when a parent presents with depression, there is a need to evaluate children in the family not only for depression but also for anxiety, as early intervention may decrease symptoms of both disorders along with the distress associated with them (Kowalenko et al., 2005; see Hirshfeld-Becker & Biederman, 2002). Second, when an adolescent client presents with depression and her or his parent has a history of depression, clinicians again should assess for anxiety in the adolescent, as the depression may be secondary to distress associated with chronic anxiety. Third, if a child presents clinically with anxiety, parent depression deserves consideration as a stressor contributing to the anxiety. Furthermore, and of importance, parental depression is related to treatment outcomes for child anxiety. For example, Berman, Weems, Silverman, and Kurtines (2000) found that higher levels of parental depressive symptoms at baseline were associated with poorer treatment outcomes for child anxiety when intervention included both a focus on the child and parental involvement. Southam-Geraw, Kendall, and Weersing (2001) found similar results for child anxiety outcome when the focus of treatment was on the child with some parental collaboration. Thus, to maximize effective treatment for child anxiety, parental depression should be assessed and, if necessary, targeted as part of the intervention.

Fourth, as several lines of research studies suggest that anxiety precedes child depression in some children, there is a need not only for intervention for anxiety when a parent is depressed but also for prevention of future depression in children with anxiety disorders. Although empirically supported treatments for anxiety and depression share many components (e.g. psychoeducation, cognitive restructuring) (Norton, Hayes, & Hope, 2004), they also differ in important ways. For example, exposure is an important component of the intervention package only for child anxiety treatment (see Weisz, 2004).

As another example, there currently appear to be trends in opposite directions regarding the role of parents in intervention for child anxiety versus prevention of child depression. Although early research indicated that inclusion of parents was beneficial in the treatment of child anxiety (Barrett, Dodds, & Rapee, 1996; Silverman, Kurtines, Ginsburg, Weems, Lampkin, & Carmichael, 1999a; Silverman, Kurtines, Ginsburg, Weems, Rabian, & Serafini, 1999b), the outcome in one recent treatment study suggests that parents do not necessarily need to be included in the child's treatment sessions for change to occur. Rather, the focus is on intervention with the child with occasional parent collaboration in the form of psychoeducation (Kendall, Hudson, Gosch, Flannery-Schroeder, & Suveg, 2008). In contrast, when prevention programs for child depression are considered, early well controlled research focused on the child or adolescent with occasional parent collaboration (e.g., Clarke, Hawkins, Murphy, Sheeber, Lewinsohn, & Seeley, 1995); however, several groups of researchers have recently called for the inclusion of parents (Compas, Keller, & Forehand, in press; Collins & Dozios, 2008; Horowitz & Garber, 2006). For example, Horowitz and Garber (2006) noted that prevention programs have been overly child focused and called for increased parental involvement, and Collins and Dozios (2008) concluded that current evidence suggests that including parents in prevention programs improves their outcomes. In sum, parents have played different roles in intervention for child anxiety compared with prevention of child depression, suggesting that different approaches may be required when treating anxiety and preventing depression.

Fifth, as we noted earlier, the same parenting behaviors (e.g., withdrawal) appear to be associated with depression in parents and anxiety in children; these behaviors suggest promising targets for intervention. Targeting the parenting behaviors of depressed parents that are specifically linked to anxiety disorders in children should increase the precision and effectiveness of interventions for child anxiety when a parent has depression. Finally, an awareness of the potential link between parent depression and child anxiety should increase our sensitivity as clinicians to the broad array of difficulties children can experience when growing up in families characterized by parental depression.

Additional Readings

The referenced articles by Goodman (2007), Goodman and Tully (2006), and Cummings and Davies (1994) provide more in-depth reviews of parental depression and child behavioral and emotional problems.

Acknowledgments

The idea for this paper emerged from work on two parallel NIMH grants, R01MH069940 and R01MH069928, examining a family cognitive-behavioral preventive intervention for children of depressed parents.

References

Andrade L, Caraveo-Anduaga JJ, Berglund P, Bijl RV, De Graaf R, Vollebergh W, et al. The epidemiology of major depressive episodes: Results from the International Consortium of Psychiatric

- Epidemiology (ICPE) Surveys. *International Journal of Methods in Psychiatric Research* 2003;12:3–21. [PubMed: 12830306]
- Angold A, Costello EJ. Depressive comorbidity in children and adolescents: empirical, theoretical, and methodological issues. *American Journal of Psychiatry* 1993;150:1779–1791. [PubMed: 8238631]
- Barrett PM, Dadds MR, Rapee RM. Family treatment of childhood anxiety: A controlled trial. *Journal of Consulting and Clinical Psychology* 1996;64:333–342. [PubMed: 8871418]
- Beidel DC, Turner SM. At risk for anxiety: I. Psychopathology in the offspring of anxious parents. *Journal of the American Academy of Child and Adolescent Psychiatry* 1997;36:918–924. [PubMed: 9204669]
- Belsher G, Costello CG. Relapse after recovery from unipolar depression: A critical review. *Psychological Bulletin* 1988;104:84–96. [PubMed: 3043528]
- Berman SL, Weems CF, Silverman WK, Kurtines WK. Predictors of outcome in exposure-based cognitive and behavioral treatments for phobic and anxiety disorders in children. *Behavior Therapy* 2000;31:713–731.
- Biederman J, Petty C, Hirshfeld-Becker DR, Henin A, Faraone SV, Dang D, Jakubowski A, Rosenbaum JF. *Psychological Medicine* 2006;36:1141–1152. [PubMed: 16700966]
- Biederman J, Rosenbaum JF, Bolduc EA, Faraone SV, Hirshfeld DR. A high risk study of young children of parents with panic disorder and agoraphobia with and without comorbid major depression. *Psychiatry Research* 1991;37:333–348. [PubMed: 1891513]
- Biederman MD, Faraone SV, Hirshfeld-Becker DR, Friedman D, Robin JA, Rosenbaum JF. Patterns of psychopathology and dysfunction in high-risk children of parents with panic disorder and major depression. *American Journal of Psychiatry* 2001;158:49–57. [PubMed: 11136633]
- Boland, RJ.; Keller, MB. Course and outcome of depression. In: Gotlib, IH.; Hammen, CL., editors. *Handbook of depression*. New York, NY: Guilford Press; 2002. p. 43-60.
- Clarke DB, Smith MG, Neighbors BD, Skerlec LM. Anxiety disorders in adolescence: Characteristics, prevalence, and comorbidities. *Clinical Psychology Review* 1994;14:113–137.
- Clarke G, Hawkins W, Murphy M, Sheeber L, Lewinsohn PM, Seeley JR. Targeted prevention of unipolar depressive disorder in an at-risk sample of high school adolescents: A randomized trial of group cognitive intervention. *Journal of the American Academy of Child and Adolescent Psychiatry* 1995;34:312–321. [PubMed: 7896672]
- Cole DA, Maxwell SE. Testing mediational models with longitudinal data: Questions and tips in the use of structural equation modeling. *Journal of Abnormal Psychology* 2003;112:358–577.
- Cole DA, Peeke LG, Martin JM, Truglio R, Seroczynski AD. A longitudinal look at the relation between depression and anxiety in children and adolescents. *Journal of Consulting and Clinical Psychology* 1998;66:451–460. [PubMed: 9642883]
- Collins KA, Dozois DJA. What are the active ingredients in preventative interventions for depression? *Clinical Psychology: Science and Practice* 2008;15:313–330.
- Compas, BE.; Keller, G.; Forehand, R. Prevention of depression in families of depressed parents. In: Strauman, T.; Costanzo, PR.; Garber, J.; Abramson, LY., editors. *Preventing depression in adolescent girls*. Duke University Press; (in press)
- Costello, EJ.; Angold, A. Epidemiology. In: March, John S., editor. *Anxiety disorders in children and adolescents*. New York, NY: Guilford Press; 1995. p. 109-124.
- Crick NR, Zahn-Waxler C. The development of psychopathology in females and males: Current progress and future challenges. *Development and Psychopathology* 2003;15:719–742. [PubMed: 14582938]
- Cummings EM, Davies PT. Maternal depression and child development. *Journal of Child Psychology and Psychiatry* 1994;35:73–112. [PubMed: 8163630]
- Davies PT, Windle M. Gender-specific pathways between maternal depressive symptoms, family discord, and adolescent adjustment. *Developmental Psychology* 1997;22:657–668. [PubMed: 9232381]
- Dick DM, Rose RJ, Kaprio J. The next challenge for psychiatric genetics: Characterizing the risk associated with identified genes. *Annals of Clinical Psychiatry* 2006;18:223–231. [PubMed: 17162621]
- Elgar FJ, Mills RSL, McGrath PJ, Waschbusch DA, Brownridge DA. Maternal and paternal depressive symptoms and maladjustment: The mediating role of parent behavior. *Journal of Abnormal Child Psychology* 2007;35:943–955. [PubMed: 17577659]

- Fendrich M, Warner V, Weissman MM. Family risk factors, parental depression, and psychopathology in offspring. *Developmental Psychology* 1990;26:40–50.
- Feng X, Shaw DS, Silk JS. Developmental trajectories of anxiety symptoms among boys across early and middle childhood. *Journal of Abnormal Psychology* 2008;117:32–47. [PubMed: 18266484]
- Ginsburg GS, Schlossberg MC. Family-based treatment of childhood anxiety disorders. *International Review of Psychiatry* 2002;14:143–154.
- Goodman SH, Gotlib IH. Risk for psychopathology in the children of depressed mothers: A developmental model for understanding mechanisms of transmission. *Psychological Review* 1999;106:458–490. [PubMed: 10467895]
- Goodman, SH.; Tully, E. Depression in women who are mothers: An integrative model of risk for the development of psychopathology in their sons and daughters. In: Keyes, CLM.; Goodman, SH., editors. *Women and depression: A handbook for the social, behavioral, and biomedical sciences*. New York, NY: Cambridge University Press; 2006. p. 241–280.
- Goodman SH. Depression in mothers. *Annual Review of Clinical Psychology* 2007;3:107–135.
- Greenberg PE, Kessler RC, Birnbaum HG, Leong SA, Lowe SW, Berglund PA, et al. *Journal of Clinical Psychiatry* 2003;64:1465–1475. [PubMed: 14728109]
- Gunlicks ML, Weissman MM. Change in child psychopathology with improvement in parental depression: A systematic review. *Journal of the American Academy of Child and Adolescent Psychiatry* 2008;47:379–389. [PubMed: 18388766]
- Hammen C, Brennan P. Severity, chronicity, and timing of maternal depression and risk for adolescent offspring diagnoses in a community sample. *Archives of General Psychiatry* 2003;60:253–258. [PubMed: 12622658]
- Hammen, C.; Rudolph, KD. Childhood mood disorders. In: Mash, EJ.; Barkley, RA., editors. *Child psychopathology*. New York, NY: Guilford Press; 2003. p. 233–278.
- Hasin DS, Goodwin RD, Stinson FS, Grant BF. Epidemiology of major depressive disorder: Results from the National Epidemiological Survey on Alcoholism and Related Conditions. *Archives of General Psychiatry* 2005;62:1097–1106. [PubMed: 16203955]
- Hirshfeld-Becker D, Biederman J. Rationale and principles for early intervention with young children at risk for anxiety disorders. *Clinical Child & Family Psychology Review* 2002;5:161–172. [PubMed: 12240705]
- Horowitz JL, Garber J. The prevention of depressive symptoms in children and adolescents: A meta-analytic review. *Journal of Consulting and Clinical Psychology* 2006;74:401–415. [PubMed: 16822098]
- Kendall PC, Hudson JL, Gosch E, Flannery-Schroeder E, Suveg C. Cognitive-behavioral therapy for anxiety disordered youth: A randomized clinical trial evaluating child and family modalities. *Journal of Consulting and Clinical Psychology* 2008;76:282–297. [PubMed: 18377124]
- Kendler KS, Walters EE, Kessler RC. The prediction of length of major depressive episodes: Results from an epidemiological sample of female twins. *Psychological Medicine* 1997;27:107–117. [PubMed: 9122291]
- Kendler KS, Gardner CO, Gatz M, Pedersen NL. The sources of co-morbidity between major depression and generalized anxiety disorder in a Swedish national twin sample. *Psychological Medicine* 2007;37:453–462. [PubMed: 17121688]
- Kennedy N, Abbott R, Paykel ES. Remission and recurrence of depression in the maintenance era: Long-term outcome in a Cambridge cohort. *Psychological Medicine* 2003;33:827–838. [PubMed: 12877397]
- Kessler RC, Berglund P, Demler O, Jin R, Koretz D, Merikangas KR, et al. The epidemiology of major depressive disorder: Results from the National Comorbidity Survey Replication (NCS-R). *Journal of the American Medical Association* 2003;289:3095–3105. [PubMed: 12813115]
- Kowalenko N, Rapee RM, Simmons J, Wignall A, Hoge R, Whitefield K, et al. Short-term effectiveness of a school-based early intervention program for adolescent depression. *Clinical Child Psychology and Psychiatry* 2005;10:493–507. Special Issue: Tenth Anniversary Edition.
- Lieb R, Isensee B, Hofler M, Pfister H, Wittchen H. Parental major depression and the risk of depression and other mental disorders in offspring: A prospective-longitudinal community study. *Archives of General Psychiatry* 2002;59:365–374. [PubMed: 11926937]

- Loeber &, Farrington DP. Problems and solutions in longitudinal and experimental treatment studies of child psychopathology and delinquency. *Journal of Consulting and Clinical Psychology* 1994;62:887–900. [PubMed: 7806719]
- Lovejoy MC, Graczyk PA, O'Hare E, Neuman G. Maternal Depression and parenting behavior: A meta-analytic review. *Clinical Psychology Review* 2000;20:561–592. [PubMed: 10860167]
- McKee LG, Colletti CJM, Rakow A, Jones DJ, Forehand RL. Parenting and child externalizing behaviors: Are associations specific or diffuse? *Aggression and Violent Behavior* 2008;13:201–215. [PubMed: 19122818]
- McLeod BD, Wood JJ, Weisz JR. Examining the association between parenting and childhood anxiety: A meta-analysis. *Clinical Psychology Review* 2007;27:155–172. [PubMed: 17112647]
- McMahon SD, Grant KE, Compas BE, Thurm AE, Ey S. Stress and psychopathology in children and adolescents: Is there evidence of specificity? *Journal of Child Psychology and Psychiatry* 2003;44:107–133. [PubMed: 12553415]
- Mineka S, Watson D, Clark LA. Comorbidity of anxiety and unipolar mood disorders. *Annual Review of Psychology* 1998;49:377–412.
- Moffitt TE, Caspi A, Rutter J. Strategy for investigating interactions between measured genes and measured environments. *Archives of General Psychiatry* 2005;62:473–496. [PubMed: 15867100]
- Moffitt TE, Harrington H, Caspi A, Kim-Cohen J, Goldberg D, Gregory AM, Poulton R. Depression and generalized anxiety disorder: Cumulative and sequential comorbidity in a birth cohort followed prospectively to age 32 years. *Archives of General Psychiatry* 2008;64:651–660. [PubMed: 17548747]
- Norton PJ, Hayes SA, Hope DA. Effects of a transdiagnostic group treatment for anxiety on secondary depression. *Depression and Anxiety* 2004;20:198–202. [PubMed: 15643648]
- Ohannessian CM, Hesselbrock VM, Kramer J, Kuperman S, Bucholz KK, Schuckit MA, Nurnberger JI. The relationship between parental psychopathology and adolescent psychopathology: An examination of gender patterns. *Journal of Emotional and Behavioral Disorders* 2005;13:67–76.
- Paykel ES. Remission and residual symptomatology in major depression. *Psychopathology* 1998;31:5–14. [PubMed: 9500681]
- Pilowsky DJ, Wickramaratne PJ, Rush JA, Hughes CW, Garber J, Mallow E, et al. Children of currently depressed mothers: A STAR*D ancillary study. *Journal of Clinical Psychiatry* 2006;67:126–136. [PubMed: 16426099]
- Pine DS, Cohen P, Gurley D, Brook J, Ma Y. The risk for early-adulthood anxiety and depressive disorders in adolescents with anxiety and depressive disorders. *Archives of General Psychiatry* 1998;55:56–64. [PubMed: 9435761]
- Shanahan L, Copeland W, Costello EJ, Angold A. Specificity of putative psychosocial risk factors for psychiatric disorders in children and adolescents. *Journal of Child Psychology and Psychiatry* 2008;49:34–42. [PubMed: 18181879]
- Silverman, WK.; Ginsburg, GS. Anxiety disorders. In: Ollendick, Thomas H.; Hersen, Michel, editors. *Handbook of child psychopathology*. New York, NY: Plenum Press; 1998. p. 239–268.
- Silverman WK, Kurtines WM, Ginsburg GS, Weems CF, Lumpkin PW, Carmichael DH. Treating anxiety disorders in children with group cognitive-behavioral therapy: A randomized clinical trial. *Journal of Consulting and Clinical Psychology* 1999a;67:995–1003. [PubMed: 10596522]
- Silverman WK, Kurtines WM, Ginsburg GS, Weems CF, Rabian B, Serafini LT. Contingency management, self-control, and education support in the treatment of childhood phobic disorders: A randomized clinical trial. *Journal of Consulting and Clinical Psychology* 1999b;67:675–687. [PubMed: 10535234]
- Southam-Gerow MA, Kendall PC, Weersing VR. Examining outcome variability: Correlates of treatment response in a child and adolescent anxiety clinic. *Journal of Clinical Child & Adolescent Psychology* 2001;30:422–436.
- Thomas AM, Forehand R. The relationship between paternal depressive mood and early adolescent functioning. *Journal of Family Psychology* 1991;4:260–271.
- Weissman MM, Warner V, Wickramaratne P, Moreau D, Olfson M. Offspring of depressed parents: 10 years later. *Archives of General Psychiatry* 1997;54:932–940. [PubMed: 9337774]

- Weissman MM, Wickramaratne PJ, Nomura Y, Warner V, Verdeli H, Pilowsky DJ, et al. Family at high and low risk for depression: A 3-generation study. *Archives of General Psychiatry* 2005;62:29–36. [PubMed: 15630070]
- Weissman MM, Wickramaratne P, Nomura Y, Warner V, Pilowsky D, Verdeli H. Offspring of depressed parents: 20 years later. *American Journal of Psychiatry* 2006b;163:1001–1008. [PubMed: 16741200]
- Weissman MM, Leckman JF, Merikangas KR, Gammon D, Prusoff BA. Depression and anxiety disorders in parents and children. *Archives of General Psychiatry* 1984;41:845–852. [PubMed: 6466043]
- Weissman MM, Pilowsky DJ, Wickramaratne PJ, Talata A, Wisniewski SR, Fava M, Hughes CW. Remissions in maternal depression and child psychopathology. *Journal of the American Medical Association* 2006a;295:1389–1398. [PubMed: 16551710]
- Weisz, JR. *Psychotherapy for children and adolescents: Evidenced-based treatments and case examples*. Cambridge, U.K.: Cambridge University Press; 2004.
- Wickramaratne PJ, Weissman MM. Onset of psychopathology in offspring by developmental phase and parental depression. *Journal of the American Academy of Child & Adolescent Psychiatry* 1998;37:933–942. [PubMed: 9841243]
- Williamson DE, Forbes EE, Dahl RE, Ryan ND. A genetic epidemiologic perspective on comorbidity of depression and anxiety. *Child and Adolescent Psychiatric Clinics of North America* 2005;14:707–726. [PubMed: 16171699]