Pretaporter, a *Drosophila* protein serving as a ligand for Draper in the phagocytosis of apoptotic cells

Takayuki Kuraishi^{1,7}, Yukiko Nakagawa², Kaz Nagaosa², Yumi Hashimoto¹, Takashi Ishimoto², Takeshi Moki², Yu Fujita¹, Hiroshi Nakayama³, Naoshi Dohmae³, Akiko Shiratsuchi^{1,2,4}, Naoko Yamamoto², Koichi Ueda⁴, Masamitsu Yamaguchi⁵, Takeshi Awasaki⁶ and Yoshinobu Nakanishi^{1,2,4,*}

¹Graduate School of Medical Science, Kanazawa University, Kanazawa, Ishikawa, Japan, ²Graduate School of Natural Science and Technology, Kanazawa University, Kanazawa, Ishikawa, Japan, ³Biomolecule Characterization Team, RIKEN, Wako, Saitama, Japan, ⁴Faculty of Pharmaceutical Sciences, Kanazawa University, Kanazawa, Ishikawa, Japan, ⁵Department of Applied Biology, Kyoto Institute of Technology, Kyoto, Kyoto, Japan and ⁶Department of Neurobiology, University of Massachusetts Medical School, Worcester, MA, USA

Phagocytic removal of cells undergoing apoptosis is necessary for animal development and tissue homeostasis. Draper, a homologue of the *Caenorhabditis elegans* phagocytosis receptor CED-1, is responsible for the phagocytosis of apoptotic cells in *Drosophila*, but its ligand presumably present on apoptotic cells remains unknown. An endoplasmic reticulum protein that binds to the extracellular region of Draper was isolated. Loss of this protein, which we name Pretaporter, led to a reduced level of apoptotic cell clearance in embryos, and the overexpression of *pretaporter* in the mutant flies rescued this defect. Results from genetic analyses suggested that Pretaporter functionally interacts with Draper and the corresponding signal mediators. Pretaporter was exposed at the cell surface after the induction of apoptosis, and cells artificially expressing Pretaporter at their surface became susceptible to Draper-mediated phagocytosis. Finally, the incubation with Pretaporter augmented the tyrosine-phosphorylation of Draper in phagocytic cells. These results collectively suggest that Pretaporter relocates from the endoplasmic reticulum to the cell surface during apoptosis to serve as a ligand for Draper in the phagocytosis of apoptotic cells. *The EMBO Journal* (2009) 28, 3868–3878. doi:10.1038/emboj.2009.343; Published online 19 November 2009
Subject Categories: differentiation & death
Keywords: apoptosis; *Drosophila*; endoplasmic reticulum; phagocytosis

Introduction

Throughout the life of multi-cellular organisms, cells of particular types die and are eventually eliminated in certain places in the body and at certain developmental stages under physiological and sub-physiological conditions (Cohen, 1991; Cleveland, 1996; Jacobson *et al*, 1997; Vaux and Korsmeyer, 1999). Such ‘unwanted cells’ are mostly induced to undergo apoptosis and subjected to phagocytic elimination (Wyllie *et al*, 1980). Prompt and selective phagocytosis of apoptotic cells is prerequisite for morphogenesis, establishment of tissue functions, tissue renewal, avoidance of diseases, and effective progress of tissue functions (Savill and Fadok, 2000; Liao, 2005; Nakanishi *et al*, 2009).

There are two genetically identified signalling pathways that lead to the induction of phagocytosis of dying cells in *Caenorhabditis elegans* (Reddien and Horvitz, 2004; Kinchen and Hengartner, 2005; Mangahas and Zhou, 2005; Lettre and Hengartner, 2006): one is made up of the proteins CED-2, CED-5, and CED-12, and the other consists of CED-1, CED-6, and CED-7. The pathways converge at CED-10, a small G-protein responsible for the rearrangement of cytoskeleton in phagocytes (Kinchen *et al*, 2005). The fact that all these *C. elegans* proteins possess counterparts in *Drosophila* and mammals (Lettre and Hengartner, 2006; Kinchen and Ravichandran, 2007) suggests that these two partially redundant signalling pathways are evolutionally conserved. However, the mode of action of those proteins still remains to be clarified, and there are missing components in the pathways. The selectivity in the recognition of apoptotic cells by phagocytes is due to the specific interaction between receptors of phagocytes and their ligands present at the surface of target cells (Lauber *et al*, 2004; Ravichandran and Lorenz, 2007). Presumably, there are two phagocytosis receptors in *C. elegans*, and CED-1 is likely the one located upstream of CED-6 and CED-7 (Zhou *et al*, 2001; Yu *et al*, 2006; Venegas and Zhou, 2007) whereas the other has not been found in genetic studies. Draper (Drpr), a *Drosophila* homologue of CED-1 (Callebaut *et al*, 2003), has been shown to act as a receptor in the phagocytosis of apoptotic cells (Freeman *et al*, 2003; Manaka *et al*, 2004), and more recently molecules that act with Drpr to accomplish the engulfment and subsequent processing of apoptotic cells in *Drosophila* phagocytes were reported (Cuttell *et al*, 2008; Kurant *et al*, 2008; Ziegenfuss *et al*, 2008). However, a molecule(s) present at the surface of apoptotic cells and recognized by Drpr is yet to be identified. The study by Venegas and Zhou (2007) has implied that the membrane phospholipid phosphatidylserine, the best-characterized phagocytosis marker in mammals (Fadok *et al*, 1998; Schlegel and Williamson, 2001), could be a ligand for CED-1. In contrast, we showed that phosphatidylserine is not required in the Drpr-mediated phagocytosis of apoptotic cells (Manaka *et al*, 2004). In this study, we searched for a phagocytosis marker(s) recognized by Drpr and identified an endoplasmic reticulum (ER) protein as a strong candidate.

*Corresponding author. Graduate School of Medical Science, Kanazawa University, Shizenken, Kakuma-machi, Kanazawa, Ishikawa 920-1192, Japan. Tel.: +81 76 234 4481; Fax: +81 76 234 4480; E-mail: nakanaka@kenroku.kanazawa-u.ac.jp

⁷Present address: Global Health Institute, EPFL, CH-1015 Lausanne, Switzerland

Received: 2 June 2009; accepted: 15 October 2009; published online: 19 November 2009

Results

Isolation of an ER protein that binds to the extracellular region of Drpr

In search of a Drpr ligand(s), we adopted a biochemical approach to isolate a protein(s) that binds to Drpr. An extracellular portion of Drpr, which contained all 15 epidermal growth factor-like repeats (Callebaut *et al*, 2003), was prepared as a protein fused to glutathione S-transferase (GST) (Supplementary Figure S1A) using a vector derived from insect virus because Drpr likely undergoes a post-translational modification (Manaka *et al*, 2004). This protein, which we operationally named Drpr-GST, specifically bound to cycloheximide-treated S2 cells, a cell line established from *Drosophila* embryonic cells (Supplementary Figure S1B): treatment with cycloheximide makes S2 cells undergo typical apoptosis (see Figure 5) and susceptible to Drpr-mediated phagocytosis (Manaka *et al*, 2004). Drpr-GST was then covalently conjugated with glutathione-Sepharose, and whole-cell lysates of apoptotic S2 cells were chromatographed on this affinity matrix. The bound materials were eluted and analysed by sodium dodecyl sulphate-polyacrylamide gel electrophoresis followed by staining with silver. The gel showed a band with a molecular mass of 47 kDa that was absent in the eluates from Sepharose containing GST alone (Figure 1A). A mass spectrometric analysis revealed that this protein, which we named Pretaporter (Prtp), encoded by the as-yet uncharacterized gene *CG1837* consists of 416 amino-acid residues, including the signal peptide at the N terminus, three thioredoxin-like domains in the middle, and an ER retention motif at the C terminus (Figure 1B). When the same eluates were analysed by western blotting with an

antiserum raised against recombinant Prtp, a distinct signal was obtained at the expected position (Figure 1C). A cell-free binding assay using recombinant proteins revealed that two-thirds of Prtp to the N terminus is sufficient for the binding to Drpr-GST (Figure 1D). From these results, we concluded that Prtp is a *Drosophila* protein that binds to the extracellular region of Drpr.

We next determined the expression during development and the sub-cellular localization of this newly identified protein. Prtp was expressed throughout the development of *Drosophila* as was Drpr (Figure 1E, and Supplementary Figure S2 for specificity of antibody) and in most cell types of embryos (Figure 1F) and wing discs (Figure 4C). A cytochemical analysis of dispersed embryonic cells showed that signals derived from Prtp overlapped with those of an ER marker (Figure 1G). These results indicated that Prtp is an ER protein expressed ubiquitously throughout development.

Reduced level of apoptotic cell phagocytosis in *prtp* null mutants

To assess a function for Prtp in the Drpr-mediated phagocytosis of apoptotic cells, we established fly lines deficient in its expression. A fly line that contains a P-element inserted proximal upstream of the translation start codon of *prtp* was subjected to P-element excision, and two mutations, *prtp*^{Δ1} and *prtp*^{Δ2}, which deleted a part of the first exon of *prtp*, were isolated (Figure 2A and Supplementary Figure S3A). We also obtained *prtp*^{G19}, in which the P-element had been precisely excised. The expression of *prtp*, either mRNA or protein, was undetectable in homozygous mutants of both *prtp*^{Δ1} and *prtp*^{Δ2}, whereas *prtp*^{G19} flies showed a normal level expression of *prtp* (Supplementary Figures S3B and

Figure 1 Isolation of Pretaporter (Prtp) as a Draper (Drpr)-binding endoplasmic reticulum (ER) protein. (A) Sodium dodecyl sulphate-polyacrylamide gel electrophoresis (stained with silver) of bound materials in chromatography of whole-cell lysates of S2 cells, which were induced to undergo apoptosis by the treatment with cycloheximide, on Sepharose conjugated with the indicated proteins. The arrowhead denotes the protein band brought to mass spectrometry. Shown at the bottom are magnified views of the squared areas. (B) Predicted domain structure of Prtp (shaded, hatched, and closed boxes denote signal peptide, thioredoxin-like domain (TLD), and ER retention motif, respectively). The regions of Prtp tested for the binding to Drpr in (D) are indicated. (C) Western blotting of the samples in (A) with anti-Prtp antibody. (D) Cell-free analysis of the binding of Prtp to the extracellular region of Drpr. The indicated maltose-binding protein (MBP)-fused proteins were incubated with Drpr-glutathione S-transferase (GST), pulled down, and analysed by western blotting with anti-GST or anti-MBP antibody. (E) Expression of Prtp and Drpr analysed by western blotting with lysates of whole animals at the indicated developmental stages. (F) Immunocytochemical analysis of Prtp expression with embryos and dispersed embryonic cells at stage 16. Bars, 100 μ m. (G) Sub-cellular localization of Prtp in an embryonic cell expressing yellow fluorescent protein containing the ER retention signal KDEL (LaJeunesse *et al*, 2004). Bar, 2 μ m.

Figure 2 Requirement of Pretaporter (Prtp) in apoptotic cell phagocytosis by haemocytes and glia. (A) Generation of *prtp* null mutants. The structure of the *Drosophila* genome near *prtp* (exons are indicated by boxes with protein-coding regions shaded) and the positions of deletions in mutant flies are schematically shown. After P-element excision with *CG1837^{BC00450}*, two fly lines with deletions in the first exon of *prtp*, *prtp^{Δ1}* ($\Delta 1$) (with a deletion in the region between nucleotide positions -45 and $+354$ with the transcription start site numbered as $+1$) and *prtp^{Δ2}* ($\Delta 2$) (with a deletion in the region from -45 to $+1170$), and one line, *prtp^{G19}* (*G19*), that lost the P-element with no alteration in *prtp* were isolated. (B) Distribution (left) and ratio (right) (dispersed cells were analysed) of green fluorescent protein (GFP)-positive haemocytes in embryos of the indicated fly lines. Data are from one of two independent experiments that yielded similar results. Bar, $20\ \mu\text{m}$. (C, D) Level of phagocytosis by GFP-positive haemocytes determined with embryos (C) or dispersed embryonic cells (D) of the indicated fly lines. Signals of GFP and terminal deoxynucleotidyltransferase-mediated dUTP nick end labelling (TUNEL) are seen in blue and brown, respectively. The arrowhead points to a haemocyte containing a TUNEL-positive nucleus. The inset in (C) is a magnified view of the squared area. Bars, $10\ \mu\text{m}$ (C) and $2\ \mu\text{m}$ (D). $*P < 0.01$ (versus *G19*). (E) Level of phagocytosis by Croquemort-expressing haemocytes in embryos of the indicated fly lines. $*P < 0.01$. NS, difference not significant. (F) Level of phagocytosis of latex beads by larval haemocytes of the indicated fly lines. (G) Distribution (left) and ratio (right) (dispersed cells were analysed) of Repo-positive glia in embryos of the indicated fly lines. Data are from one of two independent experiments that yielded similar results. Bar, $20\ \mu\text{m}$. (H, I) Level of apoptotic cell phagocytosis by glia in the indicated fly lines. Embryos were analysed for the number of *in situ* nick translation-positive cells in the vicinity of the ventral nerve cord (depicted with a black line in the left panel of G) where glia are enriched (H), and dispersed embryonic cells were examined for the ratio of cells positive for both Repo (seen in blue) and TUNEL (seen in brown) against total Repo-positive cells (I). Bar, $2\ \mu\text{m}$. $*P < 0.01$ (versus *G19*). Refer to Materials and methods section for genotypes of the fly lines analysed.

S3C). The expression of *drpr* was not influenced in these three fly lines (Supplementary Figures S3B and S3C). These results indicated that two null alleles of *prtp*, *prtp^{Δ1}* and *prtp^{Δ2}*, were generated. The fly line *prtp^{G19}* was used thereafter as a control in the analysis of the mutant alleles. Either mutant fly line was viable as a homozygote, and there was no defect in their development, motility, or fertility (data not shown). We first examined whether or not a loss of *prtp* expression influences the occurrence of apoptosis during embryogenesis, which becomes apparent at the developmental stage 11 (Abrams *et al*, 1993). Whole embryos at stages 10 and 11 were subjected to immunohistochemistry with anti-activated caspase-3 antibody as well as terminal deoxynucleotidyltransferase-mediated dUTP nick end labelling

(TUNEL) for the analysis of caspase activation and DNA fragmentation, respectively. Cells containing activated caspase or fragmented DNA were absent at stage 10 and became detectable around the anterior part of embryos at stage 11 of the control flies, and this was almost the same for the mutant flies (Supplementary Figure S4). These results indicated that apoptosis occurs normally in embryos lacking Prtp. We then determined the level of phagocytosis of apoptotic cells by haemocytes and glia in embryos, which we previously reported to accomplish Drpr-mediated phagocytosis (Manaka *et al*, 2004).

Haemocytes, which were labelled with green fluorescent protein (GFP) using a specific promoter, were widely distributed in embryos as previously reported (Brückner *et al*,

2004), and the distribution (left panel in Figure 2B) and population (right panel in Figure 2B) of haemocytes in embryos remained the same irrespective of the expression of *prtp*. When those embryos were subjected to TUNEL, some of the GFP-positive haemocytes contained positively stained nuclei besides their own (left panel in Figure 2C). We thus counted the cells positive for both GFP and TUNEL as those haemocytes that had phagocytosed apoptotic cells, and determined the ratio of phagocytosing haemocytes. Only haemocytes residing near the peripheral part of embryos were analysed because individual haemocytes were hardly discriminated in the middle part. The results showed that the level of phagocytosis in either *prtp*^{Δ1} or *prtp*^{Δ2} was almost half of that in the control *prtp*^{G19} (right panel in Figure 2C) (note that the level of phagocytosis in *prtp*^{G19} was equivalent to that in the fly line *w¹¹¹⁸*). To investigate haemocytes including those residing in other parts of embryos, we mechanically disrupted embryos and analysed resulting dispersed cells. As were observed with intact embryos, GFP-positive haemocytes with and without TUNEL-stained nuclei were present in the dispersed cells, and the ratio of phagocytosing haemocytes was significantly lower in the mutant embryos than in the control (Figure 2D). To confirm the importance of Prtp, we examined whether the overexpression of *prtp* in the mutants rescues the phagocytosis phenotype. In this experiment, embryonic haemocytes were identified by immunohistochemically examining the presence of Croquemort, a marker for haemocytes (Franc *et al*, 1996, 1999) (Supplementary Figure S5). The defect in phagocytosis in *prtp*^{Δ1} was completely recovered by the ubiquitous expression of *prtp* using the *daughterless* (*da*) promoter (Figure 2E), indicating that Prtp indeed has a function in the phagocytosis of apoptotic cells. In contrast, the expression of *prtp* in a haemocyte-specific manner using the *serpent* (*srp*) promoter did not alter the level of phagocytosis (Figure 2E), suggesting that Prtp is required in apoptotic cells not haemocytes. In fact, the level of the phagocytosis of latex beads, indicative of the basal phagocytic activity of phagocytes, was the same between larval haemocytes isolated from the control and *prtp* mutant flies (Figure 2F).

We next examined the phagocytosis of apoptotic cells by glia in the *prtp* mutants. The distribution (left panel in Figure 2G) and number (right panel in Figure 2G) of glia, which were immunohistochemically visualized with antibody recognizing the glia-specific protein Repo, did not significantly differ in embryos between the control and *prtp* mutants. To determine the level of phagocytosis *in situ*, we introduced *Rep1^P*, a null allele of *dICAD* (Mukae *et al*, 2002), into *prtp*^{Δ1}, *prtp*^{Δ2}, and *prtp*^{G19}. In *Rep1^P* flies, the active form of caspase-activated DNase is not produced and, thus, DNA of apoptotic cells is degraded only after engulfment by phagocytes (Mukae *et al*, 2002). We previously showed that apoptotic cells engulfed by phagocytes in these flies are detectable by *in situ* nick translation (ISNT) (Manaka *et al*, 2004). The region in the ventral nerve cord of stage 16 embryos contains many glia that phagocytose apoptotic neurons (Abrams *et al*, 1993; Sonnenfeld and Jacobs, 1995). We therefore determined the number of ISNT signals detected in the nine abdominal segments (A1 to A9) of the ventral nerve cord in stage 16 embryos (see left panel in Figure 2G). The level of phagocytosis in the *prtp* mutants was about a half of that observed in control embryos (Figure 2H).

To further confirm a defect in the phagocytosis of apoptotic cells by glia in the *prtp* mutants, we analysed dispersed embryonic cells for the level of phagocytosis as done for the analysis of phagocytosis by haemocytes, except for the use of the glial marker Repo instead of Croquemort (left panel in Figure 2I). We found that the level of Repo-positive glia that contained TUNEL-positive nuclei was reduced in the *prtp* mutants (right panel in Figure 2I).

Furthermore, when *prtp* was overexpressed in *Rep1^P* flies that normally express *prtp*, the level of apoptotic cell phagocytosis in embryos increased and the number of apoptotic cells decreased (Supplementary Figure S6). All these results indicated that Prtp is required for the efficient phagocytosis of apoptotic cells by haemocytes and glia in *Drosophila* embryos. We previously reported that Drpr is responsible for the phagocytic removal of degenerated axons in mushroom body γ neurons by glia during metamorphosis (Awasaki *et al*, 2006). However, the timing and extent of the loss of the axons were the same between the control and *prtp* mutant (Supplementary Figure S7).

Genetic interaction of *prtp* with *drpr*, *ced-6*, and *Rac1/Rac2*

Ced-6, a *Drosophila* homologue of *C. elegans* CED-6, is important in the Drpr-mediated phagocytosis of neural axons by glia (Awasaki *et al*, 2006). More recently, Franc and co-workers reported that *drpr* genetically interacts with *ced-6* with regard to the phagocytosis of apoptotic cells (Cuttell *et al*, 2008). An immunohistochemical analysis showed that most embryonic haemocytes expressed Ced-6 (Supplementary Figure S8A). We thus genetically examined whether or not Prtp functions in the pathway where Drpr and Ced-6 are involved. To obtain a null allele of *ced-6*, we generated a fly line having a deletion in the region spanning *ced-6* by P-element excision (Supplementary Figure S8B). This mutant, *ced-6^{J26}*, lacked the expression of Ced-6 with no change in the expression of Drpr and Prtp (Supplementary Figure S8C). We found that embryonic haemocytes of this *ced-6* null mutant showed a lower level of phagocytosis than those of wild-type flies (Figure 3A), confirming the involvement of Ced-6 in the phagocytosis of apoptotic cells (Cuttell *et al*, 2008). We then examined phagocytosis in the *prtp* null mutant with additional mutation on *drpr* (*drpr*^{Δ5}, see Supplementary Figure S2C) or *ced-6*, but the loss of either Drpr or Ced-6 did not further decrease the level of phagocytosis (Figure 3A), suggesting that Prtp and Drpr, and Prtp and Ced-6 act in the same pathway. We next examined the occurrence of genetic interaction between *prtp* and *drpr*, and *prtp* and *ced-6*. A heterozygous mutant for each of the three genes, which had reduced expression of the corresponding proteins (top panel of Figure 3B), showed a level of phagocytosis comparable to that observed with control *w¹¹¹⁸* flies (bottom panel of Figure 3B). However, the level of phagocytosis by embryonic haemocytes of a *trans*-heterozygous mutant for *prtp* and either *drpr* or *ced-6* was significantly lower than that by haemocytes of the control flies (bottom panel of Figure 3B). These results indicated that *prtp* genetically interacts with *drpr* and *ced-6*. Taken together, Prtp, Drpr, and Ced-6 are likely to function in the same signalling pathway for the induction of phagocytosis of apoptotic cells.

We extended genetic studies to examine the possible interaction of *prtp* with other *Drosophila* genes homologous

Figure 3 Genetic interaction of *prtp* with *drpr*, *ced-6*, and *Rac1/Rac2*. (A) Level of phagocytosis by Croquemort-expressing haemocytes in embryos of the indicated fly lines. The data with control (+ / +) and *prtp^{Δ1}* flies are duplications of those in Figure 2E. NS, difference not significant (versus *prtp^{Δ1}*). (B) Levels of proteins (top, western blotting) and phagocytosis by green fluorescent protein-positive haemocytes (bottom), determined with dispersed embryonic cells of the indicated fly lines. *prtp^{Δ1} hemi* is a hemizygote lacking *prtp* expression. **P* < 0.005 (versus *prtp^{Δ1} / +*). (C) Level of phagocytosis by green fluorescent protein-positive haemocytes in embryos of the indicated fly lines. **P* < 0.05 (versus *prtp^{Δ1} / +*). (D) Levels of phagocytosis, Croquemort-positive haemocytes, and terminal deoxynucleotidyltransferase-mediated dUTP nick end labeling-positive apoptotic cells determined with dispersed embryonic cells of the indicated fly lines. Refer to Materials and methods section for genotypes of the fly lines analysed.

to those consisting of the engulfment pathways in *C. elegans*. An analysis of genetic interaction for *mbc* (*CED-5* homologue) and *Rac1* and *Rac2* (*CED-10* homologues) was carried out because null mutants for *mbc* (Nolan *et al*, 1998) and both *Rac1* and *Rac2* (Hakeda-Suzuki *et al*, 2002) show developmental defects in embryos. We found that embryos of *trans*-heterozygotes of *prtp^{Δ1}* and *mbc^{C1}* (null mutant of *mbc*) showed the level of phagocytosis by haemocytes comparable to that with a heterozygous mutant for each gene (Figure 3C). In contrast, the level of phagocytosis in embryos lacking one copy of *prtp*, *Rac1*, and *Rac2* was lower than that in a heterozygous mutant for *prtp*, or *Rac1* and *Rac2* (Figure 3C), further suggesting the involvement of Prtp in the pathway corresponding to CED-1/CED-6/CED-10 in *C. elegans*. To our surprise, a lack of the expression of *elmo* (*CED-12* homologue) did not seem to affect phagocytosis in embryos (Figure 3D), indicating that ELMO does not have an important function in the phagocytosis of apoptotic cells in embryos. We were unable to analyse a *Drosophila* gene homologous to *CED-2* because of a technical difficulty. These results collectively suggested that Prtp is not involved in the engulfment pathway corresponding to CED-2/CED-5/CED-12 in *C. elegans*.

We previously reported that Calreticulin, another protein residing in the ER, serves as a marker for phagocytosis in *Drosophila* embryos (Kuraishi *et al*, 2007). We thus geneti-

cally examined the relationship between *prtp* and *calreticulin* as to the phagocytosis of apoptotic cells. When cells dispersed from embryos were analysed, the ratio of phagocytosing haemocytes obtained from a double mutant for *prtp* and *calreticulin* was about 70% of that from the *prtp* single mutant (Supplementary Figure S9A). Furthermore, the level of phagocytosis in embryos of *trans*-heterozygotes of *prtp^{Δ1}* and a *calreticulin* null allele was almost the same as that in embryos of flies lacking one copy of *prtp* or *calreticulin* (Supplementary Figure S9B), suggesting that there was no genetic interaction between these two genes. Taken together, it seems likely that Prtp and Calreticulin have independent functions in the phagocytosis of apoptotic cells by embryonic haemocytes.

Relocation of Prtp to the cell surface during apoptosis

We next examined a possible change in the sub-cellular localization of Prtp during apoptosis. S2 cells before and after the induction of apoptosis by the treatment with cycloheximide were examined by immunocytochemistry under conditions without membrane permeabilization. We found that the occurrence of apoptosis rendered S2 cells positive for the surface expression of Prtp (Figure 4A). To confirm this *in vivo*, we analysed cells dispersed from embryos, in which apoptosis was induced by the overexpression of Reaper and Hid, for the externalization of Prtp. Most dispersed

Figure 4 Relocation of Pretaporter (Prtp) to the cell surface during apoptosis. **(A)** Surface expression of Prtp in S2 cells treated with cycloheximide (Apoptotic) or left untreated (Viable). Phase-contrast (bottom) and fluorescence (top) views of the same microscopic fields are shown. The inset is a magnified view of the squared area. Bar, 10 μ m. **(B)** Surface expression of Prtp in embryonic cells undergoing apoptosis. Apoptosis was induced in embryos of *UAS-rpr UAS-hid/+; da-GAL4/tub-GAL80^{ts}* flies by incubation at 29°C for 11 h after egg laying. Cells obtained from those embryos were analysed for the exposure of Prtp (using anti-Prtp antibody) and phosphatidylserine (PS) (using fluorocholesterol-labelled annexin V) under membrane-non-permeabilizing conditions. Incubation with the probes was carried out in the absence and presence of EDTA, which chelates Ca^{2+} necessary for the binding of annexin V to PS. Phase-contrast and fluorescence views of the same microscopic fields are shown in each row. Bar, 5 μ m. **(C)** Surface expression of Prtp in apoptotic cells of wing discs. Wing discs, in which apoptosis was locally induced, were immunohistochemically analysed for the expression of Prtp and tubulin with (Total) and without (Extracellular) membrane permeabilization. Apoptotic cells were detected by *in situ* nick translation (Apoptosis). Fluorescence views of the same microscopic fields are shown in each row. The insets in the top row are magnified views of the corresponding squared area. Bar, 50 μ m.

embryonic cells with externalized phosphatidylserine, indicative of apoptosis, were also positive for the binding of anti-Prtp antibody (Figure 4B). We then extended this type of analysis using wing discs where only a few cells undergo apoptosis in normal development (Milán *et al*, 1997). Spatio- and temporal-specific apoptosis was induced in the discs by the expression of *reaper* and *hid* under the *engrailed* promoter, and these discs were analysed for externalized Prtp (Figure 4C). Apoptosis examined by ISNT was evident in a portion of the disc where the *engrailed* promoter is presumed to be active (Tabata *et al*, 1995), and the same region simultaneously became positive for the surface expression of Prtp (first row): note that most Prtp-positive cells were also positive for the ISNT signal. Positive staining with anti-Prtp antibody was not due to the permeabilization of plasma membranes, because antibody that recognizes tubulin did not bind to the region of apoptosis unless the discs were membrane permeabilized (second and fourth rows). In addition, ubiquitous expression of Prtp was observed after the permeabilization of membranes (third row). These results indicated that apoptosis causes the exposure of Prtp at the cell surface. We next determined the timing of Prtp externalization during apoptosis using S2 cells. The data suggested that the externalization of Prtp occurs at almost the same time as the fragmentation of DNA, and after the loss of phospholipid asymmetry in the plasma membrane and the cleavage of pro-caspase-3 (Figure 5A). Furthermore, the surface expression of Prtp in S2 cells was not observed when caspases were inhibited during the induction of apoptosis

(Figure 5B). Taken together, these results show that Prtp relocates from the ER to the cell surface during the process of apoptosis.

Role of Prtp as a ligand for Drpr in the phagocytosis of apoptotic cells

The results obtained so far suggested that Prtp acts as a ligand for Draper in the phagocytosis of apoptotic cells. To obtain data more directly supporting this idea, we carried out a series of experiments. First, we asked if the surface expression of Prtp makes particles more effective targets for phagocytosis. Recombinant Prtp fused to GST was covalently attached to latex beads, and these particles were tested as targets for I(2)mbn cells, a *Drosophila* cell line derived from larval haemocytes, which acquire the ability to phagocytose apoptotic S2 cells in a Drpr-dependent manner after the treatment with 20-hydroxyecdysone (Manaka *et al*, 2004). We found that Prtp-coated latex beads were more efficiently phagocytosed by hormone-treated I(2)mbn cells than those coated with GST alone (Figure 6A). Furthermore, the enhancement of phagocytosis by Prtp was completely suppressed when Drpr expression in I(2)mbn cells was inhibited by RNA interference. This indicated that the presence of Prtp at the surface of target particles renders Drpr-mediated phagocytosis more effective. We next confirmed this observation using cells instead of artificial targets. To make the ER protein Prtp expressed at the cell surface, we manipulated cDNA so that the ER retention motif of Prtp was replaced with an amino-acid sequence for the attachment of

Figure 5 Timing of Pretaporter (Prtp) externalization and other biochemical changes during apoptosis. (A) S2 cells induced to undergo apoptosis by the treatment with cycloheximide were examined for the occurrence of biochemical changes associated with apoptosis at the indicated time points. Phase-contrast and fluorescence views of the same microscopic fields in each analysis are shown. Bar, 10 μ m. The biochemical changes examined were externalization of Prtp (analysed by the binding of anti-Prtp antibody), externalization of phosphatidylserine (PS) (analysed by the binding of annexin V), activation of pro-caspase-3 (casp-3) (immunochemically analysed with anti-activated caspase-3 antibody), and fragmentation of DNA (analysed by *in situ* nick translation). (B) S2 cells were induced to undergo apoptosis by the treatment with cycloheximide in the presence and absence of the pan-caspase inhibitor z-VAD-fmk and examined for the occurrence of Prtp externalization. Bar, 10 μ m. A full-colour version of this figure is available at *The EMBO Journal* Online.

the glycosylphosphatidylinositol anchor (Supplementary Figure S10A). S2 cells transfected with this DNA became positive for the surface exposure of Prtp (left panel of Figure 6B) and were more efficiently phagocytosed by hormone-treated l(2)mbn cells than those transfected with the vector alone or left untransfected (right panel of Figure 6B). This enhancement seemed Drpr dependent because RNA interference-mediated inhibition of Drpr expression in l(2)mbn cells reduced phagocytosis (Supplementary Figure S10B). Drpr is presumed to undergo tyrosine phosphorylation at the intracellular region upon activation (Mangahas and Zhou, 2005; Ziegenfuss *et al*, 2008). We thus next tested if the binding of Prtp induces the phosphorylation of Drpr. To do so, l(2)mbn cells were maintained in culture containers coated with recombinant Prtp, and their whole-cell lysates were subjected to immunoprecipitation with anti-Drpr antibody followed by western blotting with anti-phosphotyrosine antibody. The results showed that the amount of tyrosine-phosphorylated Drpr in l(2)mbn cells increased after incubation with Prtp, but not with a control protein (Figure 6C). All

the data described above collectively strengthened the idea that Prtp serves as a ligand for Drpr in the phagocytosis of apoptotic cells.

Discussion

Here, we report the identification of a *Drosophila* protein, which we have named Prtp, serving as a ligand for Drpr in the phagocytosis of apoptotic cells. The level of phagocytosis in embryos of the null mutants for *prtp* and *drpr* was almost the same and about half of that in controls. This suggests that Prtp is solely responsible for the Drpr-mediated phagocytosis by embryonic haemocytes, and that there are additional ligands and receptors for phagocytosis. Based on the data presented above, we would propose the following pathway for the induction of the Prtp-dependent, Drpr-mediated phagocytosis of apoptotic cells: Prtp relocates from the ER to the cell surface during apoptosis; Prtp binds to Drpr of haemocytes and induces tyrosine phosphorylation; Ced-6 binds to

Figure 6 Function of Pretaporter (Prtp) as a ligand for Draper (Drpr). **(A)** Phagocytosis of latex beads coated with Prtp. Latex beads coated with the indicated proteins were subjected to phagocytosis by I(2)mbn cells with and without RNA interference (dsRNA) of Drpr or Croquemort (Crq). Data are representative of three (none, Drpr) or two (Crq) independent experiments that yielded similar results. **(B)** Phagocytosis of S2 cells surface-expressing Prtp. S2 cells, which had been transfected with plasmid for the expression of glycosylphosphatidylinositol (GPI)-anchored Prtp and green fluorescent protein (GFP), were examined for the presence of GFP and Prtp (left) and the phagocytosis by I(2)mbn cells (right). Phase-contrast and fluorescent views of the same microscopic fields are shown in each row (left). Bar, 10 μ m. **(C)** Tyrosine phosphorylation of Drpr. I(2)mbn cells incubated with maltose-binding protein (MBP)-fused Prtp or β -galactosidase (β gal) were subjected to an analysis of immunoprecipitation (IP)/western blotting (Blot) using anti-Drpr and anti-phosphotyrosine (p-Tyr) antibodies.

phosphorylated Drpr and further transmits the signal leading to the activation of Rac1 and/or Rac2; and cytoskeletons are reorganized for engulfment. Additional molecules could participate in this pathway: a membrane protein named Six-microns-under reportedly acts upstream of Drpr in the phagocytosis of apoptotic neurons by glia (Kurant *et al*, 2008); another signal mediator, named Shark, located proximal downstream of Drpr has been reported (Ziegenfuss *et al*, 2008); and Undertaker, an intracellular membrane protein involved in the regulation of calcium homeostasis, was shown to function downstream of Drpr (Cuttell *et al*, 2008). In particular, Six-microns-under has been suggested to bridge glia and apoptotic neurons (Kurant *et al*, 2008). Our results suggested that Prtp still serves as a ligand for Drpr in the phagocytosis by glia that express Six-microns-under. Further investigation is necessary before we obtain a complete picture of the Prtp/Drpr-initiated signalling pathway for the phagocytosis of apoptotic cells. More molecules are involved in the phagocytosis of apoptotic cells in *Drosophila*: the receptor Croquemort (Franc *et al*, 1996, 1999), the phagocytosis marker Calreticulin (Kuraishi *et al*, 2007), and

Pallbearer, a component of ubiquitin ligase (Silva *et al*, 2007), which might be integrated into the other engulfment pathway.

Prtp normally resides in the ER, most likely in its lumen, and a portion of it seems to relocate to the cell surface during apoptosis. The ER and plasma membranes exchange their components (Levine and Rabouille, 2005), and this might be enhanced upon the induction of apoptosis. In fact, Herrmann and co-workers reported that a variety of proteins and lipids of the ER are exposed at the surface of apoptotic human cells (Franz *et al*, 2007). More recently, it was reported that the exposure of ER proteins at the surface of apoptotic mammalian cells occurs by SNARE-dependent exocytosis (Panaretakis *et al*, 2009). It is thus probable that together with other ER components Prtp moves to the cell surface during apoptosis and serves as a ligand for the phagocytosis receptor Drpr. Prtp seemed dispensable for the removal of degenerated neural axons during metamorphosis, which requires the action of Drpr (Awasaki *et al*, 2006). We speculate that Prtp is not exposed on the surface of degenerated axons because the removal of these axons occurs independently of caspases (Awasaki *et al*, 2006), and that another ligand for Drpr exists at the surface of degenerated axons. On the other hand, Drpr appears to serve as a receptor in the phagocytosis of bacteria (Cuttell *et al*, 2008; Hashimoto *et al*, 2009). Taken together, it is likely that Drpr is a multi-ligand receptor for phagocytosis responsible for the maintenance of tissue homeostasis through the removal of degenerated own cells and invading microbial pathogens.

A counterpart for Prtp in *C. elegans* could be a ligand for CED-1, but there seems no complete homologue of *prtp* in its genome. Our data suggested that the presence of two thiorodoxin-like domains is sufficient for the binding of Prtp to Drpr. Therefore, a *C. elegans* protein containing such a structure, PDI for example, is a candidate for the CED-1 ligand. In contrast, the mammalian homologue of Prtp seems to exist; a mouse protein called ERp46 (Knoblauch *et al*, 2003) and human proteins belonging to the TXNDC5 family (Nissom *et al*, 2006) possess a domain composition similar to that of Prtp. It is important to examine if *C. elegans* PDI and mammalian ERp46 and TXNDC5 act as ligands for CED-1 and its mammalian homologue MEGF10 (Hamon *et al*, 2006), respectively, in the phagocytosis of apoptotic cells.

Materials and methods

Fly stocks and cell culture

The following fly lines were used in this study: *w*¹¹¹⁸, CG1837^{BG00450} (Bloomington Drosophila Stock Center), *ced-6*^{EY11592} (Bloomington Drosophila Stock Center), *P{seq-EYFP-ER}*3 (LaJeunesse *et al*, 2004), *Rep1*^P (Mukae *et al*, 2002), *srpHemo-GAL4 UAS-srcEGFP* (Brückner *et al*, 2004), *en-GAL4* (Brand and Perrimon, 1993), *da-GAL4* (Giebel *et al*, 1997), *tub-GAL80^{ts20}* (McGuire *et al*, 2003), *UAS-cGFP S65T* (a gift from B. Dickson), *UAS-rpr UAS-hid* (Zhou *et al*, 1997), *drpr*^{Δ5} (Freeman *et al*, 2003; MacDonald *et al*, 2006), *201Y-GAL4* (Yang *et al*, 1995), *mbc*^{C1} (Rushton *et al*, 1995), *elmo*^{Δ5} (Bianco *et al*, 2007), *Rac1*¹¹¹ (Ng *et al*, 2002), *Rac2*^Δ (Ng *et al*, 2002), and *Crc*^{S114307} (Prokopenko *et al*, 2000). To determine fly genotypes based on the expression of β -galactosidase, we balanced *drpr*^{Δ5} *mbc*^{C1}, and *Rac1*¹¹¹ *Rac2*^Δ over TM3-*ftzlacZ*, and balanced *elmo*^{Δ5} over CyO-*ftzlacZ*. The alleles of *prtp* analysed were generated by imprecise excision of the P-element with CG1837^{BG00450}. After standard mitotic recombination over *w*¹¹¹⁸ chromosomes to remove possible second-site mutations, the P-element was mobilized by a Δ 2–3 transposase source. About 200 candidate excision events were

screened by genomic polymerase chain reaction amplifying the *prtp* locus (see Figure 2A and Supplementary Figure S3A), and two deletion lines, *prtp*^{Δ1} and *prtp*^{Δ2}, were isolated. A strain after precise excision of the P-element, *prtp*^{G19}, was also obtained and used as a control. A null allele for *ced-6*, *ced-6*^{Δ26}, was similarly generated by imprecise excision of the P-element with *ced-6*^{EY11592} (see Supplementary Figures S8B and S8C). To obtain flies containing an extra *prtp* expressed with the *GAL4-UAS* system, we inserted the entire coding region of Prtp cDNA isolated from the *Drosophila* EST clone LD24756 (provided by Berkeley Drosophila Genome Project and National Institute of Genetics) into pUAST (Brand and Perrimon, 1993), and we generated flies transgenic with the resultant DNA according to a standard procedure (Spradling, 1986; Robertson *et al*, 1988). Three fly lines carrying the transgene on the second or third chromosome were established, and one with it on the third chromosome was used for inter-crossing with *da-GAL4* (for ubiquitous expression) or *srpHemo-GAL4 UAS-srcEGFP* (for haemocyte-specific expression). Other fly lines used were generated through mating of the existing lines. l(2)mbn cells and S2 cells were maintained at 25°C with Schneider's *Drosophila* medium (Invitrogen) containing 10% (v/v) heat-inactivated fetal bovine serum. l(2)mbn cells were used as phagocytes after they were incubated in the presence of 20-hydroxyecdysone (Sigma-Aldrich) (1 μM) for 2–5 days, essentially as described previously (Manaka *et al*, 2004). S2 cells were induced to undergo apoptosis by incubation with cycloheximide (1.5 μg/ml) for 24 h, as described previously (Manaka *et al*, 2004). To isolate larval haemocytes, we incubated haemolymph collected from wandering larvae on glass slides according to a published method (Pearson *et al*, 2003), and cells attaching to the slides were collected and used as haemocytes.

Affinity chromatography and mass spectrometry

The region spanning amino-acid residues 143–798 (with the N terminus numbered 1) of Drpr-RB (based on the nomenclature in FlyBase), which probably corresponds to Drpr isoform I (Freeman *et al*, 2003), was expressed as a fusion protein with GST in Sf21 insect cells using the *Drosophila* EST clone GH03529 (provided by Berkeley Drosophila Genome Project and National Institute of Genetics) and a baculovirus-based DNA vector (Invitrogen), and the purified recombinant protein (Drpr-GST) was immobilized onto glutathione-Sepharose (Amersham Biosciences) using the cross-linking reagent DSP (Pierce). Apoptotic S2 cells were lysed with PIPES buffer (pH 6.5) containing 0.1 M NaCl, 20 mM KCl, 20 mM MgSO₄, 10 mM CaCl₂, and 1% (w/v) CHAPS, and the lysates were applied onto the affinity matrix. The proteins retained on the matrix were eluted with 0.1 M Tris-glycine buffer (pH 2.7) containing 2 M NaCl, 1% CHAPS, 5 mM EDTA, and 5 mM EGTA, and analysed either by sodium dodecyl sulphate–polyacrylamide gel electrophoresis followed by staining with silver or western blotting with anti-Prtp antibody. To identify a Drpr-binding protein, we excised the protein band and in-gel-digested with *Achromobacter* protease I, and analysed the resulting peptides by LC-MS/MS using DiNa nano-flow liquid chromatograph (KYA Technologies, Tokyo) and Esquire HCT ion-trap mass spectrometer (Bruker Daltonics) with a nano-electrospray ionization probe.

Assays for phagocytosis

In situ analysis of phagocytosis was carried out with whole embryos by immunohistochemistry, TUNEL, ISNT, or a combination thereof, as described in the text. Ten intact embryos or dispersed cells obtained from about 200 embryos were analysed in each experiment. An analysis of phagocytosis with dispersed embryonic cells was carried out as described previously (Manaka *et al*, 2004) except that a fluorochrome-based detection of signals was substituted by an enzyme-linked immunoassay, which modification made this assay highly reproducible, that is, results obtained from repeated experiments do not deviate over 10%. Phagocytosis reactions *in vitro* with l(2)mbn cells as phagocytes were carried out as described previously (Manaka *et al*, 2004). To prepare latex beads containing Prtp, we incubated fluorescein isothiocyanate-labelled latex beads (Φ = 1.7 μm; Polysciences) with GST-fused Prtp, which had been expressed in and purified from *Escherichia coli*, in the presence of the cross-linking reagent EDAC (Polysciences). For the surface expression of Prtp in S2 cells, a DNA coding for the amino-acid residues 1–412 (with the N terminus numbered 1) of Prtp that lacks the ER retention motif was linked with a DNA corresponding to the amino-acid residues 606–626 of Dally (Takeo *et al*, 2005) by

overlapping polymerase chain reaction using Dally cDNA (a gift from H Nakato) (see Supplementary Figure S10A). The resultant DNA was inserted into pUAST and introduced into S2 cells by the calcium/phosphate method together with pAct5C-GAL4 and pUAST-GFP (a gift from M Miura). Those S2 cells were mixed with 20-hydroxyecdysone-treated l(2)mbn cells (at 3:1), and the mixture was incubated at 25°C for 1 h. The samples were washed and examined by microscopy after staining with haematoxylin, and l(2)mbn cells containing extra S2 cell nuclei were counted as those cells that had accomplished phagocytosis. To examine the basal phagocytic activity of haemocytes, we carried out phagocytosis reactions using haemocytes isolated from wandering larvae as phagocytes and fluorescein isothiocyanate-labelled latex beads as targets.

Other materials and methods

Anti-Prtp antibody was raised by immunizing rats with GST-fused Prtp that had been expressed in *E. coli* and purified to homogeneity. The generation of anti-Drpr (Manaka *et al*, 2004), anti-Croquemort (Manaka *et al*, 2004), and anti-Ced-6 (Awasaki *et al*, 2006) antibodies was reported previously. Anti-Fasciclin II monoclonal antibody was provided by A Nose. Anti-Repo mouse monoclonal antibody 8D12 and anti-β-galactosidase monoclonal antibody (used for fly genotyping in the experiments shown as Figure 3) were obtained from Developmental Studies Hybridoma Bank. Anti-GFP monoclonal antibody and anti-GFP polyclonal antibody were purchased from Roche and Molecular Probes, respectively. Anti-α-tubulin and anti-human activated caspase-3 antibodies were purchased from Sigma-Aldrich and Cell Signaling, respectively. z-VAD-fmk was obtained from R&D Systems. Varying portions of Prtp were expressed in *E. coli* as proteins fused with maltose-binding protein (MBP) and affinity purified to homogeneity using amylose-Sepharose (New England BioLabs). These proteins were incubated with Drpr-GST, precipitated with amylose-Sepharose, and analysed for the association of Drpr-GST by western blotting with anti-GST (Upstate) or anti-MBP (New England BioLabs) antibody. MBP-fused β-galactosidase was similarly prepared and used as a negative control. To examine the phosphorylation of Drpr, we maintained l(2)mbn cells (1 × 10⁷) in culture containers (Φ = 6 cm) that had been coated with MBP-fused Prtp (full length, 0.3 mg) or β-galactosidase at 25°C for 5.5 h, washed with phosphate-buffered saline, and lysed with buffer consisting of 40 mM Tris-HCl (pH 7.5), 2% CHAPS, 0.15 M NaCl, and 1 mM EDTA. The lysates were analysed by immunoprecipitation with anti-Drpr antibody followed by western blotting with anti-phosphotyrosine (clone RC20; BD Biosciences) or anti-Drpr antibody. Western blotting, RNA interference, cytochemistry with fluorochrome-labelled annexin V, and ISNT were carried out as described previously (Manaka *et al*, 2004). TUNEL was performed according to the manufacturer's instructions (Chemicon). Immunohistochemistry with cultured cells and embryos was carried out as described previously (Manaka *et al*, 2004; Kuraishi *et al*, 2007). To induce apoptosis in wing discs, we kept late third-instar larvae of *UAS-rpr UAS-hid*/+; *en-GAL4/tub-GAL80*^{ts} at 29°C for 12 h that had been maintained at 20°C. The discs isolated from those larvae were immunohistochemically analysed for the presence of Prtp and α-tubulin with and without membrane permeabilization according to the method of Strigini and Cohen (2000). An analysis of axon pruning was carried out as described previously (Awasaki *et al*, 2006). Reverse transcription-mediated polymerase chain reaction was conducted according to a standard procedure with primers containing sequences of mRNA for Prtp and Drpr.

Genotypes of the fly lines analysed

In Figures 2B–D, and F: *w*¹¹¹⁸*prtp*^{G19}/*w*¹¹¹⁸*prtp*^{G19}; *srpHemo-GAL4 UAS-srcEGFP/srpHemo-GAL4 UAS-srcEGFP* (G19), *w*¹¹¹⁸*prtp*^{Δ1}/*w*¹¹¹⁸*prtp*^{Δ1}; *srpHemo-GAL4 UAS-srcEGFP/srpHemo-GAL4 UAS-srcEGFP* (Δ1), and *w*¹¹¹⁸*prtp*^{Δ2}/*w*¹¹¹⁸*prtp*^{Δ2}; *srpHemo-GAL4 UAS-srcEGFP/srpHemo-GAL4 UAS-srcEGFP* (Δ2).

In Figure 2E: *w*¹¹¹⁸*prtp*^{G19}/*w*¹¹¹⁸*prtp*^{G19} (+/+), *w*¹¹¹⁸*prtp*^{Δ1}/*w*¹¹¹⁸*prtp*^{Δ1} (*prtp*^{Δ1}), *prtp*^{Δ1}/*prtp*^{Δ1}; *da-GAL4*/+ (*prtp*^{Δ1}; *da-GAL4*/+), *prtp*^{Δ1}/*prtp*^{Δ1}; *srpHemo-GAL4 UAS-srcEGFP*/+ (*prtp*^{Δ1}; *srp-GAL4*/+), *prtp*^{Δ1}/*prtp*^{Δ1}; *da-GAL4/UAS-prtp* (*prtp*^{Δ1}; *da-GAL4/UAS-prtp*), and *prtp*^{Δ1}/*prtp*^{Δ1}; *srpHemo-GAL4 UAS-srcEGFP*/+; *UAS-prtp*/+ (*prtp*^{Δ1}; *srp-GAL4*/+; *UAS-prtp*/+).

In Figure 2G and I: *w*¹¹¹⁸*prtp*^{G19}/*w*¹¹¹⁸*prtp*^{G19} (G19), *w*¹¹¹⁸*prtp*^{Δ1}/*w*¹¹¹⁸*prtp*^{Δ1} (Δ1), and *w*¹¹¹⁸*prtp*^{Δ2}/*w*¹¹¹⁸*prtp*^{Δ2} (Δ2).

In Figure 2H: $w^{1118}prtp^{G19}/w^{1118}prtp^{G19}; Rep1^P/Rep1^P$ (G19), $w^{1118}prtp^{\Delta1}/w^{1118}prtp^{\Delta1}; Rep1^P/Rep1^P$ ($\Delta1$), and $w^{1118}prtp^{\Delta2}/w^{1118}prtp^{\Delta2}; Rep1^P/Rep1^P$ ($\Delta2$).

In Figure 3A: $w^{1118}prtp^{G19}/w^{1118}prtp^{G19}$ (+/+), $w^{1118}prtp^{\Delta1}/w^{1118}prtp^{\Delta1}$ ($prtp^{\Delta1}$), $w^{1118}prtp^{\Delta1}/w^{1118}prtp^{\Delta1}; drpr^{\Delta5}/drpr^{\Delta5}$ ($prtp^{\Delta1}; drpr^{\Delta5}$), and $w^{1118}prtp^{\Delta1}/w^{1118}prtp^{\Delta1}; ced-6^{J26}/ced-6^{J26}$ ($prtp^{\Delta1}; ced-6^{J26}$).

In Figure 3B: w^{1118}/w^{1118} (+/+), $w^{1118}prtp^{\Delta1}/w^{1118}prtp^{\Delta1}$ ($prtp^{\Delta1}/+$), $w^{1118}/w^{1118}; drpr^{\Delta5}/+$ ($drpr^{\Delta5}/+$), $w^{1118}/w^{1118}; ced-6^{J26}/+$ ($ced-6^{J26}/+$), $w^{1118}prtp^{\Delta1}/w^{1118}; drpr^{\Delta5}/+$ ($prtp^{\Delta1}/+; drpr^{\Delta5}/+$), $w^{1118}prtp^{\Delta1}/w^{1118}; ced-6^{J26}/+$ ($prtp^{\Delta1}/+; ced-6^{J26}/+$), and $w^{1118}prtp^{\Delta1}/Y$ ($prtp^{\Delta1}hemi$) for western blotting, and $w^{1118}/w^{1118}; srpHemo-GAL4$ UAS-*srcEGFP*/+ (+/+), $w^{1118}prtp^{\Delta1}/w^{1118}; srpHemo-GAL4$ UAS-*srcEGFP*/+ ($prtp^{\Delta1}/+$), $w^{1118}/w^{1118}; srpHemo-GAL4$ UAS-*srcEGFP*/+; $drpr^{\Delta5}/+$ ($drpr^{\Delta5}/+$), $w^{1118}/w^{1118}; ced-6^{J26}/srpHemo-GAL4$ UAS-*srcEGFP* ($ced-6^{J26}/+$), $w^{1118}prtp^{\Delta1}/w^{1118}; srpHemo-GAL4$ UAS-*srcEGFP*/+; $drpr^{\Delta5}/+$ ($prtp^{\Delta1}/+; drpr^{\Delta5}/+$), $w^{1118}prtp^{\Delta1}/w^{1118}; ced-6^{J26}/srpHemo-GAL4$ UAS-*srcEGFP* ($prtp^{\Delta1}/+; ced-6^{J26}/+$), and $w^{1118}prtp^{\Delta1}/Y; srpHemo-GAL4$ UAS-*srcEGFP*/+ ($prtp^{\Delta1}hemi$) for phagocytosis analysis.

In Figure 3C: $prtp^{G19}/prtp^{G19}; srpHemo-GAL4$ UAS-*srcEGFP*/+ (+/+), $srpHemo-GAL4$ UAS-*srcEGFP*/+; $red^1e^1mbc^{C1}/+$ ($mbc^{C1}/+$), $srpHemo-GAL4$ UAS-*srcEGFP*/+; $Rac^{J11}Rac2^{\Delta}FRT2A/+$ ($Rac^{J11}Rac2^{\Delta}/+$), $w^{1118}prtp^{\Delta1}/w^{1118}prtp^{G19}; srpHemo-GAL4$ UAS-*srcEGFP*/+ ($prtp^{\Delta1}/+$), $prtp^{\Delta1}/+; srpHemo-GAL4$ UAS-*srcEGFP*/+; $red^1e^1mbc^{C1}/+$ ($prtp^{\Delta1}/+; mbc^{C1}/+$), $prtp^{\Delta1}/+; srpHemo-GAL4$ UAS-*srcEGFP*/+; $Rac^{J11}Rac2^{\Delta}FRT2A/+$ ($prtp^{\Delta1}/+; Rac^{J11}Rac2^{\Delta}/+$), and $w^{1118}prtp^{\Delta1}/Y; srpHemo-GAL4$ UAS-*srcEGFP*/+ ($prtp^{\Delta1}hemi$).

In Figure 3D: $y w$ (+/+), $elmo^{KO}FRT40, 42/elmo^{KO}FRT40, 42$ ($elmo^{KO}$).

Data processing and statistical analysis

Data in a numerical analysis are expressed as the mean $\pm$ s.d. of the results obtained from at least three independent experiments, unless otherwise mentioned in the text. Other data are representative of at least two independent experiments that yielded similar results. For determining the level of phagocytosis, two investigators independently conducted the same experiments or the data

obtained by one investigator were re-examined in blind by another, and only the replicated results were presented in this paper. Statistical analyses were performed using the Student's *t*-test and *P*-values of <0.05 were considered significant.

Supplementary data

Supplementary data are available at *The EMBO Journal* Online (<http://www.embojournal.org>).

Acknowledgements

We thank B Dickson, M Freeman, S Hayashi, M Meister, S Nagata, and Bloomington Stock Center for the fly stocks; M Miura and H Nakato for the DNA; Berkeley Drosophila Genome Project and National Institute of Genetics for the EST clones; and A Nose and Developmental Studies Hybridoma Bank for the antibodies. We are grateful to M-O Fauvarque, S Hayashi, K Ito, K Koizumi, E Kuranaga, S Kurata, M Miura, Y Nakai, R Tajiri, L Tsuda, R Ueda, and Y Yoshioka for advice and suggestions, and M Freeman, S Hayashi, K Ito, H Kanuka, and Y-C Wu for discussion. We thank R Tatsumi for technical support. We also acknowledge the use of FlyBase. This study was supported by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (Grant Number 19370051 to YN), and in part by Grants-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (Grant Number 18570123 to AS) and from the Ministry of Education, Culture, Sports, Science and Technology-Japan (Grant Number 18057009 to AS). TK was a recipient of the pre-doctoral fellowship from the Japan Society for the Promotion of Science. All the authors performed the experiments and analysed the data. TK and YN designed the study. YN supervised the study. TK, YN, and TA prepared the paper.

Conflict of interest

The authors declare that they have no conflict of interest.

References

- Abrams JM, White K, Fessler LI, Steller H (1993) Programmed cell death during *Drosophila* embryogenesis. *Development* **117**: 29–43
- Awasaki T, Tatsumi R, Takahashi K, Arai K, Nakanishi Y, Ueda R, Ito K (2006) Essential role of the apoptotic cell engulfment genes *draper* and *ced-6* in programmed axon pruning during *Drosophila* metamorphosis. *Neuron* **50**: 855–867
- Bianco A, Poukkula M, Cliffe A, Mathieu J, Luque CM, Fulga TA, Rørth P (2007) Two distinct modes of guidance signalling during collective migration of border cells. *Nature* **448**: 362–365
- Brand AH, Perrimon N (1993) Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Development* **118**: 401–415
- Brückner K, Kockel L, Duchek P, Luque CM, Rørth P, Perrimon N (2004) The PDGF/VEGF receptor controls blood cell survival in *Drosophila*. *Dev Cell* **7**: 73–84
- Callebaut I, Mignotte V, Souchet M, Mornon J-P (2003) EMI domains are widespread and reveal the probable orthologs of the *Caenorhabditis elegans* CED-1 protein. *Biochem Biophys Res Commun* **300**: 619–623
- Cleveland DW (1996) Neuronal growth and death: order and disorder in the axoplasm. *Cell* **84**: 663–666
- Cohen JJ (1991) Programmed cell death in the immune system. *Adv Immunol* **50**: 55–85
- Cutler L, Vaughan A, Silva E, Escaron CJ, Lavine M, Van Goethem E, Eid J-P, Quirin M, Franc NC (2008) Undertaker, a *Drosophila* Junctophilin, links Draper-mediated phagocytosis and calcium homeostasis. *Cell* **135**: 524–534
- Fadok VA, Bratton DL, Frasch SC, Warner ML, Henson PM (1998) The role of phosphatidylserine in recognition of apoptotic cells by phagocytes. *Cell Death Differ* **5**: 551–562
- Franc NC, Dimarcq J-L, Lagueux M, Hoffmann J, Ezekowitz RAB (1996) Croquemort, a novel *Drosophila* hemocyte/macrophage receptor that recognizes apoptotic cells. *Immunity* **4**: 431–443
- Franc NC, Heitzler P, Ezekowitz RAB, White K (1999) Requirement for Croquemort in phagocytosis of apoptotic cells in *Drosophila*. *Science* **284**: 1991–1994
- Franz S, Herrmann K, Führnrohr B, Sheriff A, Frey B, Gaipal US, Voll RE, Kalden JR, Jäck H-M, Herrmann M (2007) After shrinkage apoptotic cells expose internal membrane-derived epitopes on their plasma membranes. *Cell Death Differ* **14**: 733–742
- Freeman MR, Delrow J, Kim J, Johnson E, Doe CQ (2003) Unwrapping glial biology: Gcm target genes regulating glial development, diversification, and function. *Neuron* **38**: 567–580
- Giebel B, Stüttem I, Hinz U, Campos-Ortega JA (1997) Lethal of Scute requires overexpression of Daughterless to elicit ectopic neuronal development during embryogenesis in *Drosophila*. *Mech Dev* **63**: 75–87
- Hakeda-Suzuki S, Ng J, Tzu J, Dietzl G, Sun Y, Harms M, Nardine T, Luo L, Dickson BJ (2002) Rac function and regulation during *Drosophila* development. *Nature* **416**: 438–442
- Hamon Y, Trompier D, Ma Z, Venegas V, Pophillat M, Mignotte V, Zhou Z, Chimini G (2006) Cooperation between engulfment receptors: the case of ABCA1 and MEGF10. *PLoS ONE* **1**: e120
- Hashimoto Y, Tabuchi Y, Sakurai K, Kutsuna M, Kurokawa K, Awasaki T, Sekimizu K, Nakanishi Y, Shiratsuchi A (2009) Identification of lipoteichoic acid as a ligand for Draper in the phagocytosis of *Staphylococcus aureus* by *Drosophila* hemocytes. *J Immunol* **138**, advance online publication, 4 November, 2009; doi:10.4049/jimmunol.0901032
- Jacobson MD, Weil M, Raff MC (1997) Programmed cell death in animal development. *Cell* **88**: 347–354
- Kinchen JM, Cabello J, Klingele D, Wong K, Feichtinger R, Schnabel H, Schnabel R, Hengartner MO (2005) Two pathways converge at CED-10 to mediate actin rearrangement and corpse removal in *C. elegans*. *Nature* **434**: 93–99

- Kinchen JM, Hengartner MO (2005) Tales of cannibalism, suicide, and murder: programmed cell death in *C.elegans*. *Curr Top Dev Biol* **65**: 1–45
- Kinchen JM, Ravichandran KS (2007) Journey to the grave: signaling events regulating removal of apoptotic cells. *J Cell Sci* **120**: 2143–2149
- Knoblach B, Keller BO, Groenendyk J, Aldred S, Zheng J, Lemire BD, Li L, Michalak M (2003) ERp19 and ERp46, new members of the thioredoxin family of endoplasmic reticulum proteins. *Mol Cell Proteomics* **2**: 1104–1119
- Kuraishi T, Manaka J, Kono M, Ishii H, Yamamoto N, Koizumi K, Shiratsuchi A, Lee BL, Higashida H, Nakanishi Y (2007) Identification of calreticulin as a marker for phagocytosis of apoptotic cells in *Drosophila*. *Exp Cell Res* **313**: 500–510
- Kurant E, Axelrod S, Leaman D, Gaul U (2008) Six-microns-under acts upstream of Draper in the glial phagocytosis of apoptotic neurons. *Cell* **133**: 498–509
- LaJeunesse DR, Buckner SM, Lake J, Na C, Pirt A, Fromson K (2004) Three new *Drosophila* markers of intracellular membranes. *Biotechniques* **36**: 784–790
- Laubert K, Blumenthal SG, Waibel M, Wesselborg S (2004) Clearance of apoptotic cells: getting rid of the corpses. *Mol Cell* **14**: 277–287
- Lettre G, Hengartner MO (2006) Developmental apoptosis in *C. elegans*: a complex CEDnario. *Nat Rev Mol Cell Biol* **7**: 97–108
- Levine T, Rabouille C (2005) Endoplasmic reticulum: one continuous network compartmentalized by extrinsic cues. *Curr Opin Cell Biol* **17**: 362–368
- Liao DJ (2005) The scavenger cell hypothesis of apoptosis: apoptosis redefined as a process by which a cell in living tissue is destroyed by phagocytosis. *Med Hypotheses* **65**: 23–28
- MacDonald JM, Beach MG, Porpiglia E, Sheehan AE, Watts RJ, Freeman MR (2006) The *Drosophila* cell corpse engulfment receptor Draper mediates glial clearance of severed axons. *Neuron* **50**: 869–881
- Manaka J, Kuraishi T, Shiratsuchi A, Nakai Y, Higashida H, Henson P, Nakanishi Y (2004) Draper-mediated and phosphatidylserine-independent phagocytosis of apoptotic cells by *Drosophila* hemocytes/macrophages. *J Biol Chem* **279**: 48466–48476
- Mangahas PM, Zhou Z (2005) Clearance of apoptotic cells in *Caenorhabditis elegans*. *Sem Cell Dev Biol* **16**: 295–306
- McGuire SE, Le PT, Osborn AJ, Matsumoto K, Davis RL (2003) Spatiotemporal rescue of memory dysfunction in *Drosophila*. *Science* **302**: 1765–1768
- Milán M, Campuzano S, García-Bellido A (1997) Developmental parameters of cell death in the wing disc of *Drosophila*. *Proc Natl Acad Sci USA* **94**: 5691–5696
- Mukae N, Yokoyama H, Yokokura T, Sakoyama Y, Nagata S (2002) Activation of the innate immunity in *Drosophila* by endogenous chromosomal DNA that escaped apoptotic degradation. *Genes Dev* **16**: 2662–2671
- Nakanishi Y, Henson PM, Shiratsuchi A (2009) Pattern recognition in phagocytic clearance of altered self. In *Target Pattern Recognition in Innate Immunity*, Kishore U (ed), pp 129–138. Austin: Landes Bioscience
- Ng J, Nardine T, Harms M, Tzu J, Goldstein A, Sun Y, Dietzl G, Dickson BJ, Luo L (2002) Rac GTPases control axon growth, guidance and branching. *Nature* **416**: 442–447
- Nissom PM, Lo SL, Lo JCY, Ong PF, Lim JWE, Ou K, Liang RC, Seow TK, Chung MCM (2006) Hcc-2, a novel mammalian ER thioredoxin that is differentially expressed in hepatocellular carcinoma. *FEBS Lett* **580**: 2216–2226
- Nolan KM, Barrett K, Lu Y, Hu K-Q, Vincent S, Settleman J (1998) Myoblast city, the *Drosophila* homolog of DOCK180/CED-5, is required in a Rac signaling pathway utilized for multiple developmental processes. *Genes Dev* **12**: 3337–3342
- Panaretakis T, Kepp O, Brockmeier U, Tesniere A, Bjorklund A-C, Chapman DC, Durchschlag M, Joza N, Pierron G, van Endert P, Yuan J, Zitvogel L, Madeo F, Williams DB, Kroemer G (2009) Mechanisms of pre-apoptotic calreticulin exposure in immunogenic cell death. *EMBO J* **28**: 578–590
- Pearson AM, Baksa K, Rämets M, Protas M, McKee M, Brown D, Ezekowitz RAB (2003) Identification of cytoskeletal regulatory proteins required for efficient phagocytosis in *Drosophila*. *Microbes Infect* **5**: 815–824
- Prokopenko SN, He Y, Lu Y, Bellen HJ (2000) Mutations affecting the development of the peripheral nervous system in *Drosophila*: a molecular screen for novel proteins. *Genetics* **156**: 1691–1715
- Ravichandran KS, Lorenz U (2007) Engulfment of apoptotic cells: signals for a good meal. *Nat Rev Immunol* **7**: 964–974
- Reddien PW, Horvitz HR (2004) The engulfment process of programmed cell death in *Caenorhabditis elegans*. *Annu Rev Cell Dev Biol* **20**: 193–221
- Robertson HM, Preston CR, Phillis RW, Johnson-Schlitz DM, Benz WK, Engels WR (1988) A stable genomic source of P element transposase in *Drosophila melanogaster*. *Genetics* **118**: 461–470
- Rushon E, Drysdale R, Abmayr SM, Michelson AM, Bate M (1995) Mutations in a novel gene, *myoblast city*, provide evidence in support of the founder cell hypothesis for *Drosophila* muscle development. *Development* **121**: 1979–1988
- Savill J, Fadok V (2000) Corpse clearance defines the meaning of cell death. *Nature* **407**: 784–788
- Schlegel RA, Williamson P (2001) Phosphatidylserine, a death knell. *Cell Death Differ* **8**: 551–563
- Silva E, Au-Yeung HW, Goethem EV, Burden J, Franc NC (2007) Requirement for a *Drosophila* E3-ubiquitin ligase in phagocytosis of apoptotic cells. *Immunity* **27**: 585–596
- Sonnenfeld MJ, Jacobs JR (1995) Macrophages and glia participate in the removal of apoptotic neurons from the *Drosophila* embryonic nervous system. *J Comp Neurol* **359**: 644–652
- Spradling AC (1986) P element-mediated transformation. In *Drosophila: A Practical Approach*, Roberts DB (ed), 2nd edn, pp 175–197. Leeds: Oxford University Press
- Strigini M, Cohen SM (2000) Wingless gradient formation in the *Drosophila* wing. *Curr Biol* **10**: 293–300
- Tabata T, Schwartz C, Gustavson E, Ali Z, Kornberg TB (1995) Creating a *Drosophila* wing *de novo*, the role of *engrailed*, and the compartment border hypothesis. *Development* **121**: 3359–3369
- Takeo S, Akiyama T, Firkus C, Aigaki T, Nakato H (2005) Expression of a secreted form of Dally, a *Drosophila* glypican, induces overgrowth phenotype by affecting action range of Hedgehog. *Dev Biol* **284**: 204–218
- Vaux DL, Korsmeyer SJ (1999) Cell death in development. *Cell* **96**: 245–254
- Venegas V, Zhou Z (2007) Two alternative mechanisms that regulate the presentation of apoptotic cell engulfment signal in *Caenorhabditis elegans*. *Mol Biol Cell* **18**: 3180–3192
- Wyllie AH, Kerr JFR, Currie AR (1980) Cell death: the significance of apoptosis. *Int Rev Cytol* **68**: 251–306
- Yang MY, Armstrong JD, Vilinsky I, Strausfeld NJ, Kaiser K (1995) Subdivision of the *Drosophila* mushroom bodies by enhancer-trap expression patterns. *Neuron* **15**: 45–54
- Yu X, Odera S, Chuang C-H, Lu N, Zhou Z (2006) *C. elegans* dynamin mediates the signaling of phagocytic receptor CED-1 for the engulfment and degradation of apoptotic cells. *Dev Cell* **10**: 743–757
- Zhou L, Schnitzler A, Agapite J, Schwartz LM, Steller H, Nambu JR (1997) Cooperative functions of the *reaper* and *head involution defective* genes in the programmed cell death of *Drosophila* central nervous system midline cells. *Proc Natl Acad Sci USA* **94**: 5131–5136
- Zhou Z, Hartwig E, Horvitz HR (2001) CED-1 is a transmembrane receptor that mediates cell corpse engulfment in *C.elegans*. *Cell* **104**: 43–56
- Ziegenfuss JS, Biswas R, Avery MA, Hong K, Sheehan AE, Yeung Y-G, Stanley ER, Freeman MR (2008) Draper-dependent glial phagocytic activity is mediated by Src and Syk family kinase signalling. *Nature* **453**: 935–939