play a role in the development of the latter during childhood. "From an evolutionary perspective, studies on the early development of language and music abilities in children are particularly interesting and have so far indicated some overlapping and possibly shared neural resources," he said. "Musical training facilitates pitch processing in both music and language, and children with specific language impairment also show impairment in music-syntactic processing."

Given that it requires a higher precision of pitch and rhythm to appreciate music than to understand speech, Dorrell believes that the only reason humans are so good at perceiving pitch is to appreciate music. "[This] strongly suggests that our ability to appreciate music has some biological purpose, since the mechanism of precise pitch perception presumably has a non-trivial implementation cost in terms of brain growth and operation, and wouldn't exist if it didn't have some value. But this doesn't get us any closer to knowing what this purpose is," he said. "The extra precision required for music perception is, however, consistent with the hypothesis that the perception of musicality represents the perception of unintentionally leaked information which the 'sender' is making no effort to send, and even perhaps trying not to send."

Scientists have just begun to scrape the surface of the biological essence of music and its connections with other cognitive domains, such as language or learning. "Interaction between music and language in bilingual children would be another fascinating area of investigation," commented Johansson. "Taking advantage of the unfortunate fact that a large part of the human population is still illiterate, it would be feasible to initiate studies on the effect on the brain of learning to read in different languages and writing systems, and on the possible interactions between local language and music traditions."

In this intellectual and scientific endeavour to understand one of our most emotive forms of expression, the tools of molecular biology—and genetics in particular—will certainly have a role, although it is too early to say how fundamental this might be. Nevertheless, regardless of the contribution of any one discipline, the result should be a deeper comprehension of the mechanisms our brain uses to decode and process sound-driven information.

REFERENCES

- Bachner-Melman R *et al* (2005) *AVPR1a* and *SLC6A4* gene polymorphisms are associated with creative dance performance. *PLoS Genet* **1**: e42
- Dorrell P (2005) What is music? Solving a scientific mystery. <http://whatismusic.info>
- Drayna D, Manichaikul A, de Lange M, Snieder H, Spector T (2001) Genetic correlates of musical pitch recognition in humans. *Science* **291**: 1969–1972
- Granot RY, Frankel Y, Gritsenko V, Lerer E, Gritsenko I, Bachner-Melman R, Israel S, Ebstein RP (2007) Provisional evidence that the arginine vasopressin 1a receptor gene is associated with musical memory. *Evol Hum Behav* **28**: 313–318
- Johansson BB (2008) Language and music: what do they have in common and how do they differ? A neuroscientific approach. *Eur Rev* **16**: 413–427
- Patel AD (2003) Language, music, syntax and the brain. *Nat Neurosci* **6**: 674–681
- Patel AD (2008) *Music, Language, and the Brain*. Oxford, UK: Oxford University Press
- Peretz I (2008) Musical disorders. From behavior to genes. *Curr Dir Psychol Sci* **17**: 329–333
- Peretz I, Cummings S, Dubé MP (2007) The genetics of congenital amusia (tone deafness): a family aggregation study. *Am J Hum Genet* **81**: 582–588
- Pulli K, Karma K, Norio R, Sistonen P, Göring HHH, Järvelä I (2008) Genome-wide linkage scan for loci of musical aptitude in Finnish families: evidence for a major locus at 4q22. *J Med Genet* **45**: 451–456
- Ross D, Choi J, Purves D (2007) Musical intervals in speech. *Proc Natl Acad Sci USA* **104**: 9852–9857
- Theusch E, Basu A, Gitschier J (2009) Genome-wide study of families with absolute pitch reveals linkage to 8q24.21 and locus heterogeneity. *Am J Hum Genet* **85**: 112–119
- Ukkola LT, Onkamo P, Raijas P, Karma K, Järvelä I (2009) Musical aptitude is associated with *AVPR1A*-haplotypes. *PLoS One* **4**: e5534

Andrea Rinaldi

doi:10.1038/embor.2009.241

Me, myself and I

The genetics and molecular biology behind self-incompatibility and the avoidance of inbreeding in plants

The emergence of sex was a huge leap for evolution. The development of sexual reproduction meant that random mutations were no longer the only source of genetic variety on which selection could act. By mixing up parental alleles, sex essentially put evolution on afterburners; it creates a rich abundance of new varieties, which are the substrate for the process of natural selection and evolution.

However, sexual reproduction came with a new set of problems. On the one hand, incest must be avoided, as reproduction between closely related individuals leads to the accumulation of detrimental traits owing to the lack of genetic variety; on the other hand, inter-species reproduction creates sterile individuals. Animals have thus evolved a range of molecular and behavioural strategies to cope with incest, while at the same time ensuring that they only reproduce with their own kind.

Plants have faced the somewhat larger challenge of overcoming their lack of mobility and resulting inability to directly select their mates. They have therefore

evolved sexual reproduction that utilizes mobile male gametes—pollen—that must be transmitted to another plant where they fertilize the oocyte, which will develop into a seed (Fig 1). Early plants released their pollen into the air, relying on the wind to carry it away to other plants. But the emergence of flowering plants—angiosperms—refined this process to commandeer insects to transport pollen from one plant to another. Yet, neither strategy solves the problem of incest, as both the wind and insects are likely to carry pollen to a flower from the same plant, or to a very closely related individual.

The emergence of self-incompatibility (SI) to prevent inbreeding and enable outcrossing was therefore a major evolutionary breakthrough for angiosperms, as it allowed them to diversify rapidly and to dominate the surface of the Earth about 150 million years ago. Charles Darwin first described SI mechanisms (McClure, 2009) and appreciated their significance, but it is only recently that the genetic and molecular details behind SI have started to emerge.

Fig 1 | Pollination, fertilization and germination in flowering plants. To overcome their lack of mobility, flowering plants release their male gametes as pollen to be delivered—usually by the wind or insects—to other members of their species. To avoid incest and to promote compatible outcrossing, various molecular and physical strategies are in place. These include early-stage checks—such as the control of pollen tube growth by the receiving flower—and late-stage processes—such as the abortion of embryos arising from fertilization by an incompatible sperm.

Amid an ongoing debate over the ancestry of angiosperms, it has become clear that SI has evolved independently at least six times, with significant differences at both the genetic and molecular levels (Hiscock & McInnis, 2003). The elucidation of these mechanisms is significant not just for understanding the evolution of angiosperms, but also for unravelling the basis of molecular signalling and recognition in general, for example in adaptive immunity in higher vertebrates.

The emergence of self-incompatibility (SI) to prevent inbreeding and enable outcrossing was therefore a major evolutionary breakthrough for angiosperms

Indeed, there are also similarities between SI mechanisms in plants and some animals, according to Bruce McClure from the University of Missouri (Columbia, MO, USA). “SI in angiosperms is not really so

different from some animal systems,” he said, citing the example of corals, which reproduce sexually as well as asexually. “You see them pumping gametes out into the sea, which actually is just like wind pollination in plants.” Most coral species are hermaphrodite, expressing both sexes, and so by pumping out gametes individuals would fertilize themselves without SI. Indeed, studies have confirmed that all hermaphroditic species of corals are either completely or almost completely self-incompatible (Carlon, 1999).

The development of molecular mechanisms to allow the pollen and stigma—the female sexual organ to which pollen attaches—from the same species to recognize one another while at the same time avoiding self-pollination, has created a dilemma for plants. According to Hugh Dickinson from the Department of Plant Sciences at Oxford University in the UK, “It goes without saying that there is a tension in the evolution of pollen–stigma interactions—selection ‘wants’ pollen and stigmas of the same species to be far more ‘compatible’ than those of other

species—so pollen and stigma ‘co-evolve’ to develop structures, recruit effective vectors to ensure reliable pollination within, and as little as possible pollination outside the species. Opposing this trend is the necessity of avoiding self-pollination.”

Some angiosperms have therefore evolved straightforward, non-molecular methods to prevent self-pollination. Various plants segregate the genders to different flowers, with both male and female flowers on the plant at the same time; others ensure that their male and female flowers are present at separate times of the season; while others only ever express either all male or all female flowers, sometimes switching sex depending on their stage of growth. But these methods are either of limited effectiveness, or, as in the case of differential timing, impose constraints on mating with compatible members of the species.

Plants have thus evolved SI at the molecular level to occur either before fertilization, or through abortion of the incestuous seed. The process takes advantage of the way in which

the pollen of flowering plants delivers its payload: siphonogamy, which is present in all seed-producing plants and allows the gametes to remain under the protection of the plant while it circumvents the need for water to facilitate the transfer of the sperm. When a pollen grain lands on the stigma of a flower, the pollen germinates and extends a pollen tube through the style towards the ovary. Once the pollen tube reaches the ovary, it enters the ovule, penetrates the megagametophyte—which produces the egg cell—and releases two sperm cells. One fertilizes the waiting egg and the other fuses with the two polar nuclei of the central cell to form the endosperm, which provides nourishment for the developing embryo.

SI could, for instance, be applied to ensure that transgenic crops can only mate with themselves...

But the endosperm is more than the plant equivalent of yolk or the placenta; it also actively mediates the growth of the developing embryo. As such, during the course of evolution, it also became a genetic guardian by aborting the progeny of parents deemed insufficiently compatible or too closely related. However, this late-stage action to prevent incompatible fertilization means that the plant has already wasted considerable resources in allowing the pollen tube access to the ovary and is now also unavailable to more suitable pollen. Such late-acting mechanisms are therefore thought to be the precursors of more advanced, efficient versions of SI, which destroy the pollen before fertilization (Allen & Hiscock, 2008).

These mechanisms rely on signalling between the growing pollen tube and the stigma that essentially negotiates compatibility between the two. In fact, the underlying signalling pathways are also recruited quite early in some late-onset SI mechanisms, according to Simon Hiscock from Bristol University in the UK. Hiscock referred to earlier work on *Narcissus triandus*, a daffodil-like plant native to southern Europe, which utilizes a late-onset SI mechanism in which ovules cease to develop when pollen from genetically similar plants has germinated on the stigma (Sage *et al*, 1999). The signalling and subsequent decision to abort takes place before the pollen tube

reaches the ovule, but it does not prevent the development of the tube. Although it is still inefficient, this mechanism does represent an advance over truly late-acting SI.

The most efficient and highly evolved SI mechanisms destroy the pollen before it even germinates. Although several early-stage SI mechanisms have been identified, they all involve signalling between the pollen tube and the gynoecium during their interaction, and they all involve the S locus—a highly polymorphic region of the plant genome that contains two sex genes: one controlling the identity of the pollen, and the other that of the gynoecium. Selective pressure has encouraged diversity around the S locus because any plant with a new allele arising through mutation will, by definition, not be recognized as ‘self’ by most other plants; thus enhancing its prospects of procreating.

Many angiosperm species have large numbers of alleles around these S-locus genes in their population. “The majority of SI systems operate on the basis of multiple alleles, so that the likelihood of an unrelated pollen grain being rejected is slim,” Dickinson explained. “More sophisticated systems have up to 60, and grasses have two loci, each allele of which forms a unique combination with the other, creating any number of permutations.” In grasses, self-compatibility is determined by alleles at the S locus and at another known as the Z locus.

In both histocompatibility and plant SI, the distinction between self and non-self is made on the basis of alleles of a particular gene or set of genes

Although it has been known for some years that alleles at the S locus—and in the case of some grasses, also at the Z locus—are at the root of more advanced SI mechanisms, the molecular details of how the relevant protein products interact are only just coming to light. SI operates in two phases: first, the gynoecium identifies proteins expressed by pollen S-locus genes. If the gynoecium identifies the pollen as self, it triggers a series of events that arrest the growth of the pollen tube and subsequently lead to its destruction. If not, the proteins expressed by genes at the

gynoecium S locus, in conjunction with other genes, encourage and stimulate the growth of the pollen tube, for example by providing cues to help it elongate in the right direction.

According to McClure, the fact that many of the same genes on the gynoecium side are involved in both rejection and acceptance explains why pollen seems to conspire in its own destruction when attempting to fertilize an ovule that is genetically close. The reason is that, in cases where pollen does not carry S-locus alleles that closely match the target ovule, the signalling process actually encourages the growth of its tube. “This is sometimes called the ‘candy apple’ hypothesis,” McClure said: the pollen is lured into conspiring with the SI process because most of the time the mechanisms actually assist its journey towards the ovule.

Two radically different genetic mechanisms for SI have evolved—gametophytic SI (GSI) and sporophytic SI (SSI)—which are distinguished by whether it is the pollen itself or its parent that exercises control over the process. In GSI, the pollen itself expresses the molecules that make up its incompatibility phenotype, from the alleles it inherited from its parent plant. In SSI, the parent expresses the incompatibility products and the pollen inherits them directly within the nucleus that produces the pollen tube. It seems that SSI is less common; so far it has only been described in detail in *Brassica*, a genus of the mustard family. GSI, conversely, is known to have evolved several times with radically different molecular mechanisms, even though the overlying genetic process is the same.

Two GSI mechanisms that have been well-studied rely on RNase and S-glycoprotein respectively. In the first case, S-RNase, expressed through the female S locus, causes the degradation of ribosomal RNA inside the pollen tube, arresting the growth of the tube and leaving the pollen grain to wither when it has progressed about one-third of the way through the style (Franklin-Tong & Franklin, 2003). On the pollen side, members of the F-box protein family seem to play a key role in compatibility. If the interaction is compatible, the proteins target the pollen's own RNases and degrade them so that pollen tube growth can continue. According to McClure, in the case of tobacco at least, the pollen normally keeps these RNases wrapped up in a membrane that is only

broken down in the event of an incompatible interaction. This releases the RNases, which then degrade the ribosomal RNA and the tube is aborted—in other words, the pollen assists in its own suicide.

The S-glycoprotein mechanism acts even more rapidly, stopping pollen growth within minutes of arrival on the stigma. This occurs when some as-yet-unknown cell membrane receptor on the male side receives an extra glycoprotein molecule from the female side. What is known is that this signalling quickly triggers an influx of calcium to the pollen tube, disrupting the calcium gradient that is essential for protein expression and tube elongation.

Several other GSI mechanisms exist but have not yet been worked out at the molecular level, according to McClure. The common thread among them is that all have evolved from the communication system that already existed between pollen and the gynoecium during the approach to fertilization. Another possible common feature is that all these SI mechanisms tend to be abandoned under certain circumstances, for example during the process of hybridization, when individuals cross from different species. Normally, the same mechanisms that protect against inbreeding also resist mating between individuals that are too distantly related for offspring to be viable or fertile, but hybridization does occur naturally and plays a vital role in creating new species in the wild.

When fertile hybrids are formed, those that continue to use SI would quickly die out as many individuals would initially be genetically similar. In these cases, it seems that selective pressure favours the abandonment of SI in favour of self-compatibility because the detrimental effects of in-breeding are initially offset by the novelty of the new plant, or its so-called “hybrid vigour” (Cook & Soltis, 2000).

The study of SI mechanisms has a range of potential applications in agriculture and horticulture. SI could, for instance, be applied to ensure that transgenic crops can only mate with themselves, as has been demonstrated by Dickinson's lab with *Brassica*. “The idea was—which I thought was pretty natty—to disassemble the SI system and reconstruct it such that only transgenic pollen would develop on transgenic stigmas,” Dickinson said. “Pollen

moving to other brassicas in the wild would be rejected, as would wild brassica pollen entering the crop.” In that way, SI could be used to prevent transgenes from spreading into wild relatives.

...SI promises to transform conventional plant breeding, while allaying some of the public fear and criticism associated with the use of transgenic crops

SI can also be applied in the creation of new varieties through conventional breeding, to prevent candidate plants crossing with themselves and thus increasing the chance of creating new desirable outcrosses, according to Noni Franklin-Tong, a professor of plant cell biology at the University of Birmingham in the UK. “In many crops that are naturally inbreeding, such as rice, wheat and rape, hand emasculating, which is time-consuming and so expensive, is often the only method available to develop hybrid varieties,” she said. “Inserting genes to make such crops self-incompatible would mean that hand emasculating would not be necessary, and so make hybrid seed production much easier and so cheaper. [...] Several patents for S-determinants in various species have been filed, but so far no-one has come up with a workable system. We are currently working on seeing if the poppy SI system will work, and preliminary results look encouraging.”

The molecular mechanisms of plant SI are also of interest for another reason, which is that they have similarities with those of histocompatibility in higher animals, according to McClure. In both histocompatibility and plant SI, the distinction between self and non-self is made on the basis of alleles of a particular gene or set of genes. With plant SI, self is rejected, whereas the opposite is the case with histocompatibility, but the underlying mechanisms are similar. There is also evidence that histocompatibility is associated with mating preferences through its link with odour (Penn & Potts, 1999). Earlier studies had established that in house mice, both males and females tend to select mates with substantially different alleles for key genes in the major histocompatibility complex (Potts *et al*, 1991).

Histocompatibility is also related to adaptive immunity, which again has similarities to plant SI. Both have the objective of eliminating undesirable target cells, and therefore seem to share some mechanisms, such as highly variable receptors to recognize equally variable ligands (Nasrallah, 2005).

The study of SI in plants thus extends beyond plants to general mechanisms of how organisms are able to distinguish between themselves and others, and how these mechanisms might have evolved. Moreover, although its application to plant breeding and agricultural biotechnologies is still at an early stage, SI could transform conventional plant breeding, while allaying some of the public fear and criticism associated with the use of transgenic crops.

REFERENCES

- Allen AM, Hiscock SJ (2008) Evolution and phylogeny of self-incompatibility systems in angiosperms. In Franklin-Tong VE (ed) *Self-Incompatibility in Flowering Plants*, pp73–101. Heidelberg, Germany: Springer
- Carlson DB (1999) The evolution of mating systems in tropical reef corals. *Trends Ecol Evol* **14**: 491–494
- Cook LM, Soltis PS (2000) Mating systems of diploid and allotetraploid populations of *Tragopogon* (Asteraceae) II. Artificial populations. *Heredity* **84**: 410–415
- Franklin-Tong VE, Franklin FCH (2003) The different mechanisms of gametophytic self-incompatibility. *Philos Trans R Soc Lond B Biol Sci* **358**: 1025–1032
- Hiscock SJ, McInnis SM (2003) The diversity of self-incompatibility systems in flowering plants. *Plant Biol* **5**: 23–32
- McClure B (2009) Darwin's foundation for investigating self-incompatibility and the progress toward a physiological model for S-RNase-based SI. *J Exp Bot* **60**: 1069–1081
- Nasrallah JB (2005) Recognition and rejection of self in plant self-incompatibility: comparisons to animal histocompatibility. *Trends Immunol* **26**: 412–418
- Penn D, Potts W (1999) The evolution of mating preferences and major histocompatibility genes. *Am Nat* **153**: 145–164
- Potts WK, Manning CJ, Wakeland EK (1991) Mating patterns in seminatural populations of mice influenced by MHC genotype. *Nature* **352**: 619–621
- Sage TL, Strumas F, Cole WW, Barrett SCH (1999) Differential ovule development following self and cross-pollination: the basis of self-sterility in *Narcissus triandrus* (Amaryllidaceae). *Am J Bot* **86**: 855–870

Philip Hunter

doi:10.1038/embor.2009.251