Published in final edited form as:

Tetrahedron Lett. 2010 February 3; 51(5): 779–782. doi:10.1016/j.tetlet.2009.12.008.

An Efficient Boric Acid Mediated Preparation of α -Hydroxyamides

J. Sravan Kumar^a, Subash C. Jonnalagadda^b, and Venkatram R. Mereddy^a

^aDepartment of Chemistry and Biochemistry, University of Minnesota Duluth, Duluth, MN 55812.

^bDepartment of Chemistry and Biochemistry, Rowan University, Glassboro, NJ 08051.

Abstract

An efficient methodology for the preparation of α -hydroxyamides via boric acid mediated addition of isonitriles on to aldehydes has been developed. The reaction of isonitriles with α -boronobenzaldehyde takes place under intramolecular catalysis conditions to provide functionalized benzoxaboroles.

Keywords

Multicomponent reactions; Isonitriles; Boric acid; Benzoxaboroles

Introduction

Boric acid is an inexpensive, non-toxic compound and it is generally considered a green material.¹ It is an excellent precursor for the preparation of various types of organoboranes and is also used as a mild Lewis acid catalyst for several organic transformations.² α -Hydroxyamides are important synthetic intermediates in organic synthesis and also serve as valuable agents in medicinal chemistry.³ The use of isonitriles in three or four component multicomponent coupling reactions such as Passerini and Ugi is well documented in the literature.⁴ The addition of isonitriles on to carbonyl compounds is usually catalyzed by strong protic acids or Lewis acids to provide α -hydroxyamides.⁵

Results and Discussion

Owing to the aforementioned advantages, we envisaged the utility of boric acid for the direct α -addition of isonitriles on to aldehydes to synthesize α -hydroxyamides. For the current study, we chose benzyl, cyclohexyl and t-butyl isonitriles (**2a-c**) and various types of aldehydes **1a-g**. We initiated the optimization of the reaction conditions with propionaldehyde **1a** with benzyl isonitrile **2a**. The reaction was studied in various solvents such as THF, CH₂Cl₂, EtOAc, DMF, DMSO, Dioxane, MeOH and water with varying amounts of boric acid (10 mol% to one equivalent). DMF as a solvent and one equivalent of boric acid were found to be optimal in obtaining cleaner reaction with good yields. Aliphatic isonitriles **2b, 2c** and aliphatic aldehydes were in general found to be more reactive and the reactions were completed in 24 hours of stirring at room temperature. Aromatic aldehydes reacted slowly and required almost 48 hours

© 2009 Elsevier Ltd. All rights reserved.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

of reaction time. Upon completion, the reaction mixture was washed with water, worked up with diethyl ether and pure products were obtained after silica gel column chromatography.

We also explored the reaction with the ketones 3-pentanone and acetophenone as representative examples. However, the reactions were found to be very sluggish even with two equivalents of boric acid and elevated temperature (80°C).

We then applied the methodology for intramolecular version of the reaction starting with *o*-formyl phenylboronic acid **5**. We envisaged that the proximity of the boronic acid to the formyl group in **5** could lead to a facile reaction with isonitrile under self catalysis mode without any additional boric acid catalysis. This reaction should provide direct access to functionalized cyclic boronic acids (benzoxaboroles) **6**. Indeed, the reaction of boronoaldehyde **5** with isonitrile **2a** in DMF at room temperature took place smoothly with out boric acid to provide the product benzoxaborole in 81% yield. Similarly, isonitriles **2b** and **2c** upon reaction with **5** provided the corresponding benzoxaboroles **6b** and **6c** respectively (Scheme 1). This class of boron compounds have extensive applications in materials chemistry and synthetic chemistry as excellent intermediates for Suzuki cross coupling reactions.⁶ Several of these compounds have also been found to exhibit important antibacterial and antifungal properties.⁷

Representative procedure for the preparation of hydroxyamide 4a

To a stirred solution of propionaldehyde **1a** (0.14 mL, 2.0 mmol) in 2mL DMF was added benzyl isonitrile **2a** (0.24 mL, 2.0 mmol), and boric acid (0.12 g, 2.0 mmol) and stirred overnight at room temperature. Upon completion (TLC), the reaction mixture was worked up with water and diethyl ether (3 × 25 mL). The combined organic layers were dried (MgSO₄), concentrated *in vacuo*, and purified by column chromatography (silica gel, hexane:acetone, 4:1) to obtain 0.30 g (78%) of hydroxy amide **4a**. ¹H NMR (500 MHz, CDCl₃): 7.22-7.32 (m, 5H), 7.18 (bs, 1H), 4.47 (t, *J* = 4.5 Hz, 1H), 4.42 (dd, *J* = 4.0, 15.0 Hz, 1H), 4.36 (dd, *J* = 4.0, 15.0 Hz, 1H), 4.05-4.08 (m, 1H), 1.78-1.90 (m, 1H), 1.62-1.72 (m, 1H), 0.95 (t, *J* = 7.5 Hz, 3H); ¹³C NMR (125 MHz, CDCl₃): 174.5, 138.2, 128.9, 127.9, 127.7, 73.2, 43.3, 28.1, 9.4; ESI-MS: 216 [(M+Na)⁺, 100%], 194 (M+H)⁺.

Representative procedure for the preparation of benzoxaborole 6b

To a stirred solution of boronoaldehyde **5** (0.3 g, 2.0 mmol) in 2mL DMF was added *tert*-butyl isonitrile **2a** (0.23 mL, 2.0 mmol), and stirred overnight at room temperature. Upon completion (TLC), the reaction mixture was worked up with water and ethyl acetate (3 × 25 mL). The combined organic layers were dried (MgSO₄), concentrated *in vacuo*, and purified by column chromatography (silica gel, hexane:ethyl acetate, 3:1) to obtain 0.30 g (65%) of benzoxaborole **6b**. (Found: C, 61.53; H, 8.10; N, 6.02 %; C₁₂H₁₆BNO₃ requires: C, 61.84; H, 6.92; N, 6.01%); ¹H NMR (500 MHz, DMSO-*d*₆): 9.33 (bs, 1H), 7.67 (d, *J* = 7.2 Hz, 1H), 7.62 (d, *J* = 7.6 Hz, 1H), 7.36 (t, *J* = 7.4 Hz, 1H), 7.26 (t, *J* = 7.4 Hz, 1H), 6.52 (s, 1H), 5.27 (s, 1H), 1.21 (s, 9H); ¹³C NMR (125 MHz, CDCl₃): 169.4, 152.3, 131.3, 130.8, 128.1, 122.9, 80.2, 51.1, 28.9; ESI-MS: 232 [(M-H)⁺, 100%].

Conclusions

In conclusion, we have developed an efficient protocol for the preparation of α -hydroxyamides via boric acid mediated addition of isonitriles on to aldehydes. The reaction in general provides good yields of the products under very mild reaction conditions. We have applied the methodology for intramolecular version to synthesize functionalized benzoxaboroles. Owing to the importance of hydroxyamides and benzoxaboroles as synthetic intermediates and also

as medicinal agents, we believe that the current methodology will find applications in organic and medicinal chemistry.

Acknowledgments

We thank the Departments of Chemistry and Biochemistry, Rowan University, and University of Minnesota Duluth for the funding. Partial support for this work was provided by research grants from the National Institutes of Health (CA129993) (VRM) and Whiteside Institute for Clinical Research (VRM).

References

1. Boric Acid General Fact Sheet-National Pesticide Information Center.
<http://npic.orst.edu/factsheets/boricgen.pdf>
- 2 (a). Shteinberg LY. Russ. J. Appl. Chem 2009;82:613–617. (b) Kondaiah GCM, Reddy LA, Babu KS, Gurav VM, Hugu KM, Bandichhor R, Reddy PP, Bhattacharya A, Anand RV. Tetrahedron Lett 2008;49:106–109. (c) Nath J, Chaudhuri MK. Green Chemistry Letters and Reviews 2008;1:223–230. (d) Tu S-J, Zhu XT, Fang F, Zhang X-J, Zhu S-L, Li T-J, Shi D-Q, Wang X-S, Ji S-J. Chin. J. Chem 2005;23:596–598. (e) Chaudhuri MK, Hussain S, Kantam ML, Neelima B. Tetrahedron Lett 2005;46:8329–8331. (f) Maki T, Ishihara K, Yamamoto H. Org. Lett 2005;7:5047–5050. [PubMed: 16235954] (g) Stapp PR. J. Org. Chem 1973;38:1433–1434.
3. Schenck HA, Lenkowski PW, Mukherjee IC, Kob SH, Stables JP, Patel MK, Brown ML. Bioorg. Med. Chem 2004;12:979–993. [PubMed: 14980611] (b) Kuduk SD, Chang RK, DiPardo RM, Di Marco CN, Murphy KL, Ransom RW, Reiss DR, Tang C, Prueksaritanont T, Pettibone DJ, Bock MG. Bioorg. Med. Chem. Lett 2008;18:5107–5110. [PubMed: 18722115]
- 4 (a). Banfi L, Riva R. Organic Reactions 2005;65:1–141. (b) Domling A, Ugi I. Angew Chem. Int. Ed 2000;39:3168–3210.
- 5 (a). Yue T, Wang M-X, Wang D-X, Masson G, Zhu J. Angew. Chem. Intl. Ed 2009;48:6717–21. (b) Yue T, Wang MX, Wang DX, Masson G, Zhu J. J. Org. Chem 2009;74:8396–8399. [PubMed: 19788173] (c) Mihara H, Xu Yingjie, Shepherd NE, Matsunaga S, Shibasaki M. J. Am. Chem. Soc 2009;131:8384–8385. [PubMed: 19485325] (d) Wang S, Wang M-X, Wang D-X, Zhu J. Eur. J. Org. Chem 2007;24:4076–4080. (e) Wang S-X, Wang M-X, Wang D-X, Zhu J. Org. Lett 2007;9:3615–3618. [PubMed: 17685533] (f) Denmark SE, Bui T. J. Org. Chem 2005;70:10190–10193. [PubMed: 16292871] (g) Denmark SE, Fan Y. J. Am. Chem. Soc 2003;125:7825–7827. [PubMed: 12823000] (h) Seebach D, Adam G, Gees T, Schiess M, Weigand W. Chem. Ber 1988;121:507–17. (i) Hagedorn I, Eholzer U. Chem. Ber 1965;98:936–940. (i).
- 6 (a). Nicolaou KC, Natarajan S, Li H, Jain NF, Hughes R, Solomon ME, Ramanjulu JM, Boody CNC, Takayanagi M. Angew. Chem. Int. Ed 1998;37:2708. (b) Nicolaou KC, Li H, Boody CNC, Ramanjulu JM, Yue TY, Natarajan S, Chu XJ, Brase S, Rubsam F. Chem. Eur. J 1999;5:2584. (c) Tan YL, White AJP, Widdowson DA, Wilhelm R, Williams DJ. J. Chem. Soc. Perkin Trans 1 2001:3269.
- 7 (a). Akama T, Baker SJ, Zhang YK, Hernandez V, Zhou H, Sanders V, Freund Y, Kimura R, Maples KR, Plattner JJ. Bioorg. Med. Chem. Lett 2009;19:2129–2132. [PubMed: 19303290] (b) Wozniack AA, Cyranski MK, Zubrowska A, Sporzynski A. J. Organomet. Chem 2009;694:3533–3541. (c) Baker SJ, Zhang YK, Akama T, Lau A, Zhou H, Hernandez V, Mao W, Alley MRK, Sanders V, Plattner JJ. J. Med. Chem 2006;49:4447–4450. [PubMed: 16854048]

Scheme 1.
Synthesis of α -Amido Benzoxaboroles

Isonitrile	Product	Yield
<p>4a</p>	78%
<p>4b</p>	74%

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
75%

4c

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
<p style="text-align: center; font-size: 2em; font-weight: bold;">4d</p>		70%

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
<p>4e</p>	<p>4f</p>	71%
<p>4f</p>	74%

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
61%

4g

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
<p>4h</p>	64%
<p>4i</p>	72%

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
<p>4j</p>	78%

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
74%
72%

4k

4l

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3

Isonitrile	Product	Yield
82%
76%

4m

4n

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
80%
65%

4o

4p

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
<p>4q</p>	68%
<p>4r</p>	67%

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
------------	---------	-------

4s

75%

4t

72%

Tetrahedron Lett. Author manuscript; available in PMC 2011 February 3.

Isonitrile	Product	Yield
78%

4u