Published in final edited form as:

Intensive Care Med. 2010 January ; 36(1): 123–130. doi:10.1007/s00134-009-1654-6.

Changes in peroxisome proliferator activated receptor-gamma activity in children with septic shock

Jennifer M. Kaplan, MD, MS, Alvin Denenberg, Marie Monaco, Marcheale Nowell, Hector Wong, MD, and Basilia Zingarelli, MD, PhD

Cincinnati Children's Hospital Medical Center and The University of Cincinnati College of Medicine, 3333 Burnet Avenue, MLC 2005, Cincinnati, OH 45229-3039

Abstract

Purpose—To assess changes in peroxisome proliferator-activated receptor- γ (PPAR γ) in peripheral blood mononuclear cells (PBMC) from critically ill children with sepsis. Additionally, to investigate the effects of sepsis on the endogenous activator of PPAR γ , 15-deoxy- $\Delta^{12,14}$ -PGJ₂ (15d-PGJ₂), and the downstream targets of PPAR γ activity, adiponectin and resistin.

Methods—Single-center, prospective case-control study in critically ill children with systemic inflammatory response syndrome, sepsis, or septic shock.

Results—PPAR γ nuclear protein expression was decreased but PPAR γ activity was increased in PBMC from children with septic shock compared to controls. The PPAR γ activity on day 1 was significantly higher in patients with a higher PRISM score compared with controls (mean 0.22 Optical Density (OD) $\pm$ SEM 0.03 vs. 0.12 OD $\pm$ 0.02; $p < 0.001$). Patients with resolved sepsis had increased levels of the endogenous PPAR γ ligand, 15d-PGJ₂, compared to patients with SIRS and septic shock (77.7 $\pm$ 21.7 pg/ml vs. 58 $\pm$ 16.5 pg/ml; $p = 0.03$). Plasma HMWA and resistin levels were increased in patients with septic shock on day 1 and were significantly higher in patients with higher PRISM scores. Non-survivors from sepsis had higher resistin levels on the first day of hospitalization compared with survivors from septic shock (660 ng/ml interquartile range (IQR) 585–833 ng/ml vs. 143 ng/ml IQR 66–342 ng/ml; $p < 0.05$).

Conclusions—Sepsis is associated with altered PPAR γ expression and activity in PBMC. Plasma adipokines correlate with risk of mortality scores in sepsis and may be useful biomarkers. Further studies are needed to understand the mechanisms underlying changes in PPAR γ in sepsis.

Keywords

sepsis; resistin; adiponectin; 15-deoxy- $\Delta^{12,14}$ -PGJ₂; PPAR γ ; cytokines

Introduction

Peroxisome proliferator-activated receptor- γ (PPAR γ) is a ligand-binding nuclear receptor and activation of PPAR γ controls the inflammatory response [1,2]. The synthetic insulin-sensitizing drugs, thiazolidinediones (TZDs), and the natural cyclopentenone prostaglandins are specific PPAR γ ligands [3,4]. 15-deoxy- $\Delta^{12,14}$ -PGJ₂ (15d-PGJ₂) inhibited the phorbol myristyl acetate-induced TNF α production in human monocytes [2]. The mechanism of cytokine inhibition in activated monocytes/macrophages occurs, in part, through repression of several inflammatory response genes, including activator protein-1 and nuclear factor- κ B (NF- κ B) [1]. 15d-PGJ₂ exerts beneficial anti-inflammatory effects, in part, through inhibition of

NF- κ B activation [5,6]. The inhibition of the inflammatory response correlates with improved survival in clinically relevant models of septic shock [7–9].

The expression, production and activity of PPAR γ are affected in inflammatory conditions. We have previously demonstrated that in the hypodynamic phase of septic shock PPAR γ expression was downregulated on the endothelium of thoracic aortas in rats [7]. Furthermore, the sepsis-induced reduction in PPAR γ expression was reversed by *in vivo* treatment with PPAR γ ligands. In an experimental model of polymicrobial sepsis, Zhou et al. demonstrate that hepatic PPAR γ protein and gene expression is preserved in the early stages of sepsis and decreased in the late stages of sepsis [10]. These findings suggest that there are kinetic differences in PPAR γ expression after an inflammatory response.

Human data suggests that nuclear receptors are altered in inflammatory disease states. PPAR γ is altered in the inflammatory diseases sarcoidosis, colitis and multiple sclerosis [11–13]. In children with sepsis, glucocorticoid receptor mRNA expression was decreased in neutrophils on the first day of hospitalization but expression normalized as children recovered from their illness [14]. In adults with sepsis, PPAR γ mRNA expression is increased in T lymphocytes and neutrophils when compared to control subjects [15,16].

In the current study, we assessed the protein expression and activity of PPAR γ in peripheral blood mononuclear cells from a cohort of critically ill children with sepsis. In addition, we investigated the effects of sepsis on the endogenous activator of PPAR γ , 15d-PGJ₂, and the downstream targets of PPAR γ activity, high molecular weight adiponectin (HMWA) and resistin.

Materials and Methods

Patients and Study Design

The study was reviewed and approved by the Cincinnati Children's Hospital Medical Center Institutional Review Board. Patients admitted to the pediatric intensive care unit (PICU) with the systemic inflammatory response syndrome (SIRS), sepsis, or septic shock as defined according to the International Pediatric Sepsis Consensus Conference were considered eligible for the study [17]. Patients who recovered from their illness with sepsis and remained hospitalized but did not meet the physiologic parameters of any category of sepsis anymore were categorized as “resolved”. Control patients were obtained from children undergoing cardiac catheterization who exhibited no evidence of systemic inflammation. Patients with leukopenia [white blood cell (WBC) count < 1,000 cells/mm³] or ongoing cardiopulmonary resuscitation were excluded from the study. Control patients with evidence of a recent illness were excluded. Seventy-four patients were enrolled in the study however it was not possible to perform all experiments with all of the specimens because of the limited amount of blood obtained from each patient.

Blood Samples

After obtaining informed consent 5mL of blood was obtained within the first 24h of admission to the PICU and processed as described below. Subsequent blood samples were obtained from each patient on days 3 and 7. However not all patients provided a blood sample on day 7 because of death, lack of vascular access, or discharge from the hospital. One 5ml sample was obtained from each control patient.

Clinical Data

Clinical data were collected prospectively throughout the study period. Severity of illness at study entry was calculated using the Pediatric Risk of Mortality (PRISM) II score [18].

Isolation of Plasma and Peripheral Blood Mononuclear Cells (PBMC)

Fresh blood samples were collected in sodium citrate tubes and transported to the Critical Care Medicine laboratories on ice. Samples were centrifuged at 1500 rpm for 5 minutes and plasma was obtained and stored at -80°C . The remaining blood was carefully layered onto a Percoll gradient (Amersham Biosciences, Piscataway, NJ) and samples centrifuged at 1500 RPM for 30 minutes. After centrifugation the buffy coat was removed, washed in Hanks BSA and stored at -80°C until protein extraction.

Isolation of PBMC Confirmed by Flow Cytometry

To confirm our experimental technique of isolating PBMC using a Percoll gradient, cells were analyzed by flow cytometry. Whole blood obtained from healthy volunteers (3.5 ml) was carefully layered onto a percoll gradient and PBMC were isolated as described above. Phosphate-buffered saline (PBS) diluted whole blood (100 μl) was blocked with rat serum for 20 minutes on ice. After centrifugation to obtain a pellet, cells were incubated with saturating amounts of fluorescein isothiocyanate (FITC)-conjugated anti-CD66b murine monoclonal antibody, allophycocyanin (APC)-conjugated anti-CD14 murine monoclonal antibody or isotype control (BD Biosciences, San Jose, California). Samples were washed and resuspended in phosphate-buffered saline with 2% bovine serum albumin (BSA). Flow cytometric analysis was performed and confirmed that the percoll layer included monocytes and lymphocytes and excluded neutrophils (data not shown).

Data Analysis

Not all blood samples were tested for all analytes. Values in the figures and text are expressed as mean and standard error of the mean for parametric data and as median and interquartile range for nonparametric nature of the data. The number of subjects per group is referred to as *n*. For comparison of two groups, Student's *t*-tests were used. For non-parametric data the Mann-Whitney Rank Sum Test (MWRST) was used. When more than two groups were compared for nonparametric data the Kruskal-Wallis analysis of variance with the Dunn *post hoc* test was used. A two-tailed $p < 0.05$ was considered significant.

Results

Patient Characteristics and Plasma Cytokine Levels

There was no difference in age or body mass index (BMI) between the control group (children undergoing cardiac catheterization without signs of systemic inflammation), SIRS/sepsis, or septic shock groups on admission (Table 1). Patients in the septic shock group had a significantly higher PRISM score and number of organ failures compared to the SIRS/septic group. Patients with septic shock had higher plasma levels of $\text{TNF}\alpha$, IL-6 and IL-8 compared to the control group.

PPAR γ Nuclear Protein Expression is Decreased in PBMC from Children with Septic Shock

To determine the effect of sepsis on PPAR γ expression, PBMC were collected from patients with various categories of sepsis on day 1, 3 or 7 of their illness and protein expression evaluated by western blot analysis. Patients were grouped according to their sepsis category. Therefore multiple samples were obtained from the same patient if they fulfilled the requirement for the diagnostic category on day 1, 3, or 7 of their illness. A representative western blot demonstrates a decrease in PPAR γ protein expression in SIRS and septic shock patients compared to controls (Fig. 1A). Quantitative analysis demonstrates that PPAR γ levels were significantly decreased in patients with septic shock when compared to control patients [median 66.5 absolute intensity, interquartile range (IQR) 60–78 absolute intensity vs. median 113.5 absolute intensity, IQR 108–133; $p < 0.05$] (Fig. 1B). Patients with resolved sepsis who recovered from their illness

with sepsis and remained hospitalized but did not meet the physiologic parameters of any category of sepsis anymore we categorized as “resolved”. Although not statistically significant, patients with resolved sepsis had an increase in PPAR γ protein levels compared to septic shock. These levels were similar to levels demonstrated in control patients.

PPAR γ Activity is Increased in PBMC from Children with Septic Shock

Nuclear PPAR γ activity in PBMC from children with septic shock was determined by use of a PPAR γ transcription factor assay kit. Interestingly, contrary to PPAR γ expression findings, median PPAR γ activity levels were higher in patients with SIRS and septic shock when compared to control patients (Fig. 2). Patients with resolved sepsis had median PPAR γ activity levels similar to control patients. The PPAR γ activity level within the first day of hospitalization was significantly higher in patients with a PRISM score >10 compared with control patients (mean 0.22 OD $\pm$ SEM 0.03 vs. 0.12 OD $\pm$ 0.02; $p < 0.001$). Thus, our data suggest that while PPAR γ protein expression is decreased in septic shock the activity of PPAR γ is increased suggesting that additional mediators result in PPAR γ activation during septic shock.

The Effect of Sepsis on Plasma 15d-PGJ₂ Levels in Children with Systemic Inflammation from Sepsis

Since downregulation of protein expression did not correlate with PPAR γ activity in PBMC, we determined whether sepsis is associated with changes in levels of the endogenous activator of PPAR γ . Plasma levels of 15d-PGJ₂ were unchanged in patients with SIRS and septic shock when compared to controls. However, patients with resolved sepsis had elevated levels of 15d-PGJ₂ compared to patients with SIRS and septic shock (77.7 $\pm$ 21.7 pg/ml vs. 58 $\pm$ 16.5 pg/ml; $p = 0.03$). Thus, our data suggest that sepsis increases the endogenous ligand for PPAR γ and may be one factor increasing PPAR γ activity levels despite lower protein levels.

Both High Molecular Weight Adiponectin (HMWA) and Resistin are Increased Early in Septic Shock and Remain Elevated in Children with Resolving Sepsis

Because of the importance of PPAR γ in adipocyte proliferation we sought to measure the effects of sepsis on adipocyte proteins. We utilized HMWA plasma levels as an easily obtainable mechanism to measure an endpoint of PPAR γ activity for two reasons. First, HMWA has a PPAR response element in its promoter region. Second, levels of high molecular weight adiponectin correlate with synthetic PPAR γ agonist treatment [19]. We found that HMWA levels are increased in patients with septic shock on day 1 compared to controls (8.0 μ g/ml IQR 4.8–12.3 vs. 3.3 μ g/ml IQR 2.4–9.4; $p < 0.05$) (Fig. 3). The plasma levels of HMWA correlate with risk of mortality scores. HMWA levels were higher in patients with a PRISM score >21 (13.7 $\pm$ 4.3 μ g/ml) when compared to patients with a low PRISM score (≤ 10) (7 $\pm$ 1 μ g/ml; $p < 0.05$) (Fig. 4). Patients with resolved sepsis also have higher levels of HMWA (10.3 μ g/ml IQR 5.4–12.7 μ g/ml) when compared to controls (4.3 μ g/ml IQR 2.6–9.8 μ g/ml; $p < 0.05$).

We demonstrate that plasma resistin levels are increased in patients with septic shock on the first day of hospitalization compared to controls (Fig. 3). Furthermore, patients with septic shock had significantly higher resistin levels on the first day of hospitalization compared to patients with SIRS/sepsis (597 ng/ml IQR 273–851 ng/ml vs. 78 ng/ml IQR 60–325; $p < 0.01$). Resistin levels also correlate with risk of mortality scores and outcomes. We demonstrate that plasma levels of resistin on the first day of hospitalization from sepsis were significantly higher in patients with a PRISM score >21 when compared to patients with a PRISM score ≤ 10 (Fig. 4) (682.4 $\pm$ 171.6 ng/ml vs. 237.4 $\pm$ 62.5 ng/ml; $p < 0.05$). As evidence that resistin may be a useful biomarker, non-survivors from sepsis had significantly higher resistin levels on the first day of hospitalization compared with survivors from septic shock (660 ng/ml IQR 585–833 ng/ml vs. 143 ng/ml IQR 66–342 ng/ml; $p < 0.05$).

Discussion

Our results demonstrate that sepsis is associated with altered PPAR γ expression and activation in PBMC. First, in children with septic shock PPAR γ nuclear protein expression is decreased however PPAR γ activity is increased. This occurs most likely as a result of an increase in production of the endogenous ligand, 15d-PGJ₂. Second, PPAR γ activity in early sepsis correlates with the risk of mortality. Third, the adipokines resistin and high molecular weight adiponectin correlate with PPAR γ activity in sepsis and can be useful biomarkers in sepsis. Furthermore, these adipokines are elevated in the resolution of sepsis and may play a role in the compensatory anti-inflammatory response syndrome.

Our findings are consistent with previous animal studies which demonstrate that sepsis alters PPAR γ expression [7,20]. The current study is the first to demonstrate alterations in PPAR γ in children with sepsis. We found that PPAR γ protein expression in PBMC was decreased in children with septic shock. This is consistent with findings in PBMC from patients with multiple sclerosis, a disease with significant inflammation. Multiple sclerosis patients had a ~65% reduction in PPAR γ protein expression in PBMC compared with healthy donors [13]. Although beyond the scope of this current study a possible explanation for the decrease in PPAR γ protein expression demonstrated in the current study may occur through post-translational modifications of PPAR γ [21]. The activation function-1 domain of PPAR γ contains a consensus mitogen-activated protein kinase site and serine phosphorylation leads to inhibition of PPAR γ transactivation [22,23]. Once phosphorylated, PPAR γ becomes degraded by the ubiquitin-proteasome system [24]. These mechanistic changes may provide an explanation for our current findings and will need to be explored in further studies.

Contrary to our data, studies in adults with sepsis demonstrate an increase in PPAR γ mRNA expression in T cells and neutrophils [15,16]. Differences in PPAR γ detected in the current study and previously published data may occur through cell type differences in PPAR γ . PBMC includes a mixed population of monocytes and lymphocytes while other studies demonstrate changes in isolated neutrophils and T lymphocytes [15,16]. Further studies are needed to confirm our findings in specific cell populations.

Children with resolved sepsis have elevated 15d-PGJ₂ levels similar to levels found in the synovial fluid of rheumatoid and osteo-arthritis patients [25]. This finding confirms previous suggestions that an endogenous activator of PPAR γ is present in septic blood. 15d-PGJ₂ was found in blood from septic rats and sera from septic patients were able to activate PPAR γ [16,26]. It is not surprising that 15d-PGJ₂ is activated during the inflammatory response from sepsis. 15d-PGJ₂ is produced from arachidonic acid via cyclo-oxygenases (COX), enzymes known to be induced after LPS stimulation [27]. Therefore, 15d-PGJ₂ levels may be increased in sepsis as a compensatory mechanism and contribute to the increase in PPAR γ activity. As a result, despite a decrease in PPAR γ protein expression it is plausible to hypothesize that PBMC of septic patients has a sufficient receptor reserve, which can be still recruited by endogenous ligands.

Our data are consistent with published data demonstrating that resistin levels are increased in patients with septic shock on the first day of hospitalization [28]. We demonstrate for the first time that day one resistin levels correlate with survival outcomes and disease severity. Non-survivors from sepsis have significantly higher resistin levels on the first day of their illness compared to survivors. These findings are in agreement with data that demonstrates that resistin levels correlate with inflammatory markers [29]. Resistin levels are increased from human PBMC after treatment with LPS [28]. As part of a feedback loop, PBMC and synovial leukocytes respond to resistin by producing pro-inflammatory cytokines [30]. Unlike

adiponectin, both upregulation and downregulation of resistin gene expression occurs after TZD treatment [31,32].

Adiponectin has anti-inflammatory effects and is affected by PPAR γ agonists. Expression of the adiponectin gene is induced by PPAR γ ligands via direct binding to the peroxisome proliferator response element in the adiponectin promoter [33]. Multiple studies have demonstrated that treatment with thiazolidinediones increase adiponectin mRNA levels in adipocytes and adipose tissues [34,35]. Few studies have evaluated the effects of sepsis on plasma adipokines. Adiponectin levels were lower in polymicrobial sepsis and adiponectin knockout mice had higher inflammatory cytokine production and higher mortality after polymicrobial sepsis compared to wild-type mice [36,37]. But human studies have failed to detect alterations in plasma adiponectin levels after endotoxin injection [38,39]. One explanation for the lack of difference in adiponectin levels after endotoxin is that those subjects were not severely ill. Second, total adiponectin may not accurately correlate with inflammation. The high molecular weight form of adiponectin has been proposed as the most potent form mediating the metabolic actions of adiponectin and the form preferentially increased by TZDs [19].

We acknowledge that our study has many limitations. The major limitation of this study is the lack of power due to the small sample size. This may contribute to the small PPAR γ differences in patients with sepsis. Another limitation is the absence of information on pre-hospital administration of anti-pyretic medications, such as non-steroidal anti-inflammatory drugs (NSAIDs), which may affect PPAR γ activity and are often used in children. NSAIDs have anti-inflammatory effects through inhibition of cyclo-oxygenase activity. High concentrations of NSAIDs can activate PPAR γ similarly to the natural ligand 15d-PGJ₂ [40]. Although our current study did not include data on NSAID use, it is possible that PPAR γ activity could be affected by pre-hospital administration of this drug. Another limitation of this study is the small amount of blood volume obtained from each patient not all blood samples were tested for all analytes. We recognize that this study is descriptive and not mechanistic. Although animal studies have demonstrated alterations in PPAR γ in sepsis no other studies demonstrate changes in PPAR γ in children with sepsis. Furthermore, the results of this study provide the foundation for future studies to investigate the mechanism involved in the PPAR γ response to sepsis.

Conclusions

Sepsis is associated with altered PPAR γ expression and activity in PBMC. PPAR γ activity in sepsis correlates with plasma adipokines and risk of mortality score. However, taken together with our previously published *in vivo* studies, these findings suggest that an increase in PPAR γ activity may represent a counter-regulatory mechanism in the inflammatory response. Further studies are needed to understand the mechanistic changes altering PPAR γ expression and activity in sepsis.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We would like to thank the parents who enrolled their children into this study during an emotionally difficult time. Supported, in part, by grants K12 HD028827 (JK), T32 ES10957 (JK), R01 GM067202 (BZ), and R01 GM064619 (HW) from the NIH, Bethesda, MD, and by the Translational Research Initiative at Cincinnati Children's Hospital Medical Center (JK).

References

1. Ricote M, Li AC, Willson TM, Kelly CJ, Glass CK. The peroxisome proliferator-activated receptor-gamma is a negative regulator of macrophage activation. *Nature* 1998;391:79–82. [PubMed: 9422508]
2. Jiang C, Ting AT, Seed B. PPAR-gamma agonists inhibit production of monocyte inflammatory cytokines. *Nature* 1998;391:82–86. [PubMed: 9422509]
3. Kliewer SA, Lenhard JM, Willson TM, Patel I, Morris DC, Lehmann JM. A prostaglandin J2 metabolite binds peroxisome proliferator-activated receptor gamma and promotes adipocyte differentiation. *Cell* 1995;83:813–819. [PubMed: 8521498]
4. Palmer CN, Hsu MH, Griffin HJ, Johnson EF. Novel sequence determinants in peroxisome proliferator signaling. *J Biol Chem* 1995;270:16114–16121. [PubMed: 7608174]
5. Straus DS, Pascual G, Li M, Welch JS, Ricote M, Hsiang CH, Sengchanthalangsy LL, Ghosh G, Glass CK. 15-deoxy-delta 12,14-prostaglandin J2 inhibits multiple steps in the NF-kappa B signaling pathway. *Proc Natl Acad Sci U S A* 2000;97:4844–4849. [PubMed: 10781090]
6. Kaplan J, Cook JA, O'Connor M, Zingarelli B. Peroxisome proliferator-activated receptor gamma is required for the inhibitory effect of ciglitazone but not 15-deoxy-Delta 12,14-prostaglandin J2 on the NFkappaB pathway in human endothelial cells. *Shock* 2007;28:722–726. [PubMed: 17621259]
7. Zingarelli B, Sheehan M, Hake PW, O'Connor M, Denenberg A, Cook JA. Peroxisome proliferator activator receptor-gamma ligands, 15-deoxy-Delta(12,14)-prostaglandin J2 and ciglitazone, reduce systemic inflammation in polymicrobial sepsis by modulation of signal transduction pathways. *J Immunol* 2003;171:6827–6837. [PubMed: 14662889]
8. Kaplan JM, Cook JA, Hake PW, O'Connor M, Burroughs TJ, Zingarelli B. 15-deoxy-delta12,14-prostaglandin J2 (15D-PGJ2), a peroxisome proliferator activated receptor gamma ligand, reduces tissue leukosequestration and mortality in endotoxic shock. *SHOCK* 2005;24:59–65. [PubMed: 15988322]
9. Haraguchi G, Kosuge H, Maejima Y, Suzuki J, Imai T, Yoshida M, Isobe M. Pioglitazone reduces systematic inflammation and improves mortality in apolipoprotein E knockout mice with sepsis. *Intensive Care Med* 2008;34:1304–1312. [PubMed: 18283431]
10. Zhou M, Wu R, Dong W, Simms HH, Wang P. Hepatic peroxisome proliferator-activated receptor-gamma (PPAR-gamma) is downregulated in sepsis. *SHOCK* 2004;21:39.
11. Culver DA, Barna BP, Raychaudhuri B, Bonfield TL, Abraham S, Malur A, Farver CF, Kavuru MS, Thomassen MJ. Peroxisome proliferator-activated receptor gamma activity is deficient in alveolar macrophages in pulmonary sarcoidosis. *Am J Respir Cell Mol Biol* 2004;30:1–5. [PubMed: 14512375]
12. Han X, Osuntokun B, Benight N, Loesch K, Frank SJ, Denson LA. Signal transducer and activator of transcription 5b promotes mucosal tolerance in pediatric Crohn's disease and murine colitis. *Am J Pathol* 2006;169:1999–2013. [PubMed: 17148664]
13. Klotz L, Schmidt M, Giese T, Sastre M, Knolle P, Klockgether T, Heneka MT. Proinflammatory stimulation and pioglitazone treatment regulate peroxisome proliferator-activated receptor gamma levels in peripheral blood mononuclear cells from healthy controls and multiple sclerosis patients. *J Immunol* 2005;175:4948–4955. [PubMed: 16210596]
14. van den Akker EL, Koper JW, Joosten K, de Jong FH, Hazelzet JA, Lamberts SW, Hokken-Koelega AC. Glucocorticoid receptor mRNA levels are selectively decreased in neutrophils of children with sepsis. *Intensive Care Med* 2009;35:1247–1254. [PubMed: 19373457]
15. Reddy RC, Narala VR, Keshamouni VG, Milam JE, Newstead MW, Standiford TJ. Sepsis-induced inhibition of neutrophil chemotaxis is mediated by activation of peroxisome proliferator-activated receptor-{gamma}. *Blood* 2008;112:4250–4258. [PubMed: 18535203]
16. Soller M, Tautenhahn A, Brune B, Zacharowski K, John S, Link H, von Knethen A. Peroxisome proliferator-activated receptor gamma contributes to T lymphocyte apoptosis during sepsis. *J Leukoc Biol* 2006;79:235–243. [PubMed: 16380602]
17. Goldstein B, Giroir B, Randolph A. International pediatric sepsis consensus conference: definitions for sepsis and organ dysfunction in pediatrics. *Pediatr Crit Care Med* 2005;6:2–8. [PubMed: 15636651]

18. Pollack MM, Ruttimann UE, Getson PR. Pediatric risk of mortality (PRISM) score. *Crit Care Med* 1988;16:1110–1116. [PubMed: 3048900]
19. Pajvani UB, Hawkins M, Combs TP, Rajala MW, Doebber T, Berger JP, Wagner JA, Wu M, Knopps A, Xiang AH, Utzschneider KM, Kahn SE, Olefsky JM, Buchanan TA, Scherer PE. Complex distribution, not absolute amount of adiponectin, correlates with thiazolidinedione-mediated improvement in insulin sensitivity. *J Biol Chem* 2004;279:12152–12162. [PubMed: 14699128]
20. Zhou M, Wu R, Dong W, Jacob A, Wang P. Endotoxin downregulates peroxisome proliferator-activated receptor-gamma via the increase in TNF-alpha release. *Am J Physiol Regul Integr Comp Physiol* 2008;294:R84–R92. [PubMed: 17989144]
21. Han J, Hajjar DP, Tauras JM, Feng J, Gotto AM Jr, Nicholson AC. Transforming growth factor-beta1 (TGF-beta1) and TGF-beta2 decrease expression of CD36, the type B scavenger receptor, through mitogen-activated protein kinase phosphorylation of peroxisome proliferator-activated receptor-gamma. *J Biol Chem* 2000;275:1241–1246. [PubMed: 10625669]
22. Camp HS, Tafuri SR. Regulation of peroxisome proliferator-activated receptor gamma activity by mitogen-activated protein kinase. *J Biol Chem* 1997;272:10811–10816. [PubMed: 9099735]
23. Adams M, Reginato MJ, Shao D, Lazar MA, Chatterjee VK. Transcriptional activation by peroxisome proliferator-activated receptor gamma is inhibited by phosphorylation at a consensus mitogen-activated protein kinase site. *J Biol Chem* 1997;272:5128–5132. [PubMed: 9030579]
24. Hauser S, Adelmant G, Sarraf P, Wright HM, Mueller E, Spiegelman BM. Degradation of the peroxisome proliferator-activated receptor gamma is linked to ligand-dependent activation. *J Biol Chem* 2000;275:18527–18533. [PubMed: 10748014]
25. Shan ZZ, Masuko-Hongo K, Dai SM, Nakamura H, Kato T, Nishioka K. A potential role of 15-deoxy-delta(12,14)-prostaglandin J2 for induction of human articular chondrocyte apoptosis in arthritis. *J Biol Chem* 2004;279:37939–37950. [PubMed: 15213234]
26. Gilroy DW, Colville-Nash PR, Willis D, Chivers J, Paul-Clark MJ, Willoughby DA. Inducible cyclooxygenase may have anti-inflammatory properties. *Nat Med* 1999;5:698–701. [PubMed: 10371510]
27. Shibata T, Kondo M, Osawa T, Shibata N, Kobayashi M, Uchida K. 15-deoxy-delta 12,14-prostaglandin J2. A prostaglandin D2 metabolite generated during inflammatory processes. *J Biol Chem* 2002;277:10459–10466. [PubMed: 11786541]
28. Sunden-Cullberg J, Nystrom T, Lee ML, Mullins GE, Tokics L, Andersson J, Norrby-Teglund A, Treutiger CJ. Pronounced elevation of resistin correlates with severity of disease in severe sepsis and septic shock. *Crit Care Med* 2007;35:1536–1542. [PubMed: 17452927]
29. Kawanami D, Maemura K, Takeda N, Harada T, Nojiri T, Imai Y, Manabe I, Utsunomiya K, Nagai R. Direct reciprocal effects of resistin and adiponectin on vascular endothelial cells: a new insight into adipocytokine-endothelial cell interactions. *Biochem Biophys Res Commun* 2004;314:415–419. [PubMed: 14733921]
30. Bokarewa M, Nagaev I, Dahlberg L, Smith U, Tarkowski A. Resistin, an adipokine with potent proinflammatory properties. *J Immunol* 2005;174:5789–5795. [PubMed: 15843582]
31. Lu SC, Shieh WY, Chen CY, Hsu SC, Chen HL. Lipopolysaccharide increases resistin gene expression in vivo and in vitro. *FEBS Lett* 2002;530:158–162. [PubMed: 12387885]
32. Way JM, Gorgun CZ, Tong Q, Uysal KT, Brown KK, Harrington WW, Oliver WR Jr, Willson TM, Kliewer SA, Hotamisligil GS. Adipose tissue resistin expression is severely suppressed in obesity and stimulated by peroxisome proliferator-activated receptor gamma agonists. *J Biol Chem* 2001;276:25651–25653. [PubMed: 11373275]
33. Iwaki M, Matsuda M, Maeda N, Funahashi T, Matsuzawa Y, Makishima M, Shimomura I. Induction of adiponectin, a fat-derived antidiabetic and antiatherogenic factor, by nuclear receptors. *Diabetes* 2003;52:1655–1663. [PubMed: 12829629]
34. Maeda N, Takahashi M, Funahashi T, Kihara S, Nishizawa H, Kishida K, Nagaretani H, Matsuda M, Komuro R, Ouchi N, Kuriyama H, Hotta K, Nakamura T, Shimomura I, Matsuzawa Y. PPARgamma ligands increase expression and plasma concentrations of adiponectin, an adipose-derived protein. *Diabetes* 2001;50:2094–2099. [PubMed: 11522676]
35. Combs TP, Wagner JA, Berger J, Doebber T, Wang WJ, Zhang BB, Tanen M, Berg AH, O'Rahilly S, Savage DB, Chatterjee K, Weiss S, Larson PJ, Gottesdiener KM, Gertz BJ, Charron MJ, Scherer

- PE, Moller DE. Induction of adipocyte complement-related protein of 30 kilodaltons by PPARgamma agonists: a potential mechanism of insulin sensitization. *Endocrinology* 2002;143:998–1007. [PubMed: 11861525]
36. Teoh H, Quan A, Bang KW, Wang G, Lovren F, Vu V, Haitsma JJ, Szmitko PE, Al-Omran M, Wang CH, Gupta M, Peterson MD, Zhang H, Chan L, Freedman J, Sweeney G, Verma S. Adiponectin deficiency promotes endothelial activation and profoundly exacerbates sepsis-related mortality. *Am J Physiol Endocrinol Metab* 2008;295:E658–E664. [PubMed: 18628355]
37. Uji Y, Yamamoto H, Tsuchihashi H, Maeda K, Funahashi T, Shimomura I, Shimizu T, Endo Y, Tani T. Adiponectin deficiency is associated with severe polymicrobial sepsis, high inflammatory cytokine levels, and high mortality. *Surgery* 2009;145:550–557. [PubMed: 19375615]
38. Keller P, Moller K, Krabbe KS, Pedersen BK. Circulating adiponectin levels during human endotoxaemia. *Clin Exp Immunol* 2003;134:107–110. [PubMed: 12974762]
39. Anderson PD, Mehta NN, Wolfe ML, Hinkle CC, Pruscino L, Comiskey LL, Tabita-Martinez J, Sellers KF, Rickels MR, Ahima RS, Reilly MP. Innate immunity modulates adipokines in humans. *J Clin Endocrinol Metab* 2007;92:2272–2279. [PubMed: 17374708]
40. Lehmann JM, Lenhard JM, Oliver BB, Ringold GM, Kliewer SA. Peroxisome proliferator-activated receptors alpha and gamma are activated by indomethacin and other non-steroidal anti-inflammatory drugs. *J Biol Chem* 1997;272:3406–3410. [PubMed: 9013583]

Fig. 1.

Nuclear PPAR γ protein expression is decreased in PBMC in septic shock. (A) Representative western blot of nuclear PPAR γ protein expression in PBMC from two patients with various categories of sepsis on day 1 or 3 and in four control patients. (B) Box and whisker plot of PPAR γ nuclear protein levels from PBMC by western blot analysis based on densitometric analysis of the absolute intensity. Patients were grouped according to their sepsis category and therefore multiple samples were obtained from the same patient if they fulfilled the requirement for the diagnostic category on day 1,3, or 7 of their illness. The vertical box represents the 25th percentile (bottom line), median (middle line), and 75th percentile (top line) values

Fig. 2. Nuclear PPAR γ activity is increased in PBMC in septic shock. Box and whisker plot of nuclear PPAR γ activity levels from PBMC. Patients were grouped according to their sepsis category. The vertical box represents the 25th percentile (bottom line), median (middle line), and 75th percentile (top line) values, while the error bars represent the 10th and 90th percentile values. The dots represent values outside the 10th and 90th percentile. *p<0.05 vs control

Fig. 3. Plasma high molecular weight adiponectin (HMWA) and resistin levels are increased in sepsis on the first day of hospitalization. Box and whisker plot of day one plasma resistin levels grouped by sepsis category. The vertical box represents the 25th percentile (bottom line), median (middle line), and 75th percentile (top line) values, while the error bars represent the 10th and 90th percentile values. The dots represent values outside the 10th and 90th percentile. *p<0.05 vs control

Fig. 4.

Plasma HMWA and resistin levels are increased in children with an increased pediatric risk of mortality score (PRISM). Box and whisker plot showing levels of plasma HMWA and resistin categorized by severity using the admission PRISM score. The vertical box represents the 25th percentile (bottom line), median (middle line), and 75th percentile (top line) values, while the error bars represent the 10th and 90th percentile values. The dots represent values outside the 10th and 90th percentile. *p < vs PRISM? 10

Table 1

Clinical characteristics and plasma cytokine levels of the study cohort.

Day One Median (IQR)	Control (n=27)	SIRS/Sepsis (n=22)	Septic Shock (n=25)
Age (yrs)	6 (3.4–9.5)	7.3 (4–10.7)	6.1 (2.1–14.3)
BMI (kg/m ²)	17.4 (16.4–20.5)	17.7 (16.1–19.6)	17.3 (16–21.4)
PRISM	NA	6.5 (3–11)	11 (7.8–18.3) ^b
Number of Organ Failures	NA	0 (0–0.5)	2 (1–2) ^c
PICU LOS	NA	4 (2–11)	6 (3–12)
TNF α (pg/ml)	3.7 (2.6–6) (n=24)	6.3 (3.3–22.1) ^a (n=21)	17 (7.8–28) ^{ab} (n=24)
IL-6 (pg/ml)	3.8 (2–11.6) (n=24)	38.1 (19.4–96.8) ^a (n=21)	571 (86.6–3100) ^{ab} (n=24)
IL-8 (pg/ml)	1.2 (1–2.4) (n=24)	10.6 (6.3–33.9) ^a (n=21)	30.7 (11.4–88.7) ^{ab} (n=24)

^a p<0.05 vs control by Kruskal-Wallis ANOVA with Dunn post hoc test.^b p<0.05 vs SIRS/sepsis by *t* test.^c p=0.001 vs SIRS/sepsis by MWRST