Published in final edited form as:

Tetrahedron Lett. 2010 February 10; 51(6): 921–923. doi:10.1016/j.tetlet.2009.12.029.

Halogenated catechols from cycloaddition reactions of η^4 -(2-ethoxyvinylketene)iron(0) complexes with 1-haloalkynes

Jimmy Truong^a, Vioela Caze^a, Ravish K. Akhiani^a, Gayatribahen K. Joshi^a, Lazaros Kakalis^b, Nikita Matsunaga^a, and Wayne F. K. Schnatter^{*,a}

^aLong Island University, Department of Chemistry and Biochemistry, 1 University Plaza, Brooklyn, New York 11201, USA

^bRutgers, The State University of New Jersey, University Heights, Newark, New Jersey, 07102, USA

Abstract

1-chloroalkynes and 1-bromohexyne undergo cycloaddition reactions with ethoxyvinylketeneiron (0) complexes to form chloro and bromocatechols. With most substituents, the halogen is incorporated *ortho* to the phenolic hydroxyl group regioselectively. With chloroethyne, chlorohexyne, and methyl chloropropiolate, the reverse regioselection is observed. *Ab initio* calculations reveal that the products are, in most cases, nearly isoenergetic, which indicates that the intermediate ketene-alkyne adduct geometry must be important in determining the product distribution.

The efficient, regiospecific synthesis of highly substituted aromatic rings represents an important challenge in organic synthesis.¹ One convergent approach is the [4+2] cycloaddition of vinylketenes² with alkynes, however vinylketenes are usually unstable and undergo [2+2] cycloaddition upon reaction with alkynes as well as themselves. The parent compound, vinylketene (buta-1,3-dienone) dimerizes in a hetero [4+2] cycloaddition.³ Masked vinylketenes have been used to carry out this transformation, but often require a deprotection step.⁴ Danheiser used silicon stabilized vinylketenes in Diels-Alder cycloaddition reactions with alkynes.⁵ Electrocyclic ring closures of dienyl ketenes, generated *in situ* from cyclobutenones, have been developed by Moore,^{6a} Danheiser,^{6b} and Liebeskind.^{6c} Danheiser also reported a formal [4+2] reaction of 2-silylvinylketenes with ynolates to form monosilylated resorcinol derivatives.⁷

Transition metals, through their ability to stabilize reactive intermediates, offer powerful alternatives for the synthesis of highly substituted aromatic systems. The reaction of Fischer carbene complexes with alkynes to form hydroquinones has been extensively studied and modified.⁸ A wealth of evidence indicates that this important transformation proceeds through the intermediacy of vinylketene complexes. Chromium promoted variants of the electrocyclization of dienylketenes have also been exploited by Merlic^{9a,b} and Wulff,^{9c} who first isolated an η^4 -vinylketene chromium complex from the reaction of a Fischer carbene complex with an alkyne.¹⁰ Liebeskind reported the synthesis of phenols from alkynes and

*Corresponding author. Tel.: +1 718 488-1453; fax: +1 718 488-1465; Wayne.Schnatter@liu.edu.

Supplementary data: Supplementary data (detailed experimental procedures, spectroscopic data, NMR spectra, and details of *ab initio* calculations associated with this article can be found), in the online version, at doi:

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

cobalt-complexed vinylketenes, which were prepared from cyclobutenones,¹¹ and subsequently discovered a nickel catalyzed variation.¹² Recently, Kondo and Mitsudo reported a rhodium catalyzed reaction.¹³ These methods use vinylketenes generated from cyclobutenones, limiting their utility. The availability of isolable η^4 -vinylketene iron complexes from a variety of sources¹⁴ provides a promising solution to the cycloaddition problem given the low cost of iron.¹⁵ One noteworthy advantage of iron over chromium in benzannulation reactions is the fact that the product arenes are not complexed to the metal as they often are with chromium. Gibson prepared η^4 -complexed vinylketene iron(0) complexes from η^4 -vinylketone-iron(0) complexes,¹⁶ however the major products of reactions of alkynes with these complexes were alkyne insertion products, which formed cyclopentenediones, and, rarely, phenols.¹⁷ Treatment of the insertion products with FeCl₃ resulted in furan-3-(2*H*)-ones and reactions with CO produced cyclopentenediones and butenolides.¹⁸

Our early studies indicated that both electron poor (i.e. ester, ketone, and trifluoromethyl substituted) and electron rich alkynes (i.e. ynol ethers) readily undergo cycloadditions with vinylketene iron complexes to form catechol monoethyl ethers upon reaction with **1a** (Table 1, R¹ = Ph).¹⁹ Tam recently reported ruthenium catalyzed [2+2] cycloadditions^{20a} of norbornenes with alkynyl halides, and rhodium catalyzed intramolecular [4+2] cycloadditions^{20b} of haloalkyne tethered dienes, which inspired us to examine the reactions of terminal haloalkynes with vinylketene iron(0) complexes. Our results are summarized in Table 1.

Initially, we compared the reactions of 1-iodohexyne, 1-bromohexyne, and 1-chlorohexyne with **1a**. Heating **1a** with iodoheptyne for 96 h in DMF at 120 °C provided mostly unreacted iodoheptyne with two putative cycloadducts (*m/z* = 396), which were detected by GC-MS. Bromohexyne was more promising, giving **3a** in 5% yield after refluxing in THF for 24 h. Heating **1a** with bromohexyne in DMF at reflux for 48 h produced the bromocatechol **3a** in 28% yield. A NOESY experiment was used to verify the structure. Reaction of one equivalent of chlorohexyne with **1a** in THF produced **4f** in 72% yield. Cyclotrimers were not a significant contributor to the mass balance of the reactions, although detectable quantities of cyclotrimerization products were found with GC-MS in the Single Ion Monitoring (SIM) mode.

We focused our subsequent efforts in this preliminary study on 1-chloroalkynes. For chloroethyne (entry 2), chlorohexyne (entry 7), and methyl chloropropiolate (entry 10), the chlorine of the major product was incorporated *meta* to the phenolic hydroxyl group. An intermediate was isolated from the reaction of chloroethyne with **1a**, which exhibited proton and carbon NMR spectra consistent with the alkyne insertion product, (Scheme 1, compound **6**, R¹ = Ph, R² = Et, R³ = H). Injection (300 °C) into the GC-MS provided a peak with retention time and mass coincident with **4b**. Thermolysis of a sample at 110 °C for 24 h in THF resulted in complete conversion to **4b**. In the other compounds studied to date the chlorine was found *ortho* to the phenolic hydroxyl. Reaction with chloroethynyl cyclopropane (entry 4) didn't produce any ring opening product, suggesting that a slow radical reaction is unlikely. Although most reactions were carried out in refluxing THF, reaction with *tert*-butylethynyl chloride at 90 °C (entry 6) produced both isomers of the cycloadduct in a 1:4 ratio, indicating that regioselection is temperature dependent. GC-MS analysis of aliquots taken from a reaction of **1a** with chlorohexyne (entry 7) revealed that formation of the product **4f** was detectable after a few minutes at room temperature.

The reaction with methyl chloropropiolate (entry 10) was complicated by dechlorination resulting in a lower yield of halocatechol. GC-MS studies are consistent with the observation that the dechlorination occurs *after* incorporation of the chloroalkyne. Reaction of **1a** with methyl propiolate provided large amounts of the two cyclotrimers, neither of which was not observed in this reaction, suggesting that dechlorination may not be occurring prior to reaction

with the vinylketene complex. GC-MS analysis of aliquots taken from this reaction showed initial formation of the desired product soon after the reaction was started. The dechlorinated catechol was not detected until after 24 hours and then increased at the expense of the desired product. Trace amounts of dechlorination product were detected (GC-MS) with chloroethynyltrimethylsilane, chlorohexyne, and chloroethyne.

Complex **1b** behaved similarly, but it was noted that under reaction conditions similar to those used with the phenyl complex, unreacted **1b** was isolated along with pyrocatechols even after extended reaction times. Again, the regioselectivity appears to be controlled primarily by steric effects. The regioselectivity exhibited in this study is most often precisely the opposite exhibited by the Dötz benzannulation⁸ of chromium carbene complexes: the more sterically demanding substituent is incorporated *meta* rather than *ortho* to the inserted carbonyl group which forms the phenol.

A mechanism is suggested in Scheme 1. Formation of a coordinatively unsaturated iron complex through decomplexation of one of the π bonds of the diene leads to formation of the η^2 alkyne complex.²¹ It seems most likely that the regioselectivity is established when the η^2 -complex forms the insertion product. Analogy with the work of Gibson^{17, 18, 22} suggests that insertion occurs first at the ketene carbonyl rather than at the terminal allylic carbon of the vinylketene. Reductive elimination to form a cyclohexadienone, followed by tautomerization provides the catechol.

In an attempt to understand the regioselectivity of the products, *ab initio* calculations are performed for various alkynes and some of the products. The Löwdin population analysis on the chloroalkyne at the various levels of the theory shows the general trend for the alkyne carbon connected to R² is electron-rich or electron-deficient in accordance with the electronegativity difference between chlorine and the R² group. Therefore, the product distribution in Table 1 cannot be understood by the alkyne electronic structure. The thermodynamic stability of the products alone does not explain the product distribution, either. As shown in Table 1, using the MP2/6-31G(d,p) level of theory including the zero-point energy from the RHF/6-31G(d,p) level, R² = H, *cyclo*-Pr, and Ph substituted model products are nearly isoenergetic. In the case of R² = SiMe₃, **4h** is more stable than **3h** by 3.5 kcal/mol. The largest difference among the calculated products is R² = COOH, and **3i** is 6 kcal/mol more stable than **4i**. The major products of the latter two substitutions are in fact opposite of what thermodynamic stability of the products should be. From these considerations, it is likely that steric factors in the formation of reactive intermediates play a more important role in the formation of the observed products. Hence the ketene-alkyne adduct geometry must be important in determining the product distribution, and we are currently investigating this aspect computationally. We are actively probing the electronic and steric parameters that influence the regioselectivity.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We acknowledge financial support through an Intramural Research Grant to N.M. and W.F.K.S. and a grant to W.F.K.S. from the NIH (SC2GM082276). The NSF provided funds for the purchase of a GC-MS (CHE-0840432). We also thank the Department of Chemistry and Biochemistry of Rutgers University, Newark for access to instrumentation. We thank Mr. William A. Helwig, Mr. Ruhul Q. Chowdhury, Ms. Kerry McKenzie, Mr. Dean Aquino, and Ms. Jeunesse Lewis (Harlem Childrens Society intern) for assistance.

References

1. An entire issue of *Tetrahedron* **2008**, *64*, 757–968 focuses on the use of benzannulation and cycloaddition reactions to form aromatic rings.
2. For a comprehensive review of ketene chemistry see: Tidwell, TT. Ketenes. Wiley; New York: 1995.
3. Trahanovsky WS, Surber BW, Wilkes MC, Preckel MM. *J Am Chem Soc* 1982;104:6779–6781. and references therein.
4. Corey EJ, Kozikowski AP. *Tetrahedron Lett* 1975;16:2389–2392.
5. (a) Loebach JL, Bennett DM, Danheiser RL. *J Org Chem* 1998;63:8380–8389. (b) Danheiser RL, Sard H. *J Org Chem* 1980;45:4810–4812.
6. (a) Moore HW, Decker OHW. *Chem Rev* 1986;86:821–830. (b) Sun L, Danheiser RL, Brisbois RG, Kowalczyk JJ, Miller RF. *J Am Chem Soc* 1990;112:3093–3100. (c) Liebeskind LS. *J Org Chem* 1995;60:8194–8203.
7. Austin WF, Zhang Y, Danheiser RL. *Org Lett* 2005;7:3905–3908. [PubMed: 16119928]
8. For a comprehensive review of the Dötz benzannulation see: Waters ML, Wulff WD. *Org React* 2008;70:121–623., and for a general updated review see Dötz KH, Stendal J Jr. *Chem Rev* 2009;109:3227–3274. [PubMed: 19642645]
9. (a) Merlic CA, Aldrich CC, Albaneze-Walker J, Saghatelian A, Mammen J. *J Org Chem* 2001;66:1297–1309. [PubMed: 11312960] (b) Merlic C, Xu D, Gladstone BG. *J Org Chem* 1993;58:538–545. (c) Rawat M, Wulff WD. *Org Lett* 2004;6:329–332. [PubMed: 14748585]
10. Anderson BA, Wulff WD, Rheingold AL. *J Am Chem Soc* 1990;112:8615–8617.
11. (a) Huffman MA, Liebeskind LS, Pennington WT. *Organometallics* 1992;11:255–266. (b) Huffman MA, Liebeskind LS. *J Am Chem Soc* 1990;112:8617–8618.
12. Huffman MA, Liebeskind LS. *J Am Chem Soc* 1991;113:2771–2772.
13. Kondo T, Niimi M, Nomura M, Wada K, Mitsudo T. *Tetrahedron Lett* 2007;48:2837–2839.
14. For a comprehensive review of vinylketene complexes see: Gibson SE, Peplow MA. *Adv Organomet Chem* 1999;44:275–353..
15. For recent reviews of iron chemistry in organic synthesis see: Sherry BD, Fürstner A. *Acc Chem Res* 2008;41:1500–1511. [PubMed: 18588321] and Bolm C, Legros J, Le Pailh J, Zani L. *Chem Rev* 2004;104:6217–6254. [PubMed: 15584700] .
16. Alcock NW, Richards CJ, Thomas SE. *Organometallics* 1991;10:231–238.
17. Morris KG, Saberi SP, Thomas SE. *J Chem Soc; Chem Commun* 1993:209–211.
18. Saberi SP, Salter MM, Slawin AMZ, Thomas SE, Williams DJ. *J Chem Soc; Perkin Trans* 1994;1:167–171.
19. Akhani RK, Rehman A, Schnatter WFK. *Tetrahedron Lett* 2009;50:930–932. [PubMed: 20046214] and Darbasie ND, Schnatter WFK, Warner KF, Manolache N. *Tetrahedron Lett* 2006;47:963–966..
20. (a) Villeneuve K, Ridell N, Jordan RW, Tsui GC, Tam W. *Org Lett* 2004;6:4543–4546. [PubMed: 15548071] and Allan A, Villeneuve K, Cockburn N, Fatila E, Riddell N, Tam W. *Eur J Org Chem* 2008:4178–4192.. (b) Yoo WJ, Allen A, Villeneuve K, Tam W. *Org Lett* 2005;7:5853–5856. [PubMed: 16354083]
21. Diphenylketene-alkyne Ir(I) complexes have been shown to undergo cyclization to form iridabenzopyrans: Lo HC, Grotjahn DB. *J Am Chem Soc* 1997;119:2958–2959.
22. Benyunes SA, Gibson (née Thomas) SE, Peplow MA. *Tetrahedron: Asymmetry* 1997;8 and references therein.

Scheme 1.

Proposed mechanism of formation of chlorocatechols.

Table 1

Cyclization of Vinylketeneiron(0) Complexes with 1-haloalkynes^a

Entry	Complex	R^2	X	Product (%)	ΔE^b (kcal/mol)
1	1a	Bu	Br	3a (28) ^c	
2	1a	H	Cl	4b (33) ^d	3b 1.1
3	1a	Cl	Cl	3c (45) ^d	
4	1a	cyc-Pr	Cl	3d (62)	4d 0.2
5	1a	<i>t</i> -Bu	Cl	3e (55)	
6	1a	<i>t</i> -Bu	Cl	3e (44)	
				4e (11) ^e

Entry	Complex	R ²	X	Product (%)	ΔE ^b (kcal/mol)
7	1a	Bu	Cl	4f (72)	
8	1a	Ph	Cl	3g (62)	4g-0.8
9	1a	SiMe ₃	Cl	3h (42)	4h-3.5
10	1a	COOMe	Cl	3i (14)	3i 5.6
				4i (23) ^f	
11	1b	Bu	Cl	4j (49)	
12	1b	Ph	Cl	3k (43)	

^a Unless otherwise stated all reactions were carried out with 0.3-1.5 mmol of vinylketene complex with 1.0 equivalent of haloalkyne in 3-15 mL THF heated at reflux for 24-48 h.

^b Positive (negative) value indicates **3** (**4**) is more stable.

^cDMF, reflux 48 h.

^dVolatile alkynes were used in excess as solutions that were heated at 70° C in pressure vessels equipped with Young valves.

^eReaction was carried out at 90°C.

^fDechlorinated product, **4i** with X = H, was also obtained in 39 % yield.