Published in final edited form as:

Biochem Biophys Res Commun. 2010 January 1; 391(1): 426–430. doi:10.1016/j.bbrc.2009.11.074.

Toll-Like Receptor 2 Is Required for Opioids-Induced Neuronal Apoptosis

Yi Li^{1,2}, Hui Li^{3,1}, Yi Zhang¹, Xiuli Sun¹, Gregory A. Hanley⁴, Ying Zhang⁵, Shenggang Sun³, Ying Peng², and Deling Yin^{1,*}

¹Department of Internal Medicine, College of Medicine, East Tennessee State University, Johnson City, TN 37614

²Department of Neurology, the Second Affiliated Hospital, Sun Yat-sen University, Guangzhou 510120, China

³Department of Neurology, Uion Hospital, Huazhong University of Science and Technology Wuhan, 430022, China

⁴Department of Laboratory Animal Resources, College of Medicine, East Tennessee State University, Johnson City, TN 37614

⁵Department of Molecular Microbiology and Immunology, Johns Hopkins University, Baltimore, MD 21205

Abstract

Toll-like receptor 2 (TLR2), a key immune receptor in the TLR family, is widely expressed in various systems, including the immune and nervous systems and plays a critical role in controlling innate and adaptive immune responses. We previously reported that opioids inhibit cell growth and trigger apoptosis. However, the underlying mechanism by which TLR2 mediates apoptosis in response to opioids is not yet known. Here we show that chronic morphine treatment in primary neurons dramatically increased the expression of TLR2 at both the messenger RNA and protein levels. In addition, TLR2 deficiency significantly inhibited chronic morphine-induced apoptosis in primary neurons. Activation of caspase-3 after morphine treatment is impaired in TLR2 deficient primary neurons. Moreover, morphine treatment failed to induce an increased level of phosphorylated glycogen synthase kinase 3 beta (GSK3 β) in TLR2 deficient primary neurons, suggesting an involvement of GSK3 β in morphine-mediated TLR2 signaling. These results thus demonstrate that opioids prime neurons to undergo apoptosis by inducing TLR2 expression. Our data suggest that inhibition of TLR2 is capable of preventing opioids-induced damage to neurons.

Keywords

TLR2; morphine; apoptosis; neurons

© 2009 Elsevier Inc. All rights reserved.

*Corresponding author: Deling Yin. Department of Internal Medicine, College of Medicine, East Tennessee State University, Johnson City, TN 37614. Tel: 423 439 8826. Fax: 423 439 6387. yin@etsu.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Opioids have been widely applied in clinics as one of the most potent pain relievers for centuries. However, chronic use of opioids as clinical therapies produces benefits as well as severe side effects. Opioids, including morphine, cause apoptosis in various systems. We have previously reported that morphine promotes apoptosis both *in vitro* and *in vivo* [1–3]. In central nervous system (CNS), opioids induced neuronal apoptosis [4]. Although opioid receptors play critical roles in the processes of opioids-induced effects, the antagonists of opioid receptors can only partially block the effects of opioids [5]. Thus, the specific cellular and molecular mechanisms underlying on opioids-induced apoptosis still need to be defined.

Toll-like receptors (TLRs) are well known as recognition of pathogens in the innate immune system aimed as defending the survival of the host. Thirteen TLRs have been identified [6]. TLRs and their functions have been established in immune cells. However, the functional role of TLRs in the CNS remains unclear. Growing evidence demonstrated that neurons express some TLRs, including TLR2, TLR4, and TLR9 [7]. Neuronal TLRs play pivotal roles in brain injuries and functional deficits [7,8]. TLR2 was identified as a key immune receptor in TLRs family with a large repertoire of ligands. Many classes of microorganisms, as well as the bacterial cell wall components peptidoglycan and lipoteichoic, have been found to activate TLR2. Activation of TLR2 signaling triggers activation of proapoptotic signals, and causes cell death in various systems [7,8]. Caspase activities increased significantly in TLR2 signaling activated cells [9,10].

Recent evidence suggests that there is cross-talk between TLR signaling and glycogen synthase kinase 3 (GSK3), a crucial regulator of many cellular functions including cell survival and apoptosis [9,11]. GSK3 is a serine/threonine kinase that refers to two isoforms- GSK3 α and GSK3 β [9,11]. It's considered that GSK3 promotes the mitochondrial intrinsic apoptotic signaling cascade induced by a diverse array of insults [9,11,12]. On mechanisms tightly regulating the activities of two isoforms of GSK3, the most well-defined mechanism is the phosphatidylinositol 3-kinase (PI3K)/Akt signaling pathway. GSK3 activity is inhibited through PI3K/Akt signaling by phosphorylation of serine-9 in GSK3 β or serine-21 in GSK3 α [9,12].

In present study we found that TLR2 is required for morphine-induced neuronal cell death and apoptosis. Furthermore, Morphine failed to induce an increased level of phosphorylated GSK3 β in TLR2 deficient primary neurons, suggesting an involvement of GSK3 β in morphine-mediated TLR2 signaling.

Materials and Methods

Reagents

Morphine sulfate was obtained from Sigma-Aldrich. Cell culture medium, horse serum, B27 supplement and reagents for neuron cell culture were purchased from Invitrogen Corporation. The Quantative PCR kit was purchased from Invitrogen Corporation. The polyclonal anti-cleaved caspase-3, caspase-3, p-Ser9-GSK3 β , total-GSK3 β , p-Akt, and GAPDH antibodies were purchased from Cell Signaling Technology. The monoclonal TLR2 antibody was obtained from Santa Cruz Biotechnology, Inc.

Animals

Toll-like receptor 2 knockout (TLR2 KO) mice on a C57BL/6 background and wild type C57BL/6 (WT) mice were obtained from the Jackson Laboratory and were maintained in the Division of Laboratory Animal Resources at East Tennessee State University (ETSU), a facility accredited by the Association for the Assessment and Accreditation of Laboratory Animal Care

International (AAALAC). All aspects of the animal care and experimental protocols were approved by the ETSU Committee on Animal Care. Pregnancy was confirmed by the presence of vaginal plug and this was considered as gestational day 0 (E0).

Primary cortical neuron culture

The method used for preparing primary cortical neuron cultures followed the procedure described in our previous publication with a slight modification [13]. Briefly, pregnant mice on E16 were anesthetized with carbon dioxide and killed by cervical dislocation. Meninges-free cortices were dissected from the 16-day-old fetuses and triturated into homogeneous solution. The cells were plated with a density of 1×10^6 /well on 6-well-plates precoated with 100ug/ml poly-D-lysine. The cells were cultured with neurobasal medium supplemented with 5% horse serum, 2% B27, 1% glutamax and 50u/ml penicillin/streptomycin, and were kept in a tissue culture incubator at 37°C in 5% CO₂. Seven-day-old cortical neurons were subjected to morphine treatment with different concentrations within 6 days.

Determination of cell death

Neuronal cell density was observed directly with a microscope during drug treatment. At the end point of each group, neurons in some wells were stained with propidium iodide (PI) solution as described previously [2] and were observed with a fluorescent microscope (Motic Company). To determine the change of cell death in wells, five areas in each well were randomly chosen for cell counting. Data were normalized from control-treated cells and shown as a percentage of the controls.

Detection of apoptosis by TUNEL assay

Terminal deoxynucleotidyl transferase biotin-dUTP nick end labeling (TUNEL) assay was detected as described previously [3]. Neurons treated with or without 15μM morphine for 6 days underwent DNA fragmentation detection using an in situ apoptosis detection kit (Roche) according to the manufacturer's instructions. Cells were observed using a fluorescent microscope (Olympus).

Real-time quantitative reverse transcription polymerase chain reaction (RT-PCR)

The real-time PCR detection technique was performed as described in our previous publication [14]. Briefly, first-strand cDNA was synthesized from 1μg of total RNA using a Reaction Ready™ first strand cDNA synthesis kit (SuperArray Bioscience Corporation). After incubation at 70 °C for 3 min and cooling down to 37 °C for 10 min, RT cocktail was added to the annealing mixture and further incubated at 37 °C for 60 min. 2μl of 1:2 diluted cDNA was subjected to real-time quantitative PCR using Bio-Rad iCycler iQ Multicolor Real-Time PCR Detection System (Bio-Rad). PCR was performed in a 25μl volume using SYBR GreenER qPCR Super Mix for iCycler (Invitrogen). All primers were purchased from SuperArray Bioscience Corporation. All PCR assays were performed in triplicate. The reaction conditions were: 50°C for 2 min, 95°C for 8 min 30s, followed by 40 cycles of 95°C for 15s, 60°C for 60s, and 72°C for 30s. Glyceraldehyde-3-phosphate dehydrogenase (GAPDH) was amplified from all samples on each plate as housekeeping genes. The reaction mixtures without template cDNA were used as negative controls.

Western blot analysis

Western blotting was performed as described in our previous publication using several antibodies [9,12]. Briefly, experimental cells were harvested and lysed in RIPA Lysis Buffer. The lysates were separated by 10% SDS-PAGE then transferred to a nitrocellulose membrane (Bio-Rad). The membrane was then incubated at room temperature in blocking solution for 1 h, followed with the blocking solution containing first antibody overnight at 4°C. Caspase-3,

p-Ser9-GSK3 β , total-GSK3 β , p-Akt, or GAPDH was detected with antibody specific to p-Ser9-GSK3 β , total-GSK3 β , p-Akt, or GAPDH. After washing three times with TBS for 5min, the blot was incubated with a second antibody. The blot was again washed three times with TBS before being exposed to the SuperSignal West Dura Extended Duration substrate (Pierce).

Immunohistochemistry

Primary neuron cells were fixed and immunostained as described in our previous publication, with slightly modification [13]. Briefly, for TLR2 immunostaining, fixed neurons were immunostained using anti-TLR2 Ab and Alexa Fluor 488 (Invitrogen). For cleaved Caspase-3 and NeuN double immunostaining, fixed neurons were immunostained using anti-NeuN and anti-cleaved caspase-3 antibodies, followed by Alexa Fluor 488 or Alexa Fluor 594 conjugated secondary antibodies. For p-Ser9-GSK3 β , fixed neurons were immunostained using anti-p-Ser9-GSK3 β Ab and Alexa Fluor 594 (Invitrogen). Confocal images were captured using a Leica TCS SP2 Laser Scanning Confocal Microscope. Fluorescent images were captured using a fluorescent microscope (Olympus).

Statistical analysis

All data were represented as means $\pm$ SEM. The data were analyzed using one-way analysis of variance (ANOVA) followed by Bonferroni tests to determine where differences among groups existed. Differences were considered statistically significant for values of $p < 0.05$.

Results

Increased TLR2 expression in wild type neurons following chronic morphine treatment

Although it is well established that several opioid receptor isoforms are expressed on neurons, the mechanisms by which opioids influence neuronal function remains to be elucidated [15]. To investigate the mechanisms, we treated primary mouse cortical neurons from wild type mice with morphine sulfate at 15 μ M for different periods of time and examined the expression of TLR2 by quantitative real time RT-PCR and immunohistochemistry. TLR2 expression in mRNA level was significantly increased following 15 μ M morphine treatment for 4 days (Fig. 1A). Morphine-induced TLR2 expression was also detected at the protein level (Fig. 1B). Our results demonstrated that chronic morphine treatment significantly increases TLR2 expression.

A deficiency of TLR2 is resistant to morphine-induced neuronal apoptosis

It has been reported that cortical neurons generated a significant reduction in cell density following 15 μ M opioid treatment for 7 days [4]. To investigate whether TLR2 plays a role in opioids-induced neuronal cell death, we treated primary mouse cortical neurons from TLR2 knockout mice and wild type mice with morphine at different concentrations for 6 days and assessed cell death using PI staining. We found that chronic morphine treatment induced cell death in a dose-dependent manner in wild type neurons, but not in TLR2 deficient neurons (Fig. 2A). Dead cell rate showed no significant differences among 5 μ M, 15 μ M morphine treated TLR2 deficient neurons and untreated neurons. We next evaluated the effect of TLR2 on morphine-induced primary neuronal apoptosis using a TUNEL technique. A significant number of cells in the wild type neurons after morphine treatment were undergoing apoptosis, whereas only a few apoptotic cells were detected in the TLR2 deficient neurons following morphine treatment (Fig. 2B). Therefore, primary neurons with a deficiency of TLR2 are resistant to morphine-induced neuronal apoptosis.

Morphine causes caspase-3 activation in wild type neurons but not in TLR2 deficient neurons

The cleaved caspase-3 acts as an active and lethal protease at the most distal stage of the apoptotic pathways, and is the major caspase involved in neuronal apoptosis [16]. Thus, we

examined caspase-3 activation in TLR2 deficient and wild type primary neurons after morphine treatment. Morphine treatment at 15 μ M for 6 days activated caspase-3 in wild type neurons to a greater increase compared to TLR2 deficient neurons (Fig. 3A). These results were verified by double-immunofluorescent staining of active caspase-3 and neurons (Fig. 3B). Morphine treatment at 15 μ M for 6 days resulted in a dramatic increase of cleaved caspase-3 immunoreactive colocalization in wild type primary neurons, but not in TLR2 deficient neurons. Our results suggested a proapoptotic role for TLR2 in opioids-mediated neuronal apoptosis.

TLR2 deficiency in neurons blocks morphine-enhanced phosphorylation of GSK3 β

We have previously reported that TLRs activate GSK3 β signaling pathways [12]. To substantiate that opioids activate TLR2 signaling, we investigated the regulation of GSK3 β in morphine-activated TLR2 signaling. As shown in Fig. 4A, GSK3 β was phosphorylated on the inactivating residue serine-9 and phosphorylation significantly increased after 15 μ M morphine treatment for 6 days in wild type neurons. A deficiency of TLR2 in primary neurons strongly suppressed chronic morphine-induced GSK3 β serine-9 phosphorylation (Fig. 4A). The changes of GSK3 β serine-9 phosphorylation were verified by immunohistochemistry. Morphine treatment resulted in a dramatic increase of GSK3 β serine-9 phosphorylation on protein level in wild type primary cortical neurons, but not in TLR2 deficient primary cortical neurons (Fig. 4B). Since phosphorylation of serine-9 is inhibitory for GSK3 β activity, these results suggested that morphine decreases GSK3 β activity through a TLR2-dependent mechanism.

Akt is an important physiologic mediator of the PI3K pathway [2,17]. Activated Akt phosphorylates several downstream targets of the PI3K pathway including GSK3 [2,17]. Therefore, we determined if morphine induces Akt phosphorylation through a TLR2-dependent pathway. We found no alterations in the protein level of phosphorylated Akt in wild type neurons and TLR2 deficient neurons following morphine treatment (Fig. 4C), suggesting no role of Akt in morphine suppression of TLR2-mediated GSK3 β activity.

Discussion

The data presented herein demonstrated for the first time, to our knowledge, a key role for neuronal TLR2 in the induction of morphine-mediated apoptosis. Our studies show that chronic morphine treatment regulates TLR2 signaling and results in neuronal apoptosis. It is widely accepted that TLR2 signaling mainly activates inflammatory responses, including proinflammatory cytokines [18]. Most studies evaluated the effects of TLR2 on neuronal cell death and apoptosis through immune cell mediated mechanisms [19]. We found in this study that deficiency of TLR2 activity in primary neurons blocked morphine-induced neuronal apoptosis. We identified here that TLR2 signaling is activated by opioids in neurons without the presence of immune cells. This provides a possible target for controlling of morphine caused neurotoxicity.

We have previously reported that TLRs activate phosphorylation of GSK3 β [12]. GSK3 activities cause diverse regulation of TLR-mediated production of pro- and anti-inflammatory cytokines [17,18]. Selective agonists of TLRs, mediate phosphorylation of GSK3 β on Ser9, and result in GSK3 β inactivation. In agreement with these results, our data showed that morphine decreased GSK3 β activity through a TLR2-dependent mechanism. Akt activates several downstream targets, such as GSK3 β . Interestingly, we found no role of Akt in morphine suppression of TLR2-mediated GSK3 β activity. It remains to be determined whether other kinases phosphorylate GSK3 β . Furthermore, scaffolding proteins provide one possible mechanism on interactions between these kinases. Our previous results also reveal that a scaffolding protein β -arrestin 2 is involved in TLR2-mediated apoptosis [10].

In our study, we found that chronic morphine decreases GSK3 β activity. These evidences lead us to propose that GSK3 β activity protects from opioids-induced neuronal apoptosis. We reported previously that morphine modulates apoptosis through cell death receptor Fas, the other major member of the death receptor family [1]. Apoptosis signaling induced by Fas was found to be regulated by the anti-apoptotic effect of GSK3 [20]. Further investigations should be performed to understand the mechanisms on GSK3 activities associated with opioids-mediated apoptosis. We showed that morphine induces neuronal cell death and apoptosis in a dose dependent manner. These data are consistent with the findings of Svensson and Hu [4, 5]. Furthermore, we found that in the deficiency of TLR2, morphine induced effects on neuronal cell death and caspase-3 activity, were counteracted. Thus, a caspase-3-dependent pathway is involved in TLR2 signaling activated by chronic morphine effects. Evidences have already shown that activation of caspase-3 is a terminal event leading to neuronal apoptosis upon chronic exposure to morphine, and there may be changes in intracellular proapoptotic elements such as Bax, Fas, and antiapoptotic elements such as Bcl-2 in response to chronic morphine effects [5]. The cleaved caspase-3 acts as an active and lethal protease at the most distal stage of the apoptotic pathways, and is the major caspase involved in neuronal apoptosis [16]. The extrinsic pathway, one of the two identified major caspase-3-activating pathways, involves cell death receptors [21]. In our study, TLR2 deficiency in neurons reversed the caspase-3 activity increasing effect induced by chronic morphine treatment. We identified that downstream of TLR2, caspase-3 as a crucial factor for morphine-induced neuronal apoptosis.

In summary, our studies reveal that TLR2 plays a critical role in opioids-induced neuronal apoptosis and an involvement of GSK3 β . Further understanding the mechanisms mediated by TLRs may provide insights into potential therapeutic interventions opioid-related neurotoxicity.

Acknowledgments

This work was supported by National Institutes of Health (NIH) grant DA020120 and East Tennessee State University Research Development Committee (ETSU RDC) 0048 to D. Yin. This work was also supported in part by ETSU RDC grant 07-026M to G. Hanley.

References

1. Yin D, Mufson RA, Wang R, Shi Y. Fas-mediated cell death promoted by opioids. *Nature* 1999;397:218. [PubMed: 9930695]
2. Yin D, Woodruff M, Zhang Y, Whaley S, Miao J, Ferslew K, Zhao J, Stuart C. Morphine promotes Jurkat cell apoptosis through pro-apoptotic FADD/P53 and anti-apoptotic PI3K/Akt/NF-kappaB pathways. *J.Neuroimmunol* 2006;174:101–107. [PubMed: 16529824]
3. Yin D, Tuthill D, Mufson RA, Shi Y. Chronic restraint stress promotes lymphocyte apoptosis by modulating CD95 expression. *J.Exp.Med* 2000;191:1423–1428. [PubMed: 10770807]
4. Svensson AL, Bucht N, Hallberg M, Nyberg F. Reversal of opiate-induced apoptosis by human recombinant growth hormone in murine foetus primary hippocampal neuronal cell cultures. *Proc.Natl.Acad.Sci.U.S.A* 2008;105:7304–7308. [PubMed: 18474854]
5. Hu S, Sheng WS, Lokensgard JR, Peterson PK. Morphine induces apoptosis of human microglia and neurons. *Neuropharmacology* 2002;42:829–836. [PubMed: 12015209]
6. Doyle SL, O'Neill LA. Toll-like receptors: from the discovery of NFkappaB to new insights into transcriptional regulations in innate immunity. *Biochem.Pharmacol* 2006;72:1102–1113. [PubMed: 16930560]
7. Tang SC, Arumugam TV, Xu X, Cheng A, Mughal MR, Jo DG, Lathia JD, Siler DA, Chigurupati S, Ouyang X, Magnus T, Camandola S, Mattson MP. Mattson. Pivotal role for neuronal Toll-like receptors in ischemic brain injury and functional deficits. *Proc.Natl.Acad.Sci.U.S.A* 2007;104:13798–13803. [PubMed: 17693552]

8. Sabroe I, Dower SK, Whyte MK. The role of Toll-like receptors in the regulation of neutrophil migration, activation, and apoptosis. *Clin.Infect.Dis* 2005;41:S421–S426. [PubMed: 16237641]
9. Chen L, Zhang Y, Sun X, Li H, LeSage G, Javer A, Zhang X, Wei X, Jiang Y, Yin D. Synthetic resveratrol aliphatic acid inhibits TLR2-mediated apoptosis and an involvement of Akt/GSK3beta pathway. *Bioorg.Med.Chem* 2009;17:4378–4382. [PubMed: 19477653]
10. Li Y, Sun X, Zhang Y, Huang J, Hanley G, Ferslew KE, Peng Y, Yin D. Morphine promotes apoptosis via TLR2, and this is negatively regulated by beta-arrestin 2. *Biochem.Biophys.Res.Commun* 2009;378:857–861. [PubMed: 19071087]
11. Guha M, Mackman N. The phosphatidylinositol 3-kinase-Akt pathway limits lipopolysaccharide activation of signaling pathways and expression of inflammatory mediators in human monocytic cells. *J.Biol.Chem* 2002;277:32124–32132. [PubMed: 12052830]
12. Zhang Y, Zhang Y, Miao J, Hanley G, Stuart C, Sun X, Chen T, Yin D. Chronic restraint stress promotes immune suppression through toll-like receptor 4-mediated phosphoinositide 3-kinase signaling. *J.Neuroimmunol* 2008;204:13–19. [PubMed: 18814920]
13. Lv X, Su L, Yin D, Sun C, Zhao J, Zhang S, Miao J. Knockdown of integrin beta4 in primary cultured mouse neurons blocks survival and induces apoptosis by elevating NADPH oxidase activity and reactive oxygen species level. *Int.J.Biochem.Cell Biol* 2008;40:689–699. [PubMed: 18006359]
14. Yin D, Zhang Y, Stuart C, Miao J, Zhang Y, Li C, Zeng X, Hanley G, Moorman J, Yao Z, Woodruff M. Chronic restraint stress modulates expression of genes in murine spleen. *J.Neuroimmunol* 2006;177:11–17. [PubMed: 16814870]
15. Stucky CL, Gold MS, Zhang X. Mechanisms of pain. *Proc.Natl.Acad.Sci.U.S.A* 2001;98:11845–11846. [PubMed: 11562504]
16. Hua F, Ma J, Ha T, Xia Y, Kelley J, Williams DL, Kao RL, Browder IW, Schweitzer JB, Kalbfleisch JH, Li C. Activation of Toll-like receptor 4 signaling contributes to hippocampal neuronal death following global cerebral ischemia/reperfusion. *J.Neuroimmunol* 2007;190:101–111. [PubMed: 17884182]
17. Fukao T, Koyasu S. PI3K and negative regulation of TLR signaling. *Trends Immunol* 2003;24:358–363. [PubMed: 12860525]
18. Akira S, Takeda K. Toll-like receptor signalling. *Nat.Rev.Immunol* 2004;4:499–511. [PubMed: 15229469]
19. Carpentier PA, Duncan DS, Miller SD. Glial toll-like receptor signaling in central nervous system infection and autoimmunity. *Brain Behav.Immun* 2008;22:140–147. [PubMed: 17920811]
20. Song L, Zhou T, Jope RS. Lithium facilitates apoptotic signaling induced by activation of the Fas death domain-containing receptor. *BMC.Neurosci* 2004;5:20. [PubMed: 15157283]
21. Yakovlev AG, Ota K, Wang G, Movsesyan V, Bao WL, Yoshihara K, Faden AI. Differential expression of apoptotic protease-activating factor-1 and caspase-3 genes and susceptibility to apoptosis during brain development and after traumatic brain injury. *J.Neurosci* 2001;21:7439–7446. [PubMed: 11567033]

Figure 1.

Morphine induces the expression of TLR2 in wild type neurons. **A**, Quantitative real time RT-PCR analysis of TLR2 expression showed a significant increase in wild type primary cortical neurons treated with morphine for 4 days. Neurons were treated with 15 μM morphine for 16 h, 2 d and 4 d, respectively. * $p < 0.05$ vs. untreated neurons. Results represent mean ± s.e.m. of three independent experiments. **B**, Immunostaining in wild type neurons showed a dramatic increase of TLR2-immunoreactive neurons after 15 μM morphine treatment for 6 days. Scale bars, 75 μm.

Figure 2.

A deficiency of TLR2 is resistant to morphine-induced neuronal apoptosis. **A**, Bar graph showed the percentage of neuronal cell death in wild type (WT) and TLR2 knockout (TLR2 KO) neurons identified with PI staining. WT and TLR2 KO neurons were treated with 5 μM, 15 μM, or 50 μM morphine for 6 days, respectively. Results represent mean ± s.e.m. of three independent experiments. * $p < 0.05$ vs. untreated WT neurons. ** $p < 0.05$ vs. WT neurons treated with 15 μM morphine. *** $p < 0.05$ vs. WT neurons treated with 50 μM morphine.

B, Representative light microscopic images showed TUNEL-positive neurons. WT and TLR2 KO neurons were treated with or without 15 μM morphine for 6 days. The dark brown color showed TUNEL-positive neuronal apoptosis (Red arrow head). Magnification 200 ×.

Figure 3.

Morphine causes caspase-3 activation in wild type neurons but not in TLR2 knockout neurons. **A**, Total and active cleaved caspase-3 in protein levels were determined by western blot. WT and TLR2 KO neurons were treated with morphine at 5 μM or 15 μM for 6 days. Representative results of the level of cleaved caspase-3 and total caspase-3 are shown at the left of each pane. Data are representative of three independent experiments. * $p < 0.05$ vs. untreated WT neurons. ** $p < 0.05$ vs. WT neurons treated with 15 μM morphine. **B**, WT and TLR2 KO neurons were treated with or without 15 μM morphine for 6 days and immunostained with antibodies to cleaved caspase-3 (Red) and NeuN (neuronal marker) (Green). Merged images displayed co-

localizations of cleaved caspase-3 and NeuN (Yellow) (White arrow head). Scale bars, 18.75 μm . Data are representative of three separate experiments.

Figure 4.

TLR2 deficiency in neurons blocks morphine-enhanced GSK3 β serine-9 phosphorylation. **A**, Phosphorylated serine-9 GSK3 β (p-Ser9-GSK3 β) and total GSK3 β in protein levels were examined by western blot. WT and TLR2 KO neurons were treated with 5 μ M or 15 μ M morphine for 6 days, respectively. Data are representative of three independent experiments. **B**, WT and TLR2 KO neurons were treated with or without 15 μ M morphine for 6 days and immunostained with p-Ser9-GSK3 β antibody. **C**, Phosphorylated Akt (p-Akt) was determined by western blot. We treated WT and TLR2 KO primary neurons with or without 15 μ M morphine for 6 days and examined for the level of phosphorylated Akt (p-Akt). Data are representative of three independent experiments.