Published in final edited form as:

J Phys Condens Matter. 2007 July 18; (28): 285223. doi:10.1088/0953-8984/19/28/285223.

Folding and unfolding of a non-fluorescent mutant of green fluorescent protein

Beata Wielgus Kutrowska¹, Marta Narczyk¹, Anna Buszko¹, Agnieszka Bzowska¹, and Patricia L. Clark²

Beata Wielgus Kutrowska: beata@biogeo.uw.edu.pl; Marta Narczyk: ; Anna Buszko: ; Agnieszka Bzowska: ; Patricia L. Clark: pclark1@nd.edu

¹ Department of Biophysics, Institute of Experimental Physics, University of Warsaw, Zwirki & Wigury 93, 02-089 Poland ² Department of Chemistry & Biochemistry, University of Notre Dame, 251 Nieuwland Science Hall, Notre Dame, IN 46556-5670 USA

Abstract

Green fluorescent protein (GFP), from the Pacific jellyfish *A. victoria*, has numerous uses in biotechnology and cell and molecular biology as a protein marker because of its specific chromophore, which is spontaneously created after proper protein folding. After formation, the chromophore is very stable and remains intact during protein unfolding, meaning that the GFP unfolding process is not the reverse of the original folding reaction; i.e., the principles of microscopic reversibility do not apply. We have generated the mutant S65T/G67A-GFP, which is unable to form the cyclic chromophore, with the goal of investigating the folding, unfolding and competing aggregation of GFP under fully reversible conditions. Our studies have been performed in the presence of GdnHCl. The GFP conformation was monitored using intrinsic tryptophan fluorescence, and fluorescence of bis-ANS. Light scattering was used to follow GFP aggregation. We conclude from these fluorescence measurements, that S65T/G67A-GFP folding is largely reversible. During equilibrium folding, the first step is formation of molten globule, prone to aggregation.

1. Introduction

There is still very little known concerning the general rules that govern β -sheet formation. In contrast to α -helical proteins, the formation of β -sheet structure requires creation of interaction between amino acids that are far away in the sequence of polypeptide chain [1]. Alternatively, this more complex topological search problem can also lead to misfolded β -sheet structure, such as the amyloid structures connected to the neurodegenerative diseases like Alzheimer's, Huntington's or prion diseases [2]. On average, proteins with mainly β -sheet structure are more prone to aggregation than α -helical proteins [3]. Yet β -sheet proteins have persisted through evolution. In light of this, it is important to understand the forces that drive the formation of native β -sheet structure, and likewise, the forces that govern the formation of misfolded, potentially toxic, β sheet aggregates.

One representative all- β -sheet protein is green fluorescent protein (GFP) from the Pacific jellyfish *A. victoria*. GFP is well known for its biotechnological relevance as a protein marker [4] because of its intrinsic chromophore, which is spontaneously created after proper protein folding. Chromophore biosynthesis involves residues Ser65, Thr66, and Gly67 in wild-type GFP [5]. For efficient chromophore formation, close proximity of the backbone atoms of amino acids 65 and 67 is required, and achieved via the conserved glycine residue at position 67 (any other amino acid imposes too much steric hindrance) [6]. After formation, the chromophore is very stable and remains intact during protein denaturation [7], meaning that the GFP unfolding

process is not the reverse of the original folding reaction; i.e., the principles of microscopic reversibility do not apply. GFP chromophore fluorescence is quenched upon protein denaturation, presumably by the interaction of chromophore with molecular oxygen [8], or by exposing the chromophore to the aqueous solvent.

The all β structure of GFP suggests it should be an excellent model to investigate the formation of β -sheet structure, but the presence of chromophore may alter the unfolding and refolding properties. Because of this, we have generated the mutant S65T/G67A-GFP, which is unable to form the cyclic chromophore, with the goal of investigating the folding, unfolding, and competing aggregation of GFP under fully reversible conditions. Our studies have been performed in the presence of guanidinium hydrochloride (GdnHCl). The GFP conformation was monitored using intrinsic tryptophan fluorescence, 1,1'-bis(4-anilino-5-naphthalenesulphonic acid) (bisANS) fluorescence, and light scattering.

2. Materials and methods

2.1 Construction of the mutant S65T/G67A-GFP and protein preparation

A plasmid encoding S65T-GFP with polyhistidine tag was a gift from Roger Tsien. The mutant S65T/G67A-GFP was prepared using Stratagene QuikChange Mutagenesis Kit according to the manufacturer's direction. S65T/G67A-GFP was overexpressed in *E. coli* strain BL21 (DE3). The protein was recovered from inclusion bodies by dissociating in 6 M GdnHCl. The unfolded S65T/G67A-GFP with polyhistidine tag was purified by metal affinity chromatography. All buffer using during purification contained GdnHCl with 6-molar concentration. The protein was stored in the unfolded state at 4°C in 50 mM phosphate buffer pH 8 with 6 M GdnHCl and 300 mM NaCl. Folding was initiated by diluting S65T/G67A-GFP into buffers with decreasing concentration of GdnHCl. For the unfolding experiments the first step was folding of the protein into the buffer without GdnHCl. The created aggregates were removed by centrifugation and the unfolding was measured by diluting S65T/G67A-GFP into buffers with increasing concentration of GdnHCl. All buffers consisted of 50 mM mixture of Na_2HPO_4 and NaH_2PO_4 , and 300 mM NaCl. As GFP contains two cysteine residues (reduced), 1 mM DTT was included in all buffers to prevent the formation of intra- or inter-molecular disulfide bonds. The protein concentration was 1 μM .

2.2 Chemicals

All chemicals were of guaranteed reagent grade. Guanidinium hydrochloride (GdnHCl), NaCl, dithiothreitol (DTT) and buffer salts were purchased from Roth, and 1,1'-bis(4-anilino-5-naphthalenesulphonic acid) (bis-ANS) was from Molecular Probes. All solutions were prepared using deionized and filtered water from a Millipore water purification system.

2.3 Spectroscopic measurements

All spectrometric measurements were performed at 20°C in 50 mM phosphate buffer pH 8.0, 300 mM NaCl, 1 mM DTT, GdnHCl concentration ranging from 0–6 M. The concentration of protein was 1 μM .

Absorbance measurements were performed on a Uvikon spectrophotometer (Kontron, Austria). Protein concentration was determined using a molar extinction coefficient of $\epsilon_{280\text{nm}} = 22\,140\text{ M}^{-1}\text{cm}^{-1}$ calculated from the amino acids sequence, which is identical (within error) of the value presented by Tsien ($\epsilon_{280\text{nm}} = 22\,000\text{ M}^{-1}\text{cm}^{-1}$) for wild type GFP [6].

Steady-state S65T/G67A-GFP fluorescence data were collected on an LS-50 Perkin-Elmer spectrofluorimeter with excitation at 280 nm, emission at 340 nm. Fluorescence intensity was measured 15 minutes after adding protein to buffer. The quartz cuvette has path length 4 mm

for excitation and 10 nm for emission. The spectral bandwidth was 2.5 nm for excitation and emission.

The population of intermediate state was followed by bisANS fluorescence. Fluorescence emission intensity (excitation 420 nm, emission at 490 nm) was measured 10 minutes after adding protein to buffer, and again after the removal of aggregates.

Protein aggregation was monitored by light scattering at 640 nm, (with the excitation and emission wavelengths at 640 nm, chosen after [9]), 10 minutes after adding protein to buffer, and again after removal of aggregates. Under these conditions, fluorescence of amino acids (Phe, Tyr and Trp) is not observed.

The removal of aggregates was performed by 30 minutes of centrifugation at 11 500×g. The light scattering of a sample without bis-ANS, after centrifugation, was equivalent to the scattering of the buffer. In the case where the light scattering of a mixture of native and aggregated protein was measured, the aggregates were not removed from the sample containing folded and aggregated protein created during folding. Alternatively, aggregates were added to samples of native protein.

The corresponding buffer spectrum was subtracted from each protein spectrum. Each point represents the average from three independent measurements. Errors were calculated as standard deviations from the mean.

GFP folding was initiated by diluting the protein from high to low concentration of GdnHCl. Unfolding was performed by increasing the concentration of GdnHCl.

All data analysis was performed with Microcal Origin.

3. Results and discussion

3.1 Tryptophan fluorescence as a probe of GFP folding and unfolding

GFP contains only one tryptophan (and eleven tyrosines) but, as with other Trp-containing proteins [10], tryptophan fluorescence dominates the emission spectrum upon excitation at 280 or 295 nm, and is sensitive to the environment of the Trp chromophore, in contrast to Tyr fluorescence, which is not particularly sensitive to the surrounding environment. We collected the fluorescence emission intensity at 340 nm after excitation at either 280 or 295 nm. In both cases, a similar trend of fluorescence emission changes as a function of [GdnHCl] was observed, but the absolute intensity was higher for excitation at 280 nm. As excitation at 280 nm results in more efficient total protein fluorescence excitation, this wavelength was used for all measurements.

During folding, S65T/G67A-GFP remains unfolded at concentrations down to 3 M GdnHCl. At lower [GdnHCl], the fluorescence emission increased, reaching a maximum at 0.6 M GdnHCl, consistent with the burial of the Trp residue in a hydrophobic environment (figure 1). From 0.1–0.3 M GdnHCl, emission intensity decreased slightly; at 0 M GdnHCl, the intensity is ~97% of the maximal intensity.

During unfolding at low concentrations of GdnHCl, the fluorescence intensity increased slightly, reaching a maximum at 0.4 M GdnHCl. At higher GdnHCl concentrations, the fluorescence decreases dramatically, reaching a plateau at concentrations higher than 3 M. The midpoint of the folding/unfolding transition is 1.5 ± 0.2 M GdnHCl. The similarity of the folding and unfolding titrations indicates that the folding of this mutant is largely reversible (figure 1). Nevertheless, the non-linear dependence of the fluorescence emission intensity at

low [GdnHCl] means that it is difficult to accurately calculate a free energy for GFP folding using standard techniques [11].

3.2 Probing exposed hydrophobic surface area using bis-ANS

Bis-ANS is a polarity-sensitive fluorescent probe that is minimally fluorescent in polar environment, such as aqueous solution, but its fluorescence emission intensity increases in nonpolar environments. In the common protein folding intermediate termed a molten globule, the hydrophobic side chains of amino acids can create nonpolar surfaces accessible to external agents and can effect binding interactions with hydrophobic probes, such as bis-ANS. The increase in fluorescence emission intensity upon bis-ANS binding has been used to estimate the population of folding intermediates during protein folding reactions [12,13].

The fluorescence emission of bis-ANS as a function of the folded state of GFP is shown in figure 2. The fluorescence was measured before and after removing aggregates by centrifugation. At GdnHCl concentrations down to 2.0 M, the protein remains unfolded and no bis-ANS fluorescence was observed. At lower concentrations of denaturant, before removing aggregates, the fluorescence of bis-ANS increased, indicating the presence of exposed hydrophobic surface area in the partially folded and aggregated protein.

Bis-ANS fluorescence was measured again after the removal of aggregates by centrifugation. From 3.5–0.6 M GdnHCl, bis-ANS fluorescence intensity after centrifugation decreases uniformly (fig. 2). This is clearly seen on the inset, which presents the multiplication of the data for all concentrations of denaturant after centrifugation (figure 2) by a constant value. In the range 3.5 – 0.6 M GdnHCl the fluorescence changes in the same way as the fluorescence before centrifugation. In the range 0.5 – 0 M GdnHCl the decrease fluorescence is observed, caused by the removing of the aggregates. After centrifugation, there is a significant maximum of bis-ANS fluorescence intensity at GdnHCl concentrations between 0.5–0.6 M. This result suggests a maximal concentration of molten globule-like intermediate occurs at these concentrations of GdnHCl.

At lower concentrations of GdnHCl, bis-ANS fluorescence decreases, though there is a slight increase at 0 M GdnHCl. The persistent bis-ANS fluorescence intensity at GdnHCl concentrations below the threshold for S65T/G67A-GFP folding, even after the removal of aggregated material by centrifugation (see section 3.3, below), indicates that the solution contains, in addition to folded S65T/G67A-GFP, small soluble yet misfolded conformations, possibly as small oligomeric structures.

3.3 Light scattering measurements of GFP aggregates

Substantial light scattering was observed during folding of S65T/G67A-GFP to residual concentration of GdnHCl (figure 3). Light scattering was low at GdnHCl concentration down to 0.6 M, but increased dramatically at lower concentrations of GdnHCl, reaching a maximum at 0.2 M GdnHCl. These results suggest that, at intermediate concentrations of GdnHCl, GFP populates an intermediate conformation that is particularly prone to aggregation. Unfolding native (without aggregates) GFP in increasing concentration of denaturant did not result in increased light scattering, indicating aggregation does not occur during GFP unfolding (figure 3). The intensity of scattering was constant as a function of GdnHCl concentration, showing that scattering is not sensitive to GFP conformational changes during unfolding.

Observation of the light scattering during unfolding was also performed on a mixture of folded and aggregated GFP. At low concentrations of GdnHCl, light scattering was high, decreasing only above 0.9 M GdnHCl, suggesting higher concentrations of GdnHCl are required to dissociate and solubilize S65T/G67A-GFP aggregates.

4. Conclusions

As measured by tryptophan fluorescence, S65T/G67A-GFP folding is largely reversible. Upon dilution from 6 M GdnHCl, S65T/G67A-GFP remains unfolded down to 3 M GdnHCl. Between 0.6–3 M GdnHCl, a partially folded intermediate is formed. This intermediate binds bis-ANS, indicating that it resembles a classical “molten globule”-type folding intermediate. The existence of intermediate is typical for many beta-sheet proteins [14,15]. However, the complicated shape of the folding curve, especially pronounced below 0.6 M concentration of GdnHCl, is not a typical phenomenon observed during folding of other beta-sheet proteins.

The persistent bis-ANS fluorescence intensity at GdnHCl residual concentrations after the removal of aggregates indicates that the solution contains mixture of folded S65T/G67A-GFP and small misfolded conformations, which are not removed by centrifugation.

During folding, the increasing of light scattering is delayed in comparison with tryptophan and bis-ANS fluorescence. It is minimal at/above 0.6 M GdnHCl, but increases dramatically at lower concentrations of GdnHCl. These results suggest that, at equilibrium, upon gradual removal of GdnHCl, GFP first forms a molten globule structure, which next can choose a way either to an aggregation or to the creation of the native structure. These aggregates are present in solution during next steps of folding. According to the fluorescence of bis-ANS it is not possible to remove aggregates by centrifugation.

During unfolding of the native protein the aggregation is not observed. In the case of mixture of aggregated and folded protein, probably two processes are observed: unfolding of the native protein and dissociation of aggregates clearly seen at concentrations of GdnHCl higher than 0.9 M.

Acknowledgments

We are indebted to Kay Finn for preparation of the mutant S65T/G67A-GFP and excellent technical assistance. B W-K thanks to prof. Edward Darzynkiewicz for the use of his laboratory for protein purification. Supported by BW-1684/BF project from Warsaw University and an National Science Foundation CAREER Award to P.L.C.(MCB-0237945).

References

1. Capaldi AP, Radford SE. Kinetic studies of β -sheet protein folding. *Curr Opin Struct Biol* 1998;8:86–92. [PubMed: 9519300]
2. Dobson CM. Protein aggregation and its consequences for human disease. *Protein Pept Lett* 2006;13:219–27. [PubMed: 16515449]
3. Richardson JS, Richardson DC. Natural β -sheet proteins use negative design to avoid edge-to-edge aggregation. *Proc Natl Acad Sci U S A* 2002;99:2754–59. [PubMed: 11880627]
4. Lippincott-Schwartz J, Patterson GH. Development and use of fluorescent protein markers in living cells. *Science* 2003;300:87–91. [PubMed: 12677058]
5. Reid B, Fynn GC. Chromophore formation in green fluorescent protein. *Biochem* 1997;36:6786–91. [PubMed: 9184161]
6. Tsien RY. The green fluorescent protein. *Ann Rev Biochem* 1998;67:509–44. [PubMed: 9759496]
7. Ward WW, Cody CW, Hart RC, Cormier MJ. Spectrophotometric identity of the energy transfer chromophores in Renilla and Aequorea green-fluorescent proteins. *Photochem Photobiol* 1980;31:611–15.
8. Chalfie, M.; Kain, S., editors. *Green Fluorescent Protein – Properties and Applications*. New York: Willey-Liss Inc; 1998.
9. Fukuda H, Arai M, Kuwajima K. Folding of green fluorescent protein and the cycle 3 mutant. *Biochem* 2000;39:12025–32. [PubMed: 11009617]

10. Lakowicz, JR. Principles of Fluorescence Spectroscopy. 2. Vol. Chap 16. Kluwer Academic/Plenum Publishers; 1999.
11. Pace CN. Determination and analysis of urea and guanidine hydrochloride denaturation curves. *Methods Enzymol* 1986;131:266–80. [PubMed: 3773761]
12. Bino J, D'Silva PR, Lala AK. Analysis of protein folding using polarity-sensitive fluorescent probes. *Curr Sci* 2001;80:287–90.
13. Semisotniv GV, Rodionova NA, Razgulyaev OI, Uversky VN, Gripas AF, Gilmanshin RI. Study of the “molten globule” intermediate state in protein folding by a hydrophobic fluorescent probe. *Biopolymers* 1991;1:119–28.
14. Reder JS, Van Nuland NA, Thompson GS, Ferguson SJ, Dobson CM, Radford SE. A partially folded intermediate species of the beta-sheet protein apo-pseudoazurin is trapped during proline-limited folding. *Protein Sci* 2001;10:1216–24. [PubMed: 11369860]
15. Dalessio PM, Ropson IJ. Beta-sheet proteins with nearly identical structures have different folding intermediates. *Biochem* 2000;39:860–71. [PubMed: 10653629]

Figure 1.

Changes in fluorescence emission intensity for excitation wavelength 280 nm and emission measured at 340 nm during folding (grey circles) and unfolding (black squares) of S65T/G67A-GFP as a function of denaturant concentration. Measurements were conducted at 20°C in 50 mM phosphate buffer pH 8.0 with 300 mM NaCl and 1 mM DTT after ten minutes of equilibration.

Figure 2. Binding of bis-ANS to S65T/G67A-GFP monitored by fluorescence emission intensity at 490 nm (excitation at 420 nm) at various concentration of GdnHCl before (black diamonds) and after (grey diamonds) removing aggregates by centrifugation at 11 500g. Data were collected at 20°C in 50 mM phosphate buffer pH 8.0 with 300 mM NaCl and 1 mM DTT. Inset: Relative fluorescence of bis-ANS before (filled squares) and after (open squares) removing aggregates.

Figure 3.

Light scattering of S65T/G67A-GFP at 640 nm as a function of denaturant concentration during unfolding of native protein without aggregates (grey squares), unfolding the native protein in the presence of aggregates (open triangles) and during folding of S65T/G67A-GFP (black circles). Data were collected at 20°C in 50 mM phosphate buffer pH 8.0 with 300 mM NaCl and 1 mM DTT.