Alteration of A β metabolism-related molecules in predementia induced by AlCl₃ and D-galactose

Zong-Zheng Sun · Zhi-Bin Chen · Hui Jiang ·
Ling-Ling Li · Er-Guang Li · Yun Xu

Received: 30 January 2009 / Accepted: 7 May 2009 / Published online: 26 May 2009
© American Aging Association 2009

Abstract The purpose of this study was to look for alterations in β -amyloid peptide (A β) metabolism-related molecules in predementia, the early stage of Alzheimer's disease (AD). AlCl₃ (Al) and D-galactose (D-gal) were used to induce the mouse model for predementia and AD. Protein expression of β -amyloid (A β), β -secretase (BACE1), neprilysin (NEP), insulin degrading enzyme (IDE) and receptor for advanced glycation end products (RAGE) in the brain was measured. The results indicated that Al + D-gal

induced an AD-like behavioral deficit at 90 days. The period from 45 to 75 days showed no significant behavioral deficit, and we tentatively define this as predementia in this model. A significant increase in BACE1 and decreasing NEP characterized days 45–90 in the cortex and hippocampus. However, high A β occurred at day 60. IDE increased from day 60 to day 75. There was no change in RAGE. The results suggest that the observed changes in BACE1, NEP and A β in predementia might relate to a different stage of the AD-like pathology, which may be developed into useful biomarkers for the diagnosis of very early AD.

Zong-Zheng Sun and Zhi-Bin Chen contributed equally to this work.

Z.-Z. Sun · Z.-B. Chen · H. Jiang · E.-G. Li · Y. Xu (✉)
Department of Neurology, Affiliated Drum Tower Hospital,
Nanjing University Medical School,
321 Zhong Shan Road,
Nanjing, Jiangsu 210008, People's Republic of China
e-mail: xuyun20042001@yahoo.com.cn

L.-L. Li · Y. Xu
Department of Neurology, Affiliated Drum Tower Hospital
of Nanjing, School of Clinical Medicine,
Southeast University,
Nanjing, Jiangsu, People's Republic of China

E.-G. Li · Y. Xu
The State Key Laboratory of Pharmaceutical
Biotechnology, Nanjing University,
Nanjing, Jiangsu 210093, People's Republic of China

E.-G. Li · Y. Xu
Jiangsu Key Laboratory for Molecular Medicine,
Nanjing, Jiangsu 210093, People's Republic of China

Keywords Predementia · β -Amyloid · β -Secretase ·
Neprilysin

Introduction

Alzheimer's disease (AD) is the most common cause of dementia in the elderly, accounting for 65–70% of all dementia (Blennow et al. 2006). However, early AD cannot be diagnosed even if the progressing cognitive deficits affect the patient's ability to cope with the functional demands of his or her social and professional life (Hampel et al. 2008). Biomarkers are attracting increasing attention in research into early diagnosis of AD or as surrogate measures of the ongoing pathology. The major neuropathological hallmark of AD is senile plaque, which occurs with aging of the human brain many years prior to disease

onset. Senile plaque is formed by conversion of the A β peptide to amyloid in the brain (Sommer 2002).

The overproduction or impaired clearance of cerebral β -amyloid peptide (A β) leads to elevated A β levels that promote A β aggregation, oligomerization, and fibrillogenesis (Hardy and Selkoe 2002). A β is generated by the cleavage of amyloid precursor protein (APP) by β -secretase (BACE1) and γ -secretase (Vardy et al. 2005). Pathways involved in the removal of cerebral amyloid peptide include enzymatic degradation and receptor-mediated clearance of the cerebral A β . The two major endopeptidases involved in A β enzymatic degradation are insulin degrading enzyme (IDE) and neprilysin (NEP). Multiple lines of evidence support a role for IDE and NEP in A β degradation (Farris et al. 2004; Tanzi et al. 2004; Wang et al. 2006). Because of the presence of a continuous monolayer of brain endothelial cells, the blood–brain barrier (BBB) in vivo does not allow free exchange of polar solutes such as A β between brain and blood. The receptor for advanced glycation end products (RAGE)—the specialized receptor for A β at the BBB—has been implicated strongly in amyloid peptide influx back into the central nervous system (CNS; Zlokovic 2004; Donahue et al. 2006). Therefore, the increased A β load in AD patients may result from the increased influx of circulating A β across the BBB and decreased A β degradation.

Most previous studies have focused on the examination of A β and Tau in cerebrospinal fluid (CSF) as markers in AD patients. Reports on other molecules of A β generation, transportation and degradation as markers for early diagnosis of AD are limited. Our previous study showed that Al + D-galactose (D-gal) induced an AD-like behavioral deficit at 90 days (Luo et al. 2008). In this study, the mouse was treated by Al + D-gal, and tested to see if molecules associated with A β metabolism were involved in the pathological process of AD before A β deposition in brain and the learning/memory injury of mice. We tentatively define the period from 45 to 75 days as predementia in this model, because there was no significant behavioral deficit during this period. Results show that alterations in the expression of BACE1, NEP and IDE occur earlier than high expression of A β in the mouse brain and behavior changes. This suggests that detection of BACE1, NEP and IDE levels as biomarkers could be

useful for the early diagnosis of AD as well as representing potential pharmacological targets for the preservation of normal brain function and the treatment of AD.

Materials and methods

AD model

Generation of the AD model (Luo et al. 2008): Kunming strain mice (6 to 8 weeks old and 20–25 g in groups of ten) were treated with AlCl₃ 40 mg kg⁻¹ day⁻¹ and D-gal 90 mg kg⁻¹ day⁻¹, i.p., once a day for 90 days using saline water as a vehicle. Animals were treated according to the guidelines of the Regulations of Experimental Animal Administration issued by the State Committee of Science and Technology of the People's Republic of China on 14 November 1988. The experiments were carried out under the approval of the Committee of Experimental Animal Administration of Nanjing University.

Morris water maze

To evaluate the learning/memory ability of the mice, a water maze behavioral test was performed at 0, 45, 60, 75 and 90 days after the treatment, as described previously (Luques et al. 2007). A circular pool (150 cm in diameter and 45 cm deep) with walls and floor painted blank was filled with opaque water (23±1 °C). A hidden circular platform (28 cm high, 12 cm in diameter, 2 cm below the water surface, fixed position) was located in the pool away from the pool wall. The pool was conceptually divided into four equal quadrants: NW, NE, SW and SE. Mice were trained on a visible platform (cued platform) for 3 days followed by 4 days of testing for learning and memory abilities with the platform submerged (hidden platform). Each mouse was subjected to four trials a day over a 4-day period. A mouse was placed in the water facing the pool wall at one of the four quadrants at a different place each day, and allowed to swim for a maximum of 90 s to locate the hidden platform, where it was allowed to stay for 10 s. The time taken for each mouse to locate the platform in all four trials was averaged and recorded. The daily administration of D-gal and Al was continued during the water maze performance tests.

Western blotting

The mice were given anesthesia at 0, 45, 60, 75 and 90 days after D-gal/Al treatment. Protein from the hippocampus and frontal cortex was extracted and then quantified using a Coomassie Blue Fast Staining Solution according to the manufacturer's instructions. Equal amounts of protein samples were separated by SDS-PAGE and blotted onto polyvinylidene fluoride (PVDF) membranes (Xu et al. 2006). The membranes were probed with primary antibodies against A β 1–42 (1:500, AB5078P, Chemicon, Temecula, CA), BACE1 (1:500, MAB5308, Chemicon), NEP (1:500, BAF1126, R&D), IDE (1:1,000, ab25970, abcam, Cambridge, UK), RAGE (1:500, AF1179, R&D Systems, Minneapolis, MN). GAPDH (1:500, Santa Cruz Europe, Heidelberg, Germany) was used as loading control. The proteins were detected using horseradish peroxidase-conjugated anti-rabbit or anti-mouse secondary antibodies and visualized using chemiluminescence reagents provided with the ECL kit (Amersham Pharmacia Biotech, Piscataway, NJ) and exposed to film. The intensity of blots was quantified by densitometry. All assays were performed at least in triplicate in independent experiments.

Statistical analysis

SPSS 13.0 software was used to handle all data. The results were expressed as mean $\pm$ SEM. Behavioral tests were determined by analysis of variance (one-way ANOVA) and paired *t*-test. The level of proteins was analyzed by Student's *t*-test (two-tailed) or one-way ANOVA followed by post hoc Tukey test. The correlation between the expression of proteins and A β was assessed by Pearson correlation analyses. Statistical significance was set at $P < 0.05$.

Results

Behavioral results

None of the mice tested had obvious health problems (e.g., weight loss, cataracts, or toxicity reactions). Throughout the training, on day 90, mice treated with Al and D-gal took longer to reach the platform than those of the vehicle group ($P > 0.05$, Fig. 1). There

Fig. 1 Morris water maze. Kunming strain mice were treated with AlCl₃ 40 mg kg⁻¹ day⁻¹ and D-galactose (D-gal) 90 mg kg⁻¹ day⁻¹, i.p., once a day to induce the AD model. The vehicle was saline water. Each group consisted of ten mice. The Morris water maze test was applied 0, 45, 60, 75 and 90 days after treatment. No significant difference was seen at 0, 45, 60, and 75 days in mice treated with Al/D-gal vs controls. (one-way ANOVA, * $P > 0.05$). Compared to mice treated with saline or Al/D-gal at 0 days, mice treated with Al/D-gal exhibited significantly prolonged escape latency only at 90 days (paired *t*-test, $P < 0.05$)

was no significant difference between treatment and vehicle at 0, 45, 60, and 75 days. Each group consisted of ten mice.

Expression of A β and BACE1 in predementia mouse brain

To investigate if molecules of the associated A β pathway start to be induced before the decline of cognition function in this model, A β expression was measured by western blotting at 0, 45, 60, 75 and 90 days. The results indicated that there was a tendency for increased A β expression in the cortex (Co) and hippocampus (Hi) of the treated mice brain at 45 days, but this was not significant compared to the vehicle group. In contrast, a significant increase in A β expression was observed at matched time points on days 60, 75 and 90 (Figs. 2, 3). Next, the level of BACE1 expression in the brain of the “predementia mouse” was examined. Compared to A β expression, high BACE1 appeared early. The level of BACE1 in the treatment group increased significantly from day 45 to 90 in both Co and Hi compared to control levels (Figs. 2, 4). Statistical analysis showed that the level of BACE1 correlated with the level of A β ($R = 0.9351$, $P < 0.05$ for Hi; $R = 0.9688$, $P < 0.01$ for Co). The highest level was seen on day 75, confirming that overproduction of APP protein was not induced by Al

Fig. 2 Western blotting. Proteins of A β metabolism-related molecules from mouse hippocampus (Hi) and frontal cortex (Co) with Al/D-gal were measured at different timepoints during treatment. The vehicle was saline; GAPDH was used as a loading control. All experiments were performed at least in triplicate (data are presented as the mean $\pm$ SEM of three independent experiments)

and D-gal (data not shown). In this paper, no data for vehicle only is given because there was no difference between the treatment group at day 0 and vehicle groups from day 0 to day 90, thus we used day 0 as a control.

NEP and IDE alteration in predementia mouse brain

NEP and IDE are the main degrading enzymes for A β clearance in the brain. The protein levels of NEP and IDE were measured in the hippocampus and the frontal cortex. The results indicated that the level of NEP was significantly reduced in both tissues at day 45 ($P < 0.05$) compared to that of the vehicle group, and decreased further at 60, 75 and 90 days (Figs. 2, 5). However, the level of NEP did not correlate significantly with the level of A β ($r = -0.8718$, $P > 0.05$ for Hi, $r = -0.793$, $P > 0.05$ for Co). However, as Fig. 6 shows, expression of IDE in the cortex was significantly higher in the treatment group than that in the vehicle group at days 60 and 75 ($P < 0.05$), and high expression of IDE in the hippocampus was observed only at day 60 ($P < 0.05$). There was no significant change in IDE protein between the two groups at other time points ($P > 0.05$).

RAGE level in predementia mouse brain

RAGE mediates a continuous influx of circulating A β into the brain. To confirm whether RAGE is a useful marker for early diagnosis of AD, RAGE protein levels were determined. We found that expression of RAGE in the brain was a slightly higher in comparison with the control after treatment with Al + D-gal, but this difference did not reach significance ($P < 0.05$, Figs. 2, 7).

Discussion

Current state of early diagnosis of AD

Alzheimer's disease is currently most commonly identified using clinical criteria and psychometric cognitive assessment. The Mini-Mental State Examination (MMSE) is often used as a screening test for dementia. However, the MMSE has been found to be insensitive to early AD, and new tools for diagnosing early stages of AD are being developed (Benson et al. 2005; Nasreddine et al. 2005). Furthermore, in the early stage, there is no clinical method to determine whether early stage AD patients will progress to AD—long clinical follow up is required. Recently, some longitudinal studies have used PET or MRI images to predict and monitor cognitive decline from normal aging to early AD (Fritzsche et al. 2008; Mosconi et al. 2008). Both CSF biomarker levels and atrophy as assessed by MRI are now used in the diagnostic work-up of AD (Fox et al. 2005; Wiltfang et al. 2005; Waldemar et al. 2007). Increasing numbers of reports show that the amount of A β in the brain is determined by the rate of A β generation versus clearance. On this basis, it was hypothesized that alteration in the expression of A β -metabolism-associated molecules before pathological and behavioral changes could be developed into early markers of AD progression. The main findings of the present study were: (1) accumulation of high levels of BACE1 and A β in mouse brain occurred earlier than mouse memory impairment, (2) an increase in BACE1 and decrease in NEP were detected earlier than changes in A β , (3) the level of RAGE remained constant throughout the entire experimental

Figs. 3–7 Protein expression in Aβ13/D-gal treated mice at different days during treatment (control = saline). The expression of β-amyloid peptide (Aβ), β-secretase (BACE1), neprilysin (NEP), insulin degrading enzyme (IDE), and receptor for advanced glycation end products (RAGE) protein was assessed by western blotting and the relative densities were assessed using Band Scan software. One-way ANOVA fol-

lowed by post hoc Tukey test. * $P < 0.05$ vs vehicle. Fig. 3 Aβ expression was significantly higher at 60, 75 and 90 days; Fig. 4 Alterations of BACE1 were significant from 45 to 90 days; Fig. 5 Alterations of NEP were significant from 45 to 90 days; Fig. 6 Significant changes in IDE was seen at 60 and 75 days; Fig. 7 No significant changes in RAGE were observed on any day

period. This finding suggests that, besides Aβ, Aβ-pathway-related molecules such as BACE1 and NEP could also be developed into critical biological markers for early diagnosis of AD in the future.

Alteration of Aβ and BACE1 in the predementia mouse brain

The period from day 45 to day 75 after treatment with A1 and D-gal was arbitrarily defined as the

“predementia stage” in our current model. AD is seen to appear by day 90, because the behavioral test showed that the learning/memory of mice in the treatment group decreased significantly compared to that in the vehicle group at this time. The present study found that expression of BACE1 increased from day 45, i.e., earlier than Aβ expression, which begins to increase at day 60. Both BACE1 and Aβ alteration occurred before behavior changes. BACE1 cleavage is a pre-

requisite for A β formation, and is the initiating step in A β generation. Ablation of BACE1 in transgenic AD models abolishes A β production and prevents the subsequent development of amyloid-associated pathologies (Ohno et al. 2004, 2007; McConlogue et al. 2007). BACE1 levels are elevated in both AD experimental models and in the AD patient brain (Li et al. 2004; Harada et al. 2006; Zhao et al. 2007; Tamagno et al. 2002, 2005). BACE1 activity is correlated with brain A β production in the frontal cortex (Li and Sudhof 2004; Gustaw et al. 2008). In agreement with others, our data suggests that BACE1 levels may start building early and become sustained during the course of AD development. In addition, the BACE1 elevation in AD may be actively involved in AD pathology and may occur prior to the appearance of overt neuron death (Zhao et al. 2007). Statistical analysis indicates that changes in BACE1 have a positive correlation with changes of brain A β .

NEP and IDE level in the predementia mouse brain

Although the cause of increased A β in Alzheimer's disease can be attributed to many factors, evidence is accumulating that it may be a result of a decrease in A β clearance (Selkoe 2001). A β clearance occurs primarily through the action of NEP and IDE. The deletion or disruption of either of these peptidase genes in mice leads to a significant increase in A β levels (Farris et al. 2003; Miller et al. 2003). This study observed that NEP began to decrease at day 45. NEP changes occurred 15 days prior to the A β increase in these mouse brains. NEP cleaves and removes A β and, consequently, leads to a delay in amyloid plaque deposition in the A β -mediated AD brain (Mohajeri et al. 2002). NEP levels were reduced in affected AD brains, and in the brains of aged mice, and are known to be down-regulated in areas vulnerable to A β peptide accumulation in the AD brain (Yasojima et al. 2001; Iwata et al. 2002). This study and others suggest that NEP has a strong negative correlation with A β aggregation and thus may be a suitable candidate for development into a critical marker for the diagnosis of early AD.

On the contrary, levels of IDE, the rate-limiting A β -degrading enzyme (Hama et al. 2001), were significantly higher in the treatment group than in the vehicle group at day 60 in both Co and Hi, and at

day 75 only in Co. The capacity of IDE to degrade A β has been recognized for many years (Kurochkin and Goto 1994). The cause of increased IDE may be due partly to activated astrocytes as a result of A β -triggered neuroinflammation (Leal et al. 2006). Contradictory results regarding IDE expression in the AD brain have been reported (Kim et al. 2007). Therefore IDE is not suitable as a special marker for the early diagnosis of AD since it might be involved in the inflammation reaction in the early stage of AD, and its role on A β degradation occurs after AD. The decrease in IDE happens in the late stage of AD (Tanzi et al. 2004).

RAGE expression in the predementia brain

RAGE has been characterized as a transporter of A β from the plasma to the brain across the BBB (Deane et al. 2004). RAGE is regarded as a multiligand receptor in the immunoglobulin (IgG) superfamily. RAGE binds soluble A β and mediates pathophysiologically relevant cellular responses (Stern et al. 2002). This study found no changes of RAGE levels in the brain at any timepoint in this model. In the central nervous system, RAGE is present at high levels during development, and its levels decline during maturity in normal aging (Leclerc et al. 2007). Microvascular RAGE expression was notably up-regulated in advanced AD as compared to early AD and control samples (Miller et al. 2008).

In conclusion, Al + D-gal treatment can generate high A β induction in mice brains, which serves as an animal model for predementia or AD. BACE1 and NEP alterations were apparent before changes in A β , showing positive and negative correlations with A β , respectively. Both may be developed into useful biomarkers for the early diagnosis of AD, and can also be regarded as targets for a therapeutic strategy against predementia and AD.

Acknowledgments This work was supported by the National Nature Science Foundation of China (30470612, 30670739), the Doctoral Program Foundation of the Ministry of Education of China (20060284044), the International Cooperation Program and talented man program (BZ2006045, 06-B-002, RC2007006) of Jiangsu Province of China of Jiangsu Province of China, and 973 funding from the Ministry of Science and Technology in China (2009CB21906). We thank Marilyn White and Prof. Ying-Dong Zhang for writing modification.

References

- Benson AD, Slavin MJ, Tran TT, Petrella JR, Doraiswamy PM (2005) Screening for early Alzheimer's disease: is there still a role for the mini-mental state examination? *Prim Care Companion J Clin Psychiat* 7:62–69
- Blennow K, de Leon MJ, Zetterberg H (2006) Alzheimer's disease. *Lancet* 368:387–403. doi:10.1016/S0140-6736(06)69113-7
- Deane R, Wu Z, Zlokovic BV (2004) RAGE (yin) versus LRP (yang) balance regulates alzheimer amyloid beta-peptide clearance through transport across the blood-brain barrier. *Stroke* 35:2628–2631. doi:10.1161/01.STR.0000143452.85382.d1
- Donahue JE, Flaherty SL, Johanson CE, Duncan JA 3rd, Silverberg GD, Miller MC, Tavares R, Yang W, Wu Q, Sabo E, Hovanesian V, Stopa EG (2006) RAGE, LRP-1, and amyloid-beta protein in Alzheimer's disease. *Acta Neuropathol* 112:405–415. doi:10.1007/s00401-006-0115-3
- Farris W, Mansourian S, Chang Y, Lindsley L, Eckman EA, Frosch MP, Eckman CB, Tanzi RE, Selkoe DJ, Guenette S (2003) Insulin-degrading enzyme regulates the levels of insulin, amyloid beta-protein, and the beta-amyloid precursor protein intracellular domain in vivo. *Proc Natl Acad Sci USA* 100:4162–4167. doi:10.1073/pnas.0230450100
- Farris W, Mansourian S, Leissring MA, Eckman EA, Bertram L, Eckman CB, Tanzi RE, Selkoe DJ (2004) Partial loss-of-function mutations in insulin-degrading enzyme that induce diabetes also impair degradation of amyloid beta-protein. *Am J Pathol* 164:1425–1434
- Fox NC, Black RS, Gilman S, Rossor MN, Griffith SG, Jenkins L, Koller M (2005) Effects of Abeta immunization (AN1792) on MRI measures of cerebral volume in Alzheimer disease. *Neurology* 64:1563–1572. doi:10.1212/01.WNL.0000159743.08996.99
- Fritzsche KH, von Wangenheim A, Abdala DD, Meinzer HP (2008) A computational method for the estimation of atrophic changes in Alzheimer's disease and mild cognitive impairment. *Comput Med Imaging Graph* 32:294–303. doi:10.1016/j.compmedimag.2007.12.006
- Gustaw KA, Garrett MR, Lee HG, Castellani RJ, Zagorski MG, Prakash A, Siedlak SL, Zhu X, Perry G, Petersen RB, Friedland RP, Smith MA (2008) Antigen-antibody dissociation in Alzheimer disease: a novel approach to diagnosis. *J Neurochem* 106:1350–1356. doi:10.1111/j.1471-4159.2008.05477.x
- Hama E, Shirohata K, Masumoto H, Sekine-Aizawa Y, Aizawa H, Saido TC (2001) Clearance of extracellular and cell-associated amyloid beta peptide through viral expression of neprilysin in primary neurons. *J Biochem* 130:721–726
- Hampel H, Burger K, Teipel SJ, Bokde AL, Zetterberg H, Blennow K (2008) Core candidate neurochemical and imaging biomarkers of Alzheimer's disease. *Alzheimers Dement* 4:38–48. doi:10.1016/j.jalz.2007.08.006
- Harada H, Tamaoka A, Ishii K, Shoji S, Kametaka S, Kametani F, Saito Y, Murayama S (2006) Beta-site APP cleaving enzyme 1 (BACE1) is increased in remaining neurons in Alzheimer's disease brains. *Neurosci Res* 54:24–29. doi:10.1016/j.neures.2005.10.001
- Hardy J, Selkoe DJ (2002) The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297:353–356. doi:10.1126/science.1072994
- Iwata N, Takaki Y, Fukami S, Tsubuki S, Saido TC (2002) Region-specific reduction of a beta-degrading endopeptidase, neprilysin, in mouse hippocampus upon aging. *J Neurosci Res* 70:493–500. doi:10.1002/jnr.10390
- Kim M, Hersh LB, Leissring MA, Ingelsson M, Matsui T, Farris W, Lu A, Hyman BT, Selkoe DJ, Bertram L, Tanzi RE (2007) Decreased catalytic activity of the insulin-degrading enzyme in chromosome 10-linked Alzheimer disease families. *J Biol Chem* 282:7825–7832. doi:10.1074/jbc.M609168200
- Kurochkin IV, Goto S (1994) Alzheimer's beta-amyloid peptide specifically interacts with and is degraded by insulin degrading enzyme. *FEBS Lett* 345:33–37. doi:10.1016/0014-5793(94)00387-4
- Leal MC, Dorfman VB, Gamba AF, Frangione B, Wisniewski T, Castano EM, Sigurdsson EM, Morelli L (2006) Plaque-associated overexpression of insulin-degrading enzyme in the cerebral cortex of aged transgenic tg2576 mice with Alzheimer pathology. *J Neuropathol Exp Neurol* 65:976–987. doi:10.1097/01.jnen.0000235853.70092.ba
- Leclerc E, Fritz G, Weibel M, Heizmann CW, Galichet A (2007) S100B and S100A6 differentially modulate cell survival by interacting with distinct RAGE (receptor for advanced glycation end products) immunoglobulin domains. *J Biol Chem* 282:31317–31331. doi:10.1074/jbc.M703951200
- Li Q, Sudhof TC (2004) Cleavage of amyloid-beta precursor protein and amyloid-beta precursor-like protein by BACE 1. *J Biol Chem* 279:10542–10550. doi:10.1074/jbc.M310001200
- Li R, Lindholm K, Yang LB, Yue X, Citron M, Yan R, Beach T, Sue L, Sabbagh M, Cai H, Wong P, Price D, Shen Y (2004) Amyloid beta peptide load is correlated with increased beta-secretase activity in sporadic Alzheimer's disease patients. *Proc Natl Acad Sci USA* 101:3632–3637. doi:10.1073/pnas.0205689101
- Luo Y, Niu F, Wang L, Cao W, Zhang X, Lv Z, Zhang B, Xu Y (2008) Altered expression of Abeta metabolism-associated molecules from d-galactose/AICl(3) induced mouse brain. *Mech Ageing Dev* . doi:10.1016/j.mad.2008.12.005
- Luques L, Shoham S, Weinstock M (2007) Chronic brain cytochrome oxidase inhibition selectively alters hippocampal cholinergic innervation and impairs memory: prevention by ladostigil. *Exp Neurol* 206:209–219. doi:10.1016/j.expneurol.2007.04.007
- McConlogue L, Buttini M, Anderson JP, Brigham EF, Chen KS, Freedman SB, Games D, Johnson-Wood K, Lee M, Zeller M, Liu W, Motter R, Sinha S (2007) Partial reduction of BACE1 has dramatic effects on Alzheimer plaque and synaptic pathology in APP Transgenic Mice. *J Biol Chem* 282:26326–26334. doi:10.1074/jbc.M611687200
- Miller BC, Eckman EA, Sambamurti K, Dobbs N, Chow KM, Eckman CB, Hersh LB, Thiele DL (2003) Amyloid-beta peptide levels in brain are inversely correlated with insulin activity levels in vivo. *Proc Natl Acad Sci USA* 100:6221–6226. doi:10.1073/pnas.1031520100

- Miller MC, Tavares R, Johanson CE, Hovanesian V, Donahue JE, Gonzalez L, Silverberg GD, Stopa EG (2008) Hippocampal RAGE immunoreactivity in early and advanced Alzheimer's disease. *Brain Res* 1230:273–280. doi:[10.1016/j.brainres.2008.06.124](https://doi.org/10.1016/j.brainres.2008.06.124)
- Mohajeri MH, Wollmer MA, Nitsch RM (2002) Abeta 42-induced increase in neprilysin is associated with prevention of amyloid plaque formation in vivo. *J Biol Chem* 277:35460–35465. doi:[10.1074/jbc.M202899200](https://doi.org/10.1074/jbc.M202899200)
- Mosconi L, De Santi S, Li J, Tsui WH, Li Y, Boppana M, Laska E, Rusinek H, de Leon MJ (2008) Hippocampal hypometabolism predicts cognitive decline from normal aging. *Neurobiol Aging* 29:676–692. doi:[10.1016/j.neurobiolaging.2006.12.008](https://doi.org/10.1016/j.neurobiolaging.2006.12.008)
- Nasreddine ZS, Phillips NA, Bedirian V, Charbonneau S, Whitehead V, Collin I, Cummings JL, Chertkow H (2005) The Montreal Cognitive Assessment, MoCA: a brief screening tool for mild cognitive impairment. *J Am Geriatr Soc* 53:695–699. doi:[10.1111/j.1532-5415.2005.53221.x](https://doi.org/10.1111/j.1532-5415.2005.53221.x)
- Ohno M, Sametsky EA, Younkin LH, Oakley H, Younkin SG, Citron M, Vassar R, Disterhoft JF (2004) BACE1 deficiency rescues memory deficits and cholinergic dysfunction in a mouse model of Alzheimer's disease. *Neuron* 41:27–33. doi:[10.1016/S0896-6273\(03\)00810-9](https://doi.org/10.1016/S0896-6273(03)00810-9)
- Ohno M, Cole SL, Yasvoina M, Zhao J, Citron M, Berry R, Disterhoft JF, Vassar R (2007) BACE1 gene deletion prevents neuron loss and memory deficits in 5XFAD APP/PS1 transgenic mice. *Neurobiol Dis* 26:134–145. doi:[10.1016/j.nbd.2006.12.008](https://doi.org/10.1016/j.nbd.2006.12.008)
- Selkoe DJ (2001) Clearing the brain's amyloid cobwebs. *Neuron* 32:177–180. doi:[10.1016/S0896-6273\(01\)00475-5](https://doi.org/10.1016/S0896-6273(01)00475-5)
- Sommer B (2002) Alzheimer's disease and the amyloid cascade hypothesis: ten years on. *Curr Opin Pharmacol* 2:87–92. doi:[10.1016/S1471-4892\(01\)00126-6](https://doi.org/10.1016/S1471-4892(01)00126-6)
- Stern D, Yan SD, Yan SF, Schmidt AM (2002) Receptor for advanced glycation endproducts: a multiligand receptor magnifying cell stress in diverse pathologic settings. *Adv Drug Deliv Rev* 54:1615–1625. doi:[10.1016/S0169-409X\(02\)00160-6](https://doi.org/10.1016/S0169-409X(02)00160-6)
- Tamagno E, Bardini P, Obbili A, Vitali A, Borghi R, Zaccheo D, Pronzato MA, Danni O, Smith MA, Perry G, Tabaton M (2002) Oxidative stress increases expression and activity of BACE in NT2 neurons. *Neurobiol Dis* 10:279–288. doi:[10.1006/nbdi.2002.0515](https://doi.org/10.1006/nbdi.2002.0515)
- Tamagno E, Parola M, Bardini P, Piccini A, Borghi R, Guglielmotto M, Santoro G, Davit A, Danni O, Smith MA, Perry G, Tabaton M (2005) β -Site APP cleaving enzyme up-regulation induced by 4-hydroxynonenal is mediated by stress-activated protein kinases pathways. *J Neurochem* 92:628–636. doi:[10.1111/j.1471-4159.2004.02895](https://doi.org/10.1111/j.1471-4159.2004.02895)
- Tanzi RE, Moir RD, Wagner SL (2004) Clearance of Alzheimer's Abeta peptide: the many roads to perdition. *Neuron* 43:605–608
- Vardy ER, Catto AJ, Hooper NM (2005) Proteolytic mechanisms in amyloid-beta metabolism: therapeutic implications for Alzheimer's disease. *Trends Mol Med* 11:464–472. doi:[10.1016/j.molmed.2005.08.004](https://doi.org/10.1016/j.molmed.2005.08.004)
- Waldemar G, Dubois B, Emre M, Georges J, McKeith IG, Rossor M, Scheltens P, Tariska P, Winblad B (2007) Recommendations for the diagnosis and management of Alzheimer's disease and other disorders associated with dementia: EFNS guideline. *Eur J Neurol* 14:e1–e26. doi:[10.1111/j.1468-1331.2006.01605.x](https://doi.org/10.1111/j.1468-1331.2006.01605.x)
- Wang YJ, Zhou HD, Zhou XF (2006) Clearance of amyloid-beta in Alzheimer's disease: progress, problems and perspectives. *Drug Discov Today* 11:931–938. doi:[10.1016/j.drudis.2006.08.004](https://doi.org/10.1016/j.drudis.2006.08.004)
- Wiltfang J, Lewczuk P, Riederer P, Grunblatt E, Hock C, Scheltens P, Hampel H, Vanderstichele H, Iqbal K, Galasko D, Lannfelt L, Otto M, Esselmann H, Henkel AW, Kornhuber J, Blennow K (2005) Consensus paper of the WFSBP Task Force on Biological Markers of Dementia: the role of CSF and blood analysis in the early and differential diagnosis of dementia. *World J Biol Psychiatry* 6:69–84. doi:[10.1080/15622970510029786](https://doi.org/10.1080/15622970510029786)
- Xu Y, Zhang W, Klaus J, Young J, Koerner I, Sheldahl LC, Hurn PD, Martinez-Murillo F, Alkayed NJ (2006) Role of cocaine- and amphetamine-regulated transcript in estradiol-mediated neuroprotection. *Proc Natl Acad Sci USA* 103:14489–14494. doi:[10.1073/pnas.0602932103](https://doi.org/10.1073/pnas.0602932103)
- Yasojima K, Akiyama H, McGeer EG, McGeer PL (2001) Reduced neprilysin in high plaque areas of Alzheimer brain: a possible relationship to deficient degradation of beta-amyloid peptide. *Neurosci Lett* 297:97–100. doi:[10.1016/S0304-3940\(00\)01675-X](https://doi.org/10.1016/S0304-3940(00)01675-X)
- Zhao J, Fu Y, Yasvoina M, Shao P, Hitt B, O'Connor T, Logan S, Maus E, Citron M, Berry R, Binder L, Vassar R (2007) Beta-site amyloid precursor protein cleaving enzyme 1 levels become elevated in neurons around amyloid plaques: implications for Alzheimer's disease pathogenesis. *J Neurosci* 27:3639–3649. doi:[10.1523/JNEUROSCI.4396-06.2007](https://doi.org/10.1523/JNEUROSCI.4396-06.2007)
- Zlokovic BV (2004) Clearing amyloid through the blood-brain barrier. *J Neurochem* 89:807–811. doi:[10.1111/j.1471-4159.2004.02385.x](https://doi.org/10.1111/j.1471-4159.2004.02385.x)