Published in final edited form as:

Org Lett. 2008 August 7; 10(15): 3247. doi:10.1021/ol801076w.

One-pot synthesis of Oligosaccharides by Combining Reductive Openings of Benzyldiene Acetals and Glycosylations

Yusuf Vohra, Mahalakshmi Vasan, Andre Venot, and Geert-Jan Boons*

Complex Carbohydrate Research Center, The University of Georgia, 315 Riverbend Road, Athens, GA 30602, USA

Abstract

Combining triflic acid promoted glycosylations of trichloroacetimidates with reductive opening of benzyldiene acetals with triflic acid and triethylsilane as one-pot procedures provides easy access to a wide range of di- and branched trisaccharides.

Protein- and lipid-bound saccharides are ubiquitous in biological systems involved in many important molecular processes such as fertilization, embryogenesis, neuronal development, hormone activities, and the proliferation of cells and their organization into specific tissues.^{1, 2} These interactions are also important in health science and are involved in the invasion and attachment of pathogens, inflammation, metastasis, blood group immunology, and xenotransplantation.³⁻⁵ A major obstacle to advances in glycobiology is the lack of pure and structurally well-defined carbohydrates and glycoconjugates. These compounds are often found in low concentrations and in micro-heterogeneous forms, greatly complicating their isolation and characterization. In many cases, well-defined oligosaccharides can only be obtained by organic synthesis.⁶⁻⁹

Although the methods for oligosaccharide synthesis have improved considerably during the past decade, the construction of complex carbohydrates remains a significant challenge due to the combined demands of elaborate procedures for glycosyl donor and acceptor preparation and the requirements of regio- and stereoselectivity in glycoside bond formation. To streamline the preparation of complex oligosaccharides, one-pot multi-step approaches for selective monosaccharide protection and oligosaccharide assembly are being pursued, which do not require intermediate work-up and purification step and hence speed-up the process of chemical synthesis considerably.^{10,11} For example, the observation that acetal formation, regioselective reductive opening of arylidene acetals, reductive etherification and acetylation can be catalyzed by triflic acid (TfOH) or trimethylsilyl triflate (TMSOTf) made it possible to program these

giboons@ccrc.uga.edu.

Supporting Information **Available** Experimental procedures and ¹H and ¹³C NMR spectra. This information is available free of charge via the Internet at <http://pubs.acs.org>.

reactions in tandem by sequential addition of reagents to give easy access to a wide variety of selective protected monosaccharide building blocks. Furthermore, many laboratories have demonstrated that chemoselective, orthogonal and iterative glycosylation strategies, which exploit differential reactivities of anomeric leaving groups, allow several selected glycosyl donors to react in a specific order resulting in a single oligosaccharide product.¹²⁻¹⁷

To further streamline the process of oligosaccharide assembly, we report here a strategy whereby a regioselective opening of a benzylidene acetal and glycosylations are combined in a one-pot multi-step synthetic procedure. The attraction of the approach is that it makes it possible to assemble branched oligosaccharides by a one-pot procedure; a task that cannot readily be accomplished by chemoselective, orthogonal and iterative glycosylations. In this respect, differential reactivities of hydroxyls have been exploited for the synthesis of branched oligosaccharides by one-pot procedures¹⁸⁻²⁴ but the scope of this approach is limited because of the need of exceptional high regioselectivities. Trichloroacetimidates were selected as glycosyl donors because their activation requires only catalytic TfOH or TMSOTf.²⁵ Furthermore, TfOH combined with triethylsilane (Et₃SiH) was employed for the opening of a 4,6-*O*-benzylidene acetals^{26,27} because these conditions provide in general high regioselectivities and excellent yields, and furthermore it was anticipated that these reaction conditions would be compatible with TfOH mediated glycosylations of trichloroacetimidates.

Thus, a mixture of trichloroacetimidate **1**²⁸ and benzylidene acetal protected glycosyl acceptor **4**²⁹ in dichloromethane at 0 °C was treated with a catalytic amount of TfOH. After a reaction time of 30 min., TLC and MALDI-MS analysis indicated consumption of the starting material and the formation of the expected disaccharide. The reaction mixture was cooled to -78 °C followed by addition of TfOH (1.8 eq.) and Et₃SiH (2.0 eq.). After stirring for 30 minutes at -78 °C, the reaction was quenched by the addition of triethylamine and methanol, and purification by silica gel column chromatography gave disaccharide **8** in a yield of 70%. As expected, only a β-galactoside was formed due to the presence of a 2,5-difluorobenzyl group³⁰ at C-2 of the galactosyl donor **1**.²⁸ This protecting group is an excellent neighboring group participant and unlike benzoyl and pivaloyl esters can be cleaved under mild conditions using a catalytic amount of sodium methoxide in methanol.

In the next set of experiments, the sequence of reactions was repeated, however, in this case the fucosyl donor **2**³¹ was employed instead of **1**. Fortunately, disaccharide **9** was isolated in an overall yield of 74% as a single regio- and stereoisomer demonstrating that the methodology is compatible with acid sensitive fucosides. It was also found that fucosylation of 2-azido-containing glycosyl acceptor **7**³² followed by a regioselective opening of the benzylidene acetal using standard conditions of the resulting disaccharide led to the facile formation of disaccharide **10**.

It has been established that the regioselective ring opening of 4,6-*O*-benzylidene acetals can be reversed to provide saccharides with a free C-6 hydroxyl by employing Cu(OTf)₂ in combination with borane³³ instead of TfOH and Et₃SiH. Furthermore, it was anticipated that trichloroacetimidates can be activated by Cu(OTf)₂ and thus it may be possible to glycosylate the C-3 hydroxyl of **4** followed by a regioselective opening of the benzylidene acetal of the resulting product to give a disaccharide having a C-6 hydroxyl. Indeed, a Cu(OTf)₂ (0.15 eq.) mediated glycosylation of glucosyl donor **1** with acceptor **4** gave, after a reaction time of 2h at room temperature, a disaccharide, which was treated with borane-THF complex to provide after stirring for an additional 4 h at ambient temperature, disaccharide **11** in a yield of 45%. In our efforts to improve the reaction yield, it was found that prolonged reaction times led to decomposition and formation of by-products.

Next, attention was focused on a reaction sequence whereby a regioselective benzylidene acetal opening is followed by glycosylation. Thus, Et₃SiH and TfOH were added to a cooled (-78 °C) solution of compound **5** in dichloromethane and after a reaction time of 30 min, glycosyl donor **1** or **2** was added followed by gradually raising the temperature to 0 °C over a period of 30 min. Standard work-up and purification by silica gel column chromatography gave regio- and stereo-isomerically pure disaccharides **12** and **13**, respectively, in yields of 72 and 70%.

Recently, we demonstrated that glycosylations with glycosyl donors modified at C-2 with a (*S*)-(phenylthiomethyl)benzyl moiety give exclusively α -anomeric selectivity due to neighboring group participation resulting in an intermediate *trans*-fused 1,2-sulfonium ion.^{34,35} To explore the possibility of a one-pot procedure involving benzylidene acetal opening and glycosylation with such a donor, compound **5**³⁶ was treated with TfOH and Et₃SiH followed by subsequent addition of **3**³⁴ and DTBMP to give **14** as only the α -anomer.

Next, we explored the possibility of synthesizing branched trisaccharides by a reaction sequence involving glycosylation, reductive benzylidene acetal opening followed by another glycosylation (Scheme 2). Thus, a mixture of trichloroacetimidate **2** and benzylidene protected glycosyl acceptor **4** in dichloromethane at 0 °C was treated with a catalytic amount of TfOH. After a reaction time of 30 min., the reaction mixture was cooled to -78 °C followed by addition of TfOH (1.8 eq.) and Et₃SiH (2.0 eq.) and after a further reaction time of 30 min, galactosyl donor **1** was added and the reaction mixture was allowed to warm to 0 °C over a period of 30 min, to give after standard work-up and purification by silica gel column chromatography, trisaccharide **15** in a yield of 63%. A similar procedure gave trisaccharide **16** by a glycosylation of **1** with **4**, followed by reductive opening of the benzylidene acetal and glycosylation of the resulting acceptor with fucosyl donor **2**. Furthermore, a TfOH promoted glycosylation of 2-azido-2-deoxyglucoside **7** with **2** followed by benzylidene acetal opening with TfOH/Et₃SiH gave a disaccharide acceptor, which was glycosylated with galactosyl donor **1** to give protected Lewis^x trisaccharide **17** in an excellent yield of 67%. Removal of anomeric TDS protecting group of **17** followed by conversion of the resulting lactol into leaving group provides an opportunity to prepare more complex oligosaccharides. Finally, it was demonstrated that the methodology can also be employed for the preparation of 2,4-branched trisaccharides by glycosylation of acceptor **6**³⁷ with fucosyl donor **2** to give a 2,4-linked disaccharide, which was subjected to Et₃SiH/TfOH to regioselectively open the benzylidene to afford a glycosyl acceptor having a C-4 hydroxyl. Addition of galactosyl donor **1** to the latter compound followed by standard work-up and purification by silica gel column chromatography gave trisaccharide **18** in 60% yield.

In conclusion, it has been demonstrated that one pot-procedures involving reductive opening of benzylidene acetals and glycosylations can give easy access to a wide range of di- and trisaccharides. It is to be expected that the utility of the methodology can be further extended by combining acid catalyzed reductive etherifications and acetylations with glycosylations. Also the use of a thioglycoside as the initial acceptor may provide an opportunity to prepare more complex structures by employing the resulting thioglycoside product as a glycosyl donor.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by a grant from the NIH NIGM (R01GM065248).

References

1. Kobata A. *Eur J Biochem* 1992;209:483–501. [PubMed: 1358608]
2. Varki A. *Cell* 2006;126:841–845. [PubMed: 16959563]
3. Dube DH, Bertozzi CR. *Nat Rev Drug Discov* 2005;4:477–488. [PubMed: 15931257]
4. Walsh G, Jefferis R. *Nature Biotech* 2006;24:1241–1252.
5. Opdenakker G, Rudd PM, Ponting CP, Dwek RA. *Faseb Journal* 1993;7:1330–1337. [PubMed: 8224606]
6. Davis BG. *J Chem Soc Perkin Trans* 1999;1:3215–3237.
7. Demchenko AV. *Synlett* 2003:1225–1240.
8. Boons GJ. *Contemp Org Synth* 1996;3:173–200.
9. Seeberger PH, Werz DB. *Nature* 2007;446:1046–51. [PubMed: 17460666]
10. Francais A, Urban D, Beau JM. *Angew Chem Int Ed* 2007;46:8662–8665.
11. Wang CC, Lee JC, Luo SY, Kulkarni SS, Huang YW, Lee CC, Chang KL, Hung SC. *Nature* 2007;446:896–899. [PubMed: 17443183]
12. Boons GJ. *Tetrahedron* 1996;52:1095–1121.
13. Codee JDC, Litjens R, van den Bos LJ, Overkleeft HS, van der Marel GA. *Chem Soc Rev* 2005;34:769–782. [PubMed: 16100617]
14. Wang YH, Ye XS, Zhang LH. *Org Biomol Chem* 2007;5:2189–2200. [PubMed: 17609746]
15. Wang YH, Zhang LH, Ye XS. *Combinatorial Chemistry & High Throughput Screening* 2006;9:63–75. [PubMed: 16454688]
16. Douglas NL, Ley SV, Lucking U, Warriner SL. *J Chem Soc Perkin Trans* 1998;1:51–65.
17. Koeller KM, Wong CH. *Chem Rev* 2000;100:4465–4493. [PubMed: 11749355]
18. Mong KKT, Wong CH. *Angew Chem Int Ed* 2002;41:4087–4090.
19. Takahashi T, Adachi M, Matsuda A, Doi T. *Tetrahedron Lett* 2000;41:2599–2603.
20. Tanaka H, Adachi M, Takahashi T. *Chem Eur J* 2005;11:849–862.
21. Valverde S, Garcia M, Gomez AM, Lopez JC. *Chem Comm* 2000:813–814.
22. Yamada H, Kato T, Takahashi T. *Tetrahedron Lett* 1999;40:4581–4584.
23. Yamada H, Harada T, Takahashi T. *J Am Chem Soc* 1994;116:7919–7920.
24. Yamada H, Harada T, Miyazaki H, Takahashi T. *Tetrahedron Lett* 1994;35:3979–3982.
25. Jung KH, Muller M, Schmidt RR. *Chem Rev* 2000;100:4423–4442. [PubMed: 11749353]
26. Wang CC, Lee JC, Luo SY, Fan HF, Pai CL, Yang WC, Lu LD, Hung SC. *Angew Chem Int Ed* 2002;41:2360–2362.
27. Sakagami M, Hamana H. *Tetrahedron Lett* 2000;41:5547–5551.
28. Cato D, Buskas T, Boons GJ. *J Carbohydr Chem* 2005;24:503–516.
29. Kim S, Chang H, Kim WJ. *J Org Chem* 1985;50:1751–1752.
30. Sjolín P, Kihlberg J. *J Org Chem* 2001;66:2957–2965. [PubMed: 11325260]
31. Rencurosi A, Lay L, Russo G, Caneva E, Poletti L. *J Org Chem* 2005;70:7765–7768. [PubMed: 16149811]
32. Muller T, Hummel G, Schmidt RR. *Liebigs Ann Chem* 1994:325–329.
33. Shie CR, Tzeng ZH, Kulkarni SS, Uang BJ, Hsu CY, Hung SC. *Angew Chem Int Ed* 2005;44:1665–1668.
34. Kim JH, Yang H, Park J, Boons GJ. *J Am Chem Soc* 2005;127:12090–12097. [PubMed: 16117550]
35. Kim JH, Yang H, Boons GJ. *Angew Chem Int Ed* 2005;44:947–949.
36. Ogawa T, Katano K, Matsui M. *Tetrahedron* 1980;36:2727–2733.
37. Methyl 4,6-O-benzylidene-3-O-benzyl- α -D-glucopyranoside (6): To a solution of methyl 4,6-benzylidene- α -D-glucopyranoside (50 mg, 0.18 mmol) in DCM (1.0 mL) was added benzaldehyde (22 μ L, 0.21 mmol) and triethylsilane (34 μ L, 0.21 mmol), and the mixture was stirred under an atmosphere of argon for 30 min. The mixture was cooled (-78°C) and TfOH (0.3 eq) was added. The reaction mixture was quenched after 30 min. with pyridine (50 μ L), diluted with DCM (5 mL), and washed with sat. aq. NaHCO_3 (3 mL) and brine (3 mL). The organic layer was dried (MgSO_4),

concentrated in vacuo and purified by silical gel column chromatography (hexane/ethylacetate, 4/1, v/v) to give 6 as a white solid in 50% yield.

Scheme 1.

One-pot Synthesis of Disaccharides by Glycosylation and Benzylidene Acetal Opening

Scheme 2.

One-pot Synthesis of Trisaccharides by Glycosylations and Benzylidene Acetal Opening