Published in final edited form as:

ACS Chem Biol. 2009 December 18; 4(12): 1039. doi:10.1021/cb9001848.

Selective Estrogen Receptor Modulator Delivery of Quinone Warheads to DNA Triggering Apoptosis in Breast Cancer Cells

Kuan-wei Peng, Huali Wang, Zhihui Qin, Gihani T. Wijewickrama, Meiling Lu, Zhican Wang, Judy L. Bolton, and Gregory R. J. Thatcher*

Department of Medicinal Chemistry and Pharmacognosy (M/C 781), College of Pharmacy, University of Illinois at Chicago, 833 South Wood Street, Chicago, Illinois 60612-7231

Abstract

Estrogen exposure is a risk factor for breast cancer and estrogen oxidative metabolites have been implicated in chemical carcinogenesis. Oxidation of the catechol metabolite of estrone (4-OHE) and the β -naphthohydroquinone metabolite of equilenin (4-OHEN) gives *o*-quinones that produce ROS and damage DNA by adduction and oxidation. To differentiate hormonal and chemical carcinogenesis pathways in estrogen receptor positive ER(+) cells, catechol or β -naphthohydroquinone warheads were conjugated to the selective estrogen receptor modulator (SERM) desmethylarrozifene (DMA). ER binding was retained in the DMA conjugates; both were antiestrogens with submicromolar potency in mammary and endometrial cells. Cytotoxicity, apoptosis, and caspase-3/7 activation were compared in ER(+) and ER(-)MDA-MB-231 cells, and production of ROS was detected using a fluorescent reporter. Comparison was made to DMA, isolated warheads, and a DMA-mustard. Conjugation of warheads to DMA increased cytotoxicity accompanied by induction of apoptosis and activation of caspase-3/7. Activation of caspase-3/7, induction of apoptosis, and cytotoxicity were all increased significantly in ER(+) cells for the DMA conjugates. ROS production was localized in the nucleus for conjugates in ER(+) cells. Observations are compatible with β -naphthohydroquinone and catechol groups being concentrated in the nucleus by ER binding, where oxidation and ROS production result, concomitant with caspase-dependent apoptosis. The results suggest DNA damage induced by catechol estrogen metabolites can be amplified in ER(+) cells independent of hormonal activity. The novel conjugation of quinone warheads to an ER-targeting SERM gives ER-dependent, enhanced apoptosis in mammary cancer cells of potential application in cancer therapy.

INTRODUCTION

Observations from clinical trials and epidemiological studies support the hypothesis that estrogen contributes to breast cancer and may be a causative agent (1,2). The large prospective Women's Health Initiative Study confirmed that the increased risk of breast cancer in postmenopausal women is strongly linked to exposure to estrogen replacement therapy (ERT) agents that contain conjugated endogenous and equine estrogens (3,4). The collective evidence supports the concept of: (1) an hormonal, proliferative, anti-apoptotic contribution to breast cancer from estrogens (5,6); together with (2), a contribution from chemical carcinogenesis resulting from the genotoxic and mutagenic actions of estrogen oxidative metabolites (7,8). In human breast tissue, DNA adducts of equine estrogens have been reported (9). The 4-hydroxy catechol metabolite (4-OHE) of endogenous estrogen oxidation by cytochrome P450 has been

*Corresponding author. Department of Medicinal Chemistry and Pharmacognosy, College of Pharmacy, University of Illinois at Chicago, 833 South Wood Street, Chicago, Illinois 60612-7231. Fax: 312 996 7107 thatcher@uic.edu.

Supporting Information Available: This material is available free of charge via the Internet.

proposed to be carcinogenic via formation of DNA-reactive, electrophilic *o*-quinones that also generate DNA-damaging, reactive oxygen species (ROS) (Figure 1a) (10–12).

The relative importance of contributions from hormonal carcinogenesis versus chemical carcinogenesis pathways remains unknown, as do the relative contributions to chemical carcinogenesis from DNA alkylation versus DNA oxidation. A further related question is the contribution of the equine estrogen oxidative metabolite, 4-hydroxyequilenin (4-OHEN), since autooxidation and facile redox cycling differentiates 4-OHEN from 4-OHE (Fig. 1a) (13,14). Autooxidation of 4-OHEN yields a naphthoquinone that induces DNA damage including single strand breaks, oxidized bases, apurinic sites, and formation of cyclic DNA adducts, both *in vitro* and *in vivo* (9,15–18). Furthermore, 4-OHEN rapidly induces DNA oxidation and DNA strand breaks with selectivity for ER(+) cells, compatible with a “Trojan horse” mechanism whereby estrogen receptor (ER) binds and concentrates 4-OHEN in the nucleus proximal to ER-binding DNA sequences (17).

Any approach to these unknowns is hindered by the difficulty of differentiating hormonal from chemical effects in ER(+) cells, which may be further confounded by the estrogenic regulation of cellular antioxidant systems (19). To circumvent this obstacle, a selective estrogen receptor modulator (SERM) was conjugated via a chemical linker to the reactive chemical moieties of 4-OHE and 4-OHEN, a catechol and a β -naphthohydroquinone, respectively (Figures 1b,c). Owing to the known autooxidation of 4-OHEN, both of these *o*-quinone-delivery warheads were protected as diacetates to facilitate cell permeation and enhance intracellular bioavailability. The delivery unit selected was desmethylarrozifene (DMA), a benzothiophene SERM with high ER binding affinity and well established antiestrogenic activity in ER(+) breast cancer cells (20,21). As depicted in figure 2, these quinone prodrug with ER delivery (QPED) conjugates are anticipated to reveal the capacity of *o*-quinone warheads to damage DNA selectively in ER(+) cells, in the absence of confounding estrogenic activity, and hence to provide information on potential contributions to chemical carcinogenesis of 4-OHE and 4-OHEN.

Since QPEDs are ER-targeted agents of DNA damage, they may represent lead compounds for selective chemotherapy of estrogen-sensitive tumors including breast, endometrial, and ovarian cancers. In addition to providing DNA damage, QPEDs are expected to retain the antiestrogenic activity of the parent SERM. To minimize side effects on healthy cells, chemotherapeutics can be targeted to molecular abnormalities needed for tumor growth and carcinogenesis. In breast cancer, such targets include growth factor receptors, and the ER: 75% of tumors in postmenopausal women and 50% in premenopausal women express ER (22,23). Traditional chemotherapeutics that damage DNA by adduct formation include platinum complexes and mustards, both of which have been conjugated to estrogen itself (23–25). Examples of quinones with anticancer activity that form DNA adducts are known; mostly derived from natural products, for example, β -lapachone (26,27).

To our knowledge this is the first report of synthetic conjugation of quinone warheads to an ER-targeting SERM leading to DNA damage. To provide comparison with the QPEDs, a third DMA conjugate was prepared containing a nitrogen mustard, in addition to both masked catechol warheads without DMA delivery units (Figure 1c). Antiestrogenic activity of QPEDs was retained, cytotoxicity was selective towards ER(+) cells via induction of apoptosis, and ROS production was observed to be amplified in the nucleus of ER(+) cells.

RESULTS AND DISCUSSION

ER binding and antiestrogenic activity

SERMs provide excellent scaffolds for bioconjugation, because the extended side chain that disrupts ER helix 12, responsible for antiestrogenic activity, extends to the extremity of the ER ligand binding site. DMA is an active metabolite of the preclinical benzothiophene SERM arzoxifene that is itself an improved version of the SERM raloxifene, used clinically for post-menopausal indications and breast cancer chemoprevention (28,29). DMA is a potent SERM: in the standard estradiol radioligand competitive binding assay, the measured IC_{50} for ER α and ER β is < 10 nM (21). Thus, extension of the 3-position side chain of DMA has been reported to produce conjugates that retain antiestrogenic activity; for example, conjugation of DMA with azide, a large biotin linker group, and a fluorophore linker group, yielded compounds with IC_{50} for ER α of 14.8, 9.4, and 28.7 nM, respectively (30,31). Hence a similar strategy was adopted for DMAcatDA, DMAnapDA, and DMA-mustard (Scheme 1). In the competitive binding assay, IC_{50} for the DMA-mustard was measured as 37 ± 11 nM and 67 ± 20 nM, for ER α and ER β respectively (Table 1).

Gene expression elicited by ligand binding to ER is dependent on receptor dimerization, formation of complexes with co-regulator proteins, and binding of these complexes to DNA. Therefore, cell based assays provide a more complete picture of estrogenic/antiestrogenic activity, in addition to revealing cell membrane permeation. Two human ER α (+) cell systems routinely used for profiling classical estrogenic activity are MCF-7 breast cancer cells transiently transfected with an ERE-luciferase reporter and the endometrial Ishikawa cell line that incorporates an endogenous estrogen sensitive alkaline phosphatase activity. For comparison, DMA-biotin and DMA-fluorophore conjugates demonstrated IC_{50} values of 9 nM and 29 nM in MCF-7 cells, and 7 nM and 23 nM in Ishikawa cells, respectively (32). Expectedly, estrogenic activity was not observed for QPEDs and these compounds antagonized the effects of estradiol (E_2 , 1 nM) in both cell systems (Table 1). Given the different cellular contexts and ER coregulators, relative and absolute potency can vary with different ligands. In MCF-7 cells, QPEDs were less potent antiestrogens with IC_{50} for DMAcatDA, DMAnapDA, and DMA-mustard measured as 129 nM, 502 nM, and 103 nM respectively; whereas in Ishikawa cells, IC_{50} for DMAcatDA, DMAnapDA, and DMA-mustard was 10.3, 101, and 17.6 nM, respectively (Figure 3; Table 1). Tethering of a catechol diacetate (catDA) or a β -naphthohydroquinone diacetate (napDA) warhead to the DMA SERM scaffold reduced antiestrogenic activity relative to DMA, but the QPEDs retained submicromolar potency in cell culture. The control compounds, catDA and napDA, did not show any antiestrogenic activity at 1 μ M.

Cytotoxicity in ER(+) versus ER(-) cells

In order to examine the selective toxicity of QPEDs toward ER(+) cells compared to ER(-) cells, cytotoxicity was evaluated by the sulforhodamine B (SRB) colorimetric assay, a widely used method for in vitro cytotoxicity screening using cell density determination based on the measurement of cellular protein content (33). Although comparisons are often made between two breast cancer cell lines, ER(+) MCF-7 cells and ER(-) MDA-MB-231 cells, the comparison of ER(+) S30 cells with MDA-MB-231 cells provides an identical cellular background, since S30 cells are MDA-MB-231 cells stably transfected with wild type ER α . The most dramatic observation of cytotoxicity towards these human mammary epithelial cancer cells was in the comparison of catDA with DMAcatDA (Figures 4a,c): the catechol warhead showed no significant toxicity at the concentrations studied (≤ 20 μ M), whereas the corresponding QPED was cytotoxic; in ER(+) cells DMAcatDA was a relatively potent cytotoxin with $LC_{50} = 1.6$ μ M. In contrast, the napDA warhead was cytotoxic in S30 cells, and DMAnapDA was only a fourfold more potent cytotoxin than the napDA (Figure 4a; Table 1).

Both QPEDs were equally potent cytotoxins in S30 cells. DMA itself was cytotoxic at higher concentrations; however, no cytotoxicity was observed for the DMA-mustard below solubility limits.

DMA was toxic at higher concentrations comparable to those reported previously for benzothiophene SERMs; however, cytotoxicity in breast cancer cells was not dependent on ER status (Figure 4a; Table 1). Similarly, LC₅₀ values obtained for napDA were also independent of ER status (Figure 4c; Table 1). However, the cytotoxicity of both QPEDs was enhanced in ER(+) S30 cells relative to the ER(−) background MDA-MB-231 cells; DMAcatDA was 5-fold more toxic, and DMAnapDA toxicity was enhanced 2.5-fold.

Localization of cellular ROS production

The cell permeant fluorescence reporter, CM-H₂DCFDA, was used to identify and localize ROS generation in S30 and MDA-MB-231 cells. The same method was reported previously to demonstrate the rapid generation of ROS by 4-OHEN, which was observed to be localized in the nucleus of the ER(+) cell line, but not localized to a cell compartment in MDA-MB-231 cells (17). CM-H₂DCF is sensitive to oxidation by oxidizing oxygen radicals including hydroxyl radicals formed from Fenton reactions of H₂O₂ (34), and is useful for localization of these cellular ROS, in combination with nuclear dyes, using confocal fluorescence microscopy. In ER(−) MDA-MB-231 cells, after incubation for 1 h with QPEDs and the catechol warheads, a general increase in cellular DCF fluorescence was observed, but this was not localized to the nucleus (Figure 5b). This experiment was designed to observe ROS localization, rather than quantify ROS production, therefore it is unsafe to speculate on relative amounts of ROS from each agent. However, the same experiment carried out in ER(+) S30 cells definitively demonstrated selective increases in DCF fluorescence in the nuclei of QPED treated cells (Figure 5a). In these cells, ROS detected from treatment with napDA and catDA were not localized, but ROS production from DMAnapDA and DMAcatDA was clearly localized in the nuclei as shown by overlay of images with stained nuclei. Curiously, cells treated with DMAcatDA produced a more intense localization within nuclear organelles, possibly nucleoli. These observations clearly support the ability of the ER to bind and concentrate QPEDs in the cell nucleus, leading to enhanced ROS generation in the nucleus.

Induction of apoptosis

Oxidative stress is implicated in numerous cellular events including apoptosis, providing a cytotoxic pathway that has been proposed to be triggered by numerous chemotherapeutic agents, including ROS-generating flavonoids, such as the experimental anticancer agent aminoflavone (35). Since QPEDs were shown to have enhanced cytotoxicity in ER(+) cells and to produce ROS selectively in the nucleus of these cells, it was reasonable to investigate apoptosis as the cause of cell death. In apoptotic cells, the membrane phospholipid phosphatidylserine (PS) is translocated from the inner to the outer surface of the plasma membrane; exposed PS binds to the phospholipid-binding protein annexin V, providing a reliable method for quantifying the percentage of cells undergoing apoptosis using flow cytometry. Cells were analyzed in this way after treatment with compound (5 μM) or vehicle for 18 h (Figure 6a). A higher concentration of napDA (9 μM, corresponding to LC₅₀) was also studied since this compound was cytotoxic at higher concentration in both cell types. The data was analyzed to quantify the total apoptotic cells (viable and non-viable) as a percentage of cell count (Figure 6b). An excellent correlation was observed between percentage cells in apoptosis and cytotoxicity for treatment groups, in that all compounds cytotoxic at 5 or 9 μM induced apoptosis. More importantly, napDA induced apoptosis with no dependence on cellular ER status, whereas both QPEDs induced significantly more apoptosis in ER(+) cells compared to ER(−) cells: DMAcatDA and DMAnapDA induced 48% and 24% apoptosis, respectively, in S30 cells compared to 18% and 10% apoptosis in MDA-MB-231 cells. The

control compounds, catDA and napDA at 5 μM showed no significant apoptosis; however, napDA at 9 μM induced apoptosis significantly in both ER α (+) and ER(−) cells.

Induction of caspase-3/7 activity

Induction of cellular oxidative stress can lead to activation of caspase 3 and 7 as mediators of apoptotic cell death (36). Oxidative DNA damage is one trigger for caspase activation, although not the only mechanism by which oxidative stress can induce apoptosis. Caspase-3/7 activity was measured using Z-DEVD-aminoluciferin as substrate after incubation of cells with compounds under the same conditions used for detection of apoptosis (Figure 6c). After 18 h incubation, caspase activity for all QPED treated cells was observed to be significantly elevated. DMAcatDA and DMA-napDA (5 μM) induced increases of 9-fold and 5-fold in caspase activity, respectively, in S30 cells, compared to 7-fold and 4-fold activation in MDA-MB-231 cells, respectively. To confirm these results, the caspase-3/7 selective inhibitor, Ac-DEVD-CHO, was shown to block activation by both QPEDs. Interestingly at the single time point studied, napDA (9 μM) induced apoptosis, but did not show induction of caspase-3/7 activity. Importantly, caspase-3/7 activation by both QPEDs was significantly greater in ER(+) cells compared to ER(−) cells.

QPEDs as chemotherapeutic agents

The traditional DNA alkylating agents, platinum complexes and nitrogen mustards, form DNA cross-links and are applied widely in chemotherapy. In part to improve upon the activity of cisplatin in breast cancer, both platins and mustards have been conjugated to estrogens to exploit ER affinity to broaden and enhance chemotherapeutic activity (37). A number of groups have tethered platin warheads at the 3, 16, and 17 positions of the steroid backbone of estradiol/estrone, showing weaker antiproliferative activity in MCF-7 cells than parent platin and/or a lack of selective cytotoxicity between ER(+) and ER(−) cells (38,39). For example, a recent study by Berubé, varying linker length in such conjugates, reported for the most potent compound, IC₅₀ values of 21 μM and 14 μM in MCF-7 and MDA-MB-231 cells, respectively, compared to 18 μM for cisplatin itself (40). The same group has linked a platin to a benzopyran SERM scaffold yielding interesting conjugates with highly variable toxicity, but no selectivity for MCF-7 cells over MDA-MB-231 cell killing. Essigmann and Croy have reported conjugates in which the 7 α position of estradiol is tethered to ethylenediamine platinum dichloride (Pt(en)Cl₂) or a nitrogen mustard, retaining ER binding affinity (23–25). The most recent estradiol-(Pt(en)Cl₂) conjugate reported by this group showed modest selectivity against MCF-7 (LC₅₀ ~ 12 μM) versus MDA-MB-231 cells (LC₅₀ ~ 16 μM), and more impressively the conjugate was able to kill ER(+) human ovarian cancer cells (LC₅₀ ~ 25 μM) that did not respond to a control warhead (25). Reported selective cytotoxicity for the estradiol-7 α -mustard conjugate was 2.6-fold higher in MCF-7 cells compared to MDA-MB-231 cells (23,24).

Three models have been proposed to justify the development of platin and mustard conjugates as nuclear receptor targeted chemotherapeutic agents. Firstly, the preferential uptake of these conjugates by the nuclear receptor and sequestration in the case of ER in ER-positive cells and tissues, enhancing ER-positive cancer selectivity. Secondly, the formation of DNA adducts that entrap the large nuclear receptor complex is proposed to physically block DNA repair enzyme function. Thirdly, “hijacking” of the nuclear receptor might prevent receptor mediated transcription, which in the case of ER would require very high affinity conjugates to compete with endogenous estrogens. The QPEDs target and bind the ER carrying a payload of masked *o*-quinone warheads that would be anticipated to give DNA-adducts *in simile* with the known capacity of 4-OHE, 4-OHEN, and other *o*-quinone forming agents to covalently modify DNA nucleobases and form stable adducts (16,41,42). The relative contributions of DNA oxidation versus DNA adduction towards the cytotoxic actions QPED in breast cancer cells remains to be determined. However, the observations herein show that QPEDs are selective cytotoxins

towards ER(+) cells and in these cells, selectively generate ROS in the nucleus. The potency of QPEDs and the selectivity towards ER(+) cells both compare favorably with platin and mustard conjugates described above which are in development as chemotherapeutic agents.

Mechanism of action

Both QPEDs studied were cytotoxic towards human breast cancer cells in which these agents induced production of ROS, apoptosis, and executor caspase activity. These observations support a mechanism whereby induction of cellular oxidative stress by QPEDs induced apoptotic cell death mediated by caspase activation, a pathway for which there is substantial precedent. QPEDs deliver a quinone warhead after oxidative bioactivation of the catechol that is itself unmasked by non-specific esterase bioactivation. Protection of the warhead as a catechol diacetate both enhances membrane permeation and prevents quinone formation in the cell culture medium, which would lead to reduced intracellular bioavailability. Several well studied anti-infective and anticancer agents are quinones that induce apoptosis; these are considered to act as DNA alkylating agents, and also to generate semiquinones and ROS that can react with DNA, in addition to depleting glutathione, and perturbing cellular energy and calcium homeostasis (43). The cytotoxicity of such quinones in hypoxic cells and a general requirement for reductive bioactivation has led to the suggestion that DNA alkylation is the dominant contributor (43,44). Nevertheless, the mechanism by which the quinone anticancer agents induce apoptosis is not fully defined and may involve triggering of mitochondrial pathways or gene regulation of pro-apoptotic signals; one study on a series of such compounds in MCF-7 cells reported that cytotoxicity correlated with induction of apoptosis and elevation of Bcl-xL/Bcl-2-associated death protein (BAD) (45).

Cellular GSH depletion represents one mechanism by which quinone anticancer agents may induce apoptosis. Total cellular GSH was measured in ER(+) and ER(-) cells as a function of time, after incubation with QPEDs and control compounds, by a kinetic assay using glutathione reductase/NADPH as a GSSG reducing system in concert with 5,5'-dithiobis(2-nitrobenzoic acid). The pro-apoptotic QPEDs and napDA were observed significantly to deplete cellular GSH by at least 50% after 2 h incubation, whereas catDA gave less than 20% depletion at this time point, and no significant depletion was observed for DMA and catDA (all agents 5 μ M; Supplemental Figure 1). Furthermore, GSH depletion measured for all compounds at the 2 h timepoint showed concentration dependence (Supplemental Figure 2). The observed GSH depletion correlated qualitatively with apoptosis, however, no dependence was observed on cellular ER status. That is: (i) efficacy and potency for GSH depletion by QPEDs were not significantly different in S30 versus MDA-MB-231 cells; and (ii), DMA napDA was not significantly more potent than napDA in either cell line. Therefore, whereas GSH depletion may contribute to apoptosis, it is not the mechanism that mediates selective apoptosis of ER (+) cells.

The differences in activity of the monocyclic quinone producing agents, catDA and DMAcatDA are most dramatic. Oxidative bioactivation of DMAcatDA may involve chelation of chromatin-bound Cu ions by the catechol, however, further studies are needed, in particular to vary the QPED linker length. Nevertheless, the results presented herein do validate the QPED design concept and encourage further research towards ER-targeted anticancer agents.

Implications for estrogen carcinogenesis

Evidence supporting estrogens as mammary carcinogens derives from clinical trials and epidemiological studies (1–3). In the hamster kidney tumor model, a variety of estrogens, including estrone and equilenin were full carcinogens (4). Both hormonal and chemical carcinogenesis pathways are expected to contribute to cancer in estrogen-responsive tissues (5–8,46). Despite the significant differences in the oxidative metabolism of human and equine

estrogens, there has been sparse attention to the consequences of these differences for cancer in women. The extended conjugation of the naphtho-*o*-quinone of 4-OHEN provides stabilization of the oxidized form in comparison to the benzo-*o*-quinone of 4-OHE. Since 4-OHE and 4-OHEN are both estrogenic, separating hormonal from chemical pathways in ER(+) cells requires development of compounds such as the QPEDs. ER-dependent activity was observed only for the QPEDs; not for any constituent parts, neither DMA nor warheads. ROS production in the cell nucleus, activation of caspase-3/7, induction of apoptosis, and cytotoxicity were all increased significantly in ER(+) cells. This observation indicates that DNA damage induced by catechol estrogen metabolites can be amplified in ER(+) cells independent of hormonal activity. Further studies will examine differences between the types of DNA damage induced by the β -naphthoquinone and benzoquinone groups of QPEDs, which model the reactive functional groups of 4-OHEN and 4-OHE, respectively.

METHODS

Synthesis

Synthetic details and characterization are described in Supporting information.

Transfection & ERE-luciferase assays

Cells were cultured in estrogen-free media for 4 days before transfection. Cells were transfected with 2 μ g of the pERE-luciferase plasmid, which contains three copies of the *Xenopus laevis* vitellogenin A2 ERE upstream of fire fly luciferase (a gift from Dr. V. C. Jordan). To normalize transfection efficiency, pRL-TK plasmid (1 μ g, Promega) was co-transfected. Cells (5×10^6) in serum-free media were transfected by electroporation in a 0.4-cm cuvette (Bio-Rad Laboratories) at a voltage of 0.320 kV and a high capacitance of 950 μ F in a GenePulser X-cell (Bio-Rad Laboratories). The cells were resuspended in estrogen-free media, transferred to 12-well plates immediately after electroporation, and incubated overnight. The cells were treated with the appropriate compounds with 1 nM E₂ for 18 h and ERE activation measured as described previously (47). Briefly, the luciferase activities in cell lysates were measured using Dual Luciferase Assay system (Promega) with a FLUOstar OPTIMA (BMG LABTECH) and data were calculated as relative luciferase activity which is the firefly luciferase reading divided by the *Renilla* luciferase reading.

Antiestrogenic activity in Ishikawa cells

The procedure of Pisha and Pezzuto was used as described previously (47). Briefly, Ishikawa cells (5×10^4 cells/mL) were incubated overnight with estrogen free media in 96 well plates. Test samples (with 1 nM 17 β -estradiol) and appropriate controls were added. The day 0 control did not contain any additional estradiol. The cells were incubated with a total volume of 200 μ L/well at 37 °C for 4 days. The cells were washed three times with PBS and lysed by freeze-thawing in the presence of 0.1 M Tris, pH 8.0. Enzyme activity was measured by reading the liberation of *p*-nitrophenol from 1 μ M *p*-nitrophenylphosphate at 340 nm every 15 s for 16–20 readings with an ELISA reader (Bio-Tek Instrument). The maximum slope of the lines generated by the kinetic readings were calculated using a Kinecalc computer program (Bio-Tek Instrument). For antiestrogenic activity, the reduction in percent induction as compared to the DMSO control was determined as follows: [(slope sample – slope cells)/(slope DMSO-slope cells)] $\times$ 100.

Cytotoxicity assay

Cells were plated (7×10^4 cells/mL) in 96 well plates. The following day, cells were treated with the compound for 18 h. After the incubation period, cells were fixed to the plastic substratum by the addition of cold 50% aqueous trichloroacetic acid. The plates were incubated

at 4°C for 1 h, washed with H₂O, and air-dried. The trichloroacetic acid-fixed cells were stained by the addition of 0.4% (w/v) SRB, dissolved in 1% acetic acid for 30 min. Free SRB solution was removed by washing with 1% aqueous acetic acid. The plates were air-dried, and the bound dye was solubilized by the addition of 10 mM unbuffered Tris base, pH 10. The plates were placed on a shaker for 5 min, and the absorption was determined at 515 nm. Finally, the absorbance obtained with each of the treatment procedures was averaged and was expressed as a percentage, relative to the 0 h control (33).

Localization of ROS by CM-H₂DCFDA oxidation

Both S30 and MDA-MB-231 cells (4.5×10^4 cells/well) were grown on a sterile Nunc™ chambered cover glass and incubated for 48 h at 37 °C with 5% CO₂ in phenol red free-MEME medium supplemented with 10% stripped FBS medium. These cells were then treated with 1 μM QPEDs, control compounds, or 0.5% DMSO for 1 h. After treating for 30 min, S30 and MDA-MB-231 cells were labeled with 10 μM CM-H₂DCFDA for 30 min at 37°C with 5% CO₂. CM-H₂DCFDA treated cells were rinsed with phosphate buffered saline (PBS) to remove the unincorporated dye and 10 mg/ml Hoechst-33258 was added to the cells to see nuclear staining. Imaging was performed with a Zeiss LSM510 laser-scanning confocal microscope with the detector gain adjusted to eliminate the background autofluorescence. The fluorescence signal from CM-DCF was monitored with a 488 nm argon/krypton laser and a 530 nm band pass filter. Hoechst nuclear staining signal was monitored with a 345nm UV laser and 420nm band pass filter. Water immersion objective (A × 63-1.2 numerical aperture) was used for all experiments. Images were analyzed using the analysis tool provided in Zeiss biophysical software package.

Apoptosis analysis

S30 and MDA-MB-231 cells (5×10^5) were seeded in 6-well plates for 24 h before treatment. Cells were treated with DMSO, QPEDs or control compounds for 18 h. Cell death was measured by flow cytometry using the Annexin-PE Apoptosis Detection Kit I (BD Bioscience) according to the manufacturer's protocol. Cells were washed twice with cold PBS and then suspended in 1× binding buffer. Cells were stained using 5 μL PE annexin V (for apoptotic status) and 5 μL 7-amino-actinomycin (7-AAD; for cell viability status) for 15 min at RT in the dark. After adding 200 μL of 1× binding buffer, cells were analyzed by flow cytometry within 1 h. Annexin V-positive cells were scored as total apoptotic cells as a percentage of all cells.

Measurement of caspase-3/7 activity

The activities of caspase-3/7 in S30 cells were measured as recommended by the kit manufacturer (Promega). Briefly, the DMSO vehicle or compound treated cells (7×10^4 cells/mL) were incubated in a 96-well plate for 18 h at 37 °C and then incubated with 100 μL Caspase-Glo® 3/7 reagent or with a combination of caspase-3/7 substrate and 10 μM inhibitor (Ac-DEVD-CHO) at room temperature for 30 min. The luminescence of each well was measured at using a FLUOstar OPTIMA (BMG LABTECH).

Determination of GSH in Cells

Experimental details and two figures presenting data observed are provided in Supporting information.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Abbreviations

Arzoxifene	[[6-hydroxy-3-[4-[2-(1-piperidiny)-ethoxy]phenoxy]-2-(4-methoxyphenyl)]benzo[b]thiophene)
catDA	catechol diacetate
CM-H ₂ DCFDA	5-(and-6)-chloromethyl-2',7'-dichloro-dihydrofluorescein diacetate
DMA	desmethylarzoxifene
DMAcatDA	3-(3-(2-(1-(2-(4-(6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophen-3-yloxy)phenoxy)ethyl)piperidin-4-yl)ethylamino)-3-oxopropyl)-1,2-phenylene diacetate
DMA-mustard	4-(4-(bis(2-chloroethyl) amino)phenyl)-N-(2-(1-(2-(4-(6-hydroxy-2-(4-hydroxyphenyl)benzo[b]thiophen-3-yloxy) phenoxy)ethyl)piperidin-4-yl)ethyl)butanamide
DMAnapDA	7-(3-(2-(1-(2-(4-(6-acetoxy-2-(4-acetoxyphenyl)benzo[b]thiophen-3-yloxy)phenoxy)ethyl)piperidin-4-yl)ethylamino)-3-oxopropyl)naphthalene-1,2-diyl diacetate
ER	estrogen receptor
ERT	estrogen replacement therapy
ERE	estrogen response element
4-OHE	4-hydroxyestrone
4-OHEN	4-hydroxyequilenin
napDA	β-naphthohydroquinone diacetate
QPED	quinone prodrug with ER delivery
SERM	selective estrogen receptor modulator
SRB	sulforhodamine B.

Acknowledgments

This work was supported by National Institutes of Health (NIH) Grants CA130037 and CA102590. Ping Yao and Johann Sohn are thanked for technical assistance with estrogenic assays.

REFERENCES

1. Colditz GA, Hankinson SE, Hunter DJ, Willett WC, Manson JE, Stampfer MJ, Hennekens C, Rosner B, Speizer FE. The use of estrogens and progestins and the risk of breast cancer in postmenopausal women. *N. Engl. J. Med* 1995;332:1589–1593. [PubMed: 7753136]
2. Key T, Appleby P, Barnes I, Reeves G. Endogenous sex hormones and breast cancer in postmenopausal women: reanalysis of nine prospective studies. *J. Natl. Cancer Inst* 2002;94:606–616. [PubMed: 11959894]
3. Rossouw JE, Anderson GL, Prentice RL, LaCroix AZ, Kooperberg C, Stefanick ML, Jackson RD, Beresford SA, Howard BV, Johnson KC, Kotchen JM, Ockene J. Risks and benefits of estrogen plus progestin in healthy postmenopausal women: principal results from the Women's Health Initiative randomized controlled trial. *Jama* 2002;288:321–333. [PubMed: 12117397]
4. Li JJ, Li SA, Oberley TD, Parsons JA. Carcinogenic activities of various steroidal and nonsteroidal estrogens in the hamster kidney: relation to hormonal activity and cell proliferation. *Cancer Res* 1995;55:4347–4351. [PubMed: 7671246]

5. Henderson BE, Feigelson HS. Hormonal carcinogenesis. *Carcinogenesis* 2000;21:427–433. [PubMed: 10688862]
6. Yager JD, Davidson NE. Estrogen carcinogenesis in breast cancer. *N. Engl. J. Med* 2006;354:270–282. [PubMed: 16421368]
7. Bolton JL, Thatcher GRJ. Potential mechanisms of estrogen quinone carcinogenesis. *Chem. Res. Toxicol* 2008;21:93–101. [PubMed: 18052105]
8. Gaikwad NW, Yang L, Muti P, Meza JL, Pruthi S, Ingle JN, Rogan EG, Cavalieri EL. The molecular etiology of breast cancer: evidence from biomarkers of risk. *Int. J. Cancer* 2008;122:1949–1957. [PubMed: 18098283]
9. Embrechts J, Lemiere F, Van Dongen W, Esmans EL, Buytaert P, Van Marck E, Kockx M, Makar A. Detection of estrogen DNA-adducts in human breast tumor tissue and healthy tissue by combined nano LC-nano ES tandem mass spectrometry. *J Am Soc Mass Spectrom* 2003;14:482–491. [PubMed: 12745217]
10. Liehr JG. Is estradiol a genotoxic mutagenic carcinogen? *Endocr. Rev* 2000;21:40–54. [PubMed: 10696569]
11. Bolton JL, Pisha E, Zhang F, Qiu S. Role of quinoids in estrogen carcinogenesis. *Chem. Res. Toxicol* 1998;11:1113–1127. [PubMed: 9778307]
12. Cavalieri EL, Stack DE, Devanesan PD, Todorovic R, Dwivedy I, Higginbotham S, Johansson SL, Patil KD, Gross ML, Gooden JK, Ramanathan R, Cerny RL, Rogan EG. Molecular origin of cancer: catechol estrogen-3,4-quinones as endogenous tumor initiators. *Proc. Natl. Acad. Sci. U S A* 1997;94:10937–10942. [PubMed: 9380738]
13. Zhang F, Chen Y, Pisha E, Shen L, Xiong Y, van Breemen RB, Bolton JL. The major metabolite of equilin, 4-hydroxyequilin, autoxidizes to an *o*-quinone which isomerizes to the potent cytotoxin 4-hydroxyequilenin-*o*-quinone. *Chem. Res. Toxicol* 1999;12:204–213. [PubMed: 10027800]
14. Shen L, Pisha E, Huang Z, Pezzuto J, Krol E, Alam Z, van Breemen R, Bolton J. Bioreductive activation of catechol estrogen-ortho-quinones: aromatization of the B ring in 4-hydroxyequilenin markedly alters quinoid formation and reactivity. *Carcinogenesis* 1997;18:1093–1101. [PubMed: 9163701]
15. Okamoto Y, Chou PH, Kim SY, Suzuki N, Laxmi YR, Okamoto K, Liu X, Matsuda T, Shibutani S. Oxidative DNA damage in XPC-knockout and its wild mice treated with equine estrogen. *Chem. Res. Toxicol* 2008;21:1120–1124. [PubMed: 18447394]
16. Wang Z, Edirisinghe P, Sohn J, Qin Z, Geacintov NE, Thatcher GRJ, Bolton JL. Development of a liquid chromatography electrospray ionization tandem mass spectrometry method for analysis of stable 4-hydroxyequilenin-DNA adducts in human breast cancer cells. *Chem Res Toxicol* 2009;22:1129–1136. [PubMed: 19368368]
17. Wang Z, Wijewickrama GT, Peng KW, Dietz BM, Yuan L, van Breemen RB, Bolton JL, Thatcher GRJ. Estrogen Receptor {alpha} Enhances the Rate of Oxidative DNA Damage by Targeting an Equine Estrogen Catechol Metabolite to the Nucleus. *J Biol Chem* 2009;284:8633–8642. [PubMed: 19158089]
18. Zhang F, Swanson SM, van Breemen RB, Liu X, Yang Y, Gu C, Bolton JL. Equine estrogen metabolite 4-hydroxyequilenin induces DNA damage in the rat mammary tissues: formation of single-strand breaks, apurinic sites, stable adducts, and oxidized bases. *Chem. Res. Toxicol* 2001;14:1654–1659. [PubMed: 11743748]
19. Mobley JA, Brueggemeier RW. Estrogen receptor-mediated regulation of oxidative stress and DNA damage in breast cancer. *Carcinogenesis* 2004;25:3–9. [PubMed: 14514655]
20. Yu B, Dietz BM, Dunlap T, Kastrati I, Lantvit DD, Overk CR, Yao P, Qin Z, Bolton JL, Thatcher GRJ. Structural modulation of reactivity/activity in design of improved benzothiophene selective estrogen receptor modulators: induction of chemopreventive mechanisms. *Mol Cancer Ther* 2007;6:2418–2428. [PubMed: 17876041]
21. Overk CR, Peng KW, Asghodom RT, Kastrati I, Lantvit DD, Qin Z, Frasor J, Bolton JL, Thatcher GRJ. Structure-activity relationships for a family of benzothiophene selective estrogen receptor modulators including raloxifene and arzoxifene. *ChemMedChem* 2007;2:1520–1526. [PubMed: 17654759]

22. Jordan VC. Chemoprevention of breast cancer with selective oestrogen-receptor modulators. *Nat Rev Cancer* 2007;7:46–53. [PubMed: 17186017]
23. Mitra K, Marquis JC, Hillier SM, Rye PT, Zayas B, Lee AS, Essigmann JM, Croy RG. A rationally designed genotoxin that selectively destroys estrogen receptor-positive breast cancer cells. *J Am Chem Soc* 2002;124:1862–1863. [PubMed: 11866593]
24. Sharma U, Marquis JC, Nicole Dinaut A, Hillier SM, Fedeles B, Rye PT, Essigmann JM, Croy RG. Design, synthesis, and evaluation of estradiol-linked genotoxicants as anti-cancer agents. *Bioorg Med Chem Lett* 2004;14:3829–3833. [PubMed: 15203171]
25. Kim E, Rye PT, Essigmann JM, Croy RG. A bifunctional platinum(II) antitumor agent that forms DNA adducts with affinity for the estrogen receptor. *J Inorg Biochem* 2009;103:256–261. [PubMed: 19054566]
26. McDonald RW, Bunjobpon W, Liu T, Fessler S, Pardo OE, Freer IK, Glaser M, Seckl MJ, Robins DJ. Synthesis and anticancer activity of nordihydroguaiaretic acid (NDGA) and analogues. *Anticancer Drug Des* 2001;16:261–270. [PubMed: 12375879]
27. Pardee AB, Zou Zhi Ling AB, Li CJ. Cancer Therapy with beta-Lapachone. *Current Cancer Drug Targets* 2002;2:227. [PubMed: 12188909]
28. Liu H, Bolton JL, Thatcher GRJ. Chemical modification modulates estrogenic activity, oxidative reactivity, and metabolic stability in 4'-F-DMA, a new benzothiophene selective estrogen receptor modulator. *Chem Res Toxicol* 2006;19:779–787. [PubMed: 16780356]
29. Qin Z, Kastrati I, Ashgdom RT, Lantvit DD, Overk CR, Choi Y, van Breemen RB, Bolton JL, Thatcher GRJ. Structural modulation of oxidative metabolism in design of improved benzothiophene selective estrogen receptor modulators. *Drug Metab Dispos* 2009;37:161–169. [PubMed: 18936111]
30. Yu B, Qin Z, Wijewickrama GT, Edirisinghe P, Bolton JL, Thatcher GR. Comparative methods for analysis of protein covalent modification by electrophilic quinoids formed from xenobiotics. *Bioconjug Chem* 2009;20:728–741. [PubMed: 19301905]
31. Liu H, Qin Z, Thatcher GRJ, Bolton JL. Uterine peroxidase-catalyzed formation of diquinone methides from the selective estrogen receptor modulators raloxifene and desmethylated arzoxifene. *Chem Res Toxicol* 2007;20:1676–1684. [PubMed: 17630709]
32. Yu B, Qin Z, Wijewickrama GT, Edirisinghe P, Bolton JL, Thatcher GRJ. Comparative methods for analysis of protein covalent modification by electrophilic quinoids formed from xenobiotics. *Bioconjug Chem* 2009;20:728–741. [PubMed: 19301905]
33. Skehan P, Storeng R, Scudiero D, Monks A, McMahon J, Vistica D, Warren JT, Bokesch H, Kenney S, Boyd MR. New colorimetric cytotoxicity assay for anticancer-drug screening. *J Natl Cancer Inst* 1990;82:1107–1112. [PubMed: 2359136]
34. LeBel CP, Ischiropoulos H, Bondy SC. Evaluation of the probe 2',7'-dichlorofluorescein as an indicator of reactive oxygen species formation and oxidative stress. *Chem Res Toxicol* 1992;5:227–231. [PubMed: 1322737]
35. McLean L, Soto U, Agama K, Francis J, Jimenez R, Pommier Y, Sowers L, Brantley E. Aminoflavone induces oxidative DNA damage and reactive oxidative species-mediated apoptosis in breast cancer cells. *Int J Cancer* 2008;122:1665–1674. [PubMed: 18059023]
36. Ueda S, Masutani H, Nakamura H, Tanaka T, Ueno M, Yodoi J. Redox control of cell death. *Antioxid Redox Signal* 2002;4:405–414. [PubMed: 12215208]
37. Cepeda V, Fuertes MA, Castilla J, Alonso C, Quevedo C, Perez JM. Biochemical mechanisms of cisplatin cytotoxicity. *Current Medicinal Chemistry - Anti-Cancer Agents* 2007;7:3–18.
38. Gandolfi O, Blum J, Mandelbaum-Shavit F. Antitumor steroidal-cis-platinum(II)-o-catecholato conjugates: preliminary evaluation on breast cancer MCF-7 cells. *Inorganica Chimica Acta* 1984;91:257–261.
39. Jackson A, Davis J, Pither RJ, Rodger A, Hannon MJ. Estrogen-derived steroidal metal complexes: agents for cellular delivery of metal centers to estrogen receptor-positive cells. *Inorg Chem* 2001;40:3964–3973. [PubMed: 11466055]
40. Provencher-Mandeville J, Descoteaux C, Mandal SK, Leblanc V, Asselin E, Berube G. Synthesis of 17beta-estradiol-platinum(II) hybrid molecules showing cytotoxic activity on breast cancer cell lines. *Bioorg Med Chem Lett* 2008;18:2282–2287. [PubMed: 18356047]
41. Liehr JG. Genotoxic effects of estrogens. *Mutat Res* 1990;238:269–276. [PubMed: 2160607]

42. Bolton JL, Thatcher GRJ. Potential mechanisms of estrogen quinone carcinogenesis. *Chem Res Toxicol* 2008;21:93–101. [PubMed: 18052105]
43. Begleiter A. Clinical applications of quinone-containing alkylating agents. *Front Biosci* 2000;5:E153–E171. [PubMed: 11056078]
44. Gutierrez PL. The role of NAD(P)H oxidoreductase (DT-Diaphorase) in the bioactivation of quinone-containing antitumor agents: a review. *Free Radic Biol Med* 2000;29:263–275. [PubMed: 11035255]
45. Tudor G, Gutierrez P, Aguilera-Gutierrez A, Sausville EA. Cytotoxicity and apoptosis of benzoquinones: redox cycling, cytochrome c release, and BAD protein expression. *Biochem Pharmacol* 2003;65:1061–1075. [PubMed: 12663042]
46. Nandi S, Guzman R, Yang J. Hormones and mammary carcinogenesis in mice, rats, and humans: a unifying hypothesis. *Proc. Natl. Acad. Sci. U S A* 1995;92:3650–3657. [PubMed: 7731959]
47. Overk CR, Peng KW, Asghodom RT, Kastrati I, Lantvit DD, Qin Z, Frasor J, Bolton JL, Thatcher GR. Structure-activity relationships for a family of benzothiophene selective estrogen receptor modulators including raloxifene and arzoxifene. *ChemMedChem* 2007;2:1520–1526. [PubMed: 17654759]

Figure 1.

a) Figurative scheme of ER-targeting, DMA delivery system bound to ER α ; the R group is the payload containing the linker and warhead; the warhead requires bioactivation by (i) non-specific esterase activity, and (ii) oxidation to an *o*-quinone. b) The human estrogen oxidative metabolite 4-OHE and the equine estrogen oxidative metabolite 4-OHEN are further oxidized to *o*-quinones generating superoxide and ROS. 4-OHEN autoxidizes and redox cycles, whereas 4-OHE requires metal ion or oxidase catalysis of oxidation. (c) Detailed chemical structures of study compounds: NapDA is the diacetate masked warhead of DMAnapDA, which mimics 4-OHEN; whereas catDA is the masked warhead of DMAcatDA, which mimics 4-OHE.

Figure 2.

The DMAcatDA and DMAnapDA molecules are designed as quinone prodrugs with ER delivery (QPEDs). QPEDs are anticipated to be unmasked by esterase activity in the cytoplasm, to bind to ERα, and to be concentrated in the nucleus proximal to ERE sequences of DNA. Oxidation of the unmasked catechol to a quinone will lead to ROS generation (detected by DCF fluorescence), redox cycling, and DNA damage via electrophilic (covalent) and oxidative mechanisms. The evidence presented herein, shows that the consequence is apoptosis via caspase activation. R = H, Ac.

Figure 3.

Antiestrogenic activity of QPEDs compared to DMA-mustard and DMA in MCF-7 and Ishikawa cells (blue squares DMAAnapDA; red circles DMAcatDA; green inverted triangles DMA-mustard; black triangles DMA). a) Concentration-response curves for inhibition of estrogen-induced ERE-luciferase activity in MCF-7 cells. Cells were transfected with ERE-luciferase plasmid for 24 h, and then treated with different concentrations of drug in the presence of E_2 (1nM) for 18 h. b) Concentration-response curve for inhibition of estrogen-induced alkaline phosphatase activity in Ishikawa cells. Cells were seeded in 96-well plates overnight in estrogen-free medium prior to treatment with compounds in the presence of E_2 (1

nM) for 4 d. Data showing mean and s.e.m. for three separate cultures was fitted to a sigmoidal curve.

Figure 4.

ER(+) selective cytotoxicity of QPEDs in breast cancer cells. a) Comparison in ER(+) S30 cells of DMAAnapDA (blue squares); napDA (purple squares); DMAcatDA (red circles); catDA (orange circles), DMA (black triangles); DMA-mustard green inverted triangles; blue arrow shows increased toxicity of napDA conjugated to DMA; red arrow shows enhanced toxicity of catDA when conjugated to DMA. b) Comparison of DMAAnapDA (blue squares); napDA (purple squares); DMA (black triangles); S30 cells (solid symbols); ER(-) MDA-MB231 cells (open symbols); the blue arrow shows left shift of DMAAnapDA toxicity in ER(+) versus ER(-) cells. c) Comparison of DMAcatDA (red circles); catDA (orange circles), DMA (black triangles); S30 cells (solid symbols); MDA-MB231 cells (open symbols); the red arrow shows

left shift of DMAcatDA toxicity in ER(+) versus ER(−) cells. Cells were seeded in 96 well plates for 24 h then treated with compounds for 18 h before cell count was measured as a percentage from the ratio of cells in compound-treated versus DMSO-treated wells using SRB assay. Data show mean and s.e.m. for three separate cultures.

Figure 5. Nuclear localization of CM-DCF fluorescence after treatment of a) S30 and b) MDA-MB231 cells with QPEDs and control compounds. Cells were incubated with compounds for 1h and followed by treatment with H_2DCFDA and Hoechst nuclear dye for 10 min before confocal fluorescence microscopy images were taken. CM-DCF fluorescence is indicative of ROS production and overlay with nuclear staining shows localization of ROS production; arrows indicate co-localization of ROS and nuclear staining. Images are representative of several experiments.

Figure 6.

Apoptosis and caspase activity induced by QPEDs and control compounds in human breast cancer cells. a) Analysis of S30 cells and MDA-MB231 cells treated with DMSO vehicle or compounds (5 μ M or as indicated) for 18h by flow cytometry (FACS) using annexin V labeling. Data shown are representative results. b) Annexin V positive cells were scored as apoptotic and expressed as percentage of total cells: S30 cells (solid columns); MDA-MB231 cells (striped columns). c) Caspase 3/7 activity induced in S30 (solid) and MDA-MB231 (striped columns) after treatment with DMSO vehicle or compounds (5 μ M unless stated) for 18 h, followed by 30 min incubation with caspase-3/7 substrate (Z-DEVD-aminoluciferin) with or without caspase-3/7 inhibitor (INH, Ac-DEVE-CHO, 10 μ M). Data presented as fold induction compared to DMSO. In (B) and (C), data show mean and s.d. of three independent experiments analyzed by ANOVA with Newman-Keuls post test (***) $p < 0.001$; * $p < 0.05$; ** $p < 0.01$ vs. DMSO control).

Table 1

Antiestrogenic and cytotoxic potency of QPEDs and controls in cell culture.

	ER α -binding IC ₅₀ (nM)	MCF-7 cell antiestrogenic IC ₅₀ (nM)	Ishikawa cell antiestrogenic IC ₅₀ (nM)	S30 cell cytotoxicity LC ₅₀ (μ M)	MDA-MB-231 cytotoxicity LC ₅₀ (μ M)
DMA	7.8 $\pm$ 1.9	1.1 $\pm$ 0.1	0.1 $\pm$ 0.1	23.0 $\pm$ 0.8	23.5 $\pm$ 0.5
DMAcatDA	N.D.	129 $\pm$ 23	10.3 $\pm$ 1.6	1.6 $\pm$ 0.5	7.8 $\pm$ 0.3
DMAanapDA	N.D.	502 $\pm$ 95	101 $\pm$ 5.9	3.2 $\pm$ 0.7	8.2 $\pm$ 2.5
DMA-mustard	37 $\pm$ 11	103 $\pm$ 8	17.6 $\pm$ 2.6	> 40	> 40
catDA	N.D.	> 10,000	> 10,000	> 40	> 40
napDA	N.D.	> 10,000	> 10,000	8.8 $\pm$ 1.4	8.9 $\pm$ 0.9
chlorambucil	N.D.	> 10,000	> 10,000	> 40	> 40

Data show means and s.d. N.D. not determined.