Published in final edited form as:

Brain Res. 2008 October 15; 1235: 153. doi:10.1016/j.brainres.2008.06.053.

Uncovering genes for cognitive (dys)function and predisposition for alcoholism spectrum disorders: A review of human brain oscillations as effective endophenotypes

Madhavi Rangaswamy^{*} and **Bernice Porjesz**

Henri Begleiter Neurodynamics Laboratory, Department of Psychiatry and Behavioral Sciences, SUNY Downstate Medical Center

Abstract

Brain oscillations provide a rich source of potentially useful endophenotypes (intermediate phenotypes) for psychiatric genetics, as they represent important correlates of human information processing and are associated with fundamental processes from perception to cognition. These oscillations are highly heritable, are modulated by genes controlling neurotransmitters in the brain, and provide links to associative and integrative brain functions. These endophenotypes represent traits that are less complex and more proximal to gene function than either diagnostic labels or traditional cognitive measures, providing a powerful strategy in searching for genes in psychiatric disorders. These intermediate phenotypes identify both affected and unaffected members of an affected family, including offspring at risk, providing a more direct connection with underlying biological vulnerability. Our group has utilized heritable neurophysiological features (i.e., brain oscillations) as endophenotypes, making it possible to identify susceptibility genes that may be difficult to detect with diagnosis alone. We have discussed our findings of significant linkage and association between brain oscillations and genes in GABAergic, cholinergic and glutamatergic systems (*GABRA2*, *CHRM2*, and *GRM8*). We have also shown that some oscillatory indices from both resting and active cognitive states have revealed a common subset of genetic foci that are shared with the diagnosis of alcoholism and related disorders. Implications of our findings have been discussed in the context of physiological and pharmacological studies on receptor function. These findings underscore the utility of quantitative neurophysiological endophenotypes in the study of the genetics of brain function and the genetic diathesis underlying complex psychiatric disorders.

Keywords

Brain Oscillations; Genetics; EEG; Endophenotypes; *CHRM2*; *GABRA2*; Theta; Beta

1. Introduction

An important focus of electrophysiological studies is to understand how large neuronal assemblies organize and function in the brain and the possible mechanisms for cooperation

^{*}Correspondence and reprint requests should be addressed to: Madhavi Rangaswamy, Ph.D., Henri Begleiter Neurodynamics Laboratory, SUNY Downstate Medical Center, Box 1203, 450 Clarkson Ave., Brooklyn, NY 11203, U.S.A., Phone: 718 270 2024; Fax: 718 270 4081, Madhavi.rangaswamy@downstate.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

that form the basis for all sensory, cognitive, and motor processes in health and disease. Not only do electrophysiological indices provide a means to delineate the millisecond-by-millisecond structure and progression of information processing in the brain, but they also provide vital clues to points of aberration that may underlie a predisposition or vulnerability, especially to psychiatric conditions, and the extent of damage caused by the disease progression. Brain regions generally display a combination of delta (1–3 Hz), theta (4–7 Hz), alpha (8–12 Hz), beta (13–30 Hz), and gamma (> 30 Hz) rhythms, as has also been seen in corticothalamic network operations (Buzsaki and Draguhn, 2004; Steriade, 2001; Steriade, 2006). Furthermore, several rhythms can coexist in the same area or interact among different structures, and adjacent frequency bands within the same neuronal network are associated with different brain states (Steriade, 2001; Varela et al., 2001). The challenge of refining the exact combination of time and frequency elements corresponding to specific brain states is aided by new methods of analyses, such as the Continuous Wavelet Transform (Grossman and Morlet, 1984), Matching Pursuits (Mallat, 1993), and most recently the S-Transform (Stockwell, 1996), among others.

Traditional physiological models of brain function inherent in the event-related potential (ERP) methodologies often conceptualize electrophysiological spontaneous activity as internal noise (Tomberg and Desmedt, 1999). Functional neuroimaging studies have shown how the baseline of spontaneous neuronal activity is organized in multiple highly specific functional anatomical networks, known as resting state networks (Damoiseaux et al., 2006; De Luca et al., 2006). The hemodynamic changes in these resting state networks have been shown to fluctuate at frequencies between 0.01 and 0.1 Hz, and strongly overlap with sensory–motor and cognitive networks that are commonly modulated during active behavioral tasks (Corbetta et al., 2000; Fox et al., 2005; Fox et al., 2006; Mazoyer et al., 2001). Hence, recent models propose that spontaneous activity may play an important functional role by providing constraints or opportunity that could be important to communication in sensory, cognitive, or motor activity networks (Engel et al., 2001; Llinas, 1988; Mantini et al., 2007; Varela et al., 2001).

Until recently, ERPs have been the basic electrophysiological indices of cognition and have provided valuable insights into human brain processes. There is now a substantial literature which suggests that some ERP features arise from oscillatory changes due to sensory and/or cognitive processes which influence the dynamics of ongoing electroencephalographic (EEG) rhythms of different frequency bands (Basar-Eroglu and Basar, 1991; Basar-Eroglu et al., 2001; Demiralp et al., 2001; Karakas et al., 2000a; Schurmann et al., 1995; Schurmann et al., 2001; Yordanova and Kolev, 1996). Sensory, cognitive and motor activities are associated with synchronization and enhancement of EEG activity, where random resting EEG oscillations become organized, amplified and coupled, thus giving rise to an ‘evoked’ (phase-locked) or ‘induced’ (non-phase-locked) rhythmicity. Event-related oscillations (EROs) have been proposed to arise in part from superpositioning or from a process-related ‘partial-phase resetting’ occurring in different EEG frequency bands in response to sensory or cognitive stimulation (Basar, 1980; Basar, 1999; Makeig et al., 2002). These oscillations thus provide links to associative and integrative brain functions. Specific frequency rhythms of oscillatory responses have been attributed to underlie various cognitive processes, as follows: delta, signal detection and decision-making (Basar et al., 1999; Schurmann et al., 2001); theta, conscious awareness, recognition memory, and episodic retrieval (Basar et al., 2001b; Doppelmayr et al., 1998; Gevins et al., 1998; Klimesch et al., 2001; Klimesch et al., 1994); slow alpha, attribution of attentional resources (Basar et al., 1997; Klimesch, 1997; Klimesch et al., 1998); fast alpha, semantic memory and stimulus processing (Klimesch, 1997; Klimesch et al., 1997a; Klimesch et al., 1997b; Klimesch et al., 1994); beta and gamma, sensory integrative processes (Basar-Eroglu et al., 1996a; Basar-Eroglu et al., 1996b; Basar et al., 2001a; Schurmann et al., 1997). The more recent models evaluating brain function using oscillations stress the importance of

phase reorganization of these task-relevant oscillations in the understanding of cognitive brain function (see review, Klimesch et al., 2007).

The two new trends in cognitive neuroscience discussed in this section make it possible to study neural network dynamics in the human brain in health and disease; they have therefore, strongly contributed to the study of predisposition and brain dysfunction in psychiatric populations (Banaschewski and Brandeis, 2007; Ford et al., 2007; Herrmann and Demiralp, 2005; Porjesz et al., 2005; van der Stelt and Belger, 2007). In the following sections, we will elaborate on the utility of studying human brain oscillations in the search for genes involved in one of the many psychiatric conditions – alcoholism – that falls in the spectrum of addictive disorders. Recent evidence also suggests that alcohol dependence is part of a spectrum of disinhibitory disorders, which include externalizing and substance use disorders (Kendler et al., 2003). We will highlight the key oscillatory indices that have emerged as significant discriminators in the context of alcoholism, and will then provide a brief overview of the current genetic analysis methods used to trace the path from EEG and EROs to genes (Figure 1). Finally, we will discuss the genes in the context of their physiological significance and the underlying genetic diathesis for alcoholism and related disorders.

2. Exploring key brain oscillations in alcoholism

The concept of “intrinsic” rather than resting activity contributing a large proportion of the measured activity is gaining prominence (Raichle et al., 2001; Raichle and Snyder, 2007). Hence, functional connectivity in the brain during these “rest” periods is likely to reflect the maintenance of dynamic equilibrium of the “intrinsic” networks (Llinas et al., 1999). Power estimates of EEG activity in different frequency bands have been demonstrated to corresponding to arousal and activity levels of the cerebral cortex (Niedermayer and Lopes Da Silva, 1999). Measures of synchronicity, such as coherence, offer a means to study the functional relation between populations of neurons/regions. EEG coherence is a measure of phase correlation (Nunez and Srinivasan, 2006; Srinivasan et al., 1998) and reflects functional cortical connectivity, possibly arising from corticocortical or corticothalamic fiber networks. Functional connectivity impairments have been reported in other psychiatric and neurological conditions, such as attention deficit hyperactivity disorder (ADHD) (Barry et al., 2002; Clarke et al., 2007), autism (Murias et al., 2007) and schizophrenia (Higashima et al., 2007; Higashima et al., 2006). In our strategy to find genes, we have focused on brain oscillations that differentiate between alcoholics and controls as well as between high risk offspring of alcoholics and low risk offspring of nonalcoholics (Table 1). In the following sections we will discuss some key findings, both current and published, pertaining to two brain states – resting and cognitive challenge (visual oddball) that have been extensively studied in our laboratory and as part of the extended Collaborative study on the Genetics of Alcoholism (COGA), with respect to understanding risk and predisposition to alcoholism and the related spectrum disorders.

2.1. EEG beta oscillations: Power

The signatures of the resting state of the brain as determined by electroencephalography (EEG) have revealed characteristic patterns in individuals with alcoholism and also those with high risk for developing alcoholism and related spectrum of externalizing conditions (Porjesz et al., 2005), review). Across most studies published in this field, increased beta power in the EEG has emerged as one such important feature; it is noted in affected (Bauer, 2001; Costa and Bauer, 1997; Propping et al., 1981; Rangaswamy et al., 2002; Winterer et al., 1998) and high risk individuals (Finn and Justus, 1999; Gabrielli et al., 1982; Pollock et al., 1995; Rangaswamy et al., 2002; Rangaswamy et al., 2004b). The low voltage alpha (LVA) EEG variant has been shown to be associated with a subtype of alcoholism that is related to anxiety disorder (Enoch et al., 1999). It has been hypothesized that altered norepinephrine levels on thalamic activity

may partly explain the connection between LVA and anxiety disorders in alcoholic women (Enoch et al., 2003).

Increased resting beta power has been reported at frontal leads in those who also have a diagnosis of antisocial personality disorder (ASP) (Bauer and Hesselbrock, 1993). Also, the beta power increase may be associated with increased vulnerability, as female high risk subjects with a larger number of affected first-degree relatives displayed significantly elevated beta power compared to those with just one affected parent (Rangaswamy et al., 2002). Thus evidence of elevated beta power provides strong support to the excitation-inhibition imbalance model proposed to underlie the predisposition to alcohol dependence (Begleiter and Porjesz, 1999).

2.2. EEG theta oscillations: Synchrony

There is evidence in the literature that not only do alcoholics manifest differences in EEG power in specific frequency bands, but they also manifest increased interhemispheric coherence (Kaplan et al., 1985; Michael et al., 1993). It has been reported that bilateral intrahemispheric coherences in alpha and beta frequency bands were increased in both long-term abstinent and non-abstinent alcoholics compared to controls (Winterer et al., 2003a). These findings were strongest for the alpha2 (10.5–12 Hz) frequency band, and were most pronounced at temporal, parietal and occipital regions, particularly when depressiveness was included as a covariate; there was no effect of length of abstinence on these findings.

Most published studies on coherence have employed monopolar pairs in the analyses, a configuration that has a considerable amount of spatial smearing. Using bipolar electrode pairs effectively results in a low resolution spatial filter that controls this issue (Cook et al., 1998), and thus the coherence measure provides a better estimate of localized activity and possibly enhances the contributions of corticothalamic and interhemispheric fiber connections. In our laboratory we have observed significant increases in resting EEG interhemispheric high theta (6–7 Hz) coherence, using the bipolar montages, in alcohol-dependent subjects when compared to normal controls (Figure 2). Peak theta band coherence was highest for posterior electrode pairs in alcohol dependent subjects, displaying a shift from a more frontocentral prominence as seen in normal control subjects. Hence increased EEG coherence (cortical synchronization) may serve as an endophenotype (i.e. biological marker) for alcoholism.

2.3. Event-related theta and delta oscillations: Power

One of the most consistent robust findings in the literature is the reduced P3 amplitude in alcoholics and in offspring at risk prior to alcohol exposure (Begleiter et al., 1984; Porjesz et al., 2005). The low P3 amplitudes coupled with the weaker and less well-organized sources in alcoholics and offspring at risk suggest inefficient allocation of resources during neural processing. This undifferentiated neurophysiological pattern suggests a level of cortical disinhibition in alcoholics and individuals at risk.

The low P3 amplitude is not only observed in abstinent alcoholics and offspring of alcoholics, but is also present in various disinhibitory conditions, such as substance abuse, ASP, conduct disorder and ADHD. Moreover, individuals with low P3 amplitudes manifest a significantly higher incidence of externalizing disorders and disinhibitory traits than those with high P3 amplitudes (Porjesz et al., 2005), review).

Several studies have demonstrated that P3 responses are primarily the outcome of theta and delta oscillations elicited during cognitive processing of stimuli (Basar-Eroglu et al., 1992; Basar et al., 1999; Karakas et al., 2000a; Karakas et al., 2000b; Yordanova and Kolev, 1996). Topographically, delta oscillation power peaks at the posterior region, while the theta power

peak is located in the fronto-central region (Karakas et al., 2000b); theta oscillations also contribute strongly to N2 components.

In a visual oddball paradigm, alcoholics manifest significantly less evoked theta and delta ERO amplitudes while processing the target stimuli (Jones et al., 2006b); these findings are most significant anteriorly for theta, and posteriorly for delta (Figure 3). In order to determine whether these deficits in theta and delta oscillations antecede the development of alcoholism we examined high-risk children of alcoholics using the same paradigm (Rangaswamy et al., 2007). The two groups in the study compared children of alcoholics and normal children in the age range of 14–17 years. These results show that the adolescent offspring of alcoholics have reduced delta and theta band ERO amplitude (underlying P3) while processing the target stimuli compared to controls. The differences were most prominent centro-parietally for theta, and parietally for delta. Interestingly, the ERO measures were superior to P3 amplitude in differentiating between high-risk and low-risk offspring. Hence, the results of these two studies indicate that decreased theta and delta EROs to target stimuli may antecede the development of alcoholism and represent a strong trait marker (Figure 3).

We have also reported theta and delta oscillatory deficits in alcoholics and high risk subjects during the performance of tasks designed to assess response inhibition (a Go-NoGo task) and error/response evaluation (a gambling task), providing more evidence that these deficient oscillatory responses may antecede the development of alcoholism. A significant reduction in delta and theta band ERO amplitude was concentrated in the frontal region for alcoholics, particularly during No-Go processing in a Go/No-Go paradigm (Kamarajan et al., 2004); in the high risk offspring the significant differences were more profound and showed a global decrease in theta and delta power (Kamarajan et al., 2006).

2.4. Event-related gamma oscillations: Power

The early phase-locked gamma has been considered to represent an important processing step related to the selection and identification of target stimuli, indicative of a top-down mechanism involved in selective attention (Basar-Eroglu et al., 1996b; Fell et al., 2003). This phase-locked gamma is larger to attended stimuli compared to unattended stimuli, particularly over frontal regions (Basar et al., 1999; Yordanova et al., 2001). Neuro-imaging studies using attentional tasks have implicated the role of fronto-parietal networks in this top-down control of selective attention (Corbetta et al., 2000; Giesbrecht et al., 2003). Some studies have found associations between stimulus induced amplitude modulations and reaction time and speed (Frund et al., 2007; Haig et al., 1999). Induced gamma oscillations have been associated with a wide range of cognitive processes and learning (Basar-Eroglu et al., 1996b; Gruber and Muller, 2006; Tallon-Baudry et al., 1999).

Our group (Padmanabhapillai et al., 2006a) has recently found that abstinent alcoholics manifest significantly less early (1–150 ms) gamma band response (28–45 Hz) in the frontal region during target processing in a visual oddball task than controls. Control subjects had significantly more gamma while processing the target when compared to their processing of the non-target stimulus. However, the alcoholics had an opposite profile, eliciting more gamma band response to non-target than to target stimuli. We have suggested that the reduction in early evoked frontal gamma band response to targets may be associated with frontal lobe dysfunction associated with a deficient top-down processing mechanism in alcoholics. Early phase-locked gamma band activity in the children of alcoholics and non-alcoholics was investigated using the same visual oddball task (Padmanabhapillai et al., 2006b). Subjects were in the age range of 14–17 and high risk children came from families who had at least one alcoholic parent. Topographically, peak gamma band response was observed in the frontal region for the target stimuli in both groups. High risk children had significantly lower gamma activity than control children, especially in the parietal regions for the target condition.

Additionally, the offspring of alcoholics showed less differentiation between target and non-target stimuli in the parietal region compared to controls, indicating difficulty in early stimulus processing. It is likely that a dysfunctional frontoparietal attentional network may underlie this deficit. There is clear evidence of deficient activations in inferior frontal and inferior parietal brain regions while high risk subjects performed the visual oddball task in an fMRI study (Rangaswamy et al., 2004a).

3. Exploring genetics of key brain oscillations and their application to alcoholism and related disorders

Alcoholism is a common, complex (non-Mendelian) disorder with contributions from both genetic and environmental influences and their interactions. As psychiatric diagnosis is dichotomous (an individual is either affected or unaffected), it is difficult to use diagnosis as the sole phenotype when studying the genetics of complex disorders, such as alcoholism. It has been suggested that, ideally, molecular genetic studies should not be performed on psychiatric diagnoses alone, which reflect distal and variable effects of genes, but on quantitative neurobiological measures or markers that reflect more proximal effects of genes involved in the genetic predisposition for developing psychiatric disorders (Tsuang and Faraone, 2000). These quantitative biological markers (endophenotypes or intermediate phenotypes) serve as covariates that correlate with the main trait of interest (diagnosis) and serve to better define that trait or its underlying genetic mechanism (Gottesman and Gould, 2003; Gottesman and Shields, 1973; Gottesman, 1972). The advantages of using quantitative neurobiological measures of risk as endophenotypes in the search for genes involved in complex disorders are that they are closer to gene action involved in the predisposition for the disorder, they are genetically simpler than clinical endpoints, and quantitative traits provide more power to localize and characterize disease susceptibility genes (Almasy, 2003; Dick et al., 2006c).

Recent evidence suggests that alcohol dependence is part of a spectrum of disinhibitory disorders, which include externalizing and substance use disorders. Many of the same genetic risk factors underlie these disinhibitory co-occurring disorders and can be explained by a small number of common underlying genetic factors (Kendler et al., 2003). These findings are in keeping with the electrophysiological findings described above (Sections 2.1 and 2.3) indicating a similar electrophysiological profile in these related disinhibitory disorders.

The identification of suitable quantitative biologic markers that are genetically transmitted could explicate the genetic factors involved in the etiology of alcoholism and related disorders, and also might elucidate the potential nature of the genetic factors. Brain function is likely to be involved in a genetic predisposition to develop alcoholism and other psychiatric disorders, and neuroelectric events may serve as excellent biological markers or endophenotypes. Understanding genetic control of brain electrical activity may provide clues about cerebral function, and may shed light on pathogenic mechanisms involved in neurological and psychiatric disorders, where impairment in brain electrical activity is apparent. Brain oscillations provide a rich source of potentially useful endophenotypes for psychiatric genetics, as they represent important correlates of human information processing and cognition.

In order to be considered as an endophenotype, there are certain criteria that must be met, the most important of which is that the trait must be heritable (Porjesz et al., 2005). The data on the heritability of EEG frequencies are quite compelling. High concordance rates in the spectral characteristics of resting eyes-closed EEG have been reported from monozygotic twin pairs compared to dizygotic twin pairs. A large twin study indicates that power in all frequency bands of the resting EEG is highly heritable: delta 76%, theta 89%, alpha 89%, and beta 86% (van Beijsterveldt and Boomsma, 1994; van Beijsterveldt et al., 1996) and EEG coherence has also

been reported to be heritable, with estimates between 50% and 70% in twin populations (Stassen et al., 1988; van Baal et al., 1998; van Beijsterveldt and Boomsma, 1994; van Beijsterveldt et al., 1998) and estimates between 40% to 60% for coherences in theta and alpha band in posterior regions reported for a sib-pair population (Chorlian et al., 2007). In order for genetic studies to be successful, it is extremely important to select a well-characterized (endo) phenotype. Therefore, genetic studies of neuroelectric endophenotypes (brain oscillations, such as EEG and those EROs underlying ERPs, including the P3 component) provide a powerful strategy for identifying underlying susceptibility genes for developing alcoholism and related disinhibitory disorders.

3.1. Exploring the gene targets: Methods

The possible approaches for finding genes associated with the variation in quantitative traits (e.g. brain oscillations) in complex diseases (e.g. psychiatric disease) fall into two main categories: (a) genome-wide studies, which include linkage studies and the more recent genome-wide association studies (GWAS), and (b) candidate gene studies, which are mostly association studies. Genome-wide studies do not require any hypotheses about gene function, while candidate gene studies are based on biological hypotheses. Both of these approaches have been successful in finding genes, and often, a successful strategy is to use both methods together; yet each has both advantages and drawbacks (For a review of these approaches see (Hirschhorn and Daly, 2005).

The most traditional method used to find genes is genome-wide linkage analysis. Linkage studies assess whether a polymorphic genetic marker from a chromosomal region can be linked to a specific trait (e.g. brain oscillation) within a family. Thus linkage analysis determines patterns of DNA sharing among family members that correlates with the phenotypic trait (Almasy and Blangero, 1998). This method serves to indicate chromosomal sharing within families, and precisely which alleles are shared can be different across families. The advantage of this approach is that the entire genome can be scanned without any prior hypotheses, and it is often successful in finding rare genes. Linkage is measured by the Logarithm of Odds (LOD) score, where a LOD score over 3 is taken as evidence of significant linkage (equivalent to p-value of 0.0001). Linkage analysis techniques have been successfully utilized to locate genes of rare Mendelian traits, where a single gene accounts for a disease. However, linkage may also be detected when multiple genes lie in the same chromosomal region, giving rise to a broad linkage peak with several candidate genes underlying the peak (see Figures 1 and 5). While linkage methods have been less successful in locating genes associated with more common complex traits (e.g. psychiatric diagnosis), which are likely the result of multiple small gene effects, studies from our laboratory (COGA) have successfully used these methods to identify genes associated with quantitative traits (electrophysiological) that have often led to genes involved in psychiatric disease. As will be seen in this review, we have successfully used this approach with brain oscillations as endophenotypes, to target chromosomal regions using linkage that contain potential candidate genes that may be involved in both brain function and diagnosis of alcohol dependence and related disorders, which is then more closely investigated using association analyses (Figure 1).

Rather than scan the entire genome, association studies test the strength of the relationship between the variants of a specific candidate gene and a trait, such as brain oscillations. Publicly available databases are used to identify single-nucleotide polymorphisms (SNPs) within and flanking candidate genes under regions of the chromosome with significant linkage peaks. The COGA strategy has sought to include SNPs in all regions, and is not just restricted to coding regions or exons. Association studies need not be family based, and can be accomplished by comparing alleles of a particular variant between cases and controls. It is important to begin the search with a set of well-characterized heritable biological quantitative measures, such as

brain oscillations, and some well-defined functional candidate gene. The selection of a candidate gene can be made in two ways: subsequent to a significant linkage finding, or determined on the basis of a priori biological facts associated with the trait under study. In the first approach, linkage analysis will identify a chromosomal region that will include a variety of potential candidate genes that may influence the trait; in the second approach, the candidates would be selected based on biological hypotheses. For example, in the case of brain oscillations, likely candidate genes could be regulatory genes that control neurochemical processes in the brain, and hence neural functions that are likely to influence brain oscillations. As will be illustrated, both linkage and association methods have successfully been used to find genes involved in brain oscillations as well as complex disorders for which they are endophenotypes. A positive association between an allele and a specific oscillation may indicate that the allele is a causative factor in the phenotype. In association studies, one allele is being tested, but it is quite possible that other nearby alleles are fully enriched in one of the tested genotype groups. This phenomenon is called linkage disequilibrium (LD). A given allele may serve as a proxy for other nearby alleles that have moved together across generations on the same chromosome.

It should be cautioned that a positive association between an allele and an oscillation may also reflect artifacts that arise from genotyping errors, as well as multiple testing. Another common source of artifact is the problem of population stratification. This refers to potential ethnic group difference in allelic frequency due primarily to effects of population of origin and geographical isolation which concentrate certain alleles in different populations. Therefore it is important to keep in mind the problem of population stratification in conducting association studies. Various methods have been implemented to deal with population stratification in conducting association studies, such as using family-based controls, or limiting analyses to ethnically homogeneous groups of individuals (e.g. only Caucasians). As will be seen in our studies, we have used all of these approaches to take these issues into account as much as possible (i.e. all association studies are performed only on Caucasians (which is the majority of our sample), and we use family based association studies (FBAT), as well as controlling for multiple comparisons using statistical methods such as false discovery rate (FDR). Since the influence of any single gene on a given oscillation is likely to be small, we must also keep in mind other potential confounding factors such as gender, age, alcohol or drug abuse, psychopathology, etc.

With the completion of the human genome and the availability of SNPs and whole genome association studies (GWAS), the possibility exists of identifying specific genes that affect human cognition and various neural oscillations that underlie them, as well as the related biological mechanisms contributing to disease. While there are more than 6 million SNPs in the human genome, very few SNPs associated with these complex traits are likely to be in coding regions of the gene, and therefore may not involve changes in the encoded protein expression, and subsequently its function (Goldberg and Weinberger, 2004; Hirschhorn and Daly, 2005). Variations in the protein coding sequences can result in changes that range from synonymous (unchanged function of the protein) to non-synonymous or missense (changes in the function of the protein) mutations. It is more likely that the majority of these functional changes will involve the regulation of transcription via promoter variants or the organization of transcription through splice-site variants. At this point in time, it is not possible to determine how many of these variations will affect brain oscillations and cognitive processes. It is most likely that multiple genes with small effects may be involved. In addition to the aforementioned role of altered gene regulation impacting on brain oscillations and cognition, special consideration needs to be given to gene–gene interactions, gene–environment interactions and potential stochastic factors.

3.2. Resting EEG beta power: *GABRA2* gene

As increased resting beta power is already observed in offspring before the onset of alcohol dependence, it antecedes the development of alcoholism and is considered to be a “trait” rather than a “state” measure. Significant linkage and linkage disequilibrium between the EEG beta frequency and a gamma-aminobutyric acid (GABA_A) receptor gene on chromosome 4 has been reported (Porjesz et al., 2002). With the use of multiple SNPs across this cluster of GABA_A receptor genes on chromosome 4, that includes *GABRA2*, *GABRA4*, *GABRB1* and *GABRG1*, we were able to specifically identify that it was variations only in the *GABRA2* receptor gene that accounts for the linkage/linkage disequilibrium findings with the beta frequency (See Figure 4; Table 1). Thus, variations in *GABRA2* (the gene encoding the alpha 2 subunit of the GABA_A receptor) affect brain oscillations and are directly involved in the level of neural excitability (balance between excitation and inhibition). There is a strong relationship between the rs279836 SNP in the *GABRA2* receptor gene and Beta 2 EEG power. It is interesting to note that individuals who are homozygous for the rarer genotype (15%) of the rs279836 SNP have significantly increased EEG beta 2 compared to individuals with all other genotypes. These individuals are more likely to manifest CNS disinhibition.

Subsequent SNP (single nucleotide polymorphism) analysis indicated that the same *GABRA2* gene associated with the EEG beta endophenotype is associated with DSM-IV diagnosis of alcohol dependence (Edenberg et al., 2004), substance dependence (Agrawal et al., 2006), ASP (Dick et al., 2006a), and childhood conduct disorder (Dick et al., 2006b). The association between *GABRA2* and alcohol dependence has been replicated by a number of groups throughout the world (Covault et al., 2004; Drgon et al., 2006; Enoch et al., 2006; Fehr et al., 2006; Lappalainen et al., 2005; Soyka et al., 2008).

It has been demonstrated that the beta rhythm is generated while maintaining the balance in networks of excitatory pyramidal cells and inhibitory interneurons, involving GABA_A action as the pacemaker (Whittington et al., 2000). Fast synaptic inhibition in the mammalian central nervous system is mediated largely by activation of GABA_A receptors (Tobler et al., 2001). GABA_A actions are a fundamental requirement for both gamma (30–80 Hz) and beta oscillations to occur, and blockade of these receptors results in loss of synchronization (Haenschel et al., 2000). Beta rhythms can synchronize over long temporal delays between more spatially distant brain loci than gamma rhythms (Kopell et al., 2000). Although the recording of electrical oscillations from a neural population reflects the firing of multiple excitatory pyramidal cells, the mechanism underlying beta and gamma oscillations depends on the firing patterns of a network of inhibitory interneurons (Faulkner et al., 1999; Kopell et al., 2000), gated by their mutually induced GABA_A action (Whittington et al., 2000).

Alcoholics and offspring at high risk manifest increased power in EEG beta oscillations, suggesting an imbalance between excitation-inhibition (CNS disinhibition). This provides a biological hypothesis relating the underlying CNS disinhibition to the genetic risk for alcohol dependence and related disorders (Begleiter and Porjesz, 1999). The involvement of the GABAergic system in alcoholism is supported by neuroimaging studies, which report specific deficits in GABA benzodiazepine receptor function in the brains of alcoholics (Abi-Dargham et al., 1998; Krystal et al., 2006; Lingford-Hughes et al., 1998) and individuals at risk (Volkow et al., 1995). Taken together, these findings suggest GABA deficits in the brains of alcoholics and individuals at risk may account for their lack of CNS inhibition (hyperexcitability) and may be involved in the predisposition to develop alcoholism.

3.3. Interhemispheric theta coherence: *GABRA2* and *CHRM2* genes

Recent studies have suggested a significant role for theta frequency coherence in normal and aberrant thalamocortical interactions (Sarnthein and Jeanmonod, 2007; Sarnthein et al.,

2005). Impairments in neural synchrony have been reported in several psychiatric conditions, including alcohol dependence, as noted in an earlier section (Section 2.2.). We conducted whole genome linkage analysis using the high theta coherence at parietal-occipital leads as the phenotype. Highly significant linkage was found in COGA on chromosome 4 in the same region spotlighted by the EEG beta power phenotype (Porjesz and Rangaswamy, 2007; Rangaswamy et al, in preparation). Family-based association analyses with the cluster of GABA_A receptor genes under this linkage peak revealed strong association with a large number of SNPs (several at $p < 0.001$) genotyped in *GABRA2* for the Caucasian-only subset. There was no significant association with other genes in the GABA_A cluster on chromosome 4 (Figure 4; Table 1).

These results suggest that *GABRA2* may indeed influence susceptibility to alcohol and drug dependence, not just by modulating level of neural excitation, but also by influencing functional connectivity of interhemispheric networks. In their model of thalamocortical dysrhythmia, Llinas and co-workers have proposed that the enhanced low-frequency (theta) oscillations in the thalamocortical module can affect the lateral inhibitory drive in the cortex and eventually result in high frequency coherent activation of cortical modules (Llinas et al., 1999). This is particularly significant in light of our genetic findings where two resting state electrophysiological signatures - Beta power (high frequency activity associated with arousal) and theta coherence (low frequency synchrony) are both linked to GABA_A receptor gene involvement.

In another study (Winterer et al., 2003c), the authors report that 3 exonic variants of the gene encoding the human GABA_B receptor on chromosome 6 modify cortical synchronization measured as scalp-recorded EEG coherence. Parietotemporal coherence showed statistical significance associated with exon 7 and the authors concluded that this exon may be functionally meaningful and impact on cortical EEG oscillations. Alpha rhythms in the scalp electrical activity are strongly influenced by the thalamus and this has been well supported by evidences from animal and human studies including the noninvasive recordings. Some authors have suggested that similar mechanisms associated with thalamocortical cell populations, may form the basis of theta (2–7 Hz) rhythms (Hughes and Crunelli, 2005). Interestingly, a study reported significant association between the exon 7 variant of the GABA_B receptor gene and EEG alpha voltage (classified as LVA or normal), for control but not alcoholic subjects (Winterer et al., 2003b). LVA in females has been associated with a genetic variant resulting in low COMT activity, which is involved in the dopaminergic system yielding low levels of norepinephrine (Enoch et al., 2003). The authors hypothesize that this finding may partly explain their observation of low LVA associated with anxiety disorders in alcoholic women.

Significant linkage on chromosome 7 was observed for the interhemispheric high theta centroparietal coherence phenotype in a whole genome linkage scan, and family based association tests revealed very significant associations with a muscarinic acetylcholine receptor (M2) gene (*CHRM2*), underlying the linkage peak on chromosome 7 (Porjesz and Rangaswamy, 2007) (Figure 5; Table 2). No significant association was observed for SNPs in another candidate gene underlying the linkage peak in this region - the *GRM8* gene.

Significant linkage and association for evoked theta band responses at the same *CHRM2* gene have been previously reported (Jones et al., 2006a; Jones et al., 2004) (see below). Both the GABAergic and cholinergic systems interact significantly in the functions of local inhibitory circuits, thus affecting network functions and influence cortical synchronization. Increased strength of GABAergic inhibition has been reported to be a likely mechanism underlying the impaired synaptic plasticity observed with M2 knockout mice. These mice demonstrate impaired behavioral flexibility and memory deficits (Seeger et al., 2004). In a recent study examining muscarinic control of long-range cortical GABAergic inhibition, the authors

suggested that M2 receptor activation produces a presynaptic inhibition of GABA release by long-range inhibitory neurons of the perirhinal cortex projecting to the entorhinal cortex (Apergis-Schoute et al., 2007).

3.4. Theta and delta event-related oscillations underlying P3: *CHRM2* and *GRM8* genes

We have reported significant linkage and association between the *CHRM2* gene on chromosome 7 and frontal theta oscillations to target stimuli in a visual oddball task; association analyses using both population based tests (Measured Genotype) and pedigree based tests (Quantitative Pedigree Disequilibrium Test, QPDT) indicate significant association of the frontal theta band ERO phenotype with several SNPs surrounding exon 4 of *CHRM2* ($p < 0.001$). Further, an examination of the slower frequency parietal-occipital delta band EROs revealed significant association with several SNPs surrounding the coding region of the *CHRM2* gene ($p < 0.01$) (Jones et al., 2006a; Jones et al., 2004). (See Figure 5; Table 2).

These findings implicate the possible role of *CHRM2* in the generation and modulation of evoked oscillations. Theta and delta EROs depend on the level of acetylcholine (muscarinic activation). M2 receptors inhibit presynaptic release of acetylcholine, leading to inhibition of irrelevant networks. Muscarinic receptors are especially concentrated in the forebrain and possibly serve to maintain the effective balance of relevant/irrelevant networks, hence, having a direct influence on P3 generation (Frodl-Bauch et al., 1999). It is proposed that the genetic underpinnings of evoked oscillations are likely to stem from regulatory genes that control the neurochemical processes of the brain and, therefore, influence neural function. The three major neurochemical substrates contributing to theta and delta rhythms and P3 involve strong GABAergic, cholinergic, and glutamatergic system interactions (Frodl-Bauch et al., 1999).

Moreover, the cholinergic muscarinic genes have a major role in memory and cognition (Calabresi et al., 1998; Comings et al., 2003). Several studies, including the COGA study, have found evidence that this gene may be involved in intelligence (Comings et al., 2003; Dick et al., 2007b; Gosso et al., 2007). In the COGA study, we found evidence of association with multiple SNPs across *CHRM2* and Performance IQ, as measured by the Wechsler Adult Intelligence Scale-Revised (WAIS-R). These results remain significant after taking into account alcohol dependence and depression diagnoses in the sample (Dick et al., 2007b).

Our results with the *CHRM2* gene and brain oscillations strongly support the role of acetylcholine in the generation of N2 (theta oscillations) and in the P3 component (delta and theta oscillations). The function of acetylcholine has been demonstrated with regard to stimulus significance (Perry et al., 1999), selective attention (Mitrofanis and Guillery, 1993), and P3 generation (Callaway, 1983). Administration of cholinergic agonists and antagonists have yielded modified memory performance, and modified P3 amplitude in humans (Dierks et al., 1994; Hammond et al., 1987; Potter et al., 2000). *In vitro* administration of moderate amounts of the muscarinic agonist carbachol in the rat hippocampus induces synchronized delta oscillations, whereas higher concentrations produced short episodes of theta oscillations, and the carbachol-induced delta rhythms were not observed concurrent with carbachol-theta (Fellous and Sejnowski, 2000; Tiesinga et al., 2001).

Recent evidence indicates that the *CHRM2* gene is not only associated with brain oscillations and cognition, but also clinical diagnosis. Significant linkage and association were reported for the *CHRM2* gene and a diagnosis of alcohol dependence and depression (Wang et al., 2004), comorbid alcohol and drug dependence (a more severe addiction profile) (Dick et al., 2007a), as well as a spectrum of externalizing disorders (Dick et al., 2008) in the COGA study. Other groups have also replicated these findings, reporting that the *CHRM2* gene predisposes to alcohol dependence, drug dependence and affective disorders (Luo et al., 2005), and major depression in women (Comings et al., 2002). Thus genes important for the expression of the

endophenotype (brain oscillations) help in identification of genes that increase the susceptibility for risk of alcohol dependence and related disorders (Begleiter and Porjesz, 2006; Dick et al., 2006c).

Under the same theta ERO linkage peak on chromosome 7 is a glutamate receptor (*GRM8*) gene that is another very likely candidate gene for modulating electrophysiological networks, particularly as the glutamatergic system is also involved in theta oscillations and P3 (Frodin-Bauch et al., 1999). Family-based association analyses of theta EROs revealed significant associations with several SNPs and theta EROs to target stimuli at frontal, central, and parietal regions (Chen et al., submitted). An interesting finding is that several SNPs were also significant for diagnosis of alcohol dependence using ICD-10 diagnostic criteria.

A 3-T proton magnetic resonance spectroscopy (1H-MRS) study has suggested the involvement of glutamatergic neurotransmission in integrative frontal-hippocampal processing (Gallinat et al., 2006) and the sensation seeking personality dimension (Gallinat et al., 2007). The study demonstrated a robust relationship between glutamate levels in the hippocampus and frontal theta activity during auditory stimulus processing. Glutamatergic neurotransmission and its neuroadaptive changes have been proposed as important molecular determinants of craving and relapse (Cornish and Kalivas, 2000; Tzschentke and Schmidt, 2000). In particular, it is suggested that a hyperglutamatergic state mediates, at least in part, alcohol relapse behavior and maintenance of alcoholism (Tsai and Coyle, 1998). Several studies have suggested the involvement of glutamate receptors – NMDA and metabotropic, in alcohol relapse (Bachteler et al., 2005; Holter et al., 2000; Krystal et al., 2003). Acamprosate, a drug used to prevent relapse in alcoholic patients (Mann et al., 2004), has been suggested to act through a suppression of a hyperglutamatergic state created by alcohol addiction (Dahchour and De Witte, 2000; Spanagel and Heilig, 2005).

The glutamate receptors belong to two subfamilies: (a) glutamate-gated ion channels, including NMDA, AMPA/kainite type receptors, and (b) metabotropic glutamate receptors, consisting of mGluRs of which intracellular actions are mediated by G-proteins. Metabotropic glutamate receptors (mGluRs) belong to three groups, mGlu I, mGlu II and mGlu III, based on signal transduction pathways and sequence homology (Schoepp, 2001). The *GRM8* gene that encodes mGluR8 spans over 800 kb, and is composed of 10 exons and 9 introns (Shigemoto et al., 1997). Prominent mGlu8 expression has been observed in several areas of limbic system, especially the hippocampus, pontine nuclei, and lateral reticular nucleus of the medulla oblongata. Strong expression for mGlu8 has also been noted in scattered cells in the deeper layers of the cerebral cortex and pyramidal cells of the piriform cortex. The mGlu8 is detected in the presynaptic active zone in both glutamatergic and GABAergic terminals in various brain regions (Ferraguti and Shigemoto, 2006). Presynaptic inhibition of synaptic transmission via activation of mGluRs occurs very widely in the brain, with involvement of group II and group III mGluRs being especially prominent. The electrophysiological studies of these receptors have shown the regulation of synaptic sensitivity via suppression of presynaptic voltage-dependent calcium channels, the activation of presynaptic K⁺ channels and the direct inhibition of regulated exocytosis (Anwyl, 1999).

In light of the theta oscillations showing strong association to both *CHRM2* and *GRM8* genes, one could speculate a synergistic genetic mechanism underlying this electrophysiological phenotype, thus opening doors to future research in this direction. Interestingly, in studies on the rat hippocampus, authors have reported that a majority of interneurons strongly immunopositive for the muscarinic M2 or the mGlu1 receptors were the primary targets of mGluR8-containing terminals (Ferraguti et al., 2005). Rare neurons coexpressing calretinin and M2 were consistently targeted by mGluR8-positive boutons. The postsynaptic interneuron

type-specific expression predicts a role in adjusting the activity of interneurons depending on the level of network activity.

4. Conclusion

This nascent field of investigation into the genetic underpinnings of the oscillatory dynamics is rapidly evolving, propelled by (a) technological advances in the field of genetic research, such as new SNP-chip technology, advanced functional studies, improved statistical genetic methods, and (b) new mathematical tools for signal analysis, improved imaging methods and the emerging idea of oscillatory interactions as units of dynamic functional brain states (Basar, 2006). Together, these advances provide a better means to understand brain function and make gene identification and understanding of gene expression imminently tenable. The recent identification of genetic loci regarding brain oscillations indicates that they are under genetic control and are modulated by genes controlling neurotransmitters in the brain. This approach has the unprecedented potential to unravel the complex interplay of various neural subsystems relevant to the generation of brain oscillations elicited under different cognitive conditions. The advent of genomics and proteomics and a fuller understanding of gene regulation will open new horizons on the critical electrical events so essential for human brain function. It is a hope that by combining these approaches we can eventually understand how much of the capacity to modulate these oscillations, in the context of a disease state, lies in control of genes and how environment influences this control.

Alcohol dependence and related disorders results from a complex interaction of changing genetic and environmental liabilities across development, with greater genetic loading for early-onset disorders. The use of quantitative brain oscillations as endophenotypes provides the power to more easily localize and characterize disease susceptibility genes than diagnostic categories. In our endeavor to study the genetic underpinnings of predisposition/risk, some key genes that have emerged from the quantitative phenotype studies are the GABA_A receptor (*GABRA2*), the metabotropic glutamate receptor (*GRM8*), and the cholinergic muscarinic receptor (*CHRM2*) genes. These results do not imply that these genes are wholly responsible for the existence of the respective brain oscillations. It is however more likely that the individual variations in the oscillations are a product of genetic variation of the *GABRA2*, *GRM8* and *CHRM2* receptor genes, and the altered oscillatory activity is its cognitive correlate. Although no functional variant affecting the electrophysiological characteristics has yet been identified at the molecular level, there is a great deal of pharmacological evidence that attests to the relevance of these receptors to aspects of cognitive functions. Moreover, these genes are known to have very complex transcriptional patterns with different types of promoters and tissue specific expression patterns that add to the complexity.

Functional studies are underway to understand the effects of observed coding and/or non-coding variants of the key genes in gene expression and protein function in animal and *in vitro* models. Data from our COGA project indicate that in the case of *GABRA2*, the regulatory differences may affect alternative splicing and alternative promoter use (Tian et al., 2005). This eventually results in several isoforms in different brain regions, possibly with differing levels of functionality. Future molecular studies examining the functionality of isoforms will allow us to come full circle in determining the brain oscillation-to-gene connections.

Understanding genetic control of brain electrical activity may provide clues about cerebral function, and may shed light on pathogenic mechanisms involved in neurological and psychiatric disorders, where impairment in brain electrical activity is apparent. Once genes are identified and understood, risk genotypes and haplotypes can be used in prospective studies of young individuals, and can lead to an improved understanding of how cognitive changes

contribute to susceptibility, which in turn can lead to the design of well-targeted prevention initiatives.

Acknowledgments

This manuscript is dedicated to the memory of Henri Begleiter (deceased April 6, 2006), Distinguished Professor of Psychiatry and Neuroscience, and founder and director of the Neurodynamics Laboratory, now renamed in his honor. His scientific vision brought together the fields of brain oscillations and genetics, culminating in the use of brain oscillations as endophenotypes, which is the subject of this manuscript. Our laboratory continues to follow the path that he envisioned, inspired by his innovative approaches to research.

The work reviewed in this manuscript was supported by NIH Grants AA008401, AA05524, AA002686 from the National Institute of Alcohol Abuse and Alcoholism.

List of abbreviations

ADHD	Attention Deficit Hyperactivity Disorder
AMPA	Alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionate glutamate receptor
ASP	Antisocial Personality Disorder
CHRM2	cholinergic muscarinic receptor type 2
CNS	Central Nervous System
COGA	Collaborative study on the Genetics of Alcoholism
EEG	electroencephalogram
ERO	Event-related oscillation
ERP	Event-related potential
FBAT	Family based Association Studies
FDR	False discovery rate
GABA	gamma aminobutyric acid
<i>GABRA2</i>	gamma-aminobutyric acid (GABA) A receptor, alpha 2
<i>GABRA4</i>	gamma-aminobutyric acid (GABA) A receptor, alpha 4
<i>GABRB1</i>	gamma-aminobutyric acid (GABA) A receptor, beta 1
<i>GABRG1</i>	gamma-aminobutyric acid (GABA) A receptor, gamma 1
GRM8	glutamate receptor, metabotropic 8
GWAS	Genomewide Association Studies
LD	Linkage disequilibrium
LOD	Logarithm of Odds
LVA	Low voltage alpha
mGluR	metabotropic glutamate receptors
ms	millisecond
NMDA	N-methyl-D-aspartate glutamate receptor
QPDT	Quantitative Pedigree Disequilibrium Test
SNP	Single-nucleotide polymorphisms

References

- Abi-Dargham A, Krystal JH, Anjilvel S, Scanley BE, Zoghbi S, Baldwin RM, Rajeevan N, Ellis S, Petrakis IL, Seibyl JP, Charney DS, Laruelle M, Innis RB. Alterations of benzodiazepine receptors in type II alcoholic subjects measured with SPECT and [123I]iomazenil. *Am J Psychiatry* 1998;155:1550–5. [PubMed: 9812116]
- Agrawal A, Edenberg HJ, Foroud T, Bierut LJ, Dunne G, Hinrichs AL, Nurnberger JI, Crowe R, Kuperman S, Schuckit MA, Begleiter H, Porjesz B, Dick DM. Association of GABRA2 with drug dependence in the collaborative study of the genetics of alcoholism sample. *Behav Genet* 2006;36:640–50. [PubMed: 16622805]
- Almasy L. Quantitative risk factors as indices of alcoholism susceptibility. *Ann Med* 2003;35:337–43. [PubMed: 12952020]
- Almasy L, Blangero J. Multipoint quantitative-trait linkage analysis in general pedigrees. *Am J Hum Genet* 1998;62:1198–211. [PubMed: 9545414]
- Anwyl R. Metabotropic glutamate receptors: electrophysiological properties and role in plasticity. *Brain Res Brain Res Rev* 1999;29:83–120. [PubMed: 9974152]
- Apergis-Schoute J, Pinto A, Pare D. Muscarinic control of long-range GABAergic inhibition within the rhinal cortices. *J Neurosci* 2007;27:4061–71. [PubMed: 17428984]
- Bachteler D, Economidou D, Danysz W, Ciccocioppo R, Spanagel R. The effects of acamprosate and neramexane on cue-induced reinstatement of ethanol-seeking behavior in rat. *Neuropsychopharmacology* 2005;30:1104–10. [PubMed: 15668725]
- Banaschewski T, Brandeis D. Annotation: what electrical brain activity tells us about brain function that other techniques cannot tell us - a child psychiatric perspective. *J Child Psychol Psychiatry* 2007;48:415–35. [PubMed: 17501723]
- Barry RJ, Clarke AR, McCarthy R, Selikowitz M. EEG coherence in attention-deficit/hyperactivity disorder: a comparative study of two DSM-IV types. *Clin Neurophysiol* 2002;113:579–85. [PubMed: 11956003]
- Basar-Eroglu C, Basar E. A compound P300-40 Hz response of the cat hippocampus. *Int J Neurosci* 1991;60:227–37. [PubMed: 1787051]
- Basar-Eroglu C, Basar E, Demiralp T, Schurmann M. P300-response: possible psychophysiological correlates in delta and theta frequency channels. A review. *Int J Psychophysiol* 1992;13:161–79. [PubMed: 1399755]
- Basar-Eroglu C, Demiralp T, Schurmann M, Basar E. Topological distribution of oddball ‘P300’ responses. *Int J Psychophysiol* 2001;39:213–20. [PubMed: 11163898]
- Basar-Eroglu C, Struber D, Kruse P, Basar E, Stadler M. Frontal gamma-band enhancement during multistable visual perception. *Int J Psychophysiol* 1996a;24:113–25. [PubMed: 8978438]
- Basar-Eroglu C, Struber D, Schurmann M, Stadler M, Basar E. Gamma-band responses in the brain: a short review of psychophysiological correlates and functional significance. *Int J Psychophysiol* 1996b;24:101–12. [PubMed: 8978437]
- Basar, E. *EEG-Brain Dynamics: Relation Between EEG and Brain Evoked Potentials*. Elsevier; Amsterdam: 1980.
- Basar, E. *Brain Function and Oscillations. Vol. II: Integrative brain function, neurophysiology and cognitive processes*. Springer Verlag; Berlin: 1999.
- Basar E. The theory of the whole-brain-work. *Int J Psychophysiol* 2006;60:133–8. [PubMed: 16563537]
- Basar E, Basar-Eroglu C, Karakas S, Schurmann M. Are cognitive processes manifested in event-related gamma, alpha, theta and delta oscillations in the EEG? *Neurosci Lett* 1999;259:165–8. [PubMed: 10025584]
- Basar E, Basar-Eroglu C, Karakas S, Schurmann M. Gamma, alpha, delta, and theta oscillations govern cognitive processes. *Int J Psychophysiol* 2001a;39:241–8. [PubMed: 11163901]
- Basar E, Schurmann M, Basar-Eroglu C, Karakas S. Alpha oscillations in brain functioning: an integrative theory. *Int J Psychophysiol* 1997;26:5–29. [PubMed: 9202992]
- Basar E, Schurmann M, Sakowitz O. The selectively distributed theta system: functions. *Int J Psychophysiol* 2001b;39:197–212. [PubMed: 11163897]

- Bauer LO. Predicting relapse to alcohol and drug abuse via quantitative electroencephalography. *Neuropsychopharmacology* 2001;25:332–40. [PubMed: 11522462]
- Bauer LO, Hesselbrock VM. EEG, autonomic and subjective correlates of the risk for alcoholism. *J Stud Alcohol* 1993;54:577–89. [PubMed: 8412148]
- Begleiter H, Porjesz B. What is inherited in the predisposition toward alcoholism? A proposed model. *Alcohol Clin Exp Res* 1999;23:1125–35. [PubMed: 10443977]
- Begleiter H, Porjesz B. Genetics of human brain oscillations. *Int J Psychophysiol* 2006;60:162–71. [PubMed: 16540194]
- Begleiter H, Porjesz B, Bihari B, Kissin B. Event-related brain potentials in boys at risk for alcoholism. *Science* 1984;225:1493–6. [PubMed: 6474187]
- Buzsaki G, Draguhn A. Neuronal oscillations in cortical networks. *Science* 2004;304:1926–9. [PubMed: 15218136]
- Calabresi P, Centonze D, Gubellini P, Pisani A, Bernardi G. Blockade of M2-like muscarinic receptors enhances long-term potentiation at corticostriatal synapses. *Eur J Neurosci* 1998;10:3020–3. [PubMed: 9758172]
- Callaway E. Presidential address, 1982. The pharmacology of human information processing. *Psychophysiology* 1983;20:359–70. [PubMed: 6356200]
- Chorlian DB, Tang Y, Rangaswamy M, O'Connor S, Rohrbaugh J, Taylor R, Porjesz B. Heritability of EEG coherence in a large sib-pair population. *Biol Psychol* 2007;75:260–6. [PubMed: 17498861]
- Clarke AR, Barry RJ, McCarthy R, Selikowitz M, Johnstone SJ, Hsu CI, Magee CA, Lawrence CA, Croft RJ. Coherence in children with Attention-Deficit/Hyperactivity Disorder and excess beta activity in their EEG. *Clin Neurophysiol* 2007;118:1472–9. [PubMed: 17502165]
- Comings DE, Wu S, Rostamkhani M, McGue M, Iacono WG, MacMurray JP. Association of the muscarinic cholinergic 2 receptor (CHRM2) gene with major depression in women. *Am J Med Genet* 2002;114:527–9. [PubMed: 12116189]
- Comings DE, Wu S, Rostamkhani M, McGue M, Iacono WG, Cheng LS, MacMurray JP. Role of the cholinergic muscarinic 2 receptor (CHRM2) gene in cognition. *Mol Psychiatry* 2003;8:10–1. [PubMed: 12556901]
- Cook IA, O'Hara R, Uijtdehaage SH, Mandelkern M, Leuchter AF. Assessing the accuracy of topographic EEG mapping for determining local brain function. *Electroencephalogr Clin Neurophysiol* 1998;107:408–14. [PubMed: 9922086]
- Corbetta M, Kincade JM, Ollinger JM, McAvoy MP, Shulman GL. Voluntary orienting is dissociated from target detection in human posterior parietal cortex. *Nat Neurosci* 2000;3:292–7. [PubMed: 10700263]
- Cornish JL, Kalivas PW. Glutamate transmission in the nucleus accumbens mediates relapse in cocaine addiction. *J Neurosci* 2000;20:RC89. [PubMed: 10899176]
- Costa L, Bauer L. Quantitative electroencephalographic differences associated with alcohol, cocaine, heroin and dual-substance dependence. *Drug Alcohol Depend* 1997;46:87–93. [PubMed: 9246556]
- Covault J, Gelernter J, Hesselbrock V, Nellissery M, Kranzler HR. Allelic and haplotypic association of GABRA2 with alcohol dependence. *Am J Med Genet B Neuropsychiatr Genet* 2004;129:104–9. [PubMed: 15274050]
- Dahchour A, De Witte P. Ethanol and amino acids in the central nervous system: assessment of the pharmacological actions of acamprosate. *Prog Neurobiol* 2000;60:343–62. [PubMed: 10670704]
- Damoiseaux JS, Rombouts SA, Barkhof F, Scheltens P, Stam CJ, Smith SM, Beckmann CF. Consistent resting-state networks across healthy subjects. *Proc Natl Acad Sci U S A* 2006;103:13848–53. [PubMed: 16945915]
- De Luca M, Beckmann CF, De Stefano N, Matthews PM, Smith SM. fMRI resting state networks define distinct modes of long-distance interactions in the human brain. *Neuroimage* 2006;29:1359–67. [PubMed: 16260155]
- Demiralp T, Ademoglu A, Istefanopulos Y, Basar-Eroglu C, Basar E. Wavelet analysis of oddball P300. *Int J Psychophysiol* 2001;39:221–7. [PubMed: 11163899]
- Dick DM, Agrawal A, Schuckit MA, Bierut L, Hinrichs A, Fox L, Mullaney J, Cloninger CR, Hesselbrock V, Nurnberger JI Jr, Almasy L, Foroud T, Porjesz B, Edenberg H, Begleiter H. Marital status, alcohol

- dependence, and GABRA2: evidence for gene-environment correlation and interaction. *J Stud Alcohol* 2006a;67:185–94. [PubMed: 16562401]
- Dick DM, Agrawal A, Wang JC, Hinrichs A, Bertelsen S, Bucholz KK, Schuckit M, Kramer J, Nurnberger J Jr, Tischfield J, Edenberg HJ, Goate A, Bierut LJ. Alcohol dependence with comorbid drug dependence: genetic and phenotypic associations suggest a more severe form of the disorder with stronger genetic contribution to risk. *Addiction* 2007a;102:1131–9. [PubMed: 17567401]
- Dick DM, Aliev F, Kramer J, Wang JC, Hinrichs A, Bertelsen S, Kuperman S, Schuckit M, Nurnberger J Jr, Edenberg HJ, Porjesz B, Begleiter H, Hesselbrock V, Goate A, Bierut L. Association of CHRM2 with IQ: converging evidence for a gene influencing intelligence. *Behav Genet* 2007b;37:265–72. [PubMed: 17160701]
- Dick DM, Aliev F, Wang JC, Grucza RA, Schuckit M, Kuperman S, Kramer J, Hinrichs A, Bertelsen S, Budde JP, Hesselbrock V, Porjesz B, Edenberg HJ, Bierut LJ, Goate A. Using Dimensional Models of Externalizing Psychopathology to Aid in Gene Identification. 2008;65:310–318.
- Dick DM, Bierut L, Hinrichs A, Fox L, Bucholz KK, Kramer J, Kuperman S, Hesselbrock V, Schuckit M, Almasy L, Tischfield J, Porjesz B, Begleiter H, Nurnberger J Jr, Xuei X, Edenberg HJ, Foroud T. The role of GABRA2 in risk for conduct disorder and alcohol and drug dependence across developmental stages. *Behav Genet* 2006b;36:577–90. [PubMed: 16557364]
- Dick DM, Jones K, Saccone N, Hinrichs A, Wang JC, Goate A, Bierut L, Almasy L, Schuckit M, Hesselbrock V, Tischfield J, Foroud T, Edenberg H, Porjesz B, Begleiter H. Endophenotypes successfully lead to gene identification: results from the collaborative study on the genetics of alcoholism. *Behav Genet* 2006c;36:112–26. [PubMed: 16341909]
- Dierks T, Frolich L, Ihl R, Maurer K. Event-related potentials and psychopharmacology. Cholinergic modulation of P300. *Pharmacopsychiatry* 1994;27:72–4. [PubMed: 7913237]
- Doppelmayr M, Klimesch W, Schwaiger J, Auinger P, Winkler T. Theta synchronization in the human EEG and episodic retrieval. *Neurosci Lett* 1998;257:41–4. [PubMed: 9857961]
- Drgon T, D'Addario C, Uhl GR. Linkage disequilibrium, haplotype and association studies of a chromosome 4 GABA receptor gene cluster: candidate gene variants for addictions. *Am J Med Genet B Neuropsychiatr Genet* 2006;141:854–60. [PubMed: 16894595]
- Edenberg HJ, Dick DM, Xuei X, Tian H, Almasy L, Bauer LO, Crowe RR, Goate A, Hesselbrock V, Jones K, Kwon J, Li TK, Nurnberger JI Jr, O'Connor SJ, Reich T, Rice J, Schuckit MA, Porjesz B, Foroud T, Begleiter H. Variations in GABRA2, encoding the alpha 2 subunit of the GABA(A) receptor, are associated with alcohol dependence and with brain oscillations. *Am J Hum Genet* 2004;74:705–14. [PubMed: 15024690]
- Engel AK, Fries P, Singer W. Dynamic predictions: oscillations and synchrony in top-down processing. *Nat Rev Neurosci* 2001;2:704–16. [PubMed: 11584308]
- Enoch MA, Schwartz L, Albaugh B, Virkkunen M, Goldman D. Dimensional anxiety mediates linkage of GABRA2 haplotypes with alcoholism. *Am J Med Genet B Neuropsychiatr Genet* 2006;141:599–607. [PubMed: 16874763]
- Enoch MA, White KV, Harris CR, Robin RW, Ross J, Rohrbaugh JW, Goldman D. Association of low-voltage alpha EEG with a subtype of alcohol use disorders. *Alcohol Clin Exp Res* 1999;23:1312–9. [PubMed: 10470973]
- Enoch MA, Xu K, Ferro E, Harris CR, Goldman D. Genetic origins of anxiety in women: a role for a functional catechol-O-methyltransferase polymorphism. *Psychiatr Genet* 2003;13:33–41. [PubMed: 12605099]
- Faulkner HJ, Traub RD, Whittington MA. Anaesthetic/amnesic agents disrupt beta frequency oscillations associated with potentiation of excitatory synaptic potentials in the rat hippocampal slice. *Br J Pharmacol* 1999;128:1813–25. [PubMed: 10588938]
- Fehr C, Sander T, Tadic A, Lenzen KP, Anghelescu I, Klawe C, Dahmen N, Schmidt LG, Szegedi A. Confirmation of association of the GABRA2 gene with alcohol dependence by subtype-specific analysis. *Psychiatr Genet* 2006;16:9–17. [PubMed: 16395124]
- Fell J, Fernandez G, Klaver P, Elger CE, Fries P. Is synchronized neuronal gamma activity relevant for selective attention? *Brain Res Brain Res Rev* 2003;42:265–72. [PubMed: 12791444]

- Fellous JM, Sejnowski TJ. Cholinergic induction of oscillations in the hippocampal slice in the slow (0.5–2 Hz), theta (5–12 Hz), and gamma (35–70 Hz) bands. *Hippocampus* 2000;10:187–97. [PubMed: 10791841]
- Ferraguti F, Klausberger T, Cobden P, Baude A, Roberts JD, Szucs P, Kinoshita A, Shigemoto R, Somogyi P, Dalezios Y. Metabotropic glutamate receptor 8-expressing nerve terminals target subsets of GABAergic neurons in the hippocampus. *J Neurosci* 2005;25:10520–36. [PubMed: 16280590]
- Ferraguti F, Shigemoto R. Metabotropic glutamate receptors. *Cell Tissue Res* 2006;326:483–504. [PubMed: 16847639]
- Finn PR, Justus A. Reduced EEG alpha power in the male and female offspring of alcoholics. *Alcohol Clin Exp Res* 1999;23:256–62. [PubMed: 10069554]
- Ford JM, Krystal JH, Mathalon DH. Neural synchrony in schizophrenia: from networks to new treatments. *Schizophr Bull* 2007;33:848–52. [PubMed: 17567628]
- Fox MD, Snyder AZ, Vincent JL, Corbetta M, Van Essen DC, Raichle ME. The human brain is intrinsically organized into dynamic, anticorrelated functional networks. *Proc Natl Acad Sci U S A* 2005;102:9673–8. [PubMed: 15976020]
- Fox MD, Snyder AZ, Zacks JM, Raichle ME. Coherent spontaneous activity accounts for trial-to-trial variability in human evoked brain responses. *Nat Neurosci* 2006;9:23–5. [PubMed: 16341210]
- Frodl-Bauch T, Bottlender R, Hegerl U. Neurochemical substrates and neuroanatomical generators of the event-related P300. *Neuropsychobiology* 1999;40:86–94. [PubMed: 10474063]
- Frund I, Busch NA, Schadow J, Korner U, Herrmann CS. From perception to action: phase-locked gamma oscillations correlate with reaction times in a speeded response task. *BMC Neurosci* 2007;8:27. [PubMed: 17439642]
- Gabrielli WF Jr, Mednick SA, Volavka J, Pollock VE, Schulsinger F, Itil TM. Electroencephalograms in children of alcoholic fathers. *Psychophysiology* 1982;19:404–7. [PubMed: 7122778]
- Gallinat J, Kunz D, Lang UE, Neu P, Kassim N, Kienast T, Seifert F, Schubert F, Bajbouj M. Association between cerebral glutamate and human behaviour: the sensation seeking personality trait. *Neuroimage* 2007;34:671–8. [PubMed: 17123835]
- Gallinat J, Kunz D, Senkowski D, Kienast T, Seifert F, Schubert F, Heinz A. Hippocampal glutamate concentration predicts cerebral theta oscillations during cognitive processing. *Psychopharmacology (Berl)* 2006;187:103–11. [PubMed: 16767420]
- Gevens A, Smith ME, Leong H, McEvoy L, Whitfield S, Du R, Rush G. Monitoring working memory load during computer-based tasks with EEG pattern recognition methods. *Hum Factors* 1998;40:79–91. [PubMed: 9579105]
- Giesbrecht B, Woldorff MG, Song AW, Mangun GR. Neural mechanisms of top-down control during spatial and feature attention. *Neuroimage* 2003;19:496–512. [PubMed: 12880783]
- Goldberg TE, Weinberger DR. Genes and the parsing of cognitive processes. *Trends Cogn Sci* 2004;8:325–35. [PubMed: 15242692]
- Gosso FM, de Geus EJ, Polderman TJ, Boomsma DI, Posthuma D, Heutink P. Exploring the functional role of the CHRM2 gene in human cognition: results from a dense genotyping and brain expression study. *BMC Med Genet* 2007;8:66. [PubMed: 17996044]
- Gottesman II, Gould TD. The endophenotype concept in psychiatry: etymology and strategic intentions. *Am J Psychiatry* 2003;160:636–45. [PubMed: 12668349]
- Gottesman II, Shields J. Genetic theorizing and schizophrenia. *Br J Psychiatry* 1973;122:15–30. [PubMed: 4683020]
- Gottesman II, Shields J. *Schizophrenia and Genetics: A Twin Study Vantage Point*. Academic Press; New York: 1972.
- Grossman A, Morlet J. Decomposition of Hardy functions into square integrable wavelets of constant shape. *SIAM J Math* 1984;15:723–736.
- Gruber T, Muller MM. Oscillatory brain activity in the human EEG during indirect and direct memory tasks. *Brain Res* 2006;1097:194–204. [PubMed: 16729980]
- Haenschel C, Baldeweg T, Croft RJ, Whittington M, Gruzelier J. Gamma and beta frequency oscillations in response to novel auditory stimuli: A comparison of human electroencephalogram (EEG) data with in vitro models. *Proc Natl Acad Sci U S A* 2000;97:7645–50. [PubMed: 10852953]

- Haig AR, De Pascalis V, Gordon E. Peak gamma latency correlated with reaction time in a conventional oddball paradigm. *Clin Neurophysiol* 1999;110:158–65. [PubMed: 10348335]
- Hammond EJ, Meador KJ, Aung-Din R, Wilder BJ. Cholinergic modulation of human P3 event-related potentials. *Neurology* 1987;37:346–50. [PubMed: 3808322]
- Herrmann CS, Demiralp T. Human EEG gamma oscillations in neuropsychiatric disorders. *Clin Neurophysiol* 2005;116:2719–33. [PubMed: 16253555]
- Higashima M, Takeda T, Kikuchi M, Nagasawa T, Hirao N, Oka T, Nakamura M, Koshino Y. State-dependent changes in intrahemispheric EEG coherence for patients with acute exacerbation of schizophrenia. *Psychiatry Res* 2007;149:41–7. [PubMed: 17140670]
- Higashima M, Takeda T, Kikuchi M, Nagasawa T, Koshino Y. Functional connectivity between hemispheres and schizophrenic symptoms: a longitudinal study of interhemispheric EEG coherence in patients with acute exacerbations of schizophrenia. *Clin EEG Neurosci* 2006;37:10–5. [PubMed: 16475479]
- Hirschhorn JN, Daly MJ. Genome-wide association studies for common diseases and complex traits. *Nat Rev Genet* 2005;6:95–108. [PubMed: 15716906]
- Holter SM, Linthorst AC, Reul JM, Spanagel R. Withdrawal symptoms in a long-term model of voluntary alcohol drinking in Wistar rats. *Pharmacol Biochem Behav* 2000;66:143–51. [PubMed: 10837854]
- Hughes SW, Crunelli V. Thalamic mechanisms of EEG alpha rhythms and their pathological implications. *Neuroscientist* 2005;11:357–72. [PubMed: 16061522]
- Jones KA, Porjesz B, Almasy L, Bierut L, Dick D, Goate A, Hinrichs A, Rice JP, Wang JC, Bauer LO, Crowe R, Foroud T, Hesselbrock V, Kuperman S, Nurnberger J Jr, O'Connor SJ, Rohrbaugh J, Schuckit MA, Tischfield J, Edenberg HJ, Begleiter H. A cholinergic receptor gene (CHRM2) affects event-related oscillations. *Behav Genet* 2006a;36:627–39. [PubMed: 16823639]
- Jones KA, Porjesz B, Almasy L, Bierut L, Goate A, Wang JC, Dick DM, Hinrichs A, Kwon J, Rice JP, Rohrbaugh J, Stock H, Wu W, Bauer LO, Chorlian DB, Crowe RR, Edenberg HJ, Foroud T, Hesselbrock V, Kuperman S, Nurnberger J Jr, O'Connor SJ, Schuckit MA, Stimus AT, Tischfield JA, Reich T, Begleiter H. Linkage and linkage disequilibrium of evoked EEG oscillations with CHRM2 receptor gene polymorphisms: implications for human brain dynamics and cognition. *Int J Psychophysiol* 2004;53:75–90. [PubMed: 15210286]
- Jones KA, Porjesz B, Chorlian D, Rangaswamy M, Kamarajan C, Padmanabhapillai A, Stimus A, Begleiter H. S-transform time-frequency analysis of P300 reveals deficits in individuals diagnosed with alcoholism. *Clin Neurophysiol* 2006b;117:2128–43. [PubMed: 16926113]
- Kamarajan C, Porjesz B, Jones K, Chorlian D, Padmanabhapillai A, Rangaswamy M, Stimus A, Begleiter H. Event-related oscillations in offspring of alcoholics: neurocognitive disinhibition as a risk for alcoholism. *Biol Psychiatry* 2006;59:625–34. [PubMed: 16213472]
- Kamarajan C, Porjesz B, Jones KA, Choi K, Chorlian DB, Padmanabhapillai A, Rangaswamy M, Stimus AT, Begleiter H. The role of brain oscillations as functional correlates of cognitive systems: a study of frontal inhibitory control in alcoholism. *Int J Psychophysiol* 2004;51:155–80. [PubMed: 14693365]
- Kaplan RF, Glueck BC, Hesselbrock MN, Reed HB Jr. Power and coherence analysis of the EEG in hospitalized alcoholics and nonalcoholic controls. *J Stud Alcohol* 1985;46:122–7. [PubMed: 3990297]
- Karakas S, Erzenin OU, Basar E. The genesis of human event-related responses explained through the theory of oscillatory neural assemblies. *Neurosci Lett* 2000a;285:45–8. [PubMed: 10788704]
- Karakas S, Erzenin OU, Basar E. A new strategy involving multiple cognitive paradigms demonstrates that ERP components are determined by the superposition of oscillatory responses. *Clin Neurophysiol* 2000b;111:1719–32. [PubMed: 11018485]
- Kendler KS, Prescott CA, Myers J, Neale MC. The structure of genetic and environmental risk factors for common psychiatric and substance use disorders in men and women. *Arch Gen Psychiatry* 2003;60:929–37. [PubMed: 12963675]
- Klimesch W. EEG-alpha rhythms and memory processes. *Int J Psychophysiol* 1997;26:319–40. [PubMed: 9203012]
- Klimesch W, Doppelmayr M, Pachinger T, Ripper B. Brain oscillations and human memory: EEG correlates in the upper alpha and theta band. *Neurosci Lett* 1997a;238:9–12. [PubMed: 9464642]

- Klimesch W, Doppelmayr M, Pachinger T, Russeger H. Event-related desynchronization in the alpha band and the processing of semantic information. *Brain Res Cogn Brain Res* 1997b;6:83–94. [PubMed: 9450602]
- Klimesch W, Doppelmayr M, Russeger H, Pachinger T, Schwaiger J. Induced alpha band power changes in the human EEG and attention. *Neurosci Lett* 1998;244:73–6. [PubMed: 9572588]
- Klimesch W, Doppelmayr M, Yonelinas A, Kroll NE, Lazzara M, Rohm D, Gruber W. Theta synchronization during episodic retrieval: neural correlates of conscious awareness. *Brain Res Cogn Brain Res* 2001;12:33–8. [PubMed: 11489606]
- Klimesch W, Sauseng P, Hanslmayr S, Gruber W, Freunberger R. Event-related phase reorganization may explain evoked neural dynamics. *Neurosci Biobehav Rev* 2007;31:1003–16. [PubMed: 17532471]
- Klimesch W, Schimke H, Schwaiger J. Episodic and semantic memory: an analysis in the EEG theta and alpha band. *Electroencephalogr Clin Neurophysiol* 1994;91:428–41. [PubMed: 7529682]
- Kopell N, Ermentrout GB, Whittington MA, Traub RD. Gamma rhythms and beta rhythms have different synchronization properties. *Proc Natl Acad Sci U S A* 2000;97:1867–72. [PubMed: 10677548]
- Krystal JH, Petrakis IL, Mason G, Trevisan L, D'Souza DC. N-methyl-D-aspartate glutamate receptors and alcoholism: reward, dependence, treatment, and vulnerability. *Pharmacol Ther* 2003;99:79–94. [PubMed: 12804700]
- Krystal JH, Staley J, Mason G, Petrakis IL, Kaufman J, Harris RA, Gelernter J, Lappalainen J. Gamma-aminobutyric acid type A receptors and alcoholism: intoxication, dependence, vulnerability, and treatment. *Arch Gen Psychiatry* 2006;63:957–68. [PubMed: 16952998]
- Lappalainen J, Krupitsky E, Remizov M, Pchelina S, Taraskina A, Zvartau E, Somberg LK, Covault J, Kranzler HR, Krystal JH, Gelernter J. Association between alcoholism and gamma-amino butyric acid alpha2 receptor subtype in a Russian population. *Alcohol Clin Exp Res* 2005;29:493–8. [PubMed: 15834213]
- Lingford-Hughes AR, Acton PD, Gacinovic S, Suckling J, Busatto GF, Boddington SJ, Bullmore E, Woodruff PW, Costa DC, Pilowsky LS, Ell PJ, Marshall EJ, Kerwin RW. Reduced levels of GABA-benzodiazepine receptor in alcohol dependency in the absence of grey matter atrophy. *Br J Psychiatry* 1998;173:116–22. [PubMed: 9850223]
- Llinas RR. The intrinsic electrophysiological properties of mammalian neurons: insights into central nervous system function. *Science* 1988;242:1654–64. [PubMed: 3059497]
- Llinas RR, Ribary U, Jeanmonod D, Kronberg E, Mitra PP. Thalamocortical dysrhythmia: A neurological and neuropsychiatric syndrome characterized by magnetoencephalography. *Proc Natl Acad Sci U S A* 1999;96:15222–7. [PubMed: 10611366]
- Luo X, Kranzler HR, Zuo L, Wang S, Blumberg HP, Gelernter J. CHRM2 gene predisposes to alcohol dependence, drug dependence and affective disorders: results from an extended case-control structured association study. *Hum Mol Genet* 2005;14:2421–34. [PubMed: 16000316]
- Makeig S, Westerfield M, Jung TP, Enghoff S, Townsend J, Courchesne E, Sejnowski TJ. Dynamic brain sources of visual evoked responses. *Science* 2002;295:690–4. [PubMed: 11809976]
- Mallat S, Zhang Z. Matching pursuits with time-frequency dictionaries. *IEEE Transactions on Signal Processing* 1993;41:3397–3415.
- Mann K, Leher P, Morgan MY. The efficacy of acamprosate in the maintenance of abstinence in alcohol-dependent individuals: results of a meta-analysis. *Alcohol Clin Exp Res* 2004;28:51–63. [PubMed: 14745302]
- Mantini D, Perrucci MG, Del Gratta C, Romani GL, Corbetta M. Electrophysiological signatures of resting state networks in the human brain. *Proc Natl Acad Sci U S A* 2007;104:13170–5. [PubMed: 17670949]
- Mazoyer B, Zago L, Mellet E, Bricogne S, Etard O, Houde O, Crivello F, Joliot M, Petit L, Tzourio-Mazoyer N. Cortical networks for working memory and executive functions sustain the conscious resting state in man. *Brain Res Bull* 2001;54:287–98. [PubMed: 11287133]
- Michael A, Mirza KA, Mukundan CR, Channabasavanna SM. Interhemispheric electroencephalographic coherence as a biological marker in alcoholism. *Acta Psychiatr Scand* 1993;87:213–7. [PubMed: 8465670]

- Mitrofanis J, Guillery RW. New views of the thalamic reticular nucleus in the adult and the developing brain. *Trends Neurosci* 1993;16:240–5. [PubMed: 7688166]
- Murias M, Webb SJ, Greenson J, Dawson G. Resting state cortical connectivity reflected in EEG coherence in individuals with autism. *Biol Psychiatry* 2007;62:270–3. [PubMed: 17336944]
- Niedermayer, E.; Lopes Da Silva, FH. *Electroencephalography: Basic Principles, Clinical Applications and Related Fields*. Williams & Wilkins; New York: 1999.
- Nunez PL, Srinivasan R. A theoretical basis for standing and traveling brain waves measured with human EEG with implications for an integrated consciousness. *Clin Neurophysiol* 2006;117:2424–35. [PubMed: 16996303]
- Padmanabhapillai A, Porjesz B, Ranganathan M, Jones KA, Chorlian DB, Tang Y, Kamarajan C, Rangaswamy M, Stimus A, Begleiter H. Suppression of early evoked gamma band response in male alcoholics during a visual oddball task. *Int J Psychophysiol* 2006a;60:15–26. [PubMed: 16019097]
- Padmanabhapillai A, Tang Y, Ranganathan M, Rangaswamy M, Jones KA, Chorlian DB, Kamarajan C, Stimus A, Kuperman S, Rohrbaugh J, O'Connor SJ, Bauer LO, Schuckit MA, Begleiter H, Porjesz B. Evoked gamma band response in male adolescent subjects at high risk for alcoholism during a visual oddball task. *Int J Psychophysiol* 2006b;62:262–71. [PubMed: 16887227]
- Perry E, Walker M, Grace J, Perry R. Acetylcholine in mind: a neurotransmitter correlate of consciousness? *Trends Neurosci* 1999;22:273–80. [PubMed: 10354606]
- Pollock VE, Earleywine M, Gabrielli WF. Personality and EEG beta in older adults with alcoholic relatives. *Alcohol Clin Exp Res* 1995;19:37–43. [PubMed: 7771661]
- Porjesz B, Almasy L, Edenberg HJ, Wang K, Chorlian DB, Foroud T, Goate A, Rice JP, O'Connor SJ, Rohrbaugh J, Kuperman S, Bauer LO, Crowe RR, Schuckit MA, Hesselbrock V, Conneally PM, Tischfield JA, Li TK, Reich T, Begleiter H. Linkage disequilibrium between the beta frequency of the human EEG and a GABAA receptor gene locus. *Proc Natl Acad Sci U S A* 2002;99:3729–33. [PubMed: 11891318]
- Porjesz B, Rangaswamy M. Neurophysiological endophenotypes, CNS disinhibition, and risk for alcohol dependence and related disorders. *ScientificWorldJournal* 2007;7:131–41. [PubMed: 17982586]
- Porjesz B, Rangaswamy M, Kamarajan C, Jones KA, Padmanabhapillai A, Begleiter H. The utility of neurophysiological markers in the study of alcoholism. *Clin Neurophysiol* 2005;116:993–1018. [PubMed: 15826840]
- Potter DD, Pickles CD, Roberts RC, Rugg MD. Scopolamine impairs memory performance and reduces frontal but not parietal visual P3 amplitude. *Biol Psychol* 2000;52:37–52. [PubMed: 10686371]
- Propping P, Kruger J, Mark N. Genetic disposition to alcoholism. An EEG study in alcoholics and their relatives. *Hum Genet* 1981;59:51–9. [PubMed: 10819022]
- Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL. A default mode of brain function. *Proc Natl Acad Sci U S A* 2001;98:676–82. [PubMed: 11209064]
- Raichle ME, Snyder AZ. A default mode of brain function: a brief history of an evolving idea. *Neuroimage* 2007;37:1083–90. discussion 1097–9. [PubMed: 17719799]
- Rangaswamy M, Jones KA, Porjesz B, Chorlian DB, Padmanabhapillai A, Kamarajan C, Kuperman S, Rohrbaugh J, O'Connor SJ, Bauer LO, Schuckit MA, Begleiter H. Delta and theta oscillations as risk markers in adolescent offspring of alcoholics. *Int J Psychophysiol* 2007;63:3–15. [PubMed: 17129626]
- Rangaswamy M, Porjesz B, Ardekani BA, Choi SJ, Tanabe JL, Lim KO, Begleiter H. A functional MRI study of visual oddball: evidence for frontoparietal dysfunction in subjects at risk for alcoholism. *Neuroimage* 2004a;21:329–39. [PubMed: 14741671]
- Rangaswamy M, Porjesz B, Chorlian DB, Wang K, Jones KA, Bauer LO, Rohrbaugh J, O'Connor SJ, Kuperman S, Reich T, Begleiter H. Beta power in the EEG of alcoholics. *Biol Psychiatry* 2002;52:831–42. [PubMed: 12372655]
- Rangaswamy M, Porjesz B, Chorlian DB, Wang K, Jones KA, Kuperman S, Rohrbaugh J, O'Connor SJ, Bauer LO, Reich T, Begleiter H. Resting EEG in offspring of male alcoholics: beta frequencies. *Int J Psychophysiol* 2004b;51:239–51. [PubMed: 14962576]
- Rangaswamy M, Porjesz B, Chorlian DB, Choi K, Jones KA, Wang K, Rohrbaugh J, O'Connor S, Kuperman S, Reich T, Begleiter H. Theta power in the EEG of alcoholics. *Alcohol Clin Exp Res* 2003;27:607–15. [PubMed: 12711923]

- Sarnthein J, Jeanmonod D. High thalamocortical theta coherence in patients with Parkinson's disease. *J Neurosci* 2007;27:124–31. [PubMed: 17202479]
- Sarnthein J, Morel A, von Stein A, Jeanmonod D. Thalamocortical theta coherence in neurological patients at rest and during a working memory task. *Int J Psychophysiol* 2005;57:87–96. [PubMed: 15982767]
- Schoepp DD. Unveiling the functions of presynaptic metabotropic glutamate receptors in the central nervous system. *J Pharmacol Exp Ther* 2001;299:12–20. [PubMed: 11561058]
- Schurmann M, Basar-Eroglu C, Basar E. Gamma responses in the EEG: elementary signals with multiple functional correlates. *Neuroreport* 1997;8:1793–6. [PubMed: 9189935]
- Schurmann M, Basar-Eroglu C, Kolev V, Basar E. A new metric for analyzing single-trial event-related potentials (ERPs): application to human visual P300 delta response. *Neurosci Lett* 1995;197:167–70. [PubMed: 8552290]
- Schurmann M, Basar-Eroglu C, Kolev V, Basar E. Delta responses and cognitive processing: single-trial evaluations of human visual P300. *Int J Psychophysiol* 2001;39:229–39. [PubMed: 11163900]
- Seeger T, Fedorova I, Zheng F, Miyakawa T, Koustova E, Gomeza J, Basile AS, Alzheimer C, Wess J. M2 muscarinic acetylcholine receptor knockout mice show deficits in behavioral flexibility, working memory, and hippocampal plasticity. *J Neurosci* 2004;24:10117–27. [PubMed: 15537882]
- Shigemoto R, Kinoshita A, Wada E, Nomura S, Ohishi H, Takada M, Flor PJ, Neki A, Abe T, Nakanishi S, Mizuno N. Differential presynaptic localization of metabotropic glutamate receptor subtypes in the rat hippocampus. *J Neurosci* 1997;17:7503–22. [PubMed: 9295396]
- Soyka M, Preuss UW, Hesselbrock V, Zill P, Koller G, Bondy B. GABA-A2 receptor subunit gene (GABRA2) polymorphisms and risk for alcohol dependence. *J Psychiatr Res* 2008;42:184–191. [PubMed: 17207817]
- Spanagel R, Heilig M. Addiction and its brain science. *Addiction* 2005;100:1813–22. [PubMed: 16367982]
- Srinivasan R, Nunez PL, Silberstein RB. Spatial filtering and neocortical dynamics: estimates of EEG coherence. *IEEE Trans Biomed Eng* 1998;45:814–26. [PubMed: 9644890]
- Stassen HH, Lykken DT, Propping P, Bomben G. Genetic determination of the human EEG. Survey of recent results on twins reared together and apart. *Hum Genet* 1988;80:165–76. [PubMed: 3169741]
- Steriade M. Impact of network activities on neuronal properties in corticothalamic systems. *J Neurophysiol* 2001;86:1–39. [PubMed: 11431485]
- Steriade M. Grouping of brain rhythms in corticothalamic systems. *Neuroscience* 2006;137:1087–106. [PubMed: 16343791]
- Stockwell RG, Mansinha L, Lowe RP. Localization of the complex spectrum: The S-transform. *IEEE Trans on Sig Proc* 1996;44:998–1001.
- Tallon-Baudry C, Kreiter A, Bertrand O. Sustained and transient oscillatory responses in the gamma and beta bands in a visual short-term memory task in humans. *Vis Neurosci* 1999;16:449–59. [PubMed: 10349966]
- Tian H, Chen HJ, Cross TH, Edenberg HJ. Alternative splicing and promoter use in the human GABRA2 gene. *Brain Res Mol Brain Res* 2005;137:174–83. [PubMed: 15950776]
- Tiesinga PH, Fellous JM, Jose JV, Sejnowski TJ. Computational model of carbachol-induced delta, theta, and gamma oscillations in the hippocampus. *Hippocampus* 2001;11:251–74. [PubMed: 11769308]
- Tobler I, Kopp C, Deboer T, Rudolph U. Diazepam-induced changes in sleep: role of the alpha 1 GABA (A) receptor subtype. *Proc Natl Acad Sci U S A* 2001;98:6464–9. [PubMed: 11353839]
- Tomberg C, Desmedt JE. The challenge of non-invasive cognitive physiology of the human brain: how to negotiate the irrelevant background noise without spoiling the recorded data through electronic averaging. *Philos Trans R Soc Lond B Biol Sci* 1999;354:1295–305. [PubMed: 10466152]
- Tsai G, Coyle JT. The role of glutamatergic neurotransmission in the pathophysiology of alcoholism. *Annu Rev Med* 1998;49:173–84. [PubMed: 9509257]
- Tsuang MT, Faraone SV. The frustrating search for schizophrenia genes. *Am J Med Genet* 2000;97:1–3. [PubMed: 10813798]

- Tzschentke TM, Schmidt WJ. Functional relationship among medial prefrontal cortex, nucleus accumbens, and ventral tegmental area in locomotion and reward. *Crit Rev Neurobiol* 2000;14:131–42. [PubMed: 11513242]
- van Baal GC, de Geus EJ, Boomsma DI. Genetic influences on EEG coherence in 5-year-old twins. *Behav Genet* 1998;28:9–19. [PubMed: 9573642]
- van Beijsterveldt CE, Boomsma DI. Genetics of the human electroencephalogram (EEG) and event-related brain potentials (ERPs): a review. *Hum Genet* 1994;94:319–30. [PubMed: 7927323]
- van Beijsterveldt CE, Molenaar PC, de Geus EJ, Boomsma DI. Heritability of human brain functioning as assessed by electroencephalography. *Am J Hum Genet* 1996;58:562–73. [PubMed: 8644716]
- van Beijsterveldt CE, Molenaar PC, de Geus EJ, Boomsma DI. Genetic and environmental influences on EEG coherence. *Behav Genet* 1998;28:443–53. [PubMed: 9926613]
- van der Stelt O, Belger A. Application of electroencephalography to the study of cognitive and brain functions in schizophrenia. *Schizophr Bull* 2007;33:955–70. [PubMed: 17363840]
- Varela F, Lachaux JP, Rodriguez E, Martinerie J. The brainweb: phase synchronization and large-scale integration. *Nat Rev Neurosci* 2001;2:229–39. [PubMed: 11283746]
- Volkow ND, Wang GJ, Begleiter H, Hitzemann R, Pappas N, Burr G, Pascani K, Wong C, Fowler JS, Wolf AP. Regional brain metabolic response to lorazepam in subjects at risk for alcoholism. *Alcohol Clin Exp Res* 1995;19:510–6. [PubMed: 7625590]
- Wang JC, Hinrichs AL, Stock H, Budde J, Allen R, Bertelsen S, Kwon JM, Wu W, Dick DM, Rice J, Jones K, Nurnberger JI Jr, Tischfield J, Porjesz B, Edenberg HJ, Hesselbrock V, Crowe R, Schuckit M, Begleiter H, Reich T, Goate AM, Bierut LJ. Evidence of common and specific genetic effects: association of the muscarinic acetylcholine receptor M2 (CHRM2) gene with alcohol dependence and major depressive syndrome. *Hum Mol Genet* 2004;13:1903–11. [PubMed: 15229186]
- Whittington MA, Traub RD, Kopell N, Ermentrout B, Buhl EH. Inhibition-based rhythms: experimental and mathematical observations on network dynamics. *Int J Psychophysiol* 2000;38:315–36. [PubMed: 11102670]
- Winterer G, Enoch MA, White KV, Saylan M, Coppola R, Goldman D. EEG phenotype in alcoholism: increased coherence in the depressive subtype. *Acta Psychiatr Scand* 2003a;108:51–60. [PubMed: 12807377]
- Winterer G, Kloppel B, Heinz A, Ziller M, Dufeu P, Schmidt LG, Herrmann WM. Quantitative EEG (QEEG) predicts relapse in patients with chronic alcoholism and points to a frontally pronounced cerebral disturbance. *Psychiatry Res* 1998;78:101–13. [PubMed: 9579706]
- Winterer G, Mahlberg R, Smolka MN, Samochowiec J, Ziller M, Rommelspacher HP, Herrmann WM, Schmidt LG, Sander T. Association analysis of exonic variants of the GABA(B)-receptor gene and alpha electroencephalogram voltage in normal subjects and alcohol-dependent patients. *Behav Genet* 2003b;33:7–15. [PubMed: 12645817]
- Winterer G, Smolka M, Samochowiec J, Ziller M, Mahlberg R, Gallinat J, Rommelspacher HP, Herrmann WM, Sander T. Association of EEG coherence and an exonic GABA(B)R1 gene polymorphism. *Am J Med Genet B Neuropsychiatr Genet* 2003c;117:51–6.
- Yordanova J, Banaschewski T, Kolev V, Woerner W, Rothenberger A. Abnormal early stages of task stimulus processing in children with attention-deficit hyperactivity disorder--evidence from event-related gamma oscillations. *Clin Neurophysiol* 2001;112:1096–108. [PubMed: 11377270]
- Yordanova J, Kolev V. Brain theta response predicts P300 latency in children. *Neuroreport* 1996;8:277–80. [PubMed: 9051795]

Figure 1. From indices of cognition to genes – An illustration of our strategy
Brain oscillations, such as theta (θ , 3–7 Hz) EROs underlying P3, are correlates of impaired cognitive brain processes in alcoholism and risk, as seen in this time-frequency representation of EEG response to target stimuli in a visual oddball task. Head plot displays differences in ERO power and topography between alcoholics and controls.
Genetic Linkage analysis scans the entire genome to assess if a polymorphic genetic marker from a chromosomal region can be linked to a quantitative trait (theta ERO power) within a family. This is a logarithm of odds (LOD) score plot of linkage for theta ERO power at Fz, Cz and Pz electrodes on chromosome 7 indicating 2 candidate genes under the linkage peak (*CHRM2*, *GRM8*).

Candidate gene studies focus on genes underlying significant linkage peaks and/or genes with relevant biological significance. This is a schematic diagram of the *CHRM2* gene indicating its coding region (gray) and exons.

Genetic Association tests are performed for each single-nucleotide polymorphism (SNP) in the candidate gene and the trait variable. SNPs are identified within and flanking the candidate *CHRM2* gene (exons and introns). Significant associations (red dots) of the trait (theta ERO power) with these SNPs indicate loci of interest.

Figure 2. Interhemispheric high theta (6–7 Hz) coherence in resting EEG

a) Bars display coherence mean and standard error at 6 interhemispheric pairs along anterior to posterior axis depicted in the head plot inset **(b)**. Alcoholics (120, blue bars) manifest significant increases in resting EEG interhemispheric high theta coherence when compared to normal controls (120, green bars) [$F = 2.59$, $p = 0.019$; MANOVA [SPSS 12.0] (Rangaswamy, Chorlian et al., in preparation).

Figure 3. S-transform derived time-frequency representations of the average instantaneous amplitude that are z-scored for each frequency

Instantaneous amplitudes were averaged across individual trials of subjects so that non-phase locked or imprecise phase locked oscillatory energy is preserved.

Top panel: Plots for the target condition at Cz electrode in 120 alcoholic (right) and 120 control (left) subjects. Headplots of the time-frequency regions of interest (TFROIs) indicate that the theta band (4–5 Hz) power at 300–500 ms shows frontal maxima while the delta band (1–3 Hz) power at 350–700 ms has posterior maxima. Note that alcoholics have weaker responses in both theta and delta bands.

Bottom panel: Plots for the target condition at Fz electrode in visual oddball task in 87 high risk (right) and 57 matched low risk (left) adolescent subjects. Head plots for the TFROI that extends from 300 to 700 milliseconds for each band revealing frontal maxima for theta band power and parietal maxima for delta band power. Note that high risk adolescents have weaker responses in both theta and delta bands to target stimuli, similar to the alcoholics.

Figure 4. Linkage plot showing maximum LOD scores with significant linkage peaks over the GABA_A receptor gene cluster on chromosome 4

(a) Red trace: Resting EEG beta band power – beta 2 (16.5–20 Hz). The dataset consists of 1553 individuals from 250 families. (b) Green trace: Resting EEG high theta band (6–7 Hz) interhemispheric coherence at parieto-occipital bipolar pairs of electrodes (P4_O2-P3_O1). The dataset consists of 1312 individuals from 251 families.

Figure 5. Linkage plot showing maximum LOD scores with significant linkage peaks on chromosome 7, over the region harboring two candidate genes: a cholinergic muscarinic receptor gene (*CHRM2*) and a glutamate receptor gene (*GRM8*)

(a) Green trace: the central midline theta (4–5 Hz) ERO band power (between 300–700 ms, P3 latency window for visual target case) on chromosome 7. The dataset consists of 1337 individuals from 253 families; (b) Blue trace: Resting EEG theta band (6–7 Hz) interhemispheric coherence at centro-parietal bipolar pairs of electrodes (C4_P4-C3_P3) on chromosome 7. The dataset consists of 1312 individuals from 251 families.

Table 1

Brain oscillations as neurophysiological phenotypes: Comparing alcoholics and controls and high risk offspring and low risk controls

Neurophysiological Phenotype	Results	Citation
<u>Resting EEG</u>		
Theta Power (3–7 Hz)	Increased theta power in Alcohol Dependent subjects when compared to Controls	Rangaswamy et al. (2003)
Beta 1&2 Power (12–20 Hz)	Increased Beta power in Alcohol Dependent subjects when compared to Controls.	Rangaswamy et al. (2002)
	Increased Beta power in male high risk subjects (12–16 Hz) and female high risk subjects (16–20 Hz).	Rangaswamy et al. (2004b)
Bipolar Theta Coherence (4–7 Hz)	Alcohol dependent subjects have significantly increased posterior interhemispheric theta band coherence when compared to Controls.	Rangaswamy, Chorlian, et al. (in preparation)
<u>Visual Oddball Task (VP3)</u>		
Theta 4–7 Hz ERO 300–500 ms	Decreased Theta power during target processing in Alcohol dependent and High risk subjects.	Jones et al. (2006b) Rangaswamy et al. (2007)
Delta 1–3 Hz ERO 300–700 ms	Decreased Delta power during target processing in Alcohol dependent and High risk subjects	Jones et al. 2006 Rangaswamy et al. 2007
Gamma 29–45 Hz ERO 0–150 ms	Decreased gamma power during target processing in Alcohol dependent and High risk subjects	Padmanabhapillai et al. (2006a, b)
<u>Gambling Task</u>		
Theta Band ERO	Decreased Theta power during outcome processing in Alcohol dependent individuals.	Kamarajan et al. (in preparation)
<u>Response Inhibition (Go/NoGo)</u>		
Delta, Theta, Alpha Band ERO	Alcohol dependent individuals had decreased delta and theta ERO power during NoGo trials. This reduction was prominent frontally.	Kamarajan et al. (2004)
	High risk subjects had significantly decreased delta, theta ERO power	Kamarajan et al. (2006)

Neurophysiological Phenotype	Results	Citation
	for Go and NoGo and an additionally decreased alpha1 (8–9 Hz) ERO power for the NoGo.	

Table 2

Summary of association results from the quantitative pedigree disequilibrium test (QPDT) for SNPs in the candidate gene *GABRA2*.

	<i>GABRA2</i>	
	Theta Coherence	EEG Beta Power
3'	rs490434	rs490434
	rs576666	rs576666
	rs531460	rs531460
	rs561779	rs561779
	rs495818	rs495818
	rs497068	rs497068
	rs572227	rs572227
3'UTR	rs573400	rs573400
Intron 9	rs541418	rs541418
	rs481311	rs481311
	rs507788	rs507788
	rs532780	rs532780
	rs548583	rs548583
	rs10938435	rs10938435
Intron 8	rs496650	rs496650
	rs540363	rs540363
	rs526752	rs526752
	rs530329	rs530329
	rs483160	rs483160
Intron 7	rs279871	rs279871
Intron 6	rs279869	rs279869
	rs279867	rs279867
	rs279866	rs279866
	rs1808851	rs1808851
Intron 5	rs279863	rs279863
	rs279861	rs279861
Exon 5	rs279858	rs279858
Intron 4	rs175931	rs175931
	rs279843	rs279843
	rs279845	rs279845
	rs279846	rs279846
	rs183961	rs183961
	rs1440130	rs1440130
	rs279826	rs279826
	rs11503016	rs11503016

	<i>GABRA2</i>	
	Theta Coherence	EEG Beta Power
Intron 3	rs279827	rs279827
	rs279828	rs279828
	rs279834	rs279834
	rs279836	rs279836
	rs279837	rs279837
	rs279841	rs279841
	rs189957	rs189957
	rs1442059	rs1442059
	rs1442061	rs1442061
	rs1442062	rs1442062
	rs11503015	rs11503015
Intron 0	rs11503014	rs11503014
	rs3756007	rs3756007
5'	rs894269	rs894269
	rs1372472	rs1372472
	rs2165607	rs2165607
	rs1545234	rs1545234

SNPs showing significant associations for the two endophenotypes (theta coherence and EEG beta power) are indicated in bold red font.

Table 3

Summary of association results from the quantitative pedigree disequilibrium test (QPDT) for SNPs in the candidate gene *CHRM2* SNPs.

	<i>CHRM2</i>		
	Theta ERO	Delta ERO	Theta Coherence
upstream of exon 1	rs1424558	rs1424558	rs1424558
	rs1424574	rs1424574	rs1424574
	rs13247260	rs13247260	rs13247260
intron 3-4	rs1424569	rs1424569	rs1424569
	rs1424387	rs1424387	rs1424387
	rs2350780	rs2350780	rs2350780
	rs978437	rs978437	rs978437
	cc785	cc785	cc785
	cc1218	cc1218	cc1218
	rs7782965	rs7782965	rs7782965
	rs7800170	rs7800170	rs7800170
	rs1455858	rs1455858	rs1455858
	rs1378646	rs1378646	rs1378646
intron 4-5	rs1824024	rs1824024	rs1824024
	rs2061174	rs2061174	rs2061174
	rs7799047	rs7799047	rs7799047
	rs2350786	rs2350786	rs2350786
exon 5	chr2ex5	chr2ex5	chr2ex5
intron 5-6	rs6948054	rs6948054	rs6948054
	rs324640	rs324640	rs324640
	rs324650	rs324650	rs324650
	rs324651	rs324651	rs324651
3'UTR	rs8191992	rs8191992	rs8191992
	rs8191993	rs8191993	rs8191993
downstream of exon 6	rs1378650	rs1378650	rs1378650
	rs1424548	rs1424548	rs1424548
	rs324656	rs324656	rs324656

SNPs showing significant associations for the three endophenotypes (theta ERO, delta ERO and theta coherence) are indicated in bold red font.