Fig 1 | Rapid NF- κ B activation by *Helicobacter pylori* is type IV secretion system (T4SS)-dependent and CagA-independent. AGS cells were infected (MOI of 100) with the P1 *H. pylori* wild-type (wt) strain, the CagA-deficient (*cagA*) or T4SS-deficient (*virB7*) isogenic mutant strains. At the indicated times after infection, whole cell lysates were prepared and analysed by SDS-PAGE and western blotting. For immunodetection, protein-specific or phosphoprotein-specific antibodies (phospho-I κ B α Ser32/36; phospho-p65 Ser536) were used, as indicated.

In our opinion, a single CagA-deficient *H. pylori* strain that is able to rapidly activate NF- κ B with a potency similar to the wild-type strain represents strong evidence against an essential requirement for CagA, irrespective of the genetic background of the strain. Although few studies suggest a contributory role of CagA in late *H. pylori*-induced NF- κ B activation (for example, Suzuki *et al* 2009), these studies address long-term

effects involving indirect processes, but not direct NF- κ B activation within the first hour of infection.

As a possible explanation for the conflicting data, we would like to refer to the publication by Fischer *et al* (2001). Therein, the authors stated that great care needs to be taken in the generation and propagation of *H. pylori* isogenic mutant strains to prevent accidental functional inactivation of the

PAI concomitant with the deletion of single genes encoded in it: a phenomenon named 'polar effect'. Functional inactivation of the PAI due to the 'polar effect' has been documented in the literature (Naumann *et al* 1999). Thus, complementation of the *cagA*-deficient *H. pylori* strains with *cagA* could have unequivocally proven the suggested function of CagA in NF- κ B activation.

We think our comment offers fair, putative reasons for the conflicting data and helps to diminish uncertainty in the scientific community.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

REFERENCES

- Censini S *et al* (1996) *Proc Natl Acad Sci USA* **93**: 14648–14653
 Fischer W *et al* (2001) *Mol Microbiol* **42**: 1337–1348
 Forst-Ludwig A *et al* (2004) *Biochem Biophys Res Commun* **316**: 1065–1072
 Lamb A *et al* (2009) *EMBO Rep* **10**: 1242–1249
 Naumann M *et al* (1999) *J Biol Chem* **274**: 31655–31662
 Neu B *et al* (2002) *Am J Gastrointest Liver Physiol* **283**: G309–G318
 Sharma SA *et al* (1995) *Infect Immun* **63**: 1681–1687
 Suzuki M *et al* (2009) *Cell Host Microbe* **5**: 23–34
 Katrin Schweitzer, Olga Sokolova, Przemyslaw M. Bozko & Michael Naumann are at the Institute of Experimental Internal Medicine, Otto von Guericke University Magdeburg, Germany.
 E-mail: naumann@med.ovgu.de
 EMBO reports (2010) **11**, 10–11.
 doi:10.1038/embor.2009.263

Response: strain-specific activation of NF- κ B by *Helicobacter pylori* CagA

We appreciate the comments from Schweitzer *et al* in this issue about our discovery that *Helicobacter pylori* cytotoxin-associated gene A (CagA) activates nuclear factor κ B (NF- κ B), as they reveal the lack of consensus on the role that CagA plays during *H. pylori* infection.

We are confident in our data for several reasons. First, we demonstrated in our studies that three independent *H. pylori* *cagA*-positive strains, but not their respective isogenic *cagA* mutants, activate NF- κ B. Second, because *cagA* is monocistronic and transcribed in a different direction than other genes comprising the *cag* pathogenicity

island (*cagPAI*; Censini *et al*, 1996), it is unlikely that our *H. pylori* isogenic *cagA* mutant strains are exerting "polar effects" on other genes within the *cagPAI*. Third, these strains have been used extensively by other groups without apparent evidence of potential polar effects due to disruption of *cagA* (for example, Brandt *et al*, 2005; Franco *et al*, 2005; Guillemin *et al*, 2002). Fourth, while many studies demonstrate the importance of the *cagPAI*-encoded type 4 secretion system (T4SS) in *H. pylori*-induced pathogenesis and inflammation (Glocker *et al*, 1998), CagA is injected into host epithelial cells via the T4SS (Odenbreit *et al*, 2000), and studies that show defects in the ability of

T4SS-deficient bacteria to activate NF- κ B cannot completely exclude the importance of CagA. Lastly, the demonstration that CagA is important for the activation of NF- κ B is not entirely unprecedented. Using an interleukin (IL) 8 promoter-reporter assay, Sharma *et al* (1998) showed the requirement of CagA for NF- κ B activation. Brandt *et al* (2005) demonstrated that ectopically expressed CagA induces the translocation of NF- κ B from the cytoplasm into the nucleus, and, in accord with a study from Kim *et al* (2006), showed that ectopic CagA expression induces IL-8 secretion from gastric epithelial cells. Furthermore, Shibata *et al* found that Mongolian gerbils infected with *cagA*-deficient *H. pylori* showed significantly decreased NF- κ B activation and NF- κ B-linked inflammation in the gastric antra of infected gerbils (2006). These data,

in conjunction with our results, demonstrate that in certain *H. pylori* strains, CagA plays an essential role in the activation of NF- κ B.

We acknowledge in our article that CagA may not be the only component that *H. pylori* uses to activate inflammation. Compelling evidence exists that under some circumstances other *H. pylori* factors can induce an inflammatory response. For example, CagA clearly does not seem to be essential for activation of NF- κ B in HEK293 cells or macrophages (Maeda *et al*, 2001; Viala *et al*, 2004). Previous studies have also convincingly demonstrated that among populations of *H. pylori* cagA-positive strains, the ability of CagA to activate NF- κ B is strain-specific (Brandt *et al*, 2005). Thus, the accompanying remarks from Schweitzer *et al* underscore the importance of considering the experimental model when studying *H. pylori*-mediated inflammation. The goal of our work was to decipher the molecular mechanism for the activation of NF- κ B by *H. pylori*. In that

context, our study provides new insight into the pathogen-effector-induced activation of the inflammatory response.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest

REFERENCES

- Brandt S *et al* (2005) *Proc Natl Acad Sci USA* **102**: 9300–9305
 Censini S *et al* (1996) *Proc Natl Acad Sci USA* **93**: 14648–14653
 Franco AT *et al* (2005) *Proc Natl Acad Sci USA* **102**: 10646–10651
 Glocker E *et al* (1998) *Infect Immun* **66**: 2346–2348
 Guillemin K *et al* (2002) *Proc Natl Acad Sci USA* **99**: 15136–15141
 Kim SY *et al* (2006) *Cell Microbiol* **8**: 97–106
 Maeda S *et al* (2001) *J Biol Chem* **276**: 44856–44864
 Odenbreit S *et al* (2000) *Science* **287**(5457): 1497–1500
 Sharma SA *et al* (1998) *J Immunol* **160**: 2401–2407
 Shibata W *et al* (2006) *J Pathol* **210**: 306–314
 Viala J *et al* (2004) *Nat Immunol* **5**: 1166–1174

Acacia Lamb is at the Department of Biochemistry, College of Medicine, University

of Illinois at Urbana-Champaign, Urbana, IL, USA.

Jiang-Dong Li is at the Department of Microbiology and Immunology, University of Rochester Medical Center, Rochester, NY, USA.

Masanori Hatakeyama is at the Department of Microbiology, University of Tokyo, Tokyo, Japan.

Richard M. Peek is at the Department of Medicine, Division of Gastroenterology, Vanderbilt University School of Medicine, Nashville, TN, USA.

Steven R. Blanke is in the Department of Microbiology, Institute for Genomic Biology, University of Illinois at Urbana-Champaign, Urbana, IL, USA.

Lin-Feng Chen is also at the Department of Biochemistry, University of Illinois at Urbana-Champaign.

E-mail: lfchen@life.uiuc.edu

EMBO reports (2010) **11**, 11–12.

doi:10.1038/embo.2009.269

The Bronze Age of science in Spain

In 2008, the Spanish government promised a new Silver Age of science that would comprise the creation of a new Ministry of Science together with major, long-awaited investments in research and development (Editorial, 2008). Unfortunately, those expectations have not translated into reality. The recently tabled 2010 national budget proposes that funding for the Ministry of Science should be cut by more than 15%, thus returning to 2006 levels. In fact, the situation will be even more serious given that the new Ministry also comprises the Health Institutes, which were previously the oversight of the Ministry of Health. Only the funding of fellowships for education at all levels will be retained. Unlike the somewhat positive outlook that Michela Bertero described in her recent *EMBO reports* article about science in Spain (Bertero, 2009), the reality is likely to be rather different, at least for the next few years.

The Ministry of Science is the main source of funding for research in Spain, as there is very little investment available from private sources or foundations. The rationale behind the serious decrease in public funds is a programme of austerity in response to the financial crisis. The Ministry of Education, which is responsible for universities and

their research, has already declared that it will reduce funding for the maintenance of laboratories and for attracting young scientists from abroad (Díez & Riviera, 2009). As an example of what is to come next year, the Spanish Research Council (www.csic.es), which depends on funding from the Ministry of Science, has reduced posts by more than 50% compared with 2008, whereas the CNRS (www.cnrs.fr)—the French equivalent—has continued to offer a large number of positions during this year. The president and former presidents of the Spanish Society of Molecular Biology have signed a manifesto to protest against this massive decrease in funding. At the same time, the Spanish Prime Minister, José Luis Rodríguez Zapatero, has claimed that the economic crisis will force Spain to develop a more competitive and innovative economy based on knowledge, so that Spain, like other European countries, can emerge from the financial turmoil stronger and better prepared for future crises.

Spain has long lagged behind similar European countries in terms of scientific research. With the exception of a few scientists from the Spanish Golden Age from the fifteenth to seventeenth centuries, and a handful from the eighteenth century, Spain lacks the long and fertile

scientific tradition evident in much of central Europe. Even in the twentieth century, the only well-known Spanish scientist was Nobel laureate Santiago Ramón y Cajal. Some historians attribute this lack of Spanish scientific pedigree to the influence that the Catholic Church exercised during the past centuries, while others note the lack of a true democratic Spanish society until the second half of the twentieth century (Sagan, 1995). Yet others, including a number of renowned Spanish philosophers, think that science is simply a strange concept to many Spaniards; the scientific method is five centuries old and was developed in countries that tended to be the enemies of Spain—essentially it ‘hasn’t caught on’ in Spanish society. Nevertheless, the Spanish people cannot afford for their government to waste more opportunities to invest in science.

In times of crisis and economic difficulty, the best strategy for recovery is to bolster investment in research and education—an area where Spain is also at the bottom of the rankings of West European countries—without leaving more basic social necessities unattended. Japan, Finland and South Korea invested heavily in science during the 1990s when their economies were floundering, and the US administration has committed an extraordinary amount of money to basic research with the argument that scientists