Published in final edited form as:

J Org Chem. 2008 May 16; 73(10): 3885. doi:10.1021/jo800383e.

Functionalization of Organotrifluoroborates: Reductive Amination

Gary A. Molander and David J. Cooper

Roy and Diana Vagelos Laboratories, Department of Chemistry, University of Pennsylvania, Philadelphia, Pennsylvania 19104-6323

Gary A. Molander: gmolandr@sas.upenn.edu; David J. Cooper:

Abstract

Herein we report the conversion of aldehyde-containing potassium and tetrabutylammonium organotrifluoroborates to the corresponding amines through reductive amination protocols. Potassium formate facilitated by catalytic palladium acetate, sodium triacetoxyborohydride, and pyridine borane have all served as effective hydride donors, reducing the initially formed imines or iminium ions to provide the corresponding amines.

Introduction

Owing to the existence of an empty p-orbital on boron, boronic acids and, to a lesser extent, boronate esters exhibit sensitivity to several classes of useful reagents commonly employed during the course of organic synthesis. Their reactivity toward oxidants, bases, nucleophiles, and organic acids limits the conditions to which these organoboron species can be exposed, and therefore to a great extent this dictates at which point the boron must be introduced during a synthesis campaign. Organotrifluoroborates, which lack an empty p-orbital and therefore are mechanistically inhibited from reacting through modes normally observed for tricoordinate organoboron compounds, fulfill all of the criteria needed for a protected boronic acid with little cost and substantial benefits.¹ Thus, the tetracoordinate nature of these species, fortified with exceptionally strong boron-fluorine bonds, provides extraordinary opportunities for manipulation of ancillary functional groups with retention of the trifluoroborate, allowing one to make or purchase a very simple organoboron compound and build molecular complexity into the molecule from the ground up. The full development of organotrifluoroborates has the

Correspondence to: Gary A. Molander, gmolandr@sas.upenn.edu.

Supporting Information **Available**. Full experimental details and copies of all pertinent NMR spectra. This material is available free of charge via the Internet at <http://pubs.acs.org>.

potential to liberate strategic planning in the construction of complex molecules as well as the design of scaffolds for diversity-oriented synthesis. As previously reported, we have demonstrated that alkene ozonolysis, Wittig reactions, dihydroxylations, and epoxidation reactions can be performed in the presence of the organotrifluoroborate unit, creating a novel entry to unique and potentially valuable organic synthons.ⁱⁱ Herein we describe the reductive aminationⁱⁱⁱ of aldehyde-containing potassium- and tetrabutylammonium organotrifluoroborates, which not only demonstrates the resistance of the carbon-boron bond to a variety of aggressive reaction conditions, but also provides a unique access to amine-substituted organotrifluoroborates.

Results and Discussion

During our investigation of methods to access tetrabutylammonium 3-trifluoroboropropanal **1** by an ozonolysis protocol it was discovered that the remarkably stable secondary ozonide **2**, formed by treatment of tetrabutylammonium 3-butenyltrifluoroborate **3** with ozone, could be treated with an amine base to generate the desired aldehyde product **1** (eq 1).^{2e,iv}

Interestingly, while the desired product was obtained in pure form when pyridine was used as the base, it was also observed that if a secondary amine was used instead of pyridine, the corresponding tertiary amine **4** was obtained in addition to the desired aldehyde (Scheme 1). We rationalized that a process closely resembling the Leuckart reaction occurred, whereby ammonium formate was liberated during the course of the reaction and served as a reducing agent for a reductive amination process.^v Although this reaction gave an inseparable mixture of the aldehyde and the amine as products, it foretold that conditions could be developed to access valuable amine-containing organoboron coupling partners in pure form.

There is precedent for the reductive amination of aldehyde-containing boronic acids and in particular 2-formylarylboronic acids and derivatives.^{vi} However, to our knowledge unless this boronic acid or structurally related *ortho*-substituted arylboronic acids are used, there exist few examples where the amine component is something other than an aniline derivative.^{vii} To fill this perceived void, a more general method for the reductive amination of aldehyde-containing organoborons was sought to provide access to nucleophilic cross-coupling partners incorporating various amine functional groups.

There was initially some concern that one of several different processes could result in unproductive side reactions resulting from reaction of the aldehyde-containing organotrifluoroborate with the amine. It was hoped that the tetracoordinate

organotrifluoroborates, in contrast to their tricoordinate boronic acid counterparts, would prevent coordination of the boron with the oxygen atom of the intermediate hemiaminal, a necessary initial step for subsequent bond formation in the Petasis boronic acid Mannich reaction (eq 2).^{viii} Although this and related types of transformations have been reported using organotrifluoroborates (eqs 3 and 4),^{ix} it has only occurred in the presence of strong fluorophiles that generate difluoroboranes in situ.^x Depending on the nature of the organotrifluoroborate and electrophile, several different mechanistic pathways have been proposed for these transformations,^{9a,b,f,g,¹¹} including the traditional Petasis boronic acid Mannich mechanism through an intermediate hemiaminal or amination (eqs 2 and 3)^{8,9b,xi} and direct intramolecular addition via an imine-complexed RBF₂ species (eq 4).^{9c}

(2)

(3)

(4)

More relevant to the current work, during the course of our studies MacMillan et al. demonstrated that potassium aryl- and heteroaryltrifluoroborates^{xii} participate as nucleophiles in amine-catalyzed 1,4-additions to enals without Lewis acid activation (although in the

presence of 1 equiv of HF) through a mechanism involving direct addition of the organotrifluoroborate to the intermediate conjugated iminium ions. This raised the possibility that related 1,2-additions might occur between organotrifluoroborates and iminium ions generated via Mannich-type reactions from the corresponding aldehydes (Scheme 2).

As it transpired none of these potentially deleterious issues were validated in our studies, as under the conditions developed there was no evidence of these alternative reaction pathways. Instead, when tetrabutylammonium 4-formylfuran-2-yl-trifluoroborate **5** was treated with piperidine and formic acid in the presence of 4Å molecular sieves a near quantitative yield of the corresponding iminium ion was isolated (eq 5).

(5)

Based on these initial observations, an in situ reductive amination was developed to provide the amine product **6** using potassium formate facilitated by a palladium catalyst (eq 6).^{xiii} This method was optimized for **5** and applied to a variety of aldehyde-containing trifluoroborates to give the corresponding piperidinyl methylamines (Table 1). To determine the scope of the method, aldehyde **7** was then transformed using two different secondary amines, providing the corresponding tertiary amines in excellent yields (Table 1, entries 5 and 6).

(6)

Unfortunately, this first reductive amination protocol suffered in several respects. Although it proved effective for access to pure amine-containing trifluoroborates in high yields when applied to the furanyl substrates **5** (83%) and 5-formylfuran-2-yl-trifluoroborate **7** (86%), the substrate scope was limited. When extended to a variety of activated and non-activated benzaldehydes, it gave either an inseparable impurity that appeared to derive from a side

reaction with DMF, or a mixture of the aldehyde starting material, the corresponding imine, and the amino product. This method was also determined to be effective almost exclusively for tetrabutylammonium trifluoroborate substrates, with only one of the potassium trifluoroborates surveyed, 4-formylfuran-2-yl-trifluoroborate **11** (74%), giving the desired product (Table 1, entry 4). Furthermore, following the survey of a variety of primary amines including aniline, benzylamine, *n*-butylamine, *tert*-butylamine, and cyclohexylamine, the method was determined to be ineffective in the installation of secondary amines into the trifluoroborate substrates examined. Thus, in an effort to expand this chemistry to include a range of potassium organotrifluoroborates and primary amines as effective partners, other protocols for the reductive amination were considered.

Sodium triacetoxyborohydride (STAB) seemed an obvious choice as a selective reducing agent. In addition to its high level of selectivity, reactions utilizing STAB with acetonitrile as a solvent have also been well developed.^{xiv} This was encouraging to us as we expected many of the iminium ion intermediates to be soluble in acetonitrile. With this in mind, a variety of conditions for the reductive amination of aldehyde **15** and cyclohexylamine using STAB were explored.

There was some anxiety concerning a single-step protocol, wherein the reduction took place on an imine or iminium ion derived in situ, because it could result in the formation of a mixture of amine and alcohol products.¹² Because such a mixture could prove difficult to separate, a two-step, one-pot reductive amination process was utilized whereby addition of the reducing agent was delayed until complete conversion to the imine or iminium ion had occurred. Thus, the overall conversion from aldehyde to amine was considered in two stages, examining first the conversion of the aldehyde to the corresponding imine. When **15** and cyclohexylamine were stirred with anhydrous methanol as a solvent, conversion to the imine was realized in 20 min without the use of acetic acid (eq 7). The second stage, reduction of the imine using STAB, was conducted in acetonitrile and proceeded smoothly using 1.4 equivalents of STAB and 1.1 equivalents of acetic acid at ambient temperature to give the desired methylaminofuran **16**. Although this new method did allow the reductive amination of the potassium salt **15**, the method demonstrated significant substrate dependence, and obtaining reasonable yields of pure material with other substrates proved difficult.

(7)

Other selective reducing agents were examined to solve this problem. Because borane amine complexes allow the use of methanol, a solvent that had proven very effective in the conversion of formyltrifluoroborates to the corresponding imines, and because these reducing agents demonstrate a high degree of selectivity for reduction of the iminium ion over the aldehyde,^{xv} use of these reagents would allow the entire sequence to be run in methanol.¹¹ After screening several boranes, including *tert*-butylamine borane, picoline borane, and triethylamine borane, pyridine borane (pyr•BH₃) was found to reduce the iminium ion generated from aldehyde **17** and cyclohexylamine most efficaciously. Furthermore, while acetic acid can be used in conjunction with these boranes to protonate the imine giving the

reactive iminium ion,^{xvi} inexpensive KHF_2 was used instead. This served double duty. Acting as an acid, the KHF_2 activated the imine. Additionally, the KHF_2 appeared to serve as a fluoride source, allowing removal of borane byproducts that otherwise contaminated the desired organotrifluoroborates.

A variety of amines were then used with **17** to determine the scope of this protocol (Table 2). We were delighted to discover that good to excellent yields could be achieved for a wide range of amines. Although application of this method to benzaldehyde **17** and *n*-butylamine gave a mixture of the aminomethyl product **18** and the corresponding internal ammonium salt, other primary amines, including cyclopropylamine and cyclohexylamine, gave the desired products in pure form (entries 5 and 7). Furthermore, the method was successfully applied to a variety of amine-containing heterocycles to give the corresponding tertiary amine products. Pyrrolidine, piperidine, morpholine, and even Boc-protected piperazine could be used to give the corresponding boron-containing amines in good yield and in high purity (Table 2, entries 2-4, 6).

Cyclohexylamine and piperidine were then used to show that the method could be applied to a variety of aryl- and heteroaryl-aldehydes (Table 3, entries 1-7). Although the majority of the aminomethyl-substituted organotrifluoroborates synthesized were easily purified by precipitation from acetone using diethyl ether, those derived from substrates with an ortho relationship between the aldehyde and trifluoroborate functional groups (**16**, **26**, and **35**), had to be further dried by azeotropically removing the water with toluene and placing the resulting powders under vacuum (0.01 torr).

Amino alcohols can be utilized as substrates in the reductive amination. Thus, it was shown that the method could be employed using the chiral *cis*-aminoindanols in reactions with aldehydes **15** and **31** (entries 8 and 9). Optimal experimental conditions in these two cases required that a single equivalent of *cis*-aminoindanol and no acid be used in the reaction mixture to obtain pure product. Thus, good yields of the chiral products **36** and **37** were obtained by the method which remained one-pot, but which was conducted in two steps to ensure that the aldehyde was completely converted the corresponding imine prior to introduction of pyridine borane.

Finally, we demonstrated that the amine-substituted organotrifluoroborates formed in these reductive aminations could be coupled in high yields to aryl halide partners. In an adaptation of conditions developed by our group, the coupled product was realized employing 3 mol % $\text{Pd}(\text{OAc})_2$ under ligandless conditions (eq 8).¹⁷ In this variation, K_2CO_3 (3.0 equiv) was utilized as the base and the reaction was performed in MeOH at 75 °C for 3 h. Under these conditions, no column chromatography was required and the coupled product was isolated in 82% yield.

(8)

In summary, protocols for the reductive amination of aldehyde-containing organotrifluoroborates using potassium formate facilitated by catalytic palladium acetate (Table 1), sodium triacetoxyborohydride (eq 2), and pyridine borane (Tables 2 and 3) have been developed. These methods offer access to valuable amine-containing organoborons and provide another example of the complementarity between organotrifluoroborates (which allow functionalization of ancillary carbonyls) and boronic acids. Use of organotrifluoroborates provides access to both trifluoroborato-substituted imines and iminium ions, where under similar reaction conditions boronic acids might be expected to undergo the Petasis boronic acid Mannich or related reactions. Reduction of the imine- and iminium ion-substituted organotrifluoroborates generated in situ affords the corresponding trifluoroborato-functionalized amines in a one-pot process. The amino-substituted organotrifluoroborates thus generated can be cross-coupled under ligandless conditions, affording high yields of a biaryl product wherein no chromatography is utilized at any point in the multi-step process.

Experimental

General Procedure for Preparation of Tetrabutylammonium Formylaryltrifluoroborates. Tetrabutylammonium 4-Formylfuran-2-yltrifluoroborate (5)

To a slurry of potassium 4-formylfuran-2-yltrifluoroborate (101 mg, 0.50 mmol) in H₂O (1 mL) and CH₂Cl₂ (5 mL) was added an aqueous solution of (*n*-Bu)₄NOH (0.32 mL, 0.50 mmol, 1.6 M in water) with stirring. The biphasic solution was stirred for 0.5 h. The layers were separated. The aqueous layer was extracted with CH₂Cl₂ (3 × 5 mL) after which the combined organic layers were washed with H₂O, dried (MgSO₄) and concentrated under high vacuum to give the product as a colorless oil (154 mg, 76%). IR = 1673 cm⁻¹; ¹H NMR (500 MHz, CDCl₃): δ 0.89 (t, *J* = 8 Hz, 12H), 1.25 – 1.36 (m, 8H), 1.45 – 1.55 (m, 8H), 3.08 (t, *J* = 9 Hz, 8H), 6.56 (s, 1H), 8.0 (s, 1H), 9.82 (s, 1H); ¹³C NMR (125.8 MHz, CDCl₃): δ 13.5, 19.5, 23.7, 58.5, 107.5, 129.1, 152.8, 185.5; ¹⁹F NMR (471 MHz, CDCl₃): δ -142.5; ¹¹B NMR (128 MHz, CDCl₃): δ -0.42; HRMS (ESI): calcd for C₅H₃BF₃O₂ [M-C₁₆H₃₆N]⁻ 163.0178, found 163.0222.

General Procedure A for the Reductive Amination of Tetrabutylammonium Formylaryltrifluoroborates using Palladium-Catalyzed Formate. Tetrabutylammonium 4-((Piperidin-1-yl)methyl)furan-2-yltrifluoroborate (6)

A test tube containing activated 4Å molecular sieves and tetrabutylammonium 4-formylfuran-2-yltrifluoroborate **5** was evacuated under vacuum (0.01 torr) for 15 min. DMF (1 mL), piperidine (60 mg, 0.71 mmol), and formic acid (28 mg, 0.61 mmol), were then added with stirring. The solution was heated under nitrogen for 5 h at 70 °C, after which potassium formate (100 mg, 1.19 mmol), and palladium acetate (5 mg, 0.02 mmol, 6 mol %) were added, and stirring was continued an additional 4 h. Heat was removed and the reaction was allowed to cool to rt. Aqueous KHF₂ (1 mL, 4.5M) was then added and the solution was stirred for 20 min. The mixture was filtered, rinsing the white solid with CH₂Cl₂ (2 × 5 mL). The filtrate was washed with H₂O, 3M NaOH, and brine, dried (MgSO₄), and the solvent was removed to give the product as a deep yellow oil (130 mg, 83%). IR = 2933, 1669 cm⁻¹; ¹H NMR (500 MHz, CDCl₃): δ 0.81 (t, *J* = 8 Hz, 12H), 1.17 – 1.26 (m, 10H), 1.35 – 1.44 (m, 12H), 2.23 (s, 4H), 2.98 (t, *J* = 9 Hz, 8H), 3.32 (s, 2H), 6.11 (s, 1H), 7.09 (s, 1H); ¹³C NMR (125.8 MHz, CDCl₃): δ 13.6, 19.5, 23.7, 24.2, 26.1, 53.9, 56.0, 58.3, 107.8, 111.7, 151.0; ¹⁹F NMR (471 MHz, CDCl₃): δ -141.2; ¹¹B NMR (128 MHz, CDCl₃): δ 0.164; HRMS (ESI): calcd for C₁₀H₁₄BF₃NO (M-C₁₆H₃₆N) 232.1121, found 232.1131.

General Procedure for Preparation of Potassium Formylaryltrifluoroborates. Potassium 4-Formylfuran-2-yltrifluoroborate (11)

To 4-formylfuran-2-ylboronic acid (5.0 g, 3.57 mmol) in MeOH (20 mL) was added dropwise a solution of KHF_2 (13.65 g, 0.175 mol) in H_2O (40 mL) using an addition funnel. The mixture was stirred for 2 h and then concentrated under high vacuum. The residual solids were extracted with 20% MeOH in acetone (4×10 mL). The combined extracts were concentrated close to the saturation point. Et_2O (20 mL) was then added until no more precipitation was observed. The solids were collected, washed with Et_2O (2×10 mL), and dried under high vacuum to afford the product as a white crystalline solid (4.78 g, 66%). mp >200 °C; IR (KBr) = 1687 cm^{-1} ; ^1H NMR (500 MHz, $\text{DMSO}-d_6$): δ 6.29 (d, $J = 20$ Hz, 1H), 8.34 (s, 1H), 9.81 (s, 1H); ^{13}C NMR (125.8 MHz, $\text{DMSO}-d_6$): δ 106.8, 128.9, 154.3, 186.3; ^{19}F NMR (471 MHz, $\text{DMSO}-d_6$): δ -140.5; ^{11}B NMR (128 MHz, $\text{DMSO}-d_6$): δ -0.225; HRMS (ESI): calcd for $\text{C}_5\text{H}_3\text{BF}_3\text{O}_2$ (M-K) 163.0178, found 163.0176.

Potassium 4-((Piperidin-1-yl)methyl)furan-2-yltrifluoroborate (12)

A test tube containing activated 4Å molecular sieves (400 mg) and potassium 4-formylfuran-2-yltrifluoroborate (192 mg, 0.95 mmol) was evacuated under vacuum (0.01 torr) for 15 min. DMF (1 mL), piperidine (212 mg, 2.49 mmol) and formic acid (69 mg, 1.5 mmol) were added with stirring, and the solution was then heated under nitrogen for 5 h at 70 °C. Palladium acetate (22 mg, 0.10 mmol, 10 mol %) and potassium formate (195 mg, 2.32 mmol) were then added and stirring was continued an additional 4 h. Heat was removed and the reaction mixture was filtered, rinsing the white solid with MeOH (3×5 mL). To the filtrate was added an aqueous solution of KHF_2 (1 mL, 4.4M) with stirring, which was continued for 1 h. The solution was then concentrated under vacuum. The resulting yellow solid was extracted with 10% MeOH/acetone (3×5 mL). This was concentrated and precipitated with Et_2O (10 mL) to give the title compound as a yellow solid (190 mg, 74%). mp >200 °C; IR = $2933, 1669\text{ cm}^{-1}$; ^1H NMR (500 MHz, acetone- d_6): δ 1.38 – 1.44 (m, 2H), 1.52 – 1.60 (m, 4H), 2.10 – 2.16 (m, 4H), 3.44 (s, 2H), 6.21 (s, 1H), 7.30 (s, 1H); ^{13}C NMR (125.8 MHz, acetone- d_6): δ 23.7, 25.1, 53.1, 53.1, 113.1, 140.8; ^{19}F NMR (471 MHz, acetone- d_6): δ -142.1; ^{11}B NMR (128 MHz, acetone- d_6): δ 0.662; HRMS (ESI): calcd for $\text{C}_{10}\text{H}_{14}\text{BF}_3\text{NO}$ (M-K) 232.1121, found 232.1129.

Potassium 3-((Cyclohexylamino)methyl)furan-2-yltrifluoroborate (16)

To a sealed test tube containing potassium 3-formylfuran-2-yltrifluoroborate (110 mg, 0.54 mmol) in MeOH (4 mL) was added cyclohexylamine (198 mg, 2.0 mmol) with stirring. Stirring was continued for 1 h, then the solvent was removed to give a yellow solid. This was taken up into acetonitrile (4 mL). Acetic acid (40 mg, 0.67 mmol) and sodium triacetoxyborohydride (150 mg, 0.71 mmol) were added and the test tube was resealed. This was stirred for 16 h at rt. KOH (168 mg, 1.5 mmol) was then added, followed by H_2O (2 mL). This was stirred for 20 min, then KHF_2 (421 mg, 5.4 mmol) was added and solution was stirred an additional 30 min. The solution was then concentrated under vacuum. The resulting yellow solid was extracted with 10% MeOH/acetone (3×5 mL). This was concentrated and precipitated with Et_2O (5 mL) to give the product as a yellow solid (135 mg, 87%). (mp >200 °C; IR (KBr) = 2939 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6): δ 1.06 – 1.31 (m, 5H), 1.60 (d, $J = 12$ Hz, 1H), 1.74 (dd, $J = 9, 4$ Hz, 2H), 1.99–2.19 (m, 3H), 2.83 – 2.87 (m, 1H), 3.95 (s, 2H), 6.28 (d, $J = 1$ Hz, 1H), 7.31 (d, $J = 1$ Hz, 1H); ^{13}C NMR (125.8 MHz, acetone- d_6): δ 24.4, 25.4, 30.6, 40.9, 55.6, 110.3, 141.4; ^{19}F NMR (471 MHz, acetone- d_6): δ -139.7; ^{11}B NMR (128 MHz, acetone- d_6): δ 1.61; HRMS (ESI): calcd for $\text{C}_{11}\text{H}_{16}\text{BF}_3\text{NO}$ (M-K) 246.1277, found 246.1281.

General Procedure B for the Reductive Amination of Potassium Formylaryltrifluoroborates using Pyridine Borane. Potassium 3-((Pyrrolidin-1-yl)methyl)phenyltrifluoroborate (20)

To a test tube containing potassium 3-formylphenyltrifluoroborate (106 mg, 0.50 mmol) and KHF_2 (156 mg, 2.00 mmol) in MeOH (2 mL) was added pyrrolidine (53 mg, 0.75 mmol). This was stirred for 2 h at rt. Pyridine borane (46 mg, 0.5 mmol) was then added and stirring was continued for 16 h. The solvent was then removed and the resulting white solid was rinsed with Et_2O (2×5 mL). The resulting solid was then extracted with 10% MeOH/acetone (3×5 mL) after which the combined extracts were concentrated and precipitated with Et_2O (10 mL) to give potassium 3-((pyrrolidin-1-yl)methyl)phenyltrifluoroborate (102 mg, 76%). (For some substrates it proved necessary to add solid KHCO_3 (50 mg, 0.5 mmol) to the MeOH/acetone extracts to neutralize the product). mp >200 °C; IR (KBr) = 2975 cm^{-1} ; ^1H NMR (500 MHz, acetone- d_6): δ 1.66 – 1.77 (m, 4H), 2.52 (s, 4H), 3.63 (s, 2H), 7.05 (d, $J = 5$ Hz, 2H), 7.37 (t, $J = 4$ Hz, 1H), 7.47 (s, 1H); ^{13}C NMR (125.8 MHz, acetone- d_6): δ 23.2, 53.7, 60.5, 126.3, 126.4, 130.7, 132.4, 135.7; ^{19}F NMR (471 MHz, acetone- d_6): δ -142.4; ^{11}B NMR (128 MHz, acetone- d_6): δ 3.95; HRMS (ESI): calcd for $\text{C}_{11}\text{H}_{14}\text{BF}_3\text{N}$ (M-K) 228.1171, found 228.1173.

N-((4'-Methoxybiphenyl-4-yl)methyl)cyclohexanamine (38) Using Ligandless Conditions

A sealed microwave vial containing potassium 4-((cyclohexylamino)methyl)phenyltrifluoroborate **34** (89 mg, 0.30 mmol), *p*-bromoanisole (58 mg, 0.31 mmol), Pd(OAc) $_2$ (2 mg, 3 mol %), and K_2CO_3 (124 mg, 0.90 mmol) was placed under vacuum (0.01 torr) for 30 min after which MeOH (1 mL) was added with stirring. The reaction mixture was heated at 75 °C for 3 h then allowed to come to rt. Water (1 mL) was added, resulting in the formation of a white precipitate. This was filtered and the white solid was then rinsed with water (2×1 mL). The white solid was taken up in EtOAc (5 mL) and the organic phase was washed with H_2O (3×3 mL), brine (1 mL), dried (MgSO_4) and filtered to give title compound **38** (73 mg, 82%) as a solid following solvent removal. mp 65-70 °C; IR (KBr) = 2918, 2845 cm^{-1} ; ^1H NMR (500 MHz, methanol- d_4): δ 1.12 – 1.26 (m, 6H), 1.57 (d, $J = 12$ Hz, 1H), 1.73 (d, $J = 9$ Hz, 2H), 1.96 (d, $J = 10$ Hz, 2H), 2.42 – 2.47 (m, 1H), 3.77 (s, 2H), 3.80 (s, 3H), 6.95 (d, $J = 8$ Hz, 2H), 7.36 (d, $J = 8$ Hz, 2H), 7.51 (dd, $J = 11, 2$ Hz, 4H); ^{13}C NMR (125.8 MHz, methanol- d_4): δ 24.7, 25.8, 32.3, 49.5, 54.3, 55.7, 113.9, 126.2, 127.5, 128.5, 133.3, 137.8, 139.6, 159.4; HRMS (ESI): calcd for $\text{C}_{20}\text{H}_{26}\text{NO}$ (M+H) 296.2014, found 296.2014.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Financial support was derived from the NIH (R01 GM35249), and unrestricted funds from Merck Research Laboratories and Amgen are gratefully acknowledged. We thank Frontier Scientific and Johnson Matthey for donations of materials used in this research.

References

- i. Molander GA, Ellis N. Acc Chem Res 2007;40:275–286. [PubMed: 17256882]
- ii. a) Molander GA, Ribagorda M. J Am Chem Soc 2003;125:11148–11149. [PubMed: 16220907] b) Molander GA, Ellis NM. J Org Chem 2006;71:7491–7493. [PubMed: 16958551] c) Molander GA, Figueroa R. Org Lett 2006;8:75–78. [PubMed: 16381571] d) Molander GA, Figueroa R. J Org Chem 2006;71:6135–6149. [PubMed: 16872197] e) Molander GA, Cooper DJ. J Org Chem 2007;72:3558–3560. [PubMed: 17408287]
- iii. a) Abdel-Magid AF, Carson KG, Harris BD, Maryanoff CA. Org Chem 1996;61:3849–3862. b) Abdel-Magid AF, Mehrman SJ. Org Process Rev Dev 2006;10:971–1031. c) Quan ML, Han Q, Fevig JM, Lam PYS, Bai S, Knabb RM, Luetttgen JM, Wong PC, Wexler RR. Bioorg Med Chem Lett

- 2006;16:1795–1798. [PubMed: 16434195] d) Baxter, EW.; Reitz, AB. Organic Reactions. Vol. 59. John Wiley and Sons, Inc.; New York: 2002.
- iv. Schwartz C, Raible J, Mott K, Dussault PH. Org Lett 2006;8:3199–3201. [PubMed: 16836365]
- v. Leuckart R. J Prakt Chem 1890;41:33.
- vi. a) Gray CW, Walker BT, Brian T, Foley RA, Houston TA. Tetrahedron Lett 2003;44:3309–3312. b) Johnson LL, Houston TA. Tetrahedron Lett 2002;43:8905–8908. c) Luvino D. Tetrahedron Lett 2006;47:9253–9256.
- vii. a) Hall DG, Gravel M, Thompson MZ, Berube C. J Org Chem 2002;67:3–15. [PubMed: 11777432] b) Martin B, Possm F, Barbier C, Carreaux F, Carboni B, Seiler N, Moulinoux JP, Delcros JG. J Med Chem 2001;44:3653. [PubMed: 11606130] c) Asano T, Nakamura H, Uehara Y, Yamamoto Y. ChemBioChem 2004;5:483–490. [PubMed: 15185372] d) Prullano L, Rohovec J, Aime S, Maschmeyer T, Prata MI, Lima J, Geraldles C, Peters J. Chem Eur J 2004;10:5205–5217.
- viii. a) Petasis NA, Akritopoulou L. Tetrahedron Lett 1993;34:583–586. b) Petasis NA, Goodman A, Zavialov IA. Tetrahedron 1997;53:16463–16470. c) Petasis NA, Zavialov IA. J Am Chem Soc 1997;119:445–446.
- ix. a) Billard T, Langlois BR. J Org Chem 2002;67:997–1000. [PubMed: 11856050] b) Tremblay-Morin JP, Raeppe S, Gaudette R. Tetrahedron Lett 2004;45:3471–3474. c) Tehrani KA, Stas S. Tetrahedron 2007;63:8921–8931. d) Li SW, Batey RA. Chem Commun 2004:1382–1383. e) Schlienger N, Bryce MR, Hansen TK. Tetrahedron Lett 2000;41:1303–1305. f) Morgan IR, Yazici A, Pyne SG. Tetrahedron 2008;64:1409–1419. g) Viera AS, Ferreira FP, Fiorante PF, Guadagnin RC, Stefani HA. Tetrahedron 2008;64:3306–3314.
- x. Kabalka GW, Venkataiah B, Dong G. Exception. Tetrahedron Lett 2004;45:729–731.
- xi. Petasis NA. Method for the Synthesis of Amines and Amino Acids with Organoboron Derivatives WO 98/00398. January 8;1998
- xii. MacMillan DWC, Lee S. J Am Chem Soc 2007;129:15438–15439. [PubMed: 18031044]
- xiii. Basu B, Jha S, Mosharef H, Das P. Synlett 2003;4:555–557.
- xiv. Abdel-Magid AF, Maryanoff CA. Org Proc Res Dev 2006;10:971–1031.
- xv. Domann MD, Guch IC, DiMare M. J Org Chem 1995;60:5995–5996.
- xvi. Pelter A, Rosser RM, Mills S. J Chem Soc Perkin Trans 1 1984;4:717–720.
17. Molander GA, Biolatto B. J Org Chem 2003;68:4302–4314. [PubMed: 12762730]

Scheme 1.

Scheme 2.

Table 1
Organotrifluoroborate Substrates Subjected to Palladium Catalyzed Reductive Amination Using Various Amines

J Org Chem. Author manuscript; available in PMC 2010 February 10.

product

% yield

83

product

% yield

86

product

% yield

70

product

% yield

74

product

% yield

92

(n-Bu)N

product

% yield

100

$(n\text{-Bu})_4\text{N}^+\text{F}_3\text{B}^-$

^a Greater than 5% impurity from DMF

Table 2
Organotrifluoroborate Substrates Subjected to Reductive Amination Using Pyridine
Borane and Various Amines

entry	aldehyde	product
1	17	18

entry	aldehyde	product
2	17

19

entry	aldehyde	product
3	17	 20
4	17	 21
5	17	 22

entry	aldehyde	product
6	17

23

entry	aldehyde	product
7	17	 24

^aProduct exists as greater than 5% internal ammonium salt

Table 3
Organotrifluoroborate Substrates Subjected to Reductive Amination Using Pyridine Borane

entry	aldehyde
1	15

entry

aldehyde

2

25

entry

aldehyde

3

4

entry

aldehyde

5

entry

aldehyde

6

33

entry

aldehyde

7

25

entry

aldehyde

8

15

entry

aldehyde

9

31

^a Gram scale reaction.

^b Product exists as greater than 5% internal ammonium salt.

^c Performed in two stages with no KHF₂.