Published in final edited form as:

Biomed Pharmacother. 2005 October ; 59(Suppl 1): S31.

Depressive mood is independently related to stroke and cardiovascular events in a community

G. Yamanaka^a, K. Otsuka^{a,b,*}, N. Hotta^{a,b}, S. Murakami^c, Y. Kubo^a, O. Matsuoka^a, E. Takasugi^{a,b}, T. Yamanaka^{a,b}, M. Shinagawa^a, S. Nunoda^a, Y. Nishimura^a, K. Shibata^{a,b}, H. Saitoh^{a,b}, M. Nishinaga^d, M. Ishine^e, T. Wada^e, K. Okumiya^f, K. Matsubayashi^g, S. Yano^h, S. Ishizukaⁱ, K. Ichihara^j, G. Cornélissen^k, and F. Halberg^k

^aDepartment of Medicine, Tokyo Women's Medical University, Medical Center East, Nishiogu 2-1-10, Arakawa, Tokyo 116-8567, Japan

^bDivision of Neurocardiology and Chronoecology, Tokyo Women's Medical University, Medical Center East, Nishiogu 2-1-10, Arakawa, Tokyo 116-8567, Japan

^cDepartment of Internal Medicine, Osaka Medical University, Osaka, Japan

^dDepartment of Gerontology, School of Medicine, Kochi University, Kochi, Japan

^eDepartment of Field Medicine, Kyoto University Graduate School of Medicine, Kyoto, Japan

^fResearch Institute for Humanity and Nature, Kyoto, Japan

^gCenter for South-East Asian Studies, Kyoto University, Kyoto, Japan

^hSorachi Health and Welfare Office, Sorachi-Godochosha, Iwamizawa, Hokkaido, Japan

ⁱA&D Co. Ltd, R&D Division, Saitama, Japan

^jDivision of Clinical Laboratory Sciences, Faculty of Health Sciences, School of Medicine, Yamaguchi University, Ube, Japan

^kHalberg Chronobiology Center, University of Minnesota, Minneapolis, MN, USA

Abstract

By means of a multivariate Cox model, we investigated the predictive value of a depressive mood on vascular disease risk in middle-aged community-dwelling people. In 224 people (88 men and 136 women; mean age: 56.8 ± 11.2 years) of U town, Hokkaido (latitude: 43.45 degrees N, longitude: 141.85 degrees E), a chronoecological health watch was started in April 2001. Consultations were repeated every 3 months. Results at the November 30, 2004 follow-up are presented herein. 7-day/24-h blood pressure (BP) and heart rate (HR) monitoring started on a Thursday, with readings taken at 30-min intervals between 07:00 h and 22:00 h and at 60-min intervals between 22:00 h and 07:00 h. Data stored in the memory of the monitor (TM-2430-15, A&D company, Japan) were retrieved and analyzed on a personal computer with a commercial software for this device. Subjects were asked to answer a self-administered questionnaire inquiring about 15 items of a depression scale, at the start of study and again after 1-2 years. Subjects with a score higher by at least two points at the second versus first screening were classified as having a depressive mood. The other subjects served as the control group.

The mean follow-up time was 1064 days, during which four subjects suffered an adverse vascular outcome (myocardial infarction: one man and one woman; stroke: two men). Among the variables

used in the Cox proportional hazard models, a depressive mood, assessed by the Geriatric Depression Scale (GDS), as well as the MESOR of diastolic (D) BP (DBP-MESOR) and the circadian amplitude of systolic (S) BP (SBP-Amplitude) showed a statistically significant association with the occurrence of adverse vascular outcomes. The GDS score during the second but not during the first session was statistically significantly associated with the adverse vascular outcome. In univariate analyses, the relative risk (RR) of developing outcomes was predicted by a three-point increase in the GDS scale (RR = 3.088, 95% CI: 1.375-6.935, $P = 0.0063$). Increases of 5 mmHg in DBP-MESOR and of 3 mmHg in SBP-Amplitude were associated with RRs of 2.143 (95% CI: 1.232-3.727, $P = 0.0070$) and 0.700 (95% CI: 0.495-0.989, $P = 0.0430$), respectively. In multivariate analyses, when both the second GDS score and the DBP-MESOR were used as continuous variables in the same model, GDS remained statistically significantly associated with the occurrence of cardiovascular death. After adjustment for DBP-MESOR, a three-point increase in GDS score was associated with a RR of 2.172 (95% CI: 1.123-4.200). Monday endpoints of the 7-day profile showed a statistically significant association with adverse vascular outcomes. A 5 mmHg increase in DBP on Monday was associated with a RR of 1.576 (95% CI: 1.011-2.457, $P = 0.0446$).

The main result of the present study is that in middle-aged community-dwelling people, a depressive mood predicted the occurrence of vascular diseases beyond the prediction provided by age, gender, ABP, lifestyle and environmental conditions, as assessed by means of a multivariate Cox model. A depressive mood, especially enhanced for 1-2 years, was associated with adverse vascular outcomes. Results herein suggest the clinical importance of repetitive assessments of a depressive mood and the need to take sufficient care of depressed subjects.

Another result herein is that circadian and circaseptan characteristics of BP variability measured 7-day/24-h predicted the occurrence of vascular disease beyond the prediction provided by age, gender, depressive mood and lifestyle, as assessed by means of a multivariate Cox model. Earlier, we showed that the morning surge in BP on Mondays was statistically significantly higher compared with other weekdays. Although a direct association between the Monday surge in BP and cardiovascular events could not be demonstrated herein, it is possible that the BP surge on Monday mornings may also trigger cardiovascular events. We have shown that depressive people exhibit a more prominent circaseptan variation in SBP, DBP and the double product (DP) compared to non-depressed subjects.

In view of the strong relation between depression and adverse cardiac events, studies should be done to ascertain that depression is properly diagnosed and treated. Chronodiagnosis and chronotherapy can reduce an elevated blood pressure and improve the altered variability in BP and HR, thus reducing the incidence of adverse cardiac events. This recommendation stands at the basis of chronomics, focusing on pre-habilitation in preference to rehabilitation, as a public service offered in several Japanese towns.

Keywords

Depressive mood; Seven-day ambulatory blood pressure; Cardiovascular diseases; Stroke

1. Introduction

Several lines of evidence suggest that clinical depression is a risk factor for cardiac morbidity and mortality in patients with coronary heart disease, especially after an acute myocardial infarction (MI). Major depression is associated with a fourfold increase in the risk of mortality during the first 6 months after an acute MI, and its prognostic significance is comparable to that of left ventricular dysfunction and a history of MI. Depression is not unusual among individuals with coronary heart disease, with studies indicating that between 15% and 22% of patients suffer from depression after a cardiac event.

Not only circadian, but also circannual, circaseptan (about 7-day) and other variations characterize the incidence of adverse vascular events, such as acute myocardial infarctions, strokes and sudden cardiac deaths, carrying important pathophysiological implications. Most studies showed an increased incidence in the morning, peaking between 06:00 and 12:00. Ambulatory blood pressure monitoring (ABPM) has become an important tool in the diagnosis and management of hypertension. Several studies have indicated that target organ damage and cardiovascular morbidity are more strongly associated with circadian BP endpoints than with office BP values. We reported that circadian and other patterns are synchronized by socio-ecologic factors, such as human lifestyles, as well as economic and environmental conditions. We demonstrated the presence of a weekly BP variation in community-dwelling subjects, including a surge in BP on Mondays, and we showed that it was most prominent in depressed people.

In 2000, we began a community-based study to longitudinally investigate the longevity and aging of a population in Hokkaido County (LILAC) [1,2], including the evaluation of the younger population aged from 40 to 74 years (Health Watch) [3]. Our goal is the prevention of cardiovascular events, notably strokes and myocardial ischemic events in a community. Herein, we examine whether a depressive mood is associated with an increased risk of stroke and cardiovascular mortality, and we estimate the predictive value of a depressive mood on the incidence of adverse vascular events, using a multivariate Cox model.

2. Methods

2.1. Subjects and Health Watch study design

The 224 subjects monitored between April 2001 and March 2004 were between 24 and 79 years of age (88 men and 136 women; mean age: 56.8 ± 11.2 years), living in U town in Hokkaido (latitude: 43.45 degrees N, longitude: 141.85 degrees E). All subjects visited and utilized the free health screening, counseling, and educational services of the U town office. Subjects with definite neurological diseases, such as Parkinsonism and stroke, and those who were too severely ill to stand without help were excluded. Lifestyle, including nutritional and sleep quality was investigated by several types of questionnaires. BP was measured at the beginning of the study in a sitting position, and brachialankle pulse wave velocity (baPWV) was measured between the right arm and ankle in a supine position, using an ABI/Form instrument (Nippon Colin Co., Ltd., Komaki, Japan). In all studies, baPWV was measured twice after at least a 5-min rest. For the assessment of the predictive value, only baPWV measures from participants showing normal ankle/brachial pressure index (ABI) values (>0.90) were used.

2.2. Ambulatory blood pressure monitoring

The 7-day/24-h BP recordings were obtained with a commercially available ambulatory BP recorder (TM-2431, A&D Company, Japan) [4]. Subjects were fitted with the recorder in the morning, usually between 10:00 and 11:30 h of the first day of monitoring. The recorder was set to take measurements at intervals of 30 min between 07:00 and 22:00 h, and of 60 min between 22:00 and 07:00 h. Subjects were instructed to follow their usual daily routine after they left the office. Subjects were instructed to remain motionless each time a reading was to be taken, and then to record their activity on a diary sheet. Stored data were retrieved and analyzed on a personal computer with commercially available software for the device (TM-2430-15, A&D Company, Japan). The data were further analyzed by sphgmochron for the entire 7-day span and separately for each day of monitoring (provided by corne001@umn.edu for all interested comers). In addition to the circadian rhythm characteristics, estimates were also obtained for mean values during 24-h, waking and sleeping spans, and for the day-night ratio (as a gauge of “dipping”) for SBP, DBP and HR, and also

for mean blood pressure ($MBP = 1/3SBP + 2/3DBP$), pulse pressure ($PP = SBP - DBP$), and the double product ($DP = SBP \times HR/100$) ("dip" is defined as the Awake-Sleep difference divided by Awake ($\times 100$), where Awake and Sleep are average values during the awake- and sleep-span, respectively).

2.3. Depression screening scale

Numerous depression rating scales are currently available, which represent a mixture of observer-rated and self-rating scales. In this investigation, subjects were asked about 15 items on a depression scale, by means of a self-administered questionnaire [5]. When the depression score was 5, subjects were considered to have a depressed mood. A depression score was obtained again 1-2 years later. Subjects with a score during the second screening higher by at least two points compared to that of the first screening were assigned to the enhanced depressive mood group (irrespective of the score at the start of study). All other subjects were assigned to the control group.

2.4. Stroke and cardiovascular mortality

The Health Watch study was started on April 12, 2001. One or two doctors of our team visited and provided repeated consultations every 3 months. They offered advice in relation to the rehabilitation of disordered functions, and healthy lifestyle modifications (promoting complete cessation of smoking, weight reduction, reduction of salt intake, moderation in the consumption of fruits and vegetables and alcohol intake). They also advised in terms of medical prescriptions for the local general practitioner.

The follow-up herein ended on November 30, 2004. During this time, nine subjects died of myocardial infarction or stroke. The follow-up time was defined as the time elapsed between the date of the first (reference) examination and the date of first cardiovascular event or death.

2.5. Statistical analysis

Results are reported as mean $\pm$ S.D. Student's *t*-tests served to compare the enhanced depressed and control groups. A *P*-value below 0.05 was considered to indicate statistical significance (and below 0.10, borderline statistical significance).

All data were analyzed with the Statistical Software for Windows (StatFlex Ver.5.0, Artec, Osaka, <http://www.statflex.net>). We used Cox's regression analysis to calculate the unadjusted and adjusted relative risks (RRs) and 95% confidence intervals (CIs) for cardiovascular death. To identify independent predictors of stroke and cardiovascular disease in relation to a depressive mood, we used a multivariate Cox regression analysis with stepwise selection, including as variables age, gender, BMI, lifestyle, sleep quality, QOL, baPWV and ABP endpoints.

The independent correlation of the enhanced depressive mood was determined by means of a logistic regression analysis. Kaplan-Meier event probability curves were computed with two groups, and the cumulative probability of events of two groups was compared by means of the log-rank test. Significance was considered at a value of $P < 0.05$.

3. Results

Reference characteristics of the 224 subjects are given in Table 1. The sample comprised 88 men and 136 women. The mean age of participants at entry was 56.8 years. The mean follow-up time was 1064 days, during which four subjects suffered from adverse vascular outcomes (myocardial infarction: one man and one woman, stroke: two men).

Table 1 lists the mean, standard deviation (S.D.), minimum and maximum of the different variables considered in this chronoeological health watch. Items include indices of environmental conditions, lifestyle, quality of sleep, QOL and 7-day ABP endpoints, such as the 7-day (from Thursday to Wednesday) and daily averages of SBP, DBP, HR, the incidence of SBP- and DBP-CHAT Ccircadian hyper-amplitude-tensionl and of SBP, DBP and HR “dipping” (SBP-dip, DBP-dip, and HR-dip), estimates of the circadian amplitude and acrophase of SBP, DBP and HR, as well as estimates of the percentage time elevation (PTE) of SBP, DBP and HR. Averages over 7 days were also computed (not shown) as SBP 1-7, DBP 1-7, pulse pressure (PP) 1-7, double product (DP) 1-7, 24-h S.D. of HR (HRSD1-7), SBP-dip 1-7, DBP-dip 1-7, HR-dip 1-7, MAPdip 1-7, PPdip 1-7, and DPdip 1-7.

Depression scores were obtained for 218 of the 224 citizens at the start of study and from 179 subjects after 1-2 years (177 subjects provided a score during both sessions). Scores obtained during the first and second sessions averaged 4.11 ± 3.11 , and 4.42 ± 3.14 , respectively. Of the 177 citizens who answered both questionnaires, 39 were assigned to the enhanced depressive mood group. A comparison of characteristics between the two groups is shown in Table 2. It can be seen that depressive subjects had a higher body mass index (BMI), they consumed smaller meals (less amount of rice), and their total serum cholesterol was higher as compared to the control subjects. The time to fall asleep was also slightly longer in the depressive group ($P = 0.051$). Kaplan-Meier curves for event-free survival revealed a significant difference between the two groups ($P = 0.0124$, log-rank test) (Fig. 1). Subjects in the control group had a better event-free survival than depressed subjects.

Among the variables used in Cox proportional hazard models, a depressive mood, assessed by the Geriatric Depression Scale (GDS), as well as DBP-MESOR and SBP-Amplitude showed a statistically significant association with the occurrence of vascular outcomes (Table 3). It should be noted that GDS during the second but not during the first session was statistically significantly associated with adverse vascular outcomes. In univariate analyses, an increase by two or three points of the GDS score during the second session was associated with a RR of 2.121 (95% CI: 1.237-3.637) or 3.088 (95% CI: 1.375-6.935), respectively ($P = 0.0063$). A 5 mmHg increase in DBP-MESOR and a 3 mmHg increase in SBP-Amplitude were associated with RRs of 2.143 (95% CI: 1.232-3.727, $P = 0.0070$) and 0.700 (95% CI: 0.495-0.989, $P = 0.0430$), respectively. In multivariate analyses, when both the second GDS score and DBP-MESOR were used as continuous variables in the same model, GDS remained statistically significantly associated with the occurrence of cardiovascular death. After adjustment for DBP-MESOR, a three-point increase in GDS score was associated with a RR of 2.172 (95% CI: 1.123-4.200).

Monday and/or Tuesday endpoints of the 7-day ABP showed a statistically significant association with adverse vascular outcomes. A 5-mmHg increase in DBP-5 or DBP-6, namely in the arithmetic mean of DBP from data collected on Mondays and Tuesdays, was associated with RRs of 1.576 (95% CI: 1.011-2.457, $P = 0.0446$) and 1.666 (95% CI: 1.052-2.639, $P = 0.0297$), respectively. A 20-unit increase in DP-5 (double product on Mondays) and a 3% increase in MAP-dip-5 were associated with RRs of 3.067 (95% CI: 1.075-8.753, $P = 0.0362$) and 0.824 (95% CI: 0.683-0.995, $P = 0.0439$), respectively. Saturday endpoints also related to adverse vascular outcomes. A 3% increase in PP-dip-3 (dipping ratio of pulse pressure on Saturday nights) and a 5-unit increase in DP-dip-3 (dipping ratio of Saturday's double product) were inversely associated with adverse vascular outcome, with RRs of 0.865 (95% CI: 0.756-0.989, $P = 0.0337$) and 0.752 (95% CI: 0.596-0.948, $P = 0.0158$), respectively.

4. Discussion

Depression is a risk factor for cardiac morbidity and mortality in patients with coronary heart disease [6-8]. Anda et al. [9] followed a cohort of 2832 persons for an average of 12.4 years. During the reference stage, 11.1% of the study cohort had a depressed affect. The adjusted relative risk of fatal CHD was 1.4 for patients with depressed affect, compared with patients who were not depressed. For non-fatal CHD, the adjusted relative risk was 1.6. For both fatal and non-fatal CHD, the increased risk associated with depression was independent of established risk factors for CHD (e.g. smoking, cholesterol concentration, family history). Barefoot and Schroll [10] reported that high scores on a measure of symptoms of depressed mood were associated with an increased risk for AMI and early mortality during a 27-year follow-up, and that the impact of a depressed affect on health did not differ between men and women. We also observed in middle-aged community-dwelling people that a depressive mood, assessed by GDS, predicted the occurrence of vascular diseases beyond the prediction provided by age, gender, ABP, lifestyle and environmental conditions, as assessed by means of a multivariate Cox model. A depressive mood, especially when enhanced for 1-2 years, was associated with adverse vascular outcomes. Results herein suggest the clinical importance of repeated assessment of a depressive mood and of taking sufficient care of depressed subjects.

Another result of the present study is that in community-dwelling people, circadian and circaseptan characteristics of BP variability derived from 7-day/24-h ABP predicted the occurrence of vascular diseases beyond the prediction provided by age, gender, depressive mood and lifestyle, as assessed by means of a multivariate Cox model.

Ambulatory 24-h blood pressure monitoring has become quite popular in recent years, offering a number of advantages from the viewpoint of both diagnosis and treatment [11]. One advantage rests on the fact that clinic blood pressure is not representative of blood pressure values outside the clinic. Results obtained herein show that blood pressure is characterized not only by a circadian but also by a weekly variation. The 24-h averages of SBP and DBP are seen to differ from one day to another in the same citizen in the same 7-day record. In middle-aged community-dwelling people, DBP-MESOR and SBP-Amplitude predicted the occurrence of vascular diseases, assessed by means of a multivariate Cox model. We found that several endpoints from data collected on Mondays predicted adverse outcomes, suggesting that circaseptan as well as circadian characteristics are associated with the occurrence of vascular disease. We reported on weekly BP characteristics earlier [12]. Most citizens in a community showed a novelty effect (the first day effect), a holiday dip, and a Monday morning surge of SBP. A Monday peak has been reported for the incidence of acute myocardial infarction, sudden cardiac death, and stroke [13-17]. Our previous studies showed that the morning BP surge on Mondays was statistically significantly higher compared with other weekdays [18]. Although a direct association between the Monday surge in BP and cardiovascular events could not be demonstrated directly, it is possible that a morning surge of BP on Mondays may trigger cardiovascular events. This possibility is supported by the fact that we found that depressive citizens had a more prominent circaseptan component for SBP, DBP and DP, compared with non-depressive subjects [19].

Results herein indicate the clinical importance of taking care of depressed subjects. In view of the strong relation between depression and adverse cardiac events, studies should be done to ascertain whether treatment of depression, especially from the standpoint of chronodiagnosis and chronotherapy, can reduce elevated blood pressure values and improve the altered variability in BP and HR, so as to reduce the incidence of adverse cardiac events. Fewer than 7 days of monitoring means a greater chance of a false diagnosis, on which a treatment decision may depend for the long term, including perhaps a potentially unnecessary treatment for decades. Less than 7 days of monitoring can be compared with taking the pulse for less than a

cardiac cycle, if the circaseptan aspect of time structures (chronomes) is viewed as one of many new spectral components, with even longer periods, including transyears characterizing series of blood pressure and heart rate covering at least 5 and up to 38 years [20]. When the ECG and EEG, as well as the blood pressure of small rodents can be monitored continuously for most of their lifetime, this study is but a small step toward universal 7-day and eventually continuous monitoring of BP and HR, or the ECG.

Acknowledgments

This study was supported by Fukuda Foundation for Medical (Grant in 2004 for the study on association between arterial stiffness and cognitive impairment in community-dwelling subjects over 70 years old).

References

- [1]. Otsuka K, Murakami S, Kubo Y, Yamanaka T, Mitsutake G, Ohkawa S, et al. Chronomics for chronoastrobiology with immediate spin-offs for life quality and longevity. *Biomed Pharmacother* 2003;57:1s–8s.
- [2]. Murakami S, Otsuka K, Yamanaka G, Kubo Y, Matsuoka O, Yamanaka T, et al. Positive impact of social intervention on disturbed neurobehavioral function in an elderly community-dwelling population: Longitudinal Investigation for Longevity and Aging in Hokkaido County (LILAC). *Biomed Pharmacother* 2004;58:45s–7s.
- [3]. Otsuka K, Mitsutake G, Yano S. Depression, quality of life, and lifestyle: chronoecological health watch in a community. *Biomed Pharmacother* 2002;56:231s–42s. [PubMed: 12653175]
- [4]. Palatini P, Frigo G, Bertolo O, et al. Validation of the A&D TM-2430 device for ambulatory blood pressure monitoring and evaluation of performance according to subjects' characteristics. *Blood Pressure Monit* 1998;3:255–9.
- [5]. Yesavage JA, Brink TL, Rose TL, Lum O, Huang V, Adey M, et al. Development of validation of a geriatric depression screening scale: a preliminary report. *J Psychiat Res* 1983;17:37–49. [PubMed: 7183759]
- [6]. Frasure-Smith N, Lesperance F, Talajic M. Depression following myocardial infarction: determine predictive variables. *J Am Med Assoc* 1993;270:1819–61.
- [7]. Frasure-Smith N, Lesperance F, Talajic M. Depression and 18-month prognosis after myocardial infarction. *Circulation* 1995;91:999–1005. [PubMed: 7531624]
- [8]. Barefoot JC, Schroll M. Symptoms of depression, acute myocardial infarction, and total mortality in a community sample. *Circulation* 1996;93:1976–80. [PubMed: 8640971]
- [9]. Anda R, Williamson D, Jones D, et al. Depressed affect, hopelessness, and the risk of ischemic heart disease in a cohort of US adults. *Epidemiology* 1993;4:285–94. [PubMed: 8347738]
- [10]. Barefoot JC, Schroll M. Symptoms of depression, acute myocardial infarction, and total mortality in a community sample. *Circulation* 1996;93:1976–80. [PubMed: 8640971]
- [11]. Cornélissen G, Schwartzkopff O, Halberg F, Otsuka K, Watanabe Y. Seven-day ambulatory monitoring for adults with hypertension and diabetes [letter]. *Am J Kidney Dis* 2001;37:878. [PubMed: 11273894]
- [12]. Shinagawa M, Otsuka K, Murakami S, et al. Seven-day (24-h) ambulatory blood pressure monitoring, self-reported depression and quality of life scores. *Blood Pressure Monit* 2002;7:69–76.
- [13]. Halberg F. Chronobiology. *Ann Rev Physiol* 1969;31:675–725. [PubMed: 4885778]
- [14]. Otsuka, K., editor. Chronome and Janus-medicine: heart rate variability (HRV) and BP variability (BPV) from a viewpoint of chronobiology and ecology. Medical Review; Tokyo: 1998. p. 213
- [15]. Halberg F, Cornélissen G, Katinas G, Hillman D, Sampson M, Schwartzkopff O. Season's Appreciations 2000. Chronomics complement, among many other fields, genomics and proteomics. *Neuroendocrinol Lett* 2001;22:53–73. [PubMed: 11393163]
- [16]. Cornélissen G, Breus TK, Bingham C, et al. Beyond circadian chronorisk: worldwide circaseptan-circasemiseptan patterns of myocardial infarctions, other vascular events, and emergencies. *Chronobiologia* 1993;20:87–115. [PubMed: 8354104]

- [17]. Cornélissen G, Tamura K, Tarquini B, et al. Differences in some circadian patterns of cardiac arrhythmia, myocardial infarction and other adverse vascular events. *Chronobiologia* 1994;21:79–88. [PubMed: 7924643]
- [18]. Murakami S, Otsuka K, Kubo Y, Shinagawa M, Yamanaka T, Ohkawa S, et al. Repeated ambulatory monitoring reveals a Monday morning surge in blood pressure in a community-dwelling population. *Am J Hypertens* 2004;17:1179–83. [PubMed: 15607626]
- [19]. Otsuka K, Yamanaka G, Shinagawa M, Murakami S, Yamanaka T, Shibata K, et al. Chronomic community screening reveals about 31% depression, elevated blood pressure and infradian vascular rhythm alteration. *Biomed Pharmacother* 2004;58:48s–55s.
- [20]. Cornélissen G, Masalov A, Halberg F, Richardson JD, Katinas GS, Sothorn RB, et al. Multiple resonances among time structures, chronomes, around and in us. Is an about 1.3-year periodicity in solar wind built into the human cardiovascular chronome? *Hum Physiol* 2004;30(2):86–92.

Fig. 1.

Kaplan-Meier event probability curves for the incidence of vascular disease.

Calculation of Kaplan-Meier curves for event-free survival revealed a significant difference between the control and depressive groups ($P = 0.0124$, log-rank test). A depression score was obtained again after 1-2 years, and when the score during the second screening had increased by more than 2 points compared to the first screening, such subjects were assigned to the enhanced depressive mood group, shown as “Depressive”, while the other subjects were assigned to the control group, shown as “Control”. It can be seen that subjects in the control group had fewer adverse vascular events (better event-free survival).

Table 1

Reference characteristics of the 224 subjects

Variables	Units	n	Mean	S.D.	Min.	Max.
Age	years	224	56.8	11.2	24	79
Gender ratio		224	0.393	0.489	0	1
BMI	kg/m ²	220	24.6	2.8	17.5	34.7
Days Follow-up	days	224	1063.7	271.6	114	1378
Outside temperature	°C	192	3.7	8.7	-16	25
Room temperature	°C	206	22.3	3.9	0	34
Temperature difference	°C	199	18.4	9.2	-5	42
Time of sunrise	h:min	218	05:53	0.043	03:54	07:30
Time of sunset	h:min	218	17:17	0.046	16:00	19:21
Sunshine duration	h:min	216	11:24	0.087	08:57	15:22
Smoking I		218	252.1	463.8	0	2720
Smoking II		217	9.88	15.16	0	55
Family history of CVD	points	219	2.19	2.16	0	8
Exercise	points	220	1.79	1.21	0	4
Alcohol	points	219	1.00	3.53	0	50
GDS I	points	218	4.11	3.11	0	13
GDS2	points	179	4.42	3.14	0	15
Subjective healthiness	%	220	65.38	17.50	19.8	100
Subjective mood	%	220	69.13	18.19	14.3	100
Family relations	%	218	79.86	19.40	7.4	100
Relations with friends and relatives	%	220	77.90	18.32	8.7	100
Financial satisfaction	%	218	62.65	24.87	0	100
Life satisfaction	%	220	70.56	21.77	0	100
Sense of happiness	%	220	74.19	19.31	3.5	100
Total score of QOL		216	496.6	107.7	118.8	692.0
Getting up		216	06:08	0.035	04:00	09:00
Going to bed		216	22:33	0.042	19:59	01:59
Duration of sleep	h:min	156	7:51	0.043	05:30	11:00
Time from getting up to going to bed		156	16:08	0.043	13:00	18:30

Variables	Units	n	Mean	S.D.	Min.	Max.
Sleep well?		188	1.21	0.41	1	2
Time to fall asleep	min	188	4.05	2.2	2	12
Duration of snoring	years	112	4.54	2.49	0	12
Not feeling sufficiently refreshed		187	1.67	0.75	1	3
Duration of not feeling refreshed	months	167	5.13	6.70	0	15
Quality of sleep		184	19.79	2.53	14	24
Condition during sleep		184	21.33	0.90	18	22
Salt intake (Summer)	g/day	169	13.57	2.08	9.37	20.27
Salt intake (Winter)	g/day	169	13.47	2.02	9.37	20.27
Amount of meal (rice)	cups	169	2.50	0.89	0.5	6
Soybean soup	cups	169	1.52	0.77	0	3.3
Vegetables (Summer)		172	9.67	5.64	0	48
Vegetables (Winter)		172	9.02	5.75	0	48
Fruit		170	4.61	3.46	0	21
R baPWV	cm/s	202	1486.6	323.5	735	2561
L baPWV	cm/s	203	1526.3	336.5	748	2718
R ABI		204	1.11	0.09	0.60	1.33
L ABI		204	1.11	0.08	0.82	1.30
CTR	%	128	48.31	5.27	37.3	62.0
SVI + RV5	Mm	151	27.99	9.03	11.6	60.0
SBP-MESOR	mmHg	223	126.66	14.75	95.2	172.6
DBP-MESOR	mmHg	223	77.49	8.44	57.6	103.5
HR-MESOR	mmHg	223	69.76	7.09	48.2	92.9
SBPdip	%	223	15.62	7.29	-7.08	33.18
DBPdip	%	223	17.17	7.17	-3.13	32.53
HR dip	%	223	16.81	7.78	-18.89	34.84
SBP CHAT		223	0.152	0.360	0	1
DBP CHAT		223	0.054	0.226	0	1
SBP Amp	mmHg	223	24.98	10.55	0.78	56.25
DBP Amp	mmHg	223	16.48	6.80	0.61	36.96
HR Amp	mmHg	223	15.80	6.30	1.17	35.74
SBP Acro	h:min	222	13:51	0.10	01:27	21:07

Variables	Units	n	Mean	S.D.	Min.	Max.
DBPAcro	h:min	222	13:51	0.08	06:11	23:40
HRAcro	h:min	222	14:10	0.09	00:25	22:38
SBP PTE	%	223	12.53	23.16	0	97.9
DBP PTE	%	223	6.97	14.90	0	91.4
HR PTE	%	223	0.56	2.31	0	18.0
SBP1	mmHg	219	129.86	16.50	90.46	186.79
SBP2	mmHg	221	128.53	15.83	93.49	182.64
SBP3	mmHg	220	127.19	14.84	89.41	172.54
SBP4	mmHg	220	126.82	15.18	93.23	167.71
SBP5	mmHg	215	127.55	15.32	95.11	176.11
SBP6	mmHg	209	127.84	15.35	96.58	175.94
SBP7	mmHg	176	127.71	15.57	90.08	173.63
DBP1	mmHg	219	79.57	9.51	53.07	104.59
DBP2	mmHg	221	78.90	9.19	56.22	106.23
DBP3	mmHg	220	78.04	8.81	50.87	97.76
DBP4	mmHg	220	77.65	8.87	50.94	103.45
DBP5	mmHg	215	78.44	9.08	56.23	111.47
DBP6	mmHg	209	78.42	9.28	54.06	108.69
DBP7	mmHg	176	78.45	9.11	57.89	105.85
Ppaver1	mmHg	222	50.51	10.49	30.36	88.94
Ppaver2	mmHg	222	49.92	9.94	31.00	81.06
Ppaver3	mmHg	222	49.50	9.79	31.46	81.95
Ppaver4	mmHg	220	49.54	9.81	30.05	79.14
Ppaver5	mmHg	216	49.41	9.78	31.42	84.93
Ppaver6	mmHg	210	49.76	9.92	33.16	83.03
Ppaver7	mmHg	176	49.65	10.2	31.03	92.04
Dpaver1	mmHg	222	91.06	15.32	56.79	157.91
Dpaver2	mmHg	222	92.48	15.69	57.03	158.19
Dpaver3	mmHg	222	91.27	14.03	63.94	145.57
Dpaver4	mHg	220	90.80	14.94	62.57	142.29
Dpaver5	mmHg	216	92.28	15.57	59.89	145.66
Dpaver6	mmHg	210	92.11	15.27	55.48	137.98

Variables	Units	n	Mean	S.D.	Min.	Max.
Dpaver7	mmHg	176	92.28	14.70	59.38	134.26
HRSD1		222	11.27	4.62	2.42	33.78
HRSD2		222	11.74	4.17	3.11	27.68
HRSD3		222	12.12	4.49	3.03	27.92
HRSD4		220	11.87	4.52	4.18	34.11
HRSD5		216	11.39	4.07	3.09	27.25
HRSD6		211	11.64	4.37	3.53	33.61
HRSD7		177	11.79	4.29	4.27	26.74
SBP dip 1	%	216	17.49	9.80	-16.66	41.44
SBP dip 2	%	210	15.21	10.35	-23.19	42.36
SBP dip 3	%	218	14.64	10.58	-22.4	40.9
SBP dip 4	%	211	15.59	10.49	-15.36	46.89
SBP dip 5	%	210	15.80	10.33	-33.1	36.29
SBP dip 6	%	200	16.89	10.03	-31.18	39.96
SBP dip 7	%	174	16.47	9.18	-13	38.41
DBP dip 1	%	216	18.46	10.71	-11.43	49.73
DBP dip 2	%	210	16.65	10.55	-17.33	43.29
DBP dip 3	%	218	16.36	10.62	-12.44	43.26
DBP dip 4	%	210	16.41	11.15	-22.51	50.65
DBP dip 5	%	209	17.82	10.18	-29.05	42.56
DBP dip 6	%	199	18.51	10.84	-24.9	39.67
DBP dip 7	%	173	17.52	10.40	-18.06	41.82
HR dip 1	%	216	14.75	12.24	-42.82	41.96
HR dip 2	%	210	16.01	11.36	-28.93	42.57
HR dip 3	%	218	17.60	10.25	-23.78	42.32
HR dip 4	%	211	16.42	12.59	-54.08	45.99
HR dip 5	%	210	17.42	10.12	-19.99	43.87
HR dip 6	%	199	17.61	10.57	-27.61	40.08
HR dip 7	%	173	17.46	10.83	-24.94	43.29
MAP dip 1	%	216	18.16	9.59	-13.85	41.81
MAP dip 2	%	210	15.99	9.92	-20.11	42.47
MAP dip 3	%	218	15.55	10.17	-13.59	42.26

Variables	Units	n	Mean	S.D.	Min.	Max.
MAP dip 4	%	211	16.13	10.39	-16.66	49.11
MAP dip 5	%	210	17.00	9.75	-30.87	39.3
MAP dip 6	%	199	17.82	9.93	-27.87	37.47
MAP dip 7	%	173	17.14	9.27	-12.66	39.07
PP dip 1	%	216	14.48	17.72	-42.37	53.11
PP dip 2	%	210	12.22	16.55	-47.26	55.36
PP dip 3	%	218	11.54	15.90	-47.8	45.64
PP dip 4	%	211	13.24	16.05	-35.93	54.82
PP dip 5	%	210	11.46	17.36	-65.85	48.88
PP dip 6	%	199	12.84	16.26	-56.39	46.5
PP dip 7	%	173	13.86	15.06	-44.22	52.95
DP dip 1	%	216	29.82	12.93	-15.58	58.08
DP dip 2	%	210	29.12	12.38	-30.23	54.21
DP dip 3	%	218	29.87	12.65	-29.7	56.7
DP dip 4	%	211	29.69	13.58	-17.23	58.52
DP dip 5	%	210	30.61	12.14	-20.4	55.15
DP dip 6	%	199	31.63	11.72	-1.94	58.3
DP dip 7	%	173	31.15	11.88	-15.43	59.39

Gender: $M=1$, $F=0$ (88 men and 136 women).

“dip” is defined as Awake-Sleep difference divided by Awake ($\times 100$), where Awake and Sleep are average values during awake- and sleep-span, respectively.

Table 2
Comparison of subject characteristics between the control and enhanced depressive mood groups

	Control			Enhanced depressive mood			t-value	P-value
	Units	N	Mean	S.D.	n	Mean	S.D.	
GDS 1	Points	138	4.34	3.13	39	3.26	2.70	0.049
GDS 2	Points	138	3.75	2.83	39	6.80	3.07	0.000
Difference of GDS scores	Points	138	-0.61	1.57	39	3.54	1.85	0.000
Age	Years	138	56.7	11.7	39	57.7	10.3	N.S.
Gender		138			39			N.S.
BMI	kg/m ²	138	24.1	2.6	39	25.2	2.9	0.029
Smoking	points	135	9.5	14.9	39	8.3	14.2	N.S.
Family Hx	points	138	2.27	2.22	39	2.18	2.02	N.S.
Alcohol	points	137	0.77	1.24	39	0.82	0.97	N.S.
Vegetables	times daily	138	2.12	0.80	39	2.27	0.77	N.S.
Subjective healthiness	%	138	65.6	17.6	39	65.0	17.5	N.S.
Subjective mood	%	138	69.8	17.9	39	65.9	19.0	N.S.
Sense of happiness	%	138	74.3	19.3	39	72.6	17.2	N.S.
Getting up Time	clock time	138	06:13	0.03	39	06:02	0.03	N.S.
Duration of Sleep	clock time	115	07:50	0.04	33	07:51	0.04	N.S.
Time for falling asleep	min	133	18.5	12.7	39	24.1	23.1	0.051
Sleep well?	points	135	1.2	0.4	39	1.3	0.5	N.S.
Sleep Quality Score	points	135	19.8	2.4	39	19.5	2.8	N.S.
Salt intake daily	g	123	13.6	2.2	39	13.3	1.7	N.S.
Rice	cups	123	2.60	0.95	39	2.20	0.64	0.016
rt baPWV	cm/s	127	1458.1	309.1	35	1520.6	335.4	N.S.
lt baPWV	cm/s	127	1500.6	330.1	35	1555.7	340.2	N.S.
rt ABI		127	1.12	0.08	35	1.09	0.08	0.080
k ABI		127	1.11	0.08	35	1.09	0.09	N.S.
ECG (SVI+RV5)	mm	98	28.0	9.3	24	29.7	10.0	N.S.
CTR	%	81	47.9	5.5	28	49.4	4.9	N.S.
Total cholesterol	mg/dl	131	204.4	30.3	37	215.5	28.0	0.046

	Control			Enhanced depressive mood			<i>t</i> -value	<i>P</i> -value
	Units	N	Mean	S.D.	n	Mean	S.D.	
HDL cholesterol	mg/dl	130	58.8	14.2	37	59.6	13.0	N.S.
Triglyceride	mg/dl	131	123.3	85.3	37	128.9	83.9	N.S.
Uric acid	mg/dl	128	5.0	1.3	35	5.0	1.6	N.S.
Serum creatinine	mg/dl	125	0.88	0.33	34	0.88	0.16	N.S.
FBS	mg/dl	128	99.5	16.5	36	96.7	12.9	N.S.
Follow up days	days	138	1085.8	227.9	39	1128.1	260.5	N.S.
Uric acid	mg/dl	128	5.0	1.3	35	5.0	1.6	N.S.
Serum creatinine	mg/dl	125	0.88	0.33	34	0.88	0.16	N.S.
FBS	mg/dl	128	99.5	16.5	36	96.7	12.9	N.S.
Follow up days	days	138	1085.8	227.9	39	1128.1	260.5	N.S.

Gender: $M = 1$, $F = 0$.

GDS scores and difference of GDS scores used for classification.

Table 3

Relative risk (RR) of adverse vascular outcome accounted for by variables investigated in chronoeological health watch

Variables	Number of data	β	S.E. (β)	z-value	P-value	RR	95% CI
Age	221	0.1163	0.0654	1.7769	0.0756		
Gender	221	1.5365	1.1548	1.3306	N.S.		
BMI	217	0.1964	0.1667	1.1780	N.S.		
Smoking	215	0.0002	0.0010	0.2238	N.S.		
Family history	216	-0.1854	0.2771	0.6692	N.S.		
Exercise	217	0.1482	0.3847	0.3852	N.S.		
Alcohol	216	0.0366	0.0860	0.4248	N.S.		
GDS 1	215	0.0329	0.1557	0.2113	N.S.		
GDS 2 (2 points)	176	0.3759	0.1376	2.7323	0.0063	2.121	1.237 3.637
GDS 2 (3 points)	176	0.3759	0.1376	2.7323	0.0063	3.088	1.375 6.935
Subjective healthiness	217	-0.0293	0.0272	1.0763	N.S.		
Subjective mood	217	-0.0261	0.0269	0.9681	N.S.		
Sense of happiness	217	-0.0050	0.0248	0.2007	N.S.		
Getting up time	213	-10.3425	13.8703	0.7457	N.S.		
Duration of sleep	153	12.1102	12.0658	1.0037	N.S.		
Time for falling asleep	185	-0.0104	0.2293	0.0452	N.S.		
Sleep well?	185	0.1778	1.1547	0.1540	N.S.		
Sleep quality score	181	-0.1019	0.1946	0.5234	N.S.		
Daily salt intake	167	-0.1249	0.2612	0.4782	N.S.		
Vegetables	170	0.0540	0.0559	0.9654	N.S.		
SBP-MESOR	220	0.0540	0.0290	1.8595	0.0630		
DBP-MESOR (5 mmHg)	220	0.1524	0.0565	2.6988	0.0070	2.143	1.232 3.727
HR-MESOR	220	-0.0020	0.0699	0.0281	N.S.		
SBP-dip	220	-0.1170	0.0607	1.9275	0.0539		
DBP-dip	220	-0.1158	0.0695	1.6661	0.0957		
HR-dip	220	0.0029	0.0659	0.0434	N.S.		
SBP-CHAT	220	-30.2348	2588781	0.0000	N.S.		
SBP-Amp (3 mmHg)	220	-0.1191	0.0589	2.0240	0.0430	0.700	0.495 0.989
DBP-Amp	220	-0.1368	0.0859	1.5937	N.S.		

Variables	Number of data	β	S.E. (β)	z-value	P-value	RR	95% CI
HR-Amp	220	-0.0030	0.0791	0.0382	N.S.		
HR-Acro	219	-3.2113	3.8626	0.8314	N.S.		
SBP-PTE	220	0.0152	0.0151	1.0058	N.S.		
HR-PTE	220	-38.8712	943727.8	0.0000	N.S.		
DBP-5 (5)	212	0.0910	0.0453	2.0085	0.0446	1.576	1.011 2.457
DBP-6 (5)	206	0.1021	0.0470	2.1740	0.0297	1.666	1.052 2.639
DP-5 (20)	213	0.0560	0.0268	2.0946	0.0362	3.067	1.075 8.753
DP-6	207	0.0457	0.0289	1.5814	N.S.		
SBP-dip-3	215	-0.0699	0.0373	1.8738	0.0610		
SBP-dip-7 (3%)	173	-0.0966	0.0485	1.9929	0.0463	0.749	0.563 0.995
MAP-dip-5 (3%)	207	-0.0644	0.0320	2.0155	0.0439	0.824	0.683 0.995
HR-dip-3	215	-0.0468	0.0372	1.2562	N.S.		
PP-dip-3 (3%)	215	-0.0485	0.0228	2.1242	0.0337	0.865	0.756 0.989
DP-dip-3 (5)	215	-0.0571	0.0237	2.4125	0.0158	0.752	0.596 0.948
DP-dip-5	207	-0.0446	0.0325	1.3745	N.S.		
Right baPWV	200	0.0015	0.0013	1.1678	N.S.		
Left baPWV	201	0.0018	0.0012	1.4543	N.S.		
Right ABI	202	-1.1412	4.8159	0.2370	N.S.		
Left ABI	202	-3.9975	5.7243	0.6984	N.S.		

"Dip" is defined as Awake-Sleep difference divided by Awake ($\times 100$), where Awake and Sleep are average values during awake- and sleep-span, respectively.