Published in final edited form as:

Int Rev Cell Mol Biol. 2009 ; 274: 69. doi:10.1016/S1937-6448(08)02002-9.

Hypothalamic neural systems controlling the female reproductive life cycle: Gonadotropin-releasing hormone, glutamate, and GABA

Jacqueline A. Maffucci¹ and Andrea C. Gore^{1,2,3}

¹Institute for Neuroscience, University of Texas, Austin, TX 78712, USA

²Division of Pharmacology and Toxicology, University of Texas, Austin, TX 78712, USA

³Institute for Cellular & Molecular Biology, University of Texas, Austin, TX 78712, USA

Abstract

The hypothalamic-pituitary-gonadal (HPG) axis undergoes a number of changes throughout the reproductive life cycle that are responsible for the development, puberty, adulthood, and senescence of reproductive systems. This natural progression is dictated by the neural network controlling the hypothalamus including the cells that synthesize and release gonadotropin-releasing hormone (GnRH) and their regulatory neurotransmitters. Glutamate and GABA are the primary excitatory and inhibitory neurotransmitters in the central nervous system, and as such contribute a great deal to modulating this axis throughout the lifetime via their actions on receptors in the hypothalamus, both directly on GnRH neurons as well as indirectly through other hypothalamic neural networks. Interactions among GnRH neurons, glutamate, and GABA, including the regulation of GnRH gene and protein expression, hormone release, and modulation by estrogen, are critical to age-appropriate changes in reproductive function. Here, we present evidence for the modulation of GnRH neurosecretory cells by the balance of glutamate and GABA in the hypothalamus, and the functional consequences of these interactions on reproductive physiology across the life cycle.

Keywords

GnRH; NMDA receptor; glutamate; GABA; hypothalamus; reproductive aging; puberty

I. Introduction

The development and maintenance of reproductive systems are necessary for the propagation of all vertebrate species. Each level of the reproductive hypothalamic-pituitary-gonadal (HPG) axis produces unique hormone(s) that regulate one another through feed-forward and feedback mechanisms. Not surprisingly, this system is remarkably complex and involves not only cross-talk among its components, but it also includes extrinsic inputs from other somatic and neural systems. This network needs to be considered in the context of the life cycle, as reproductive demands vary enormously through an organism's life from birth to death.

Correspondence: Dr. Andrea C. Gore, Division of Pharmacology & Toxicology, The University of Texas at Austin, 1 University Station A1915, Austin, TX 78712, Telephone: 512-471-3669, Fax: 512-471-5002, andrea.gore@mail.utexas.edu. Jacqueline A. Maffucci, Institute for Neuroscience, The University of Texas at Austin 1 University Station A1915, Austin, TX 78712 Telephone: 512-471-6311, Fax: 512-471-5002, maffucci@mail.utexas.edu.

This review will focus on the neural (hypothalamic) control of reproduction across the life cycle, focusing on the female rat, a model for which most information is published. Because the hypothalamic control of reproduction is highly conserved among vertebrates, many of the conclusions drawn in the current article are likely to be in place in other species. However, when relevant we will note species differences. Because of the primary importance of the hypothalamus in driving reproduction, we will specifically discuss the roles of the hypothalamic neurons that produce the neuropeptide gonadotropin-releasing hormone (GnRH), and their regulation by two opposing neurotransmitter inputs: glutamate and GABA. Glutamate and GABA are the principal excitatory and inhibitory neurotransmitters, respectively, of the entire central nervous system including the hypothalamus (van den Pol, 2003), so it is not surprising that they play key regulatory roles in the control of GnRH neurons. Numerous other neuroactive substances are also critically involved in integrating the function of GnRH neurosecretion and we refer readers to other papers on this subject (de la Iglesia and Schwartz, 2006; Gore, 2002; Malyala et al., 2005), as it is far beyond the scope of the current review to provide a comprehensive understanding of the more than fifty neurotransmitters, neurotrophic factors, and other regulatory molecules that affect the GnRH system. Here, the interactions of GnRH neurons, GABA, and glutamate in the hypothalamus will be considered at molecular, anatomical, and physiological levels. In addition, the potential role of glutamate and GABA neurons in the mediation of feedback effects of steroid hormones on HPG function will be considered. There is no question that this hypothalamic network is an integral part of the overall circuitry that is responsible for the proper development, maintenance and function of the reproductive axis in adulthood, and changes in this network may also contribute to its eventual decline through the life history.

II. The Hypothalamic-Pituitary-Gonadal (HPG) Axis

The capacity for vertebrates to reproduce is dependent upon a functional HPG axis. The hypothalamus encompasses a relatively small area of the brain at the ventral portion of the diencephalon, just below the thalamus. It contains neurons that release the decapeptide GnRH into the portal capillary system, where it is transported to the anterior pituitary gland, located just below the brain. Here, GnRH binds to its receptors on the gonadotropes and stimulates the release of the gonadotropins, luteinizing hormone (LH) and follicle-stimulating hormone (FSH). These are released systemically, and target their receptors on the gonads, where they bind and stimulate the release of steroid hormones (estrogens, progestins, and androgens in both females and males). These sex steroid hormones are transported through the general circulatory system, and as part of their activities, they feed back onto the pituitary and hypothalamus, inhibiting the release of GnRH and LH, completing a negative feedback circuit. Additionally, in females, just prior to ovulation, the estrogen-induced negative feedback changes to positive via a mechanism that is still not entirely understood but undoubtedly involves steroid-sensitive inputs to GnRH neurons. This causes stimulation of GnRH and LH, resulting in the preovulatory GnRH/LH surge that triggers ovulation in the females of species with spontaneous ovulation, including primates and rodents.

III. The Reproductive Life Cycle in Females

A. Development and the attainment of reproductive competence

Throughout the lifespan, the reproductive axis undergoes a series of changes. During late prenatal/early postnatal life, the HPG axis is hyperactive in a sexually dimorphic manner, contributing to sexual differentiation of the brain (Gore, 2008; McCarthy, 2008). Later in postnatal life, this hyperactivity declines and the HPG axis enters a quiescent period during which GnRH release decreases, resulting in a decline of gonadotropin and steroid hormone levels. During the pubertal transition, the quiescent period ends, with basal GnRH levels, GnRH pulse frequency and pulse amplitude increasing, thereby activating the HPG axis, and resulting

in the sexual maturation of the animal. This process includes the onset of folliculogenesis and ovulation in the female, spermatogenesis in the male, steroidogenesis in the gonads of both sexes, as well as the manifestation of reproductive behaviors. Although all three HPG levels become active during the pubertal period, the driving force is the hypothalamic GnRH neurons (Richter and Terasawa, 2001; Sisk and Foster, 2004). However, GnRH neurons have the ability to release the GnRH decapeptide long before puberty. Therefore, the attainment of reproductive competence involves not only molecular modifications of gene and protein expression of GnRH (Ebling and Cronin, 2000; Sisk and Foster, 2004), but also, a maturation of the hypothalamic neural circuitry that regulates the GnRH cells (Ebling and Cronin, 2000; Gore, 2002). Puberty involves changes in the excitatory and inhibitory neurotransmitters that regulate GnRH release, and of relevance to this article, is characterized by decreased GABAergic inhibition of GnRH neurons, and an enhancement of facilitatory effects of glutamatergic inputs to the GnRH system, among other neurotransmitters that modulate the timing of puberty (Clarkson and Herbison, 2006; Moguilevsky and Wuttke, 2001; Terasawa, 2005).

In female rats, once puberty has occurred, the adult animal experiences regular estrous cycles of 4–5 days that are divided into four phases: diestrus I, diestrus II, proestrus, and estrus. During diestrus I and II, GnRH and LH release are low, and estrogen (primarily estradiol) levels are at their lowest. As diestrus II transitions to proestrus, estrogen levels increase and concurrently switch from exerting a negative feedback onto the hypothalamus and pituitary, to positive feedback. This results in the GnRH/LH surge that triggers ovulation during estrus. The timing of reproductive behaviors is also coordinated by these cyclic hormonal fluctuations, with females becoming sexually receptive during this transition to estrus, the time that maximizes the likelihood of successful fertilization upon mating (Gore, 2008; McCarthy, 2008). Although this current article focuses on the rat model, a species that exhibits short estrous cycles (as do mice), other vertebrates, including non-human and human primates, undergo similar developmental processes, with the capacity to generate menstrual cycles first occurring during pubertal development. This latter process, while much more prolonged than the rodent's estrous cycle, also shares similar traits of negative and positive feedback of preovulatory GnRH/LH release (Maffucci and Gore, 2005).

Reproductive systems are subject to an array of biological clocks. A critical biological rhythm of reproduction is the pulsatile (circoral) mode of release of GnRH, and subsequently LH. In rats, pulses of GnRH/LH occur at intervals of roughly 30 minutes (Maffucci *et al.*, 2008), and in primates, the rhythm is on the order of every 1–2 hours (Terasawa, 1995; Terasawa, 2005). This mechanism is in place throughout the reproductive life of the organism, and is necessary for proper reproductive function (Belchetz *et al.*, 1978; Loucopoulos *et al.*, 1984; Wildt *et al.*, 1981). Notably, pulses of GnRH occur throughout reproductive cycles of females, and therefore are superimposed upon the longer cyclic changes in GnRH during estrous and menstrual cycles in rodents and primates, respectively. Even shorter biological rhythms of GnRH exist, as shown by electrophysiological recordings from GnRH neurons in explant cultures, with short (millisecond to second to minute) bursts of electrical activity, calcium currents, and fluxes in GnRH peptide. It has been proposed that the coordination of this activity results in the pulsatile GnRH pattern described above (Abe and Terasawa, 2005; Suter *et al.*, 2000). The reproductive axis of mammals is also subject to circadian/diurnal rhythms of 24 hours (de la Iglesia and Schwartz, 2006). Additionally, at least some mammalian species exhibit seasonal rhythms of reproduction, and even more specifically, GnRH (Ebling and Cronin, 2000).

Thus, in examining the development and maintenance of the female reproductive axis, all of these biological rhythms of reproduction must be considered. Here, we will focus on regulation of GnRH/LH pulsatile release and the preovulatory GnRH/LH surge.

B. Reproductive senescence

During aging, reproductive systems undergo declines in function. While this is most apparent in the case of women, who undergo menopause at mid-life, animals also experience partial or total reproductive failure (Maffucci and Gore, 2005). Rats have proven to be an interesting model for reproductive aging because they experience neuroendocrine changes prior to ovarian failure. At middle age (about 1 year), the rat estrous cycle becomes irregular, resulting in cycles greater than 5 days in length. As the animal ages further, it eventually undergoes a transition to an acyclic state, resulting in persistent estrus, in which no ovulation occurs. This transition is accompanied by an attenuation and decline of the preovulatory or steroid-induced GnRH/LH surge (Wise, 1984), as well as a diminution of the GnRH/LH pulsatile pattern (Maffucci *et al.*, 2008; Scarbrough and Wise, 1990). Additionally, changes to hypothalamic GnRH gene expression are seen (Bottner *et al.*, 2007; Gore *et al.*, 2002; Gore *et al.*, 2000a; Rubin *et al.*, 1997). Because these changes precede the loss of ovarian cyclicity and the capacity to ovulate, these data suggest that at least in rats, reproductive aging involves, and may even be causally linked to, a gradual failure of hypothalamic GnRH function.

However, there may be substantial species differences in the mechanisms for reproductive senescence. Women undergo follicular loss at mid-life, making it very difficult to ascertain whether there are also neuroendocrine changes at this same time, not to mention the impossibility of measuring hypothalamic hormones in humans, as this is a highly invasive procedure. Such measurements have been made in the rhesus monkey, and Gore *et al.* (Gore *et al.*, 2004) showed that pulsatile GnRH release *increases* in perimenopausal rhesus monkeys, presumably due at least in part to a loss of negative feedback from estradiol. Consistent with this finding, serum LH, FSH, and gonadotropin free- α subunit (a proxy for GnRH) increased levels of their release during the perimenopause in women and female monkeys (Downs and Urbanski, 2006; Gill *et al.*, 2002; Hall *et al.*, 2000; Woller *et al.*, 2002). However, later post-menopausal women do show a decrease in LH pulse amplitude and inter-pulse interval (Genazzani *et al.*, 1997; Hall *et al.*, 2000), suggesting immediate increases in gonadotropins due to the relief of estrogen stimulated negative feedback, but an eventual reversal of this elevation. Thus, while slight species differences may exist in hormonal profiles during this transition, likely due to the lack of ovarian degradation in the rat, overall trends in hormone release between aging rats and humans share some similarities.

In aging rats, despite these changes in pulsatile and surge release of GnRH, and aberrations in GnRH gene expression, there are actually relatively few changes that are *intrinsic* to GnRH neurons. That is, GnRH cell numbers and distributions change very little with aging (Funabashi and Kimura, 1995; Hoffman and Finch, 1986; Miller and Gore, 2002; Miller *et al.*, 1990; Rubin and King, 1994; Rubin *et al.*, 1984; Witkin, 1986). GnRH neuronal ultrastructure does appear to undergo age-associated change (Romero *et al.*, 1994), as does the association of GnRH cells with astrocytes (Cashion *et al.*, 2003). Thus, we invoke the broader GnRH neural network – the GnRH neurons together with their direct and indirect inputs from neurons, glia, and steroid hormone feedback – as an instigator of reproductive aging. This hypothesis is supported by observations of age-related changes in function of hypothalamic neurotransmitters and neurotrophic factors, and expression of their receptors and steroid hormone receptors. Some of these changes may occur directly upon GnRH neurons, others upon the more extended circuit of cells that regulates the GnRH dendrites, perikarya and neuroterminals more broadly. The net effect of these alterations with aging is a loss of drive upon GnRH release (Yin and Gore, 2006).

C. The GnRH-glutamate-GABA connection: an introduction

Glutamate and GABA are regulators of developmental-, adulthood- and age-related control of GnRH function. As the primary excitatory and inhibitory amino acid neurotransmitters,

glutamate and GABA, respectively, are widely expressed in cells throughout the central nervous system. They each have several classes of receptors, which in turn, are made up of subunits whose expression varies by developmental age, sex, and region (Brann and Mahesh, 2005; Clarkson and Herbison, 2006; Gore, 2001; Henderson, 2007). Of relevance to reproductive neuroendocrinology, GnRH neurons themselves express glutamate and GABA receptors, and this co-expression varies across the life cycle (Adams *et al.*, 1999; Bailey *et al.*, 2006; Eyigor and Jennes, 1996; Eyigor and Jennes, 1997; Gore *et al.*, 1996; Jung *et al.*, 1998; Miller and Gore, 2002; Sim *et al.*, 2000). In addition, hypothalamic regions that mediate effects of steroid hormone negative and positive feedback are abundant in glutamatergic and GABAergic receptors. Here, we provide evidence for the control of GnRH function by glutamate and GABA, and how this regulation changes across the reproductive lifespan in females.

IV. Glutamate and the HPG axis

Excitatory amino acid neurotransmitters, and their interactions with their receptors, are an extremely important mechanism of excitatory synaptic activity in the CNS. Of these, glutamate is the most abundant in the brain (Brann, 1995; Brann and Mahesh, 1994).

Immunohistochemical and *in situ* hybridization studies examining glutamate receptor localization in the hypothalamus have shown their presence in a variety of hypothalamic nuclei, including medial preoptic nucleus (MPN), median preoptic nucleus, arcuate nucleus (ARC), dorsomedial, and ventromedial hypothalamic nuclei, supraoptic nucleus, suprachiasmatic nucleus, and paraventricular nucleus (Brann, 1995; Jennes *et al.*, 2002). Additionally, immunohistochemical analysis of glutamatergic fibers in the hypothalamus has shown them to be in close proximity with GnRH perikarya in the preoptic area, and GnRH axons in the median eminence (Eyigor and Jennes, 1996; Kawakami *et al.*, 1998b).

A. Expression of glutamate receptors in the hypothalamus, and specifically on GnRH cells

There are three classes of ionotropic receptors to which glutamate binds, named after agonist binding (Table 1). GnRH neurons express all of these: N-methyl-D-aspartate (NMDA) (Gore *et al.*, 2002; Gore *et al.*, 1996; Gore *et al.*, 2000b; Gu *et al.*, 1999; Ottem *et al.*, 2002); kainate (Eyigor and Jennes, 2000), and DL- α -amino-3-hydroxy-5-methyl-4-isoxazole propionic acid (AMPA) (Bailey *et al.*, 2006; Gu *et al.*, 1999). Furthermore, these receptors are present not only on GnRH perikarya themselves, but also in a number of hypothalamic nuclei implicated in GnRH secretory control. These include anteroventral periventricular nucleus (AVPV), MPN, ARC, and median eminence (ME), the latter being an area rich in GnRH nerve terminals (Brann and Mahesh, 2005; Gu *et al.*, 1999; Yin *et al.*, 2007). Activation of ionotropic receptors by glutamate has important effects on GnRH activity and resulting reproductive function, and this likely occurs via both these direct and indirect circuits.

Conventionally, ionotropic glutamate receptors are designated as NMDA and non-NMDA (AMPA and kainate) glutamatergic subtypes. The NMDAR is believed to be a tetrameric receptor (Laube *et al.*, 1998; Mano and Teichberg, 1998; Rosenmund *et al.*, 1998), and is composed of multiple subunits (NR1, NR2a–d, NR3a–b; Table 1). In the hypothalamus, NR1, NR2a and NR2b are abundant and are co-expressed on GnRH cell bodies (Eyigor *et al.*, 2001; Gore *et al.*, 2000b; Herman *et al.*, 2000; Meeker *et al.*, 1994; Miller and Gore, 2002). Non-NMDARs form tetrameric channels which, for AMPA-type receptors, may contain any of four subunits (GluR1–4), all of which are present in the hypothalamus and are co-expressed with GnRH neurons, although GluR1–2 appear to be the most abundantly expressed (Bailey *et al.*, 2006; Eyigor *et al.*, 2001). Kainate receptors may contain any of five subunits (GluR5–7, KA1, KA2) and all of these have been identified in rat hypothalamus, with KA2 being the most common (Eyigor *et al.*, 2001; Meeker *et al.*, 1994). The KA2 subunit is also the only such subunit to be co-localized immunohistochemically with GnRH perikarya in female rats to date,

as shown by *in situ* hybridization (Eyigor and Jennes, 1996). In addition, Todman *et al.* (Todman *et al.*, 2005) assayed the mRNA contents of GnRH neurons of adult female mice with PCR amplification and oligonucleotide microarrays. Of the ionotropic receptors, they detected the presence of NR2b, NR2d, GluR1, and GluR3, as well as Grid 1 and Grid 2, receptors of a fourth and less well known ionotropic glutamate class (δ) that are not known to form functional channels in *Xenopus* oocytes or mammalian cells (Lomeli *et al.*, 1993; Yamazaki *et al.*, 1992). It should be noted that the lack of detection of other receptor subtypes does not necessarily reflect the absence of the mRNA, so these latter findings probably represent a subset of the total receptor subunit populations. As a whole, these immunohistochemical, molecular, and physiological assays indicate that GnRH perikarya have the capacity to respond directly to glutamatergic stimulation through a variety of ionotropic and metabotropic glutamate receptors.

The previous discussion has focused on the expression of glutamate receptors at the level of the preoptic area, both directly and indirectly, on GnRH cells. However, the GnRH neuroterminals also must be considered as a site of glutamatergic regulation. Using electron microscopy, Kawakami *et al.* (Kawakami *et al.*, 1998a) demonstrated in adult female rats that GnRH terminals in the median eminence co-express NR1 and KA2 immunoreactivity. Although the GnRH terminals were not found to express GluR1, GluR2/3 and GluR6/7 in that study, other, non-GnRH, terminals in the median eminence were found to co-express these receptors.

The GT 1–7 cell line, an immortalized GnRH cell line that was produced using targeted tumorigenesis technique (Liposits *et al.*, 1991; Mellon *et al.*, 1991) also provides evidence for the expression of glutamate receptor subunits on GnRH neurons. As GnRH cells are often difficult to isolate and maintain in culture, these cells offer a viable alternative to studying GnRH neuronal function, and direct effects of glutamatergic influence on the action of these neurons. However, there is some disagreement in the field about whether and which glutamate receptors are expressed on GT1 cells. For example, the expression of the NMDA receptor subunits NR1 and NR2b-2d mRNA were reported (El-Etr *et al.*, 2006; Mahesh *et al.*, 1999; Urbanski *et al.*, 1994). However, Mahesh *et al.* (Mahesh *et al.*, 1999) did not show the presence of GluR1, GluR4 and GluR6 mRNA, nor were NR1 or GluR1 protein present as determined by western blot analysis. Binding studies using AMPA, kainate, and NMDA specific antagonists suggested the absence or minimal effect of glutamate ionotropic receptors in this cell line (Mahesh *et al.*, 1999). However, recent preliminary work from our laboratory (Maffucci and Gore, unpublished) did verify the presence of the NR1 and NR2a-2d subunit mRNA using real-time PCR, and both NR1 and NR2a (but not NR2b-d) protein using Western Blot analysis. Furthermore, Garyfallou *et al.* (Garyfallou *et al.*, 2006) found GluR2 and GluR4, as well as KA2 in the GT1-7 cell line using gene microarrays and RT-PCR. The variability of these data are likely due to differences in individual cell lines, emphasizing the importance of characterization of cell lines prior to experimental use. At the very least the results suggest the potential for GT1 cells to express glutamatergic receptors under certain culturing conditions.

GT1 cells may also respond to glutamate agonists/antagonists. Incubation of these cells with AMPA, kainate, NMDA, or glutamate yielded increased GnRH release and an increase in intracellular calcium concentration (El-Etr *et al.*, 2006; Spergel *et al.*, 1994; Spergel *et al.*, 1995). Addition of the NMDAR antagonists MK-801 or AP-5 inhibited the NMDA-induced increase in GnRH release (Mahachoklertwattana *et al.*, 1994).

B. Structure-function properties of glutamate receptors

The structural properties of the ionotropic receptors determine their ability to bind and respond to glutamate. Specifically, the subunit composition of the receptor determines the kinetic properties of the channel, and thus understanding the stoichiometry of the receptor population,

as well as its distribution in various nuclei in the hypothalamus, appears to be fundamental to understanding its influence on the reproductive axis. Of relevance to this article, the stoichiometry of glutamate receptors undergoes substantial developmental changes, having potential implications for glutamate's target cells, including GnRH neurons. Moreover, the steroid hormone environment modulates the properties of glutamate receptors, making those cells that express the receptors potential targets of integration of steroid hormone influences on HPG function.

C. The glutamate-GnRH connection across the life cycle

1. Early development—The role of the NMDAR on reproduction changes throughout the female's lifespan. Some of these developmental differences may be attributable to either the absence of expression of the NMDAR in hypothalamic tissues, including on GnRH neurons, and/or the lack of a key receptor subunit that is required for the functionality of the receptor. This certainly appears to be the case during the infantile life stage. Our laboratory has quantified both gene and protein expression of NMDAR subunits in the preoptic area-anterior hypothalamus (POA-AH), a sexually dimorphic region that is important for reproductive physiology and sexual behaviors in male and female rats. Using rats across a range of early developmental ages, spanning late fetal development (day 18 of gestation) through early postnatal life (15 days of age), we reported age-, sex-, and subunit-specific differences in mRNAs of the NR1, NR2a and NR2b subunits of the NMDAR in the POA-AH (Adams *et al.*, 1999). Specifically, NR1 mRNA levels increased progressively during this developmental period, about 10-fold in males and 5-fold in females. NR2a mRNA was virtually undetectable in late embryonic life and remained extremely low in females through postnatal day (P) 15. In males, NR2a mRNA was first detectable on the day of birth and underwent a large increase from P10–P15. Finally, NR2b mRNA underwent an approximate doubling in levels from P0 to P5 in both males and females. Taken together, these results suggest that the ability of cells to synthesize NMDAR subunits changes dramatically during early life development in the POA, a finding that has implications for differential abilities of NMDARs to respond to glutamate (Adams *et al.*, 1999; Chen *et al.*, 1999; Erreger *et al.*, 2005; Vicini *et al.*, 1998). This same study evaluated whether GnRH neurons co-express the NR1 subunit during early postnatal development and did not detect any co-expression at these ages. Finally, when the NMDA receptor agonist NMA, or antagonist MK-801 was administered to these developing rats, little effect on GnRH mRNA or primary transcript RNA levels was detected (Table 2; Adams *et al.*, 1999). Thus, during the perinatal life period, the ability of glutamate to affect changes on the GnRH system is not due to direct actions via NMDARs on GnRH neurons, but rather, is either due to indirect inputs from other NMDAR expressing cells, or to non-NMDA glutamate receptors.

2. Puberty—To our knowledge, the earliest studies demonstrating an involvement of glutamatergic NMDARs on HPG function came from reports determining consequences of NMDAR agonists on the timing of puberty in male rhesus monkeys and female rats. In those experiments, prepubertal animals were treated with an NMDAR agonist, and the results showed an advancement in the timing of pubertal landmarks (Gay and Plant, 1987; Gay and Plant, 1988; Urbanski and Ojeda, 1987). Since then and as elaborated upon below, studies have investigated pubertal changes in responsiveness of GnRH neurons to glutamatergic agonists/antagonists, through measurements of serum LH, central GnRH, and/or GnRH gene expression, in coordination with evaluation of external markers of puberty such as secondary sex characteristics.

2a. Glutamate synthesis and release in the hypothalamus increases during puberty: The synthesis and release of glutamate in the POA-hypothalamus changes during the life cycle, including puberty, and this may have functional outcomes on the reproductive axis. *In vitro*

work by Bourguignon *et al.* (Bourguignon *et al.*, 1995) suggested that an increase in glutaminase activity, the enzyme responsible for conversion of glutamine into glutamate (Table 1), occurred in conjunction with the observed increase in GnRH pulsatile release accompanying pubertal onset. Furthermore, glutamate levels in the medial preoptic nucleus increased during the pubertal transition (Goroll *et al.*, 1993).

2b. Expression of glutamate receptors in the hypothalamus and on GnRH neurons

increases during puberty: NMDA and non-NMDAR expression in the hypothalamus undergo pubertal changes in mRNA and protein expression, and in the co-expression of their receptor subunits on GnRH neurons. For example, Gore *et al.* (Gore *et al.*, 1996) showed that NR1 mRNA levels in the POA-AH increased peripubertally, and using double-label immunohistochemistry, demonstrated that the co-expression of NR1 with GnRH neurons increased significantly during the time period when adult reproductive function was attained. Eyigor and Jennes (Eyigor and Jennes, 1997) used *in situ* hybridization to examine the co-expression of KA2 and NR2a with GnRH neurons in the rostral hypothalamus (medial septum-diagonal band complex and POA). Although they found no change in NR2a co-expression during the pubertal transition, KA2 was not only present in GnRH neurons, but showed a diurnal patterning throughout development, with higher expression levels in the morning from P20–P40 switching to higher expression levels in the afternoon at P45–P50 (Eyigor and Jennes, 1997).

The OVX, steroid-treated prepubertal rat has also been used as a model of glutamatergic regulation of GnRH function. Using *in situ* hybridization, expression patterns of GluR1-7 and NMDAR1 subunits in the AVPV, and changes in these patterns following estrogen or estrogen/progesterone treatment, were assessed (Gu *et al.*, 1999). The results showed differential expression of these various receptor subunits in different structures within the AVPV itself, and little evidence for the presence of kainate receptors (GluR5-7). Furthermore, an estrogen dependent increase in the GluR1 subunit and suppression of NR1 (as compared to OVX controls) was reported (Gu *et al.*, 1999) suggesting steroid regulation, a concept that will be discussed below (see subsection 3c).

2c. Functional changes in the response of the HPG axis to glutamate during puberty:

Not only does glutamate receptor expression in the hypothalamus in general and on GnRH neurons in particular change during the progression of puberty, but the ability of glutamate to affect GnRH gene expression is facilitated during puberty. Gore *et al.* (Gore *et al.*, 1996) found that administration of the glutamate agonist NMA to intact female rats enhanced the naturally occurring increase in GnRH mRNA expression that accompanies the onset of puberty (Table 2). A further study examining blockade of NMDARs or non-NMDARs during the steroid-induced LH surge in peripubertal (P28) OVX, steroid replaced females showed that MK-801 (NMDAR antagonist), but not CNQX (AMPA-kainate antagonist), caused a decrease in the GnRH mRNA levels in the rostral hypothalamus (Seong *et al.*, 1993).

The ability of NMDAR agonists and antagonists to influence the reproductive axis does not become apparent until the pubertal time period. It appears that there is a critical period ($\geq$ P21 in rats) during which administration of an NMDAR agonist will induce precocious puberty *in vivo* and increase GnRH release *in vitro* (Moguilevsky *et al.*, 1995; Smyth and Wilkinson, 1994). Prior to this age, NMDAR agonists/antagonists have relatively little effect on GnRH/LH release and the timing of puberty (c.f. Adams *et al.*, 1999). Beginning at about P21, a time period when rats first begin to undergo activation of the HPG axis, the activation of NMDARs results in the induction of precocious puberty (Brann *et al.*, 1993a; MacDonald, 1992; Smyth and Wilkinson, 1994; Urbanski and Ojeda, 1987; Urbanski and Ojeda, 1990), while NMDAR antagonists delay pubertal onset (MacDonald and Wilkinson, 1990; Meijs-Roelofs *et al.*, 1991; Urbanski and Ojeda, 1990; Veneroni *et al.*, 1990).

The functional role of non-NMDARs in this pubertal transition is less clear. Administration of kainate either systemically or directly (i.c.v.) into the third ventricle did not induce precocious puberty in immature female rats, nor did i.c.v. application of DNQX, an AMPA/kainate antagonist (Table 1), delay its onset (Brann *et al.*, 1993a). Further investigation into why this difference exists between NMDARs and non-NMDARs suggest that these receptors may show sexually dimorphic mediation of puberty. Comparison of the activity of NMDA and MK-801 versus kainate and DNQX (AMPA/kainate antagonist) on gonadotropin secretion in both male and female prepubertal rats showed that, while the NMDAR drugs affected both sexes equally, kainate stimulated LH, and DNQX inhibited LH, in males only (Pinilla *et al.*, 1998). Another study found slightly different effects on the gonadotropin response to kainate in males. While NMDA administration caused LH increases *in vivo* and GnRH *in vitro* in both males and females, kainate administration resulted in increased GnRH, but no effect on LH (Carbone *et al.*, 1992; Carbone *et al.*, 1996). These differential findings in gonadotropin release are likely due to variations in methodology, including drug dose administered or time of day administered. However, overall, these results support a role for these receptors in the functioning of the reproductive axis during this important transition, and suggest a possible sexual dimorphism with kainate receptor action. Differential regulation of gonadotropin secretion by glutamate could be the result of differential distribution, and hence alterations in the stoichiometry, of the receptors within specific regions of the hypothalamus and differential stimulation of the GnRH neuronal populations, whether via direct or indirect mechanisms. More research will be required in order to better understand the physiological influence of glutamate and its various ionotropic receptors during this important developmental time period.

It is important to point out that analyses of kainate and NMDAR binding in the hypothalamic area spanning the POA to MBH showed no change in the number of binding sites during the pubertal transition (Brann *et al.*, 1993a). This finding suggests that the effects of these receptors are via increased sensitivity to glutamate or increased sensitivity of GnRH neurons to their activity, rather than to the ability of glutamate to bind to its receptors. Changes to receptor stoichiometries during this time period also need to be considered, as it is plausible that while receptor numbers remain the same, subunit distributions differ, changing the electrical properties and hence physiological responses of these channels.

3. Adulthood

3a. Glutamatergic actions upon the GnRH/LH surge—As mentioned previously, reproductive physiology in adult females involves several rhythms of GnRH/LH release. Of primary interest to reproductive physiologists studying species with spontaneous ovulatory cycles, such as rats and primates (both human and non-human), are the patterns of GnRH/LH pulsatile release and the GnRH/LH surge that precedes ovulation. Both of these events are influenced by the action of NMDARs and non-NMDARS. With respect to the activation of GnRH gene expression and release during the surge, NMDAR activation in proestrous young adult female rats (Gore *et al.*, 2000b) led to rapidly increased GnRH gene expression in the hypothalamus (Table 2). The NMDAR antagonist MK-801 given to proestrous females not only blocked the LH surge, but also resulted in a less prominent increase in GnRH mRNA expression in the hypothalamus as compared to controls (Suzuki *et al.*, 1995). However, in OVX, steroid hormone treated females, kainate injection (i.c.v.) administered on the morning of the expected steroid induced surge did not result in upregulation of GnRH mRNA levels, nor did it induce c-Fos expression in GnRH neurons 40 minutes following injection. However, it did result in an increase in LH levels 10 minutes following treatment (Eyigor and Jennes, 2000). Together, these data suggest that kainate is working indirectly to affect LH release during this time period. Furthermore, application of AP-7 (NMDA antagonist) or DNQX (AMPA/kainate antagonist) into the third ventricle of OVX, E2 replaced females blocked the steroid-induced LH surge (Lopez *et al.*, 1990), as did administration of MK-801 to the

proestrous female (Brann and Mahesh, 1991). Administration of NMDA or NMA on the morning of proestrus resulted in an increase in LH levels shortly after (Brann and Mahesh, 1992; Luderer *et al.* 1993). Thus, activation of glutamate receptors contributes to the onset of the preovulatory LH surge, with roles being played by both NMDA and non-NMDA receptors.

3b. Glutamatergic effects on pulsatile GnRH/LH release—Both NMDAR and non-NMDAR agonists and antagonists affect the pulsatile release of GnRH and LH in a similar manner to their effects on the GnRH/LH surge. Agonists to both receptor types facilitate, and antagonists diminish/block, pulsatility. For example, pulsatile release of GnRH *in vitro* and LH release *in vivo* was diminished by administration of antagonists to NMDARs and non-NMDARs in both adult male and female rats (Arslan *et al.*, 1988; Bourguignon *et al.*, 1989b; Maffucci *et al.*, 2008; Ping *et al.*, 1994a; Ping *et al.*, 1995).

In vitro studies on explanted hypothalami have further implicated glutamate receptors as key regulatory factors of GnRH output. Application of AP-5 (NMDAR antagonist) and DNQX (non-NMDAR antagonist) to male hypothalamic retrochiasmatic explants resulted in an inhibition of GnRH output (Bourguignon *et al.*, 1989a). Analysis of NMDA or kainate application to median eminence fragments from the OVX female in the absence of estrogen demonstrated similar increases of GnRH output (Kawakami *et al.*, 1998a). Donoso *et al.* (Donoso *et al.*, 1990) found that administration of the agonist kainate to ME fragments that had been explanted from male adult rats caused a 1.8 fold increase in GnRH output *in vitro*, while NMDA required a higher concentration (20 mM) to elicit a GnRH response. It should be noted that most of these latter studies focused on GnRH nerve terminals in the median eminence, a region where kainate receptors and NMDARs are expressed (Maragos *et al.*, 1988; Unnerstall and Wamsley, 1983).

In vivo experiments further highlight the role of these receptors in adult reproductive physiology. Kainate injected directly into the medial preoptic nucleus of intact adult females on diestrus 1 resulted in increased GnRH content in OVLT and MBH, but gonadotropin output varied, with LH increasing, but not significantly, and FSH significantly decreasing (Gerendai *et al.*, 1980). Repeated injections of kainate or NMA (i.v.) to intact (Abbud and Smith, 1991) or NMA (i.v.) to OVX (Pohl *et al.*, 1989) female rats significantly increased LH release, but this elevation was not maintained with kainate. Again these findings support the potential for differential modulation of gonadotropin output based on the type of glutamate receptor activated.

3c. Steroid modulation of glutamate's effects on GnRH neurons—The steroid hormone environment has considerable effects on glutamate receptor activation of GnRH neurons and has been studied at length in adult females. This is an important area of research because GnRH neurons do not express nuclear estrogen receptor (ER) alpha (Shivers, 1983) and although they express estrogen receptor beta (ER β), this is not adequate to explain positive feedback effects of steroids upon the GnRH/LH surge (Wintermantel *et al.*, 2006). Thus, effects of estradiol and progesterone on this event are mediated to a large extent indirectly by inputs to GnRH cells. Observations that hypothalamic inputs to GnRH neurons express ER α corroborate this idea. In the case of glutamatergic regulation, immunohistochemical analyses show that steroid hormone receptors (ER α , ER β , androgen receptor) are co-expressed with NMDARs and non-NMDARs in various nuclei of the hypothalamus that are important to regulation of GnRH release (Chakraborty *et al.*, 2003a; Chakraborty *et al.*, 2003b; Diano *et al.*, 1997). Additionally, testosterone or estrogen replacement in gonadectomized adult males or females resulted in increases in GluR1 and GluR2/3 hypothalamic protein expression (Diano *et al.*, 1997). By contrast, no effect of estradiol on NR1 immunoreactive cell numbers in the AVPV was found (Chakraborty *et al.*, 2003a).

It appears that the duration of estrogen treatment, as well as the time of day that the endpoints are examined, play a role in determining actions of glutamate receptors on GnRH output. In general, estradiol enhances NMDA-induced LH release (Arias *et al.*, 1993; Gore, 2001; Maffucci *et al.*, 2008). An examination of both GnRH and NMDAR subunit mRNA expression in response to different durations of estradiol treatment suggested little effect of duration in the POA-AH (Brann *et al.*, 1993b; Gore *et al.*, 2002; Ottem *et al.*, 2002), but slight differences in the median eminence/medial basal hypothalamus (Gore *et al.*, 2002). Additionally, while NMDA administered in the morning (0700h) increased GnRH mRNA levels in OVX, estrogen treated females, MK801 administered in the afternoon (1200h) did not block the estrogen induced increase in GnRH mRNA levels that accompanies the estrogen-induced surge (Ottem *et al.*, 2002). Examination of MK-801 administration in the morning was not examined.

In focusing on AMPA receptors, Ping *et al.* (Ping *et al.*, 1997) found a differential effect of AMPA activation on GnRH release *in vitro* (MBH fragments from OVX, estradiol- or non-estradiol primed females) and LH *in vivo* (in OVX, estradiol- and non-estradiol primed females). The authors concluded that, while GnRH release increased regardless of the steroid hormone environment with AMPA administration and was reversed by administration of NBQX (AMPA specific antagonist), increases in LH were estrogen dependent. AMPA stimulated LH release in estradiol-primed females, but inhibited it in the absence of estradiol. Similar findings were seen when examining the effect of NMDA administration to OVX, steroid primed females (Arias *et al.*, 1993). Thus estradiol is a powerful regulatory factor in the modulation of this system by glutamate and its receptors.

3d. Glutamate release in the hypothalamus-preoptic area—The ability of glutamate to activate GnRH release is clearly contingent upon its release at GnRH cell bodies, terminals, and dendrites. However, relatively little is known about this phenomenon in adult rats. Measurement of glutamate release in the POA show that it is increased during the steroid hormone-induced LH surge in adult female rats as measured by push-pull perfusion (Jarry *et al.*, 1995; Ping *et al.*, 1994b) and microdialysis sampling (Neal-Perry *et al.*, 2005). These results are consistent with the proposed stimulatory role of glutamate during the GnRH/LH surge.

3e. Conclusions on effects of glutamate on GnRH neurons of adults—While these studies shed light on the role of these receptors, there are many questions yet to be answered. It appears that there may not only be differential effects of glutamate receptors on the reproductive axis, but that these effects may be mediated by the presence (or absence) of steroid hormones, and may be more or less robust depending upon the endpoint observed, including GnRH/LH pulsatility, the GnRH/LH surge, or other HPG axis responses at various time points during the cycle. The differential distribution patterns of these receptors throughout the hypothalamus suggests that specific nuclei may play different roles in regulating reproductive physiology in response to the presence (or absence) of glutamate. Thus, while we can conclusively state that these receptors play a role in the female reproductive physiology, the extent of that role, the individual contributions of each receptor, and their interplay, requires further research.

4. Reproductive Senescence

4a. Age-related changes in hypothalamic glutamate receptors—Middle-aged rats experience a reproductive decline that results in decreased fertility and fecundity. Previous studies support that some of these changes are effected at the hypothalamic level, and that excitatory amino acids are strongly implicated. Age-associated changes in expression of NMDAR and non-NMDARs in the hypothalamus have been shown that may underlie some of these functional differences. Gene expression of NR1, NR2a and NR2b subunits were measured in the POA-AH and medial basal hypothalamus (MBH) of young, middle-aged, and

aged intact animals of known cycling statuses (Gore *et al.*, 2000b). NR2a and NR2b mRNA expression in the POA-AH were higher in cycling (both young and middle-aged) than acyclic (both middle-aged and aged) animals. In the MBH there was an overall increase in NR1, NR2a, and NR2b subunit mRNA expression with age. While this study showed no significant change in NR1 gene expression in POA-AH, by contrast, Zuo *et al.* (Zuo *et al.*, 1996) found a decline in NR1 gene expression in both POA-AH and MBH regions in middle-aged versus young proestrus females using RT-PCR. The inconsistencies in findings may be due to the varied techniques used in each study, including tissue dissection, variation in times of sacrifice, mRNA extraction, and mRNA processing. However, both studies support the presence of important changes in NMDAR stoichiometry based on cycle status and age.

Ovariectomized rats have also been used to examine age-related changes in gene and protein expression of NMDAR subunits. Using young, middle-aged and old female rats that were OVX and given estradiol or vehicle treatment, Gore *et al.* (Gore *et al.*, 2002) showed significant effects of both age and hormone treatment on NMDAR subunit mRNA expression. Specifically, in POA, NR1 mRNA levels were significantly decreased by short-term (2 day) but not longer-term (2 week) estradiol. NR2a mRNA levels were also down-regulated by estradiol, particularly in middle-aged and old rats. NR2b mRNA levels were highest in old rats and treatment with estradiol increased these levels in the old group. In the MBH, NR1 mRNA levels were decreased in estradiol compared to vehicle treatment; NR2a mRNA was not affected by age or estradiol; and NR2b expression was up-regulated by 2 week estradiol treatment in young rats, and decreased by this treatment in old rats (Gore *et al.*, 2002). Therefore, these studies show the importance of regional specificity in the expression of these receptors, as results differed between the POA-AH and MBH.

To further pinpoint the anatomical site at which estradiol may regulate NMDAR expression, the number of cells expressing the NR1 protein was quantified in the AVPV of OVX, estrogen or cholesterol treated aging female rats, a region chosen for its importance in the preovulatory GnRH/LH surge (Terasawa *et al.*, 1980; Wiegand *et al.*, 1980). Results of quantitative stereology showed no effect of estradiol, but significant age-associated declines in NR1 immunoreactive cell numbers in the lateral AVPV (Chakraborty *et al.*, 2003a). Recently, we have performed stereologic analyses of NR2b subunit expression in the AVPV of OVX rats at three ages (young, middle-aged and old) and hormone treatments (Maffucci & Gore, unpublished; Figure 1). Preliminary results suggest a decrease in NR2b-immunoreactive cell density in the AVPV from young to aged animals regardless of hormone treatment (vehicle-vehicle, estradiol-vehicle, or estradiol-progesterone). Furthermore, the volume fraction of the AVPV represented by immunofluorescent cells expressing colocalized NR1 and NR2b in this region increased with age, whereas the volume fraction of either NR1 or NR2b immunofluorescent independently showed no age associated changes (Figure 1). Together, these data suggest that whereas the number of cells in the AVPV that express NR2b is decreasing with age, those cells that are remaining are more likely to co-express NR1 with NR2b. The consequence of such changes would be altered channel properties of these receptors, and hence their effects on the reproductive system (Maffucci *et al.*, unpublished).

4b. Co-expression of glutamate receptor subunits on GnRH neurons, and changes with age—By examining NMDAR protein expression changes directly on GnRH neurons of the OVLT-POA hypothalamic region, our lab has demonstrated no age-related change in NR1 co-expression, but an increase in the NR2b subunit co-expression with GnRH from young to middle-aged intact animals (Miller and Gore, 2002). Thus, there may be a change in the stoichiometry of the NMDAR on GnRH cells with aging, that may underlie some of the functional differences in the responsiveness of GnRH neurons to NMDAR agonists, as described below (see section 4c).

Co-expression of NMDAR subunits on GnRH neurons during the steroid-induced LH surge, a time when the GnRH response to glutamate agonists *in vitro* is attenuated in middle-aged females. Zuo *et al.* (1996) showed differences between middle-aged and young rats. Specifically, AMPARs (GluR1, 3–4) on GnRH neurons underwent altered expression, as compared to young animals, on proestrus (Bailey *et al.*, 2006). Specifically, GluR1/GnRH co-expression was lower in middle-aged females at 1200h than young and failed to decline at the end of the proestrus day (2000h) in the young but not the middle-aged rats. GluR3 co-expression showed a diurnal pattern that was age specific. Young animals showed higher expression on the proestrus morning whereas middle-aged animals had a higher number in the afternoon. Finally, GluR4 co-expression in young animals was elevated early in the day as compared to middle-aged females, but showed no difference on the afternoon of the surge. These changing expression patterns likely contribute to the age-associated differences in hypothalamic activity, and as discussed for NMDARs above, probably represent altered stoichiometries of non-NMDA receptors in the aging hypothalamus.

4c. Functional changes in responsiveness of GnRH neurons to glutamate with aging—Pharmacological studies addressed some of the functional consequences in changes in NMDAR stoichiometry and/or expression of NMDARs in hypothalamus and on GnRH neurons. *In vitro* application of NMDA to anterior POA-MBH fragments from young and aged OVX females resulted in increases in GnRH peptide concentrations, with young females showing a greater increase than the aged group (Arias *et al.*, 1996). In this same study, *in vivo* administration of NMDA to the POA resulted in a greater LH response of the young animals compared to aged females, and baseline LH levels prior to NMDA administration were higher in the young animals (Arias *et al.*, 1996). Zuo *et al.* (Zuo *et al.*, 1996) performed a similar study using ARC/median eminence fragments from intact, proestrus young and middle-aged females. No age-associated differences were seen in basal GnRH output, but application of NMDA caused increases in GnRH, with a significantly greater effect in young explants. Furthermore, young animals had significantly higher levels of LH than middle-aged. However, this study did not measure the effect of *in vivo* NMDA application on LH levels.

In assessing GnRH gene expression, NMA application to intact young and middle-aged female rats resulted in an increase in young, but a decrease in middle-aged animals (Gore *et al.*, 2000b). Thus, while GnRH release appears to be stimulated by NMDAR agonists, there appears to be an age-associated disconnection between release and GnRH mRNA, as well as an attenuation of the NMDA effect. Finally, a recent functional study of the role of the NR2b subunit in the reproductive senescent transition showed that inhibition of NMDARs containing the NR2b subunit with an NR2b-selective antagonist, ifenprodil, led to increases in overall LH levels and parameters of LH pulsatility in young and middle-aged OVX, E2 treated females (Maffucci *et al.*, 2008). In that study, little effect was seen on GnRH gene expression, suggesting that the action of ifenprodil is probably on the release of GnRH, independently of *de novo* GnRH biosynthesis (Table 2). Together, these results once again highlight the importance of NMDAR stoichiometry, particularly the presence or absence of an NR2b subunit, and suggests this subunit may cause inhibition of the GnRH neuron, either through indirect actions, or possibly through direct actions on GnRH neurons, as the NR2b subunit historically is associated with slower activation and deactivation channel kinetics (Chen *et al.*, 1999; Erreger *et al.*, 2005; Vicini *et al.*, 1998).

4d. Age-related changes in hypothalamic glutamate release—Glutamate levels in the POA decline during aging, as shown *in vitro* using male MPN-MBH explants (Bonavera *et al.*, 1998) and *in vivo* using intracerebral microdialysis in steroid-primed females (Neal-Perry *et al.*, 2005). Thus, beyond apparent changes to glutamate receptors themselves, there are accompanying changes in glutamate availability throughout the life cycle and it appears

that these levels are regulated, at least in part, by the steroid hormone environment (Carbone *et al.*, 1995; Jarry *et al.*, 1992).

V. Gamma-aminobutyric acid (GABA) and the HPG axis

GABA is the dominant inhibitory amino acid neurotransmitter in the brain, including the hypothalamus, where 50% of all synaptic contacts in the hypothalamus are estimated to be GABAergic (Decavel and Van den Pol, 1990). It is derived from the neurotransmitter glutamate via the enzyme glutamic acid decarboxylase (GAD; Table 3), which is found in two isoforms, GAD₆₅ and GAD₆₇, of which GAD₆₇ is viewed as the key enzyme for GABA synthesis (Asada *et al.*, 1997). This relatedness between glutamate and GABA, the fact that they are the dominant excitatory and inhibitory amino neurotransmitters in the nervous system, respectively, along with other data presented in the next section, strongly suggests a close interplay between the glutamatergic and GABAergic regulatory pathways.

As the key inhibitory amino acid in the adult hypothalamus, GABA is the main effector of inhibitory synaptic activity in this region (Decavel and Van den Pol, 1990). However, it has a very dynamic developmental progression. In addition to its role as a neurotransmitter, two of the GABA receptors (GABA_A and GABA_B; Table 3) have a large influence on the development of the central nervous system (Owens and Kriegstein, 2002). Anatomical studies of these receptors and their respective subunit compositions during this period support a developmental change, as well as a shift in their physiological responses to the neurotransmitter (Clarkson and Herbison, 2006; Moguilevsky and Wuttke, 2001). Specific to the hypothalamus, studies support GABA, particularly through its action on the GABA_AR, as affecting the establishment of sexually differentiated nuclei (McCarthy *et al.*, 2002). It is important to note that the expression of the GABA_C receptor appears to be primarily in the retina, superior colliculus, and cerebellum (Boue-Grabot *et al.*, 1998) and, although this receptor has been detected in the pituitary gland (Boue-Grabot *et al.*, 1995; Boue-Grabot *et al.*, 2000), it is not implicated in the establishment and maintenance of hypothalamic function. Taken together, these data suggest that, much like the glutamate receptors, the GABAergic system is undergoing developmental changes that have important repercussions on various neural systems, including the reproductive axis.

A. Expression of GABA receptors in the hypothalamus, and specifically on GnRH cells

GABA_A and GABA_B receptors have been localized to many nuclei of the hypothalamus, both via mRNA (Whiting *et al.*, 1997; Wisden and Seeburg, 1992) and protein (Davis *et al.*, 2000; Fenelon and Herbison, 1995; Fritschy and Mohler, 1995) detection methods. Even more specifically, the α , β , and γ subunits of the GABA_A receptor have been co-localized to hypothalamic GnRH neurons in female rats (Jung *et al.*, 1998), and ϵ subunits have been co-localized on GnRH cells in male rats (Moragues *et al.*, 2003). Ultrastructural studies have provided evidence for GABAergic synapses on GnRH neurons in the medial preoptic nucleus of the rat (Leranth *et al.*, 1985). Thus, it appears that, like glutamate receptors, GABARs can influence the GnRH neuronal network directly as well as indirectly.

The immortalized GnRH GT1-7 cells expresses GABA receptor subunits. The α and β , but not the γ subunits, of the GABA_AR have been detected (El-Etr *et al.*, 1995; Favit *et al.*, 1993; Garyfallou *et al.*, 2006; Hales *et al.*, 1992). Application of GABA to GT1 cells resulted in either a biphasic effect (an immediate stimulation of GnRH release followed by a suppression, suggested to be a result of two different GABARs working in concert) (Martinez de la Escalera *et al.*, 1994) or solely a decrease in GnRH secretion (Favit *et al.*, 1993; Martinez de la Escalera *et al.*, 1994). As with experiments testing glutamatergic responses, these differential results are likely due to differences in growing and passaging of the cell lines used in either study.

However, together, these *in vivo* and *in vitro* studies suggest a close link between GABA tone and GnRH function.

B. Structure-function properties of GABA receptors

There are three types of GABA receptors (GABARs), two of which (GABA_A and GABA_C) are ionotropic and one of which (GABA_B) is metabotropic. Both GABA_A and GABA_B appear to regulate the GnRH system. However, because the majority of research in rats has focused on the GABA_AR, we will focus our discussion predominantly on the ionotropic receptor GABA_A and mention other receptor types only when relevant.

The GABA_AR forms a pentameric structure for which sixteen subunits have been identified ($\alpha 1$ – $\alpha 6$, $\beta 1$ – $\beta 3$, $\gamma 1$ – $\gamma 3$, δ , ϵ , π , θ ; Table 3). Of these, α , β , and γ are the most common to receptor composition (Henderson, 2007). Using recombination techniques, it has been demonstrated that in order to activate a functional chloride channel, the GABA_AR must contain at least one α and one β subunit (Rabow *et al.*, 1995; Wisden and Seeburg, 1992). The composition of the GABA_AR is heterogeneous, regionally specific and changes with age. The stoichiometry of this receptor, as with glutamate receptors, is indicative of its channel properties and its specificity to various GABA_AR modulators. Additionally, specific subunits can also be modulated directly by steroid hormones, such as estrogens, having strong implications for the role of GABA_AR stoichiometry in the regulation of reproduction (McCarthy *et al.*, 1997).

C. The GABA-GnRH connection across the life cycle

1. Early development—A key component of hypothalamic development involves alterations in GABA levels, receptors and synapses. Not only do GABA subunit expression patterns in the hypothalamus change with development, but the nature of the effect of GABA undergoes a developmental switch from excitatory to inhibitory. Thus, in early development, studies suggests that GABA *stimulates* neural developmental processes, and later in development (e.g. puberty), it appears to play *inhibitory roles* including the mediation of negative feedback effects of steroids on reproductive physiology. It is also important to note that although pharmacological studies support an inhibitory effect of GABA on GnRH release during puberty, electrophysiological studies actually suggest that GABA may exert excitatory effects on GnRH electrical properties that are potentially mediated through actions of GABA through a more complex neural network (DeFazio *et al.*, 2002; Yin *et al.*, 2008; Moenter and DeFazio, 2005). These changes in GABA levels, tone, and physiology appear to be key players in the regulation of GnRH neural circuits.

Using radiolabeled dansyl chloride, a reagent that identifies primary amino groups, and hence is able to detect GABA levels, Cutler and Dudzinski (Cutler and Dudzinski, 1974) showed an increase in GABA content in the rostral hypothalamus beginning at week 1 and going through to adulthood (please note that in this study the sex of the animals examined were not identified). Davis *et al.* (Davis *et al.*, 1999) expanded on these findings using HPLC to examine individual hypothalamic nuclei and found that these increases are nucleus specific, with no change occurring in POA, but increases in VMN and ARC observed in both developing males and females from P1–P20.

GABA plays a role in the sexual differentiation of the hypothalamus. Margaret McCarthy's laboratory has shown the presence of sex and age dependent variations in GAD mRNA expression and GABA neurotransmitter levels in specific hypothalamic nuclei during the developmental time period (E18-P10) when organizational sex differences develop (McCarthy *et al.*, 2002; McCarthy *et al.*, 1997). Additionally, the GABA_AR shows very interesting changes in both structure and function through development in the hypothalamus. *In situ* hybridization studies examining the distribution of α , β , and γ subunit mRNA in male and female rats

throughout early development (P0, 1, 3, 5, 6, 7, 10, 14, 21, 35, and adult, defined as 200–250 g) suggest a switch in the expression of the $\alpha 1$ and $\alpha 2$ subunits, with $\alpha 1$ showing an increase, and $\alpha 2$ a decrease in many forebrain nuclei, including hypothalamus. Additionally, it appears that the $\beta 3$ subunit remains ubiquitously expressed throughout development (Laurie *et al.*, 1992; Zhang *et al.*, 1992; Zhang *et al.*, 1991). While these studies identified the distribution of many other receptor subunits, the developmental profiling was limited to $\alpha 1$, $\alpha 2$ and $\beta 3$. Thus, other GABA_AR subunits may change expression patterns with development, but have yet to be examined in this context.

Examination of GABA_AR protein expression of the alpha (1 and 2) subunit in both males and females from P1 through to adulthood found developmental variations in the levels of the GABA_AR- $\alpha 1$ subunit and GABA_AR- $\alpha 2$ subunit, dependent on age and the hypothalamic nucleus studied (Davis *et al.*, 2000). In the ventromedial nucleus (VMN), the GABA_AR- $\alpha 1$ subunit decreased while GABA_AR- $\alpha 2$ increased, whereas the POA showed no change in either subunit, and the arcuate nucleus showed an increase in both. A different study found that ³H-muscimol binding (a GABA_AR agonist; Table 3) in P20 animals was significantly higher as compared to earlier time points in the POA, but not VMN (Davis and McCarthy, 2000). Clark *et al.* (Clark *et al.*, 1997) detected an increase in binding in the VMN between P7–P14, with a decline occurring by P70. These are not in complete disagreement with the findings of Davis and McCarthy (Davis and McCarthy, 2000) as the latter showed a slight, but non-significant increase between P10 and P20. Together, these suggest that there is an increase in GABA_AR activity during this transitional period, and an alteration in receptor activity based on nucleus specific changes in receptor stoichiometry.

Expression of GABA subunits has also been studied specifically within developing GnRH neurons. Temple and Wray (Temple and Wray, 2005) examined embryonic day (E) 11.5 mouse nasal explants at various time points (culture day 3.5, 7, 14, and 28) to determine the expression patterns of the GABA_AR subunits $\alpha 1$ – $\alpha 4$, $\alpha 6$, and $\beta 3$ using PCR analysis, and to ascertain their changes as a function of age. Some significant changes in expression were seen, suggesting alterations in the receptor stoichiometry during development *in vitro*. At culture day 28, GABA_AR- $\alpha 2$ expression in GnRH neurons increased approximately 20–33% from previous ages, to 75%, and $\alpha 6$ decreased from 83–100% to 25% at day 28. $\alpha 1$ maintained low expression levels throughout the time course (20–38%), while $\alpha 3$ remained at high levels (50–100%). Expression of $\alpha 4$ and $\beta 3$ were variable (0–50%) but showed no significant differences. Thus, even at the embryonic stage, these subunits are modifying expression levels with development.

The excitatory nature of GABA in early development is thought to influence neuronal growth, suggesting that this excitatory mechanism is functional during a time of neuronal migration, neurite outgrowth, and differentiation of various neurons (van den Pol, 1997). To examine the role of GABA on GnRH development, mouse embryonic olfactory tissues, in which GnRH neurons are present, were cultured and incubated with bicuculline, a GABA_AR antagonist (Fueshko *et al.*, 1998). Measures of neuronal outgrowth and migration showed no effect on GnRH cell number or fiber length, but did show an increase in the distance of neuronal migration (measured by the migration of GnRH-immunopositive cells from their olfactory pits). In that same study, incubation with the GABA_AR agonist, muscimol, resulted in an inhibition of this migration (Fueshko *et al.*, 1998). A recent live imaging analysis of GnRH neuronal migration (from embryonic mouse tissue) showed that application of bicuculline resulted in an increase in the percentage of video frames showing motion, and a decrease in the percentage of frames showing turns (Bless *et al.*, 2005). This, in total, indicates that bicuculline application caused more forward motion, and less lateral motion. Together these two studies imply that the GABA_AR plays a role in the facilitation of GnRH embryonic development and migration.

2. Puberty—There has been extensive research on the potential role of GABA as a mediator of puberty, with the long-held hypothesis that a removal of GABAergic input to GnRH neurons is necessary to enable the increase in pulsatile GnRH release that drives pubertal development. Evidence for this hypothesis has been provided by a number of different techniques, although there has been some disagreement in the literature as discussed below. It is noteworthy that push-pull perfusion measuring GABA levels in the POA of prepubertal (26 and 32 days) and post-pubertal (P45) female rats demonstrated a pubertal decrease in GABA (Goroll *et al.*, 1993), consistent with a pubertal diminution of GABAergic tone on GnRH neurons.

Examination of GABA_AR gene and protein levels in GnRH neurons during pubertal development supports this trend of age-associated changes in GABA subunit expression. Jung *et al.* (Jung *et al.*, 1998) determined that the GABA_AR subunits $\alpha 1$ and 2, $\beta 2/3$, and $\gamma 2$ mRNA were all expressed within the GnRH neuron in female prepubertal (28 day) rostral hypothalamic regions (including medial septum, diagonal band of Broca, OVLT, POA, and rostral regions of the anterior hypothalamus), whereas $\gamma 1$ mRNA was absent. In addition, Sim *et al.* (Sim *et al.*, 2000) compared juvenile (P15–20) and adult (P50–70) mRNA expression of GABA subunits in GnRH neurons of female mice and showed that juvenile GnRH neurons express a much more heterogeneous population of GABA_ARs. Alpha 5, $\beta 1$, and $\gamma 2$ were the most frequently co-expressed subunits, and excepting $\alpha 4$ and $\gamma 1$, all subunits that were probed ($\alpha 1$ –5, $\beta 1$ –3, and $\gamma 1$ –3) were detected. In general, these were detected at low frequencies in the juvenile hypothalamus, suggesting a very heterogeneous population of receptors, whereas the adult hypothalamus showed much less variation. Alpha 1, $\alpha 3$, 5, $\beta 1$, $\beta 3$, and $\gamma 2$ were detected at high frequencies and specific combinations of these subunits were much more common. Additionally, more GnRH neurons co-expressed GABA_AR subunits in the adult (Sim *et al.*, 2000).

It is noteworthy that GABA_BR subunits also change protein expression patterns in the hypothalamus during development. Western blot analysis of the GABA_BR-1a and 1b splice variants in female hypothalamic tissue at P1, P4, P12, P20, P28, and P38 showed an overall decline in both variants with age (Bianchi *et al.*, 2005). Together, these findings support an anatomical change in the GABAergic system through development, both in the hypothalamus in general and on GnRH neurons in particular.

To further understand how these anatomical changes may be represented by physiological output, electrophysiological studies examining the response of hypothalamic neurons to GABA stimulation or inhibition have been essential. Recordings taken from embryonic day 15 (E15) neurons in culture show that application of GABA induces an excitatory, rather than the expected inhibitory response in early neuronal cultures. This response gradually shifts to inhibition as the cultures age (Chen *et al.*, 1995; Chen *et al.*, 1996). Such a finding is consistent with results of Han *et al.* (Han *et al.*, 2002). Here, GnRH neurons recorded from slice preparations of female immature (P10–17), peripubertal (P25–30) and adult (P36–55) female mice showed differential responses to GABA application. Specifically, immature neurons showed a depolarizing response to GABA, peripubertal neurons exhibited a variable response of depolarization, hyperpolarization, or no response, and adult neurons were consistently hyperpolarized (Han *et al.*, 2002). These findings suggest a developmental switch in the GABAergic response, in which initially, this system is stimulatory and then becomes inhibitory peri-pubertally. However, a study using similar recording techniques reported that GABA application actually depolarized GnRH neurons in adult (>42 days) female mice (DeFazio *et al.*, 2002). A reconciliation of these results has not been made to date but differences may be attributable to direct vs. indirect actions of GABA on GnRH cells (DeFazio *et al.*, 2002; Yin *et al.*, 2008; Moenter and DeFazio, 2005). These studies will be expanded upon below (section 3e).

The translation of these electrophysiological data to the whole organism is crucial to understanding the importance of the GABA regulation of GnRH function. As mentioned previously, GnRH is released in a pulsatile manner, and this mechanism is functional in early postnatal life. However, in prepubertal animals, GnRH release enters a period of relative quiescence, during which the amplitude of pulses declines substantially, and then increase during the pubertal transition (Dohler and Wuttke, 1975; Hompes *et al.*, 1982; Reiter and Grumbach, 1982; Roth *et al.*, 1997). It has been suggested that the establishment of this quiescent period early in development is likely due, at least in part, to an increase in inhibitory GABAergic activity, and that the reversal of this process in the peripubertal period enables an increase in excitatory input to GnRH neurons.

The period of the pubertal transition is one in which pharmacological properties of the GABAR undergo interesting changes. Activation of the GABA system using muscimol (a GABA_AR agonist), baclofen (a GABA_BR agonist), or aminooxyacetic acid (AOAA: a GABA-transaminase inhibitor; see Table 3) peripubertally (P23–P30) resulted in an inhibition of GnRH release *in vitro*, LH release *in vivo* and the onset of puberty (Feleder *et al.*, 1999). It appears likely that there is also a shift sometime between P20 and P30 whereby the GABAR, and more specifically the GABA_AR, changes from an excitatory to an inhibitory action on the HPG axis (Brann *et al.*, 1992; Moguilevsky *et al.*, 1991; Szwarcfarb *et al.*, 1994). As evidence, differential effects of muscimol and baclofen on GnRH *in vitro* and LH and FSH *in vivo* in prepubertal (P16) and peripubertal (P30) females have been demonstrated (Moguilevsky *et al.*, 1991). While baclofen (the GABA_BR agonist) remained inhibitory on these HPG endpoints at both ages, muscimol (a GABA_AR agonist) was stimulatory at P16, and inhibitory at P30. Brann *et al.* (Brann *et al.*, 1992), using estradiol-treated females, found that muscimol remained stimulatory on basal LH levels at P29 both *in vivo* and *in vitro* (the latter from hemipituitaries), but found no effect on GnRH release *in vitro*. These studies suggest that the pubertal period is a critical one for a shift in GABA receptor pharmacologic responses.

Analysis of the muscimol (GABA_AR) binding sites in female rats throughout development (P5–P45) showed that females have high levels of binding until the prepubertal timepoint (P15), at which point they experience a decline (Szwarcfarb *et al.*, 1994). Additionally, administration of AOAA to females resulted in a stimulation of LH levels at P12, 16, and 18 with AOAA treatment, but a significant decrease at P30. In total, these data suggest that GABA binding to the GABA_AR changes with development. However, further studies examining changes in GABA_BR binding are warranted to fully understand the physiology of these receptors, and their regulatory roles, during the pubertal period.

3. Adulthood—The role of GABA in the adult female reproductive system has revealed both stimulatory and inhibitory GABAergic mechanisms that remain controversial to this day. An early pharmacological study in the OVX and OVX steroid hormone primed female rat showed an increase in LH release following GABA administration. In OVX females (steroid hormone primed were not tested), pre-injection with bicuculline blocked the GABAergic rise in LH, suggesting an excitatory effect on this system (Vijayan and McCann, 1978). Using the GnRH GT1-7 cell line, Martinez de la Escalera *et al.* (Martinez de la Escalera *et al.*, 1994) demonstrated a biphasic effect of GABA application, during which an immediate stimulation of GnRH release was observed, followed by a tonic inhibition. The application of specific GABA_AR and GABA_BR agonists suggested that this stimulatory effect was GABA_AR mediated, results that were replicated and expanded upon by Hales *et al.* (Hales *et al.*, 1994). An experiment by Virmani *et al.* (Virmani *et al.*, 1990) used rat pituitary cells in culture, and applied GABA or muscimol either using a static incubation set-up, or a perfusion method with drug administration every 10 minutes. In the static set-up, LH release increased in response to GABA or muscimol, whereas in the perfusion experiment, LH levels progressively decreased with application of both agents. Thus, pulsatile application of GABA and GABAergic drugs

may prove more relevant in determining GABAergic function. In addition, these studies suggest direct effects of GABA on the pituitary level of the HPG axis, along with the considerable information showing hypothalamic actions of GABA on GnRH cells.

3a. Pulsatile LH release: The question of the role of GABA in the pulsatile release of LH has been extensively studied, and results have been variable. This is likely due to 1) the models examined, including whether or not rats were OVXed, for how long, and if they received hormone replacement; 2) the dose of agonist or antagonist used; 3) the method of drug delivery; and 4) the brain region examined. To follow is evidence for the pulsatile release of GABA in the hypothalamus, as well as effects of pharmacological manipulations of GABARs on GnRH/LH pulsatile release.

GABA release has been measured in the POA and MBH of adult female rats and shown to be pulsatile (Jarry *et al.*, 1992; Jarry *et al.*, 1988). However, the correlation of GABA pulses with LH (and presumably GnRH) pulses is debatable, as there is evidence for both the absence (Demling *et al.*, 1985; Jarry *et al.*, 1991) and presence (Jarry *et al.*, 1988) of such a correlation. This variability is likely due to the different techniques, or possibly different areas within the POA-AH, being probed.

The POA-AH, the site of GnRH perikarya, has been a main focus of studies examining the effects of manipulations of GABA levels or receptors on GnRH/LH pulsatility. Administration of AOAA (Table 3) to OVX female rats resulted in a decline of mean LH levels, as well as the frequency of pulsatile events and the interpulse interval. Bicuculline (GABA_A antagonist) prevented this effect, and bicuculline also increased mean LH levels (Donoso and Banzan, 1984), suggesting a role for GABA_ARs in this mechanism. Complementary to that finding, muscimol (GABA_A agonist) treatment resulted in an inhibition of LH pulses (Akema *et al.*, 1990). Furthermore, unilateral lesion of the POA resulted in a temporary (2 hour) decline in mean LH levels, LH amplitude and the number of pulses in OVX females, which recovered following a 48-hour period (Jarry *et al.*, 1993). Application of GABA (i.c.v.) into the POA-AH provided the same outcome (Jarry *et al.*, 1993). Taken as a whole, these data suggest a role of GABA in the regulation of GnRH/LH pulsatility, but support the idea that GABA is not the only neurotransmitter regulating this event, and hence even in the presence of GABA, compensatory mechanisms may be in place to recover lost pulsatile function.

Two studies examining OVX female rats found similar GABAergic effects using bilateral ablation techniques, but reported unexpected results when administering bicuculline (Herbison *et al.*, 1991; Jarry *et al.*, 1991). Bicuculline resulted in a decrease in mean LH levels (in both studies) as well as a decrease in pulse amplitude and frequency (Jarry *et al.*, 1991) or an overall absence of pulses (Herbison *et al.*, 1991). This may indicate a disinhibition of the pulsatile system, specifically in reference to the GABA_AR, whereby inhibition of this receptor causes stimulation of an inhibitory response. The interplay of this receptor with GABA_BRs likely adds another potential explanation for this mechanism. Moreover, we remind readers of the previously discussed studies showing some stimulatory effects of the GABAergic system in GnRH/LH release, which may come into play to explain the inhibitory effect of a GABA_AR antagonist on pulsatile LH release. Clearly more research on the precise mechanisms by which GABA affects the HPG axis is warranted.

The role of the GABA_BR in LH pulsatility is not as well-studied as that of the GABA_AR. To our knowledge, there is only one study examining this receptor subtype on pulsatile LH release. Administration of baclofen (GABA_BR agonist; Table 3) to OVX female rats led to a very complex response of the pulsatile properties of LH release, and the extent of these outcomes were dose dependent (Akema and Kimura, 1992). To summarize, the authors found a temporary ablation of LH pulses, followed by an increase in pulse amplitude and inter-pulse interval.

Biphasic and triphasic fluctuations in mean LH release were also seen (Akema and Kimura, 1992).

In total, this evidence suggests that both GABA_ARs and GABA_BRs are involved in a complex regulation of GnRH/LH pulsatility, and likely influence different endpoints of hormone release. It would be interesting to examine the interplay of these two receptors in the regulations of pulsatile output, as it is likely that these two are interacting, rather than acting as separate entities. This likely could explain the differential results observed with bicuculline administration, as well as the complex response to baclofen administration. Finally, researchers must strive further to examine smaller nuclei in the hypothalamus, as evidence suggests a finely defined neuroanatomy and neurochemistry in these regions that may underlie very discrete and different physiological responses.

3b. The GnRH/LH surge: The literature concerning the role of GABA in the GnRH/LH surge suggests this neurotransmitter's strong involvement in this physiological process. Studies examining GAD₆₇ mRNA levels as an indicator for GABA synthesis during the preovulatory (Grattan *et al.*, 1996; Herbison *et al.*, 1992) or steroid induced (Curran-Rauhut and Petersen, 2002) LH surge overall support region specific declines in GAD₆₇ on the afternoon of the surge (summarized in Table 4). Although these studies differ slightly in their findings for specific hypothalamic nuclei, likely due to differences in methodology, they all demonstrate an active involvement of GABA in the rostral hypothalamus in the regulation of the GnRH/LH surge. Specifically, a diminution of GABA biosynthesis precedes the generation of the large increase in GnRH that occurs during the surge. A study by Leonhardt *et al.* (Leonhardt *et al.*, 2000) contradicts these findings, suggesting that GAD₆₇ mRNA in the intact female increases during the afternoon of proestrus. However, these findings are likely due to their use of the whole POA rather than a focus on specific nuclei within this region, as well as the use of competitive RT-PCR rather than *in situ* hybridization, as was used in the former studies (Table 4). Additionally, findings by Herbison *et al.* (Herbison *et al.*, 1992) suggest that GABA synthesis in the medial preoptic nucleus and diagonal band of Broca specifically, as measured by GAD₆₇ levels, changes only during the proestrous surge, and not throughout the estrous cycle, although multiple timepoints throughout the cycle should be looked at to further develop this timeline.

In further considering the estrous cycle, examination of GABA levels in the whole hypothalamus of intact female rats showed a significant decline on the proestrus day, as compared to estrus and diestrus (Loscher *et al.*, 1992). Further, regionally specific changes in GABA levels were demonstrated using both push-pull perfusion in the POA-AH (Demling *et al.*, 1985; Jarry *et al.*, 1988) and microdialysis in the medial preoptic nucleus (Tin Tin Win *et al.*, 2004) of OVX-estradiol primed females. These studies demonstrated a decrease in GABA levels just prior to the LH surge, whereas levels in the median eminence/medial basal hypothalamus did not decline (Jarry *et al.*, 1995). It appears that the observed decline in GABA in the medial preoptic nucleus is specific to the proestrous afternoon, as very few changes were seen throughout the rest of the cycle (Mitsushima *et al.*, 2002). This decline in GABA prior to the GnRH/LH surge presumably works to disinhibit GnRH neurons, thus allowing the GnRH release to increase. This mechanism is likely to involve actions on GnRH biosynthesis and not on neurosecretion *per se*, as changes appear to be limited to POA-AH and are not seen in the median eminence.

Other enzymes involved in GABA metabolism appear to be involved in the LH surge. Administration of GABA-transaminase (GVG; Table 3), an enzyme that inhibits uptake and recycling of GABA, to intact, proestrus females resulted in an inhibition of the LH surge and the absence of ovulation (Donoso and Banzan, 1986). Application of AOAA, which inhibits GABA breakdown (Table 3) reduced the amplitude of the steroid-induced LH surge (Donoso

and Banzan, 1984), presumably by maintaining relatively high levels of GABA. However, unilateral administration of bicuculline followed by electrochemical stimulation of the medial preoptic nucleus caused a decline of LH levels and an inhibition of ovulation by a mechanism that at least in part involves opiate receptors (Taleisnik and Haymal, 1997). This suggests that GABA regulation of the GnRH/LH surge involves both direct and indirect actions on GnRH neurons.

The respective roles of the GABA_AR and the GABA_BR on the GnRH/LH surge remain largely unknown. Whereas direct and continual infusion of GABA into the medial preoptic nucleus (Herbison and Dyer, 1991) and injection into the third ventricle (Morello *et al.*, 1989) of proestrous females resulted in an inhibition of the LH surge, administration of bicuculline prior to GABA recovered it, suggesting a role for the GABA_AR in this action. Local injection of muscimol into the third ventricle (Akema *et al.*, 1990; Akema and Kimura, 1993) or medial preoptic nucleus (Seltzer and Donoso, 1992) also inhibited the estradiol-induced LH surge, and bicuculline administration reversed this effect. These studies all support a role of the GABA_AR in the LH surge regulation.

For the GABA_BR, bilateral infusion of baclofen (GABA_BR agonist) at either a low or high dose into the third ventricle of OVX, estradiol primed females resulted in a delayed (low dose) or inhibited (high dose) LH surge (Akema and Kimura, 1991). Follow-up studies from that same laboratory, in which baclofen was administered to OVX females and LH pulsatile release monitored, showed that baclofen interfered with GnRH/LH pulsatility, causing an immediate disruption in pulsatility (low dose) or a decrease in pulse frequency and increase in amplitude (high dose). The authors suggest that the GABA_BR-induced inhibition of the LH surge may be in part due to an interference with the normal pulsatile rhythmicity (Akema and Kimura, 1992).

In an attempt to understand how the GABA_A and GABA_B receptors work as a unit to regulate the surge mechanism, Adler and Crowley (Adler and Crowley, 1986) administered muscimol or baclofen to intact, estradiol + progesterone primed females, resulting in a decline in LH levels during the LH surge. This decline was not recovered by the simultaneous administration of bicuculline when administered with baclofen, but was slightly recovered in those animals receiving simultaneous muscimol (however, not to the level of control animals). Together, these studies suggest that both the GABA_AR and GABA_BR affect the action of the surge, but still leave unanswered to what extent, if any, these receptors interact to do so. Future studies should focus on revealing the roles of these receptors in the surge, specifically examining their role in the timing of the surge, its amplitude, and other physiological components.

The timing of the LH surge in rats is predictable and tightly regulated, and coordinated to the time of day as driven by the circadian pacemaker. Of relevance, GABA regulation of the GnRH circuit is dependent upon the time of day. Morello *et al.* (Morello *et al.*, 1992) found that administration of GABA into the third ventricle of proestrus females at 1100h inhibited the onset of the LH surge, whereas administration at 1300h had no effect. Further examination implicates the GABA_AR system as a timekeeper for the surge. Bicuculline administration produced similar effects as GABA, in that earlier administration (between 0900–1100h) had no effect on the pre-ovulatory LH surge, but administration at 1000–1200h caused a premature surge (Kimura and Jinnai, 1994) with higher amplitude (Kimura and Jinnai, 1994). Funabashi *et al.* (Funabashi *et al.*, 1997) examined co-expression of GnRH neurons with c-fos during the pre-ovulatory LH surge, and the effect of bicuculline administration thereupon. The bicuculline-associated advancement of the surge was replicated, and c-fos expression on GnRH neurons in rostral hypothalamic nuclei did not differ between those tissues collected in late afternoon from controls and early afternoon from bicuculline-treated animals, whereas those collected in late afternoon from controls had much lower expression levels. Thus, the

GABA_AR appears to act as a timekeeper for the GABAergic actions on the LH surge. To our knowledge, the role of the GABA_BR in this process is unknown.

3c. GnRH gene expression: Few studies have examined the role of GABA in GnRH gene regulation. Studies to date have suggested either no change, an increase, or a decrease in GnRH gene expression based on the pharmacology (GABA_AR or GABA_BR) of the agonist or antagonist used in specific studies (Bergen et al., 1991; Cho and Kim, 1997; Kang et al., 1995; Leonhardt et al., 1995). These differences are likely dependent upon differing methodologies, absence or presence of steroid hormones, and dissection of the hypothalamic region. As summarized in Table 5, in OVX rats, GABA antagonists (both GABA_A and GABA_B) tend to increase, and agonists decrease, GnRH mRNA. When OVX rats are treated with estradiol, these GABA agonists tend to increase GnRH mRNA, an effect that is opposite of that in OVX rats that are not estrogen primed. Clearly, much more work is needed to clarify these interactions among GABA, estrogens, and GnRH neurons at the level of gene expression.

3d. Steroid regulation of GABAergic systems: As mentioned previously, the steroid hormone environment is critical to consider when examining the neurotransmitter regulation of the HPG axis. There is ample evidence to suggest steroid hormone regulation of GABA, beginning with co-expression of estrogen receptors with GABAergic neurons (as labeled by a GAD-specific antibody; GAD 65 or GAD-67 not specified) in the POA-AH (Flugge *et al.*, 1986). Furthermore, the GABA_AR- α 2 subunit is co-expressed with estrogen receptor in the medial preoptic nucleus and bed nucleus of the stria terminalis, and the expression of this subunit, along with GABA_AR- β 3, is regulated in these regions by estrogen (Herbison and Fenelon, 1995).

Ovariectomy (OVX) in the adult female rat results in an increase in GnRH and LH basal levels due to the absence of estrogen-regulated negative feedback. In comparing GABA levels in the medial preoptic nucleus of OVX female rats to intact, diestrous females, as measured by push-pull perfusion, levels were significantly lower in the OVX rats, consistent with the higher GnRH release (Ondo *et al.*, 1982). Furthermore, bilateral infusion of GABA into the third ventricle of OVX females resulted in an acute increase in LH levels (within 30 minutes of this infusion); injection of bicuculline resulted in a slight and acute decrease. However, the animals treated with estrogen exhibited no such changes (McCann *et al.*, 1984). Lamberts *et al.* (Lamberts *et al.*, 1983) replicated these findings in the OVX female, showing a decline in LH release following i.c.v. injection of muscimol, but no change in those treated with estradiol. These data, together, suggest that the presence of estrogen changes the nature of action of GABA on the HPG system.

There is additional evidence that the presence of estrogen results in an increase in GABA levels in the POA-AH both *in vitro* (Herbison *et al.*, 1989) and *in vivo* (Demling *et al.*, 1985; Herbison *et al.*, 1990; Mansky *et al.*, 1982; Ondo *et al.*, 1982), suggesting that GABA may play a role in the negative feedback effects of estrogens on GnRH cells. A more recent study examining the diagonal band of Broca (DBB), VMN, medial septum, MPN and dorsomedial nucleus (all nuclei within the rostral hypothalamus), OVX, estradiol treated females showed an increase in GABA levels in the DBB and VMN as compared to OVX controls (Luine *et al.*, 1997). Furthermore examination of intact, female rats suggests a relationship between estrogen and GABA levels throughout the estrous cycle, with GABA levels being highest in the medial preoptic area during the morning of proestrus as compared to any other time in the cycle (Mitsushima *et al.*, 2002). These data were expanded to suggest such correlation is regionally specific, with GABA levels varying on proestrus in the ventro- and dorso- medial hypothalamus only, and not in the medial preoptic nucleus, diagonal band of Broca, lateral or medial septum, anterior hypothalamus (Frankfurt *et al.*, 1984).

Estrogen appears to selectively influence the enzymes involved in GABA synthesis, GAD₆₅ and GAD₆₇, and likely affects other regulatory proteins, such as GABA transporters (GAT; Table 3) responsible for removal of GABA from the synaptic cleft. Comparison of GAD₆₇ mRNA level in the POA throughout the estrous cycle in females sacrificed at noon suggest no change in expression as measured by *in situ* hybridization (Herbison *et al.*, 1992). However, as discussed previously, there are nucleus-specific changes in GAD₆₇ gene expression during the preovulatory or steroid induced LH surge (Table 4). Further, findings from Curran-Rauhut *et al.* (Curran-Rauhut and Petersen, 2002) suggest an estrogen-mediated action during this period, as *in situ* hybridization studies examining the rostral POA/OVLT showed an increase in GAD₆₇ levels on the morning of the estradiol-induced LH surge, and a decline in the afternoon, but no change in OVX females that were not treated with estradiol.

GABA transporters (GAT) provide another source of possible estrogen regulation. Of these, GAT-1, expressed on neurons) and GAT-3, localized to both neuron and glial cells, are most abundant in the POA, and 75% of GATs in this region are estimated to be GAT-1 (Herbison *et al.*, 1995). GAT-1 mRNA levels decline in a regionally-specific manner (specifically in the AVPV and medial preoptic nucleus (MPN)) and with ovariectomy, whereby long-term (7-day) estrogen replacement resulted in significantly higher expression (in the MPN only). Furthermore, GABA uptake was enhanced, an unexpected finding that suggests the presence of estrogen is increasing transporter levels, the net result of which would be a decrease in extracellular GABA levels.

In the caudal hypothalamus, Parducz *et al.* (Parducz *et al.*, 1993) reported that the presence of estrogen resulted in ultrastructural changes to the GABAergic neurons of the arcuate nucleus (ARC) in the OVX, female rat. Specifically GABA-positive synaptic contacts on the membrane and axosomatic synapses declined with estrogen treatment. A similar study using the intact female across the estrous cycle, found a decline in GABA positive axosomatic synapses, and an increase in non-GABA labeled dendritic spine synapses during the transition from the proestrous morning to the proestrous afternoon and estrus (Csakvari *et al.*, 2007). This suggests a diminution of GABAergic inhibition in this region and an increase in stimulatory synapses, thereby enabling female ovulation and reproductive behaviors during this time.

Together, these studies show multiple levels at which estrogens and GABA may interact to influence the GnRH system. Apparently, all regulatory levels of GABA synthesis, release, and uptake are affected by estrogen, the net outcome of which is an apparent decrease in GABAergic tone to the GnRH cells prior to the preovulatory or steroid-induced GnRH/LH surge. We believe that this attenuation of GABAergic input then enables stimulatory inputs to GnRH neurons to come into play to stimulate the surge; a potential candidate for this stimulation is glutamate, among other neurotransmitters. We will discuss some of the possible interactions among GABA, glutamate and GnRH in section VI.

3e. Electrophysiological studies: Electrophysiological studies focusing on the GABAergic drive on GnRH neurons have verified, without dispute, that GABA_ARs are located on GnRH neurons, that their activation results in functional physiological responses (Spergel *et al.*, 1999), and that they are active at the synaptic level (Moenter and DeFazio, 2005; Sim *et al.*, 2000; Sullivan and Moenter, 2003; Sullivan and Moenter, 2004; Sullivan and Moenter, 2005). As mentioned previously, in early development (E15-peripubertal), electrophysiological recordings of non-GnRH, hypothalamic neurons (E15) and GnRH neurons in response to GABA application suggest an excitatory rather than inhibitory neuronal response (Chen *et al.*, 1995; Chen *et al.*, 1996; Han *et al.*, 2002). However, in adult female mice, GABAergic activity was shown to be either inhibitory (Han *et al.*, 2002) or excitatory (DeFazio *et al.*, 2002; Moenter and DeFazio, 2005) on GnRH neurons, results that may differ due to several methodological differences: first, when considering the method of drug

application, Han *et al.* (Han *et al.*, 2002) applied bicuculline (GABA_AR antagonist) as a bath. DeFazio *et al.* (DeFazio *et al.*, 2002) used a rapid application of muscimol (GABA_AR agonist). The response of these cells to long-term versus short-term application of drug likely varies, as receptors become activated and possibly desensitized to long-term application. Second, the method of GnRH identification also differed. In the former study (Han *et al.*, 2002), GnRH neurons were identified using a GnRH- β -galactosidase transgenic whereas the latter study (DeFazio *et al.*, 2002) used GnRH-GFP labeled animals. It is possible that these targeting techniques focused on two different populations of GnRH neurons that differed in their expression of GABA receptors in some way. Notably, a recent study (Yin *et al.*, 2008) has also suggested excitatory effects of GABA on GnRH neurons of adult rats, consistent with work from Moenter's laboratory. Research into other CNS regions (including cortex, suprachiasmatic nucleus of the hypothalamus, and spinal neurons) support the idea that different populations of neurons may react to GABA differently, resulting in depolarization rather than the expected hyperpolarization (Gulledge, 2003; Marty and Llano 2005; Rohrbough and Spitzer, 1996; Wagner *et al.*, 1997). Third, the former study (Han *et al.*, 2002) did not block glutamatergic activity, whereas the latter study performed this blockade (DeFazio *et al.*, 2002). As this review hopes to show, glutamate and GABA are two very important neurotransmitters that act both directly and indirectly to regulate this system and its inputs, as well as to regulate one another. The absence of glutamatergic signaling in one study versus its presence in another likely affects the final activity of the GnRH neuron. In another study, Moenter and DeFazio (Moenter and DeFazio, 2005), using hypothalamic tissue slices from female mice, showed that bicuculline, when applied in the absence of glutamate inhibition, caused a marked increase in the neuronal firing rate, whereas the presence of such inhibition resulted in a decreased firing rate, rather than the expected increase attributed to relief of inhibition. This work shows the importance of examining both the individual contribution of each neurotransmitter as well as their interactions.

From these studies, it is important to recognize that there are likely different populations of GABAergic cells that respond differently to varying stimuli. The fact that GABA_ARs contain many combinations of subunits and these are differentially modulated by steroid hormones (Herbison *et al.*, 1995), that signals from other GnRH inputs (Roberts *et al.*, 2008) integrate with GABAergic signals to affect a response, and that the expression of chloride transporters on GnRH neurons varies (Leupen *et al.*, 2003, DeFazio, 2002 #321) all suggest that the expected GABAergic result may vary depending upon the target tissue and the neural environment.

4. Reproductive Senescence—As discussed in section IIIB, in many species including rats, aging is accompanied by a transition to a non-reproductive state. Part of this process involves changes to GnRH/LH pulsatile release, an attenuated and decreased GnRH/LH surge, and the change from a regular estrous cycle to a non-cycling state. Despite abundant evidence for critical roles of GABA in the regulation of these mechanisms in young adults (described in the previous sections), it is surprising how few studies have examined GABA during the age-related transition to acyclicity.

GABA turnover rates, using push-pull perfusion, were examined in intact young and aged female rats, and differences were observed in both the medial preoptic nucleus (MPN) and suprachiasmatic nucleus (SCN), regions important to the modulation and timing of the GnRH/LH surge, ovulation, and circadian rhythmicity (Harney *et al.*, 1996; Jarry *et al.*, 1995). Acyclic, aged animals showed a decline in GABA turnover, as compared to young, in the MPN, but an increase in the SCN (in persistent estrous females only). Thus changes to GABA turnover are likely affecting the regulatory mechanism of GnRH/LH release, such as pulsatile output, although further research is needed to resolve the mechanisms. Moreover, the net effect of decreased GABA turnover might be enhanced synaptic GABA. This possibility is consistent with a recent report by Neal-Perry *et al.* (2008) who showed by microdialysis that GABA

release in the POA was increased in middle-aged compared to young ovariectomized, hormone-primed rats.

In examining the GnRH/LH surge, Cashion *et al.* (Cashion *et al.*, 2004) found that middle-aged animals had overall lower expression levels of GAD₆₇ mRNA on the proestrus day, as compared to young rats, in the AVPV and organum vasculosum of the lamina terminalis (OVLT). Additionally, whereas in young animals GAD₆₇ expression in these nuclei increased in the early morning of proestrus, followed by a decline at later time points, no such fluctuations were observed in their older counterparts in the AVPV only. Thus, the biological rhythms of GABA synthesis appear to be attenuated with age. A study by Grove-Strawser *et al.* (Grove-Strawser *et al.*, 2007) support these findings, showing a decline in GAD₆₇ expression during the afternoon of proestrus in the OVLT/rostral hypothalamus of young female rats, but no such change in middle-aged females, and supporting a significantly higher expression level in middle-aged animals during the time period that LH is ascending. These data suggest a change in the synthesis of GABA such that the rhythmicity of its release, important to the timing of the pre-ovulatory surge, is interrupted, thus providing one explanation for the alteration of this mechanism during the aging transition, contributing to an attenuation of the GnRH/LH surge. Consistently, the important study of Neal-Perry *et al.* (2008) confirms that pharmacological blockade of GABA_ARs with bicuculline in middle-aged rats mitigates the attenuation of the steroid-induced LH surge.

These studies are just the beginning of numerous questions that researchers must examine to further understand the role of this inhibitory neurotransmitter in the reproductive senescent transition. Changes to GABA subunit populations, effects of agonists and antagonists, and their consequences on reproductive parameters (i.e. the preovulatory GnRH/LH surge, GnRH pulsatile release, estrogenic regulation, prolongation of estrous cycles with age) are all needed to help shed light on the GABAergic mechanisms during this important life period.

VI. Glutamate/GABA interactions

Thus far we have examined the individual contributions of GABA and glutamate to the regulation of the reproductive system throughout the life span of female rats. These data show a very complex interaction of both GABA and glutamate in the regulation of the HPG axis, and suggest that these amino acid neurotransmitters set the inhibitory and excitatory tone for neurotransmission in the hypothalamus-preoptic area (Figure 2). As such, it is important not only to examine how they act individually to regulate this system, but also how they interact with one another in this regard. Although beyond the scope of this article, we point out that this glutamate-GABA balance in the POA provides a backdrop in which other hypothalamic neurotransmitters, neuropeptides, neurotrophic factors, and other neural factors then must act to exert their respective influences upon the GnRH neural circuit. Here, we provide a brief overview of the glutamate-GABA interactions across the reproductive life cycle.

The glutamatergic and GABAergic systems are both in developmental transition during the early postnatal period (van den Pol *et al.*, 1995). Indeed, during early development, these systems probably regulate one another in the hypothalamus to ensure a proper level of excitatory/inhibitory activity (Gao *et al.*, 1998; van den Pol *et al.*, 1998). van den Pol *et al.* (van den Pol *et al.*, 1995) used patch clamp recordings to determine the response of E15 rat hypothalamic neurons to GABA and glutamate agonists. These preparations showed that, while all measured neurons responded to GABA, only a fraction of these responded to the glutamate agonists (with no response recorded for NMDA application until E17). Later studies from this laboratory support this data and further suggest that the glutamate system develops more slowly than the GABAergic system, and that GABA may actually facilitate glutamatergic depolarization during this period (E15–E18) (Gao *et al.*, 1998; van den Pol *et al.*, 1998). This

evidence suggests that during development, these systems have a very close interaction in which they regulate one another to insure a proper level of excitatory activity. The authors suggest that once established, the glutamatergic system becomes the main excitatory component and the GABAergic system switches to become the main inhibitory drive.

From P15–P30 there is a transition in hypothalamic GABAergic regulation, from a stimulatory to inhibitory action, and it appears likely that the GABA_AR is the primary target receptor for this action. Furthermore, during this time the reproductive axis is undergoing a very important change from quiescent to active, as the period from P25–P30 represents the onset of puberty. It has been suggested that the GABAergic system interacts with the glutamatergic system during this transition to regulate the activity of the reproductive axis and influence these hypothalamic hormonal changes.

Administration of AOAA, an inhibitor of GABA breakdown (Table 3) at P15 caused the expected increase in levels of GABA, and both glutamate levels *in vitro* and LH *in vivo*, followed, increasing as well. At P30, the AOAA-induced increase in GABA resulted in decline in glutamate and LH levels, presumably due to the increase in GABA levels (Carbone *et al.*, 2002). This suggests the possibility of GABAergic regulation of glutamate release, and a switch in that regulation from excitatory to inhibitory during the pubertal transition, causing a decrease in LH prepubertally, and an increase postpubertally. To further decipher the role of the GABA_AR and GABA_BR in this, muscimol or baclofen was applied to hypothalamic explants of 30-day old female rats. These treatments caused a decline in GnRH and glutamate release, suggesting that GABA, acting through both receptor types, may influence GnRH both independently as well as through an inhibition of excitatory glutamate release (Feleder *et al.*, 1999).

To determine the role of glutamate regulation of GABA activity during the prepubertal period, Scacchi *et al.* (Scacchi *et al.*, 1998) demonstrated that, while administration of the NMDAR antagonist MK-801 to prepubertal females did not significantly alter LH levels, treatment with the GABAergic agonists muscimol or baclofen increased or decreased LH levels, respectively, suggesting differential roles for the GABA_AR (excitatory) and GABA_BR (inhibitory) during this time period. The addition of MK-801 to these GABAergic treatments resulted in an inhibition of the muscimol-induced LH increase, suggesting a robust regulation of the GABA_ARs by the NMDA system. Further, treatment with NMDA caused an increase in LH, which was inhibited by MK-801, but not bicuculline, implying that at this time period, the activity of the GABA_ARs is not regulating the NMDA system (GABA_BR activity was not measured). However, a study by Pinilla *et al.* (2002) found the opposite. Administration of either bicuculline or phaclofen (a GABA_BR antagonist) in P23 females inhibited NMDA-stimulated LH levels. Although some of these results are not in agreement, possibly due to differences in methodology including different ages of experimental animals, mode of administration, specific pharmacologic agent, the bottom line of these studies is that during pubertal development, there appears to be reciprocal interactions of glutamate and GABA, with downstream effects on the GnRH neurosecretory system.

In the adult model, few studies have been undertaken to examine this interplay. Glutamate-stimulated release of GABA in various regions of the brain is well documented (de Mello *et al.*, 1988; do Nascimento and de Mello, 1985; Harris and Miller, 1989; Moran *et al.*, 1986; Pin and Bockaert, 1989). Examination of the POA specifically supported the presence of this mechanism, and further suggested that steroid hormone treatment, specifically estradiol + progesterone, results in increased glutamate-induced release of GABA (Fleischmann *et al.*, 1992) as measured from synaptosomes in culture. Using intact males, this same lab demonstrated that activation of GABA_ARs via muscimol (GABA_A agonist) application resulted in both increased GABA and glutamate release from this same *in vitro* preparation

(Fleischmann et al., 1995). An examination of a functional endpoint, such as the GnRH/LH surge, adds further insight to this observation. In OVX, estradiol treated rats given NMDA to facilitate the GnRH/LH surge, intracerebroventricular administration of muscimol (GABA_A agonist), but not baclofen (GABA_B agonist), resulted in an inhibition of the surge (Akema and Kimura, 1993). These results further suggest some interaction GABA_ARs and NMDARs, but the nuances of this interaction are not known.

Perhaps the most intriguing finding in this area is evidence for a dual-phenotype GABA/glutamate neuronal population in the AVPV of the rat (Ottem *et al.*, 2004). The AVPV is a nucleus of great importance to the execution of the GnRH/LH surge and ovulation. Ottem *et al.* (Ottem *et al.*, 2004) used both *in situ* hybridization histochemistry and immunohistochemistry to show that the majority of neurons in the AVPV that express vesicular glutamate transporter (VGlut; Table 1), a marker of glutamate, also express vesicular GABA transporter (VGAT; Table 3), a marker of GABA, and vice-versa. Additionally, these dual-phenotype neurons strongly express ER α , they have contacts on GnRH neurons, and their glutamatergic-GABAergic phenotype is estradiol-sensitive. Specifically, during the estradiol-induced LH surge, the VGAT-containing vesicles decreased while VGlut-containing vesicles increased during the afternoon transition to the surge, suggesting an increase in glutamatergic, and decrease in GABAergic, activity during this event (Ottem *et al.*, 2004). To our knowledge, this is the first model of its kind, and suggests an even more complex architecture for the regulation of the HPG reproductive axis by GABA, glutamate, and estradiol.

Finally, to our knowledge, there is a single study that has interrogated the interplay between glutamate, GABA, and GnRH neurons during reproductive aging. Using ovariectomized young and middle-aged female rats, Neal-Perry *et al.* (2008) showed: 1) extracellular glutamate is decreased, and GABA is increased, in middle-aged rats with attenuated LH surges (measured by microdialysis); 2) pharmacological blockade of the GABA_AR enhances both glutamate release and the LH surge; and 3) such treatment in concert with an inhibitor of synaptic glutamate reuptake further enhances glutamate levels and restores the LH surge. These results show a tight interaction of glutamate and GABA, along with GABAergic regulation of glutamate levels, and subsequent effects on the GnRH/LH surge (Figure 2).

VII. Conclusion

Understanding the reproductive axis, its development, maintenance, and senescence, has proven to be an important but difficult task due to the many regulatory components of this system. The complex interactions among the three levels of the hypothalamic-pituitary-gonadal axis, as well as the neuromodulatory mechanisms upstream of this axis, creates an endocrine, neural, and neuroendocrine network that is only beginning to be fully appreciated. In considering the actions of each component part of this network, it is important to consider regional specificity of action, reproductive life cycle stage, and the steroid hormone environment, as many studies demonstrate differential mechanisms of modulation due to these variables.

This review suggests that GABA, the main inhibitory neurotransmitter, and glutamate, the main stimulatory neurotransmitter, set a level of excitability in the hypothalamus that decreases or increases the likelihood that GnRH will be synthesized or released. On top of this tone, other neuropeptides and regulatory factors may modulate GnRH release in a manner appropriate to the animal's reproductive status. For example, monoaminergic neurotransmitters such as serotonin, dopamine, epinephrine, norepinephrine; neuropeptides such as kisspeptin, neuropeptide Y, vasoactive intestinal protein, and many others; neurotrophic factors in the transforming-growth factor and insulin-like growth factor families, among others; and neuroactive gases such as nitric oxide; feed into this neurocircuitry. By further understanding

the neural regulation of hypothalamic GnRH neurons, we will gain insights not only into normal reproductive functions, but better understand dysfunctions of puberty, infertility, and aging.

Literature Cited

1. Abbud R, Smith MS. Differences in the luteinizing hormone and prolactin responses to multiple injections of kainate, as compared to N-methyl-D,L-aspartate, in cycling rats. *Endocrinology* 1991;129:3254–3258. [PubMed: 1659525]
2. Abe H, Terasawa E. Firing pattern and rapid modulation of activity by estrogen in primate luteinizing hormone releasing hormone-1 neurons. *Endocrinology* 2005;146:4312–4320. [PubMed: 15976055]
3. Adams MM, Flagg RA, Gore AC. Perinatal changes in hypothalamic N-methyl-D-aspartate receptors and their relationship to gonadotropin-releasing hormone neurons. *Endocrinology* 1999;140:2288–2296. [PubMed: 10218982]
4. Adler BA, Crowley WR. Evidence for gamma-aminobutyric acid modulation of ovarian hormonal effects on luteinizing hormone secretion and hypothalamic catecholamine activity in the female rat. *Endocrinology* 1986;118:91–97. [PubMed: 3510124]
5. Akema T, Chiba A, Kimura F. On the relationship between noradrenergic stimulatory and GABAergic inhibitory systems in the control of luteinizing hormone secretion in female rats. *Neuroendocrinology* 1990;52:566–572. [PubMed: 2178232]
6. Akema T, Kimura F. The mode of GABAB receptor-mediated inhibition of the preovulatory luteinizing hormone surge in female rats. *Brain Res* 1991;562:169–172. [PubMed: 1666015]
7. Akema T, Kimura F. Modulation of pulsatile LH secretion by baclofen, a selective GABAB receptor agonist, in ovariectomized rats. *Neuroendocrinology* 1992;56:141–147. [PubMed: 1328914]
8. Akema T, Kimura F. Differential effects of GABAA and GABAB receptor agonists on NMDA-induced and noradrenaline-induced luteinizing-hormone release in the ovariectomized estrogen-primed rat. *Neuroendocrinology* 1993;57:28–33. [PubMed: 8097576]
9. Arias P, Carbone S, Szwarcfarb B, Feleder C, Rodriguez M, Scacchi P, Moguilevsky JA. Effects of aging on N-methyl-D-aspartate (NMDA)-induced GnRH and LH release in female rats. *Brain Res* 1996;740:234–238. [PubMed: 8973819]
10. Arias P, Jarry H, Leonhardt S, Moguilevsky JA, Wuttke W. Estradiol modulates the LH release response to N-methyl-D-aspartate in adult female rats: studies on hypothalamic luteinizing hormone-releasing hormone and neurotransmitter release. *Neuroendocrinology* 1993;57:710–715. [PubMed: 7690119]
11. Arslan M, Pohl CR, Plant TM. DL-2-amino-5-phosphonopentanoic acid, a specific N-methyl-D-aspartic acid receptor antagonist, suppresses pulsatile LH release in the rat. *Neuroendocrinology* 1988;47:465–468. [PubMed: 2840595]
12. Asada H, Kawamura Y, Maruyama K, Kume H, Ding RG, Kanbara N, Kuzume H, Sanbo M, Yagi T, Obata K. Cleft palate and decreased brain gamma-aminobutyric acid in mice lacking the 67-kDa isoform of glutamic acid decarboxylase. *Proc Natl Acad Sci U S A* 1997;94:6496–6499. [PubMed: 9177246]
13. Bailey JD, Centers A, Jennes L. Expression of AMPA receptor subunits (GluR1-GluR4) in gonadotrophin-releasing hormone neurones of young and middle-aged persistently oestrous rats during the steroid-induced luteinising hormone surge. *J Neuroendocrinol* 2006;18:1–12. [PubMed: 16451215]
14. Belchetz PE, Plant TM, Nakai Y, Keogh EJ, Knobil E. Hypophysial responses to continuous and intermittent delivery of hypothalamic gonadotropin-releasing hormone. *Science* 1978;202:631–633. [PubMed: 100883]
15. Bergen HT, Hejtmanick JF, Pfaff DW. Effects of gamma-aminobutyric acid receptor agonists and antagonist on LHRH-synthesizing neurons as detected by immunocytochemistry and in situ hybridization. *Exp Brain Res* 1991;87:46–56. [PubMed: 1661682]
16. Bianchi MS, Lux-Lantos VA, Bettler B, Libertun C. Expression of gamma-aminobutyric acid B receptor subunits in hypothalamus of male and female developing rats. *Brain Res Dev Brain Res* 2005;160:124–129.

17. Bless EP, Walker HJ, Yu KW, Knoll JG, Moenter SM, Schwarting GA, Tobet SA. Live view of gonadotropin-releasing hormone containing neuron migration. *Endocrinology* 2005;146:463–468. [PubMed: 15486219]
18. Bonavera JJ, Swerdloff RS, Sinha Hakim AP, Lue YH, Wang C. Aging results in attenuated gonadotropin releasing hormone-luteinizing hormone axis responsiveness to glutamate receptor agonist N-methyl-D-aspartate. *J Neuroendocrinol* 1998;10:93–99. [PubMed: 9535055]
19. Bottner M, Leonhardt S, Wuttke W, Jarry H. Changes of expression of genes related to the activity of the gonadotrophin-releasing hormone pulse generator in young versus middle-aged male rats. *J Neuroendocrinol* 2007;19:779–787. [PubMed: 17850460]
20. Boue-Grabot E, Dufy B, Garret M. Molecular diversity of GABA-gated chloride channels in the rat anterior pituitary. *Brain Res* 1995;704:125–129. [PubMed: 8750973]
21. Boue-Grabot E, Roudbaraki M, Bascles L, Tramu G, Bloch B, Garret M. Expression of GABA receptor rho subunits in rat brain. *J Neurochem* 1998;70:899–907. [PubMed: 9489708]
22. Boue-Grabot E, Taupignon A, Tramu G, Garret M. Molecular and electrophysiological evidence for a GABA_A receptor in thyrotropin-secreting cells. *Endocrinology* 2000;141:1627–1632. [PubMed: 10803570]
23. Bourguignon JP, Gerard A, Alvarez Gonzalez ML, Purnelle G, Franchimont P. Endogenous glutamate involvement in pulsatile secretion of gonadotropin-releasing hormone: evidence from effect of glutamine and developmental changes. *Endocrinology* 1995;136:911–916. [PubMed: 7867599]
24. Bourguignon JP, Gerard A, Franchimont P. Direct activation of gonadotropin-releasing hormone secretion through different receptors to neuroexcitatory amino acids. *Neuroendocrinology* 1989a;49:402–408. [PubMed: 2541362]
25. Bourguignon JP, Gerard A, Mathieu J, Simons J, Franchimont P. Pulsatile release of gonadotropin-releasing hormone from hypothalamic explants is restrained by blockade of N-methyl-D,L-aspartate receptors. *Endocrinology* 1989b;125:1090–1096. [PubMed: 2546737]
26. Brann DW. Glutamate: a major excitatory transmitter in neuroendocrine regulation. *Neuroendocrinology* 1995;61:213–225. [PubMed: 7898626]
27. Brann DW, Mahesh VB. Endogenous excitatory amino acid involvement in the preovulatory and steroid-induced surge of gonadotropins in the female rat. *Endocrinology* 1991;128:1541–1547. [PubMed: 1900231]
28. Brann DW, Mahesh VB. Excitatory amino acid regulation of gonadotropin secretion: modulation by steroid hormones. *J Steroid Biochem Mol Biol* 1992;41:847–850. [PubMed: 1348628]
29. Brann DW, Mahesh VB. Excitatory amino acids: function and significance in reproduction and neuroendocrine regulation. *Front Neuroendocrinol* 1994;15:3–49. [PubMed: 7958168]
30. Brann DW, Mahesh VB. The aging reproductive neuroendocrine axis. *Steroids* 2005;70:273–283. [PubMed: 16080236]
31. Brann DW, Zamorano P, Ping L, Mahesh V. Role of excitatory amino acid neurotransmission during puberty in the female rat. *Molecular and cellular neurosciences* 1993a;4:107–112. [PubMed: 19912913]
32. Brann DW, Zamorano PL, Chorch LP, Mahesh VB. Steroid hormone effects on NMDA receptor binding and NMDA receptor mRNA levels in the hypothalamus and cerebral cortex of the adult rat. *Neuroendocrinology* 1993b;58:666–672. [PubMed: 7907400]
33. Brann DW, Zamorano PL, Putnam-Roberts CD, Mahesh VB. Gamma-aminobutyric acid-opioid interactions in the regulation of gonadotropin secretion in the immature female rat. *Neuroendocrinology* 1992;56:445–452. [PubMed: 1335551]
34. Carbone S, Ponzo O, Szwarcfarb B, Rondina D, Reynoso R, Scacchi P, Alberto Moguilevsky J. Ontogenic modifications in the effect of the GABAergic system on the hypothalamic excitatory amino acids: its relationship with GABAergic control of gonadotrophin secretion during sexual maturation in female rats. *Brain Res Dev Brain Res* 2002;133:13–18.
35. Carbone S, Szwarcfarb B, Losada M, Moguilevsky JA. Effect of ovarian hormones on the hypothalamic excitatory amino acids system during sexual maturation in female rats. *Neuroendocrinology* 1995;61:235–242. [PubMed: 7898628]

36. Carbone S, Szwarcfarb B, Otero Losada ME, Moguilevsky JA. Effects of ovarian steroids on the gonadotropin response to N-methyl-D-aspartate and on hypothalamic excitatory amino acid levels during sexual maturation in female rats. *Endocrinology* 1992;130:1365–1370. [PubMed: 1347007]
37. Carbone S, Szwarcfarb B, Rondina D, Feleder C, Moguilevsky JA. Differential effects of the N-methyl-D-aspartate and non-N-methyl-D-aspartate receptors of the excitatory amino acids system on LH and FSH secretion. Its effects on the hypothalamic luteinizing hormone releasing hormone during maturation in male rats. *Brain Res* 1996;707:139–145. [PubMed: 8919290]
38. Cashion AB, Smith MJ, Wise PM. The morphometry of astrocytes in the rostral preoptic area exhibits a diurnal rhythm on proestrus: relationship to the luteinizing hormone surge and effects of age. *Endocrinology* 2003;144:274–280. [PubMed: 12488355]
39. Cashion AB, Smith MJ, Wise PM. Glutamic acid decarboxylase 67 (GAD67) gene expression in discrete regions of the rostral preoptic area change during the oestrous cycle and with age. *J Neuroendocrinol* 2004;16:711–716. [PubMed: 15271064]
40. Chakraborty TR, Ng L, Gore AC. Age-related changes in estrogen receptor beta in rat hypothalamus: a quantitative analysis. *Endocrinology* 2003a;144:4164–4171. [PubMed: 12933691]
41. Chakraborty TR, Ng L, Gore AC. Colocalization and hormone regulation of estrogen receptor alpha and N-methyl-D-aspartate receptor in the hypothalamus of female rats. *Endocrinology* 2003b;144:299–305. [PubMed: 12488358]
42. Chen CL, Trombley PQ, van den Pol AN. GABA receptors precede glutamate receptors in hypothalamic development; Differential regulation by astrocytes. *Journal of Neurophysiology* 1995;74:1473–1484. [PubMed: 8989386]
43. Chen G, Trombley PQ, van den Pol AN. Excitatory actions of GABA in developing rat hypothalamic neurones. *J Physiol* 1996;494(Pt 2):451–464. [PubMed: 8842004]
44. Chen N, Luo T, Raymond LA. Subtype-dependence of NMDA receptor channel open probability. *J Neurosci* 1999;19:6844–6854. [PubMed: 10436042]
45. Cho BN, Kim K. Differential effect of baclofen on hypothalamic GnRH and pituitary LH beta gene expression in steroid-treated rats. *Mol Cells* 1997;7:605–609. [PubMed: 9387146]
46. Clark AS, Robinson S, Henderson LP. Dynamics of GABA(A) receptor binding in the ventromedial hypothalamus during postnatal development in the rat. *Brain Res Dev Brain Res* 1997;103:195–198.
47. Clarkson J, Herbison AE. Development of GABA and glutamate signaling at the GnRH neuron in relation to puberty. *Mol Cell Endocrinol* 2006;254–255. 32–38.
48. Csakvari E, Hoyk Z, Gyenes A, Garcia-Ovejero D, Garcia-Segura LM, Parducz A. Fluctuation of synapse density in the arcuate nucleus during the estrous cycle. *Neuroscience* 2007;144:1288–1292. [PubMed: 17161546]
49. Curran-Rauhut MA, Petersen SL. Regulation of glutamic acid decarboxylase 65 and 67 gene expression by ovarian steroids: identification of two functionally distinct populations of GABA neurones in the preoptic area. *J Neuroendocrinol* 2002;14:310–317. [PubMed: 11963828]
50. Cutler RW, Dudzinski DS. Regional changes in amino acid content in developing rat brain. *J Neurochem* 1974;23:1005–1009. [PubMed: 4154971]
51. Davis AM, McCarthy MM. Developmental increase in (3)H-muscimol binding to the gamma-aminobutyric acid(A) receptor in hypothalamic and limbic areas of the rat: why is the ventromedial nucleus of the hypothalamus an exception? *Neurosci Lett* 2000;288:223–227. [PubMed: 10889348]
52. Davis AM, Penschuck S, Fritschy JM, McCarthy MM. Developmental switch in the expression of GABA(A) receptor subunits alpha(1) and alpha(2) in the hypothalamus and limbic system of the rat. *Brain Res Dev Brain Res* 2000;119:127–138.
53. Davis AM, Ward SC, Selmanoff M, Herbison AE, McCarthy MM. Developmental sex differences in amino acid neurotransmitter levels in hypothalamic and limbic areas of rat brain. *Neuroscience* 1999;90:1471–1482. [PubMed: 10338313]
54. de la Iglesia HO, Schwartz WJ. Minireview: timely ovulation: circadian regulation of the female hypothalamo-pituitary-gonadal axis. *Endocrinology* 2006;147:1148–1153. [PubMed: 16373412]
55. de Mello MC, Klein WL, de Mello FG. L-glutamate evoked release of GABA from cultured avian retina cells does not require glutamate receptor activation. *Brain Res* 1988;443:166–172. [PubMed: 2896053]

56. Decavel C, Van den Pol AN. GABA: a dominant neurotransmitter in the hypothalamus. *J Comp Neurol* 1990;302:1019–1037. [PubMed: 2081813]
57. DeFazio RA, Heger S, Ojeda SR, Moenter SM. Activation of A-type gamma-aminobutyric acid receptors excites gonadotropin-releasing hormone neurons. *Mol Endocrinol* 2002;16:2872–2891. [PubMed: 12456806]
58. Demling J, Fuchs E, Baumert M, Wuttke W. Preoptic catecholamine, GABA, and glutamate release in ovariectomized and ovariectomized estrogen-primed rats utilizing a push-pull cannula technique. *Neuroendocrinology* 1985;41:212–218. [PubMed: 2864651]
59. Diano S, Naftolin F, Horvath TL. Gonadal steroids target AMPA glutamate receptor-containing neurons in the rat hypothalamus, septum and amygdala: a morphological and biochemical study. *Endocrinology* 1997;138:778–789. [PubMed: 9003015]
60. do Nascimento JL, de Mello FG. Induced release of gamma-aminobutyric acid by a carrier-mediated, high-affinity uptake of L-glutamate in cultured chick retina cells. *J Neurochem* 1985;45:1820–1827. [PubMed: 2865335]
61. Dohler KD, Wuttke W. Changes with age in levels of serum gonadotropins, prolactin and gonadal steroids in prepubertal male and female rats. *Endocrinology* 1975;97:898–907. [PubMed: 1193012]
62. Donoso AO, Banzan AM. Effects of increase of brain GABA levels on the hypothalamic-pituitary-luteinizing hormone axis in rats. *Acta Endocrinol (Copenh)* 1984;106:298–304. [PubMed: 6377787]
63. Donoso AO, Banzan AM. Blockade of the LH surge and ovulation by GABA-T inhibitory drugs that increase brain GABA levels in rats. *Psychoneuroendocrinology* 1986;11:429–435. [PubMed: 3562742]
64. Donoso AO, Lopez FJ, Negro-Vilar A. Glutamate receptors of the non-N-methyl-D-aspartic acid type mediate the increase in luteinizing hormone-releasing hormone release by excitatory amino acids in vitro. *Endocrinology* 1990;126:414–420. [PubMed: 2152870]
65. Downs JL, Urbanski HF. Neuroendocrine changes in the aging reproductive axis of female rhesus macaques (*Macaca mulatta*). *Biol Reprod* 2006;75:539–546. [PubMed: 16837643]
66. Ebling FJ, Cronin AS. The neurobiology of reproductive development. *Neuroreport* 2000;11:R23–R33. [PubMed: 11095489]
67. El-Etr M, Akwa Y, Baulieu EE, Schumacher M. The neuroactive steroid pregnenolone sulfate stimulates the release of gonadotropin-releasing hormone from GT1-7 hypothalamic neurons, through N-methyl-D-aspartate receptors. *Endocrinology* 2006;147:2737–2743. [PubMed: 16513833]
68. El-Etr M, Akwa Y, Fiddes RJ, Robel P, Baulieu EE. A progesterone metabolite stimulates the release of gonadotropin-releasing hormone from GT1-1 hypothalamic neurons via the gamma-aminobutyric acid type A receptor. *Proc Natl Acad Sci U S A* 1995;92:3769–3773. [PubMed: 7731981]
69. Erreger K, Dravid SM, Banke TG, Wyllie DJ, Traynelis SF. Subunit-specific gating controls rat NR1/NR2A and NR1/NR2B NMDA channel kinetics and synaptic signalling profiles. *J Physiol* 2005;563:345–358. [PubMed: 15649985]
70. Eyigor O, Centers A, Jennes L. Distribution of ionotropic glutamate receptor subunit mRNAs in the rat hypothalamus. *J Comp Neurol* 2001;434:101–124. [PubMed: 11329132]
71. Eyigor O, Jennes L. Identification of glutamate receptor subtype mRNAs in gonadotropin-releasing hormone neurons in the rat brain. *Endocrine* 1996;4:133–139.
72. Eyigor O, Jennes L. Expression of glutamate receptor subunit mRNAs in gonadotropin-releasing hormone neurons during the sexual maturation of the female rat. *Neuroendocrinology* 1997;66:122–129. [PubMed: 9263209]
73. Eyigor O, Jennes L. Kainate receptor subunit-positive gonadotropin-releasing hormone neurons express c-Fos during the steroid-induced luteinizing hormone surge in the female rat. *Endocrinology* 2000;141:779–786. [PubMed: 10650960]
74. Favit A, Wetsel WC, Negro-Vilar A. Differential expression of gamma-aminobutyric acid receptors in immortalized luteinizing hormone-releasing hormone neurons. *Endocrinology* 1993;133:1983–1989. [PubMed: 8404645]
75. Feleder C, Ginzburg M, Wuttke W, Moguilevsky JA, Arias P. GABAergic activation inhibits the hypothalamic-pituitary-ovarian axis and sexual development in the immature female rat. Associated

- changes in hypothalamic glutamatergic and taurinerger systems. *Brain Res Dev Brain Res* 1999;116:151–157.
76. Fenelon VS, Herbison AE. Characterisation of GABAA receptor gamma subunit expression by magnocellular neurones in rat hypothalamus. *Brain Res Mol Brain Res* 1995;34:45–56. [PubMed: 8750860]
 77. Fleischmann A, Etgen AM, Makman MH. Estradiol plus progesterone promote glutamate-induced release of gamma-aminobutyric acid from preoptic area synaptosomes. *Neuropharmacology* 1992;31:799–807. [PubMed: 1356255]
 78. Fleischmann A, Makman MH, Etgen AM. GABAA receptor activation induces GABA and glutamate release from preoptic area. *Life Sci* 1995;56:1665–1678. [PubMed: 7723596]
 79. Flugge G, Oertel WH, Wuttke W. Evidence for estrogen-receptive GABAergic neurons in the preoptic/anterior hypothalamic area of the rat brain. *Neuroendocrinology* 1986;43:1–5. [PubMed: 3520371]
 80. Frankfurt M, Fuchs E, Wuttke W. Sex differences in gamma-aminobutyric acid and glutamate concentrations in discrete rat brain nuclei. *Neurosci Lett* 1984;50:245–250. [PubMed: 6149503]
 81. Fritschy JM, Mohler H. GABAA-receptor heterogeneity in the adult rat brain: differential regional and cellular distribution of seven major subunits. *J Comp Neurol* 1995;359:154–194. [PubMed: 8557845]
 82. Fueshko SM, Key S, Wray S. GABA inhibits migration of luteinizing hormone-releasing hormone neurons in embryonic olfactory explants. *J Neurosci* 1998;18:2560–2569. [PubMed: 9502815]
 83. Funabashi T, Jinnai K, Kimura F. Bicuculline infusion advances the timing of Fos expression in LHRH neurons in the preoptic area of proestrous rats. *Neuroreport* 1997;8:771–774. [PubMed: 9106764]
 84. Funabashi T, Kimura F. The number of luteinizing hormone-releasing hormone immunoreactive neurons is significantly decreased in the forebrain of old-aged female rats. *Neurosci Lett* 1995;189:85–88. [PubMed: 7609925]
 85. Gao XB, Chen G, van den Pol AN. GABA-dependent firing of glutamate-evoked action potentials at AMPA/kainate receptors in developing hypothalamic neurons. *J Neurophysiol* 1998;79:716–726. [PubMed: 9463435]
 86. Garyfallou VT, Lemos D, Urbanski HF. Expression profiling of genes encoding glutamate and GABA receptor subunits in three immortalized GnRH cell lines. *Brain Res* 2006;1086:50–54. [PubMed: 16626661]
 87. Gay VL, Plant TM. N-methyl-D,L-aspartate elicits hypothalamic gonadotropin-releasing hormone release in prepubertal male rhesus monkeys (*Macaca mulatta*). *Endocrinology* 1987;120:2289–2296. [PubMed: 3106017]
 88. Gay VL, Plant TM. Sustained intermittent release of gonadotropin-releasing hormone in the prepubertal male rhesus monkey induced by N-methyl-DL-aspartic acid. *Neuroendocrinology* 1988;48:147–152. [PubMed: 3065674]
 89. Genazzani AD, Petraglia F, Sgarbi L, Montanini V, Hartmann B, Surico N, Biolcati A, Volpe A, Genazzani AR. Difference of LH and FSH secretory characteristics and degree of concordance between postmenopausal and aging women. *Maturitas* 1997;26:133–138. [PubMed: 9089563]
 90. Gerendai I, Wenger T, Scapagnini U. Increase in the LHRH content of the organum vasculosum of the lamina terminalis and the medio-basal hypothalamus six hours after injection of kainic acid into the rat medial preoptic area. *Neurosci Lett* 1980;17:249–252. [PubMed: 6763163]
 91. Gill S, Sharpless JL, Rado K, Hall JE. Evidence that GnRH decreases with gonadal steroid feedback but increases with age in postmenopausal women. *J Clin Endocrinol Metab* 2002;87:2290–2296. [PubMed: 11994378]
 92. Gore AC. Gonadotropin-releasing hormone neurons, NMDA receptors, and their regulation by steroid hormones across the reproductive life cycle. *Brain Res Brain Res Rev* 2001;37:235–248. [PubMed: 11744089]
 93. Gore AC. Gonadotropin-releasing hormone (GnRH) neurons: gene expression and neuroanatomical studies. *Prog Brain Res* 2002;141:193–208. [PubMed: 12508571]
 94. Gore AC. Developmental programming and endocrine disruptor effects on reproductive neuroendocrine systems. *Front Neuroendocrinol* 2008;29:358–374. [PubMed: 18394690]

95. Gore AC, Oung T, Woller MJ. Age-related changes in hypothalamic gonadotropin-releasing hormone and N-methyl-D-aspartate receptor gene expression, and their regulation by oestrogen, in the female rat. *J Neuroendocrinol* 2002;14:300–309. [PubMed: 11963827]
96. Gore AC, Oung T, Yung S, Flagg RA, Woller MJ. Neuroendocrine mechanisms for reproductive senescence in the female rat: gonadotropin-releasing hormone neurons. *Endocrine* 2000a;13:315–323. [PubMed: 11216643]
97. Gore AC, Windsor-Engnell BM, Terasawa E. Menopausal increases in pulsatile gonadotropin-releasing hormone release in a nonhuman primate (*Macaca mulatta*). *Endocrinology* 2004;145:4653–4659. [PubMed: 15231708]
98. Gore AC, Wu TJ, Rosenberg JJ, Roberts JL. Gonadotropin-releasing hormone and NMDA receptor gene expression and colocalization change during puberty in female rats. *J Neurosci* 1996;16:5281–5289. [PubMed: 8757241]
99. Gore AC, Yeung G, Morrison JH, Oung T. Neuroendocrine aging in the female rat: the changing relationship of hypothalamic gonadotropin-releasing hormone neurons and N-methyl-D-aspartate receptors. *Endocrinology* 2000b;141:4757–4767. [PubMed: 11108291]
100. Goroll D, Arias P, Wuttke W. Preoptic release of amino acid neurotransmitters evaluated in peripubertal and young adult female rats by push-pull perfusion. *Neuroendocrinology* 1993;58:11–15. [PubMed: 7903429]
101. Grattan DR, Rocca MS, Strauss KI, Sagrillo CA, Selmanoff M, McCarthy MM. GABAergic neuronal activity and mRNA levels for both forms of glutamic acid decarboxylase (GAD65 and GAD67) are reduced in the diagonal band of Broca during the afternoon of proestrus. *Brain Res* 1996;733:46–55. [PubMed: 8891247]
102. Grove-Strawser D, Jimenez-Linan M, Rubin BS. Middle-aged female rats lack the dynamic changes in GAD(67) mRNA levels observed in young females on the day of a luteinising hormone surge. *J Neuroendocrinol* 2007;19:708–716. [PubMed: 17680886]
103. Gu G, Varoqueaux F, Simerly RB. Hormonal regulation of glutamate receptor gene expression in the anteroventral periventricular nucleus of the hypothalamus. *J Neurosci* 1999;19:3213–3222. [PubMed: 10191334]
104. Gullledge AT, Stuart GJ. Excitatory actions of GABA in the cortex. *Neuron* 2003;37:299–307. [PubMed: 12546824]
105. Hales TG, Kim H, Longoni B, Olsen RW, Tobin AJ. Immortalized hypothalamic GT1–7 neurons express functional gamma-aminobutyric acid type A receptors. *Mol Pharmacol* 1992;42:197–202. [PubMed: 1325030]
106. Hales TG, Sanderson MJ, Charles AC. GABA has excitatory actions on GnRH-secreting immortalized hypothalamic (GT1–7) neurons. *Neuroendocrinology* 1994;59:297–308. [PubMed: 8159279]
107. Hall JE, Lavoie HB, Marsh EE, Martin KA. Decrease in gonadotropin-releasing hormone (GnRH) pulse frequency with aging in postmenopausal women. *J Clin Endocrinol Metab* 2000;85:1794–1800. [PubMed: 10843154]
108. Han SK, Abraham IM, Herbison AE. Effect of GABA on GnRH neurons switches from depolarization to hyperpolarization at puberty in the female mouse. *Endocrinology* 2002;143:1459–1466. [PubMed: 11897704]
109. Harney JP, Scarbrough K, Rosewell KL, Wise PM. In vivo antisense antagonism of vasoactive intestinal peptide in the suprachiasmatic nuclei causes aging-like changes in the estradiol-induced luteinizing hormone and prolactin surges. *Endocrinology* 1996;137:3696–3701. [PubMed: 8756535]
110. Harris KM, Miller RJ. Excitatory amino acid-evoked release of [3H]GABA from hippocampal neurons in primary culture. *Brain Res* 1989;482:23–33. [PubMed: 2565138]
111. Henderson LP. Steroid modulation of GABAA receptor-mediated transmission in the hypothalamus: effects on reproductive function. *Neuropharmacology* 2007;52:1439–1453. [PubMed: 17433821]
112. Herbison AE, Augood SJ, McGowan EM. Expression of glutamic acid decarboxylase messenger RNA in rat medial preoptic area neurones during the oestrous cycle and after ovariectomy. *Brain Res Mol Brain Res* 1992;14:310–316. [PubMed: 1326694]

113. Herbison AE, Augood SJ, Simonian SX, Chapman C. Regulation of GABA transporter activity and mRNA expression by estrogen in rat preoptic area. *J Neurosci* 1995;15:8302–8309. [PubMed: 8613763]
114. Herbison AE, Chapman C, Dyer RG. Role of medial preoptic GABA neurones in regulating luteinising hormone secretion in the ovariectomised rat. *Exp Brain Res* 1991;87:345–352. [PubMed: 1769388]
115. Herbison AE, Dyer RG. Effect on luteinizing hormone secretion of GABA receptor modulation in the medial preoptic area at the time of proestrous luteinizing hormone surge. *Neuroendocrinology* 1991;53:317–320. [PubMed: 1645854]
116. Herbison AE, Fenelon VS. Estrogen regulation of GABAA receptor subunit mRNA expression in preoptic area and bed nucleus of the stria terminalis of female rat brain. *J Neurosci* 1995;15:2328–2337. [PubMed: 7891170]
117. Herbison AE, Heavens RP, Dyer RG. Oestrogen and noradrenaline modulate endogenous GABA release from slices of the rat medial preoptic area. *Brain Res* 1989;486:195–200. [PubMed: 2720430]
118. Herbison AE, Heavens RP, Dyer RG. Oestrogen modulation of excitatory A1 noradrenergic input to rat medial preoptic gamma aminobutyric acid neurones demonstrated by microdialysis. *Neuroendocrinology* 1990;52:161–168. [PubMed: 2125704]
119. Herman JP, Eyigor O, Ziegler DR, Jennes L. Expression of ionotropic glutamate receptor subunit mRNAs in the hypothalamic paraventricular nucleus of the rat. *J Comp Neurol* 2000;422:352–362. [PubMed: 10861512]
120. Hoffman GE, Finch CE. LHRH neurons in the female C57BL/6J mouse brain during reproductive aging: no loss up to middle age. *Neurobiol Aging* 1986;7:45–48. [PubMed: 3513038]
121. Hompes PG, Vermes I, Tilders FJ, Schoemaker J. In vitro release of LHRH from the hypothalamus of female rats during prepubertal development. *Neuroendocrinology* 1982;35:8–12. [PubMed: 6810194]
122. Jarry H, Hirsch B, Leonhardt S, Wuttke W. Amino acid neurotransmitter release in the preoptic area of rats during the positive feedback actions of estradiol on LH release. *Neuroendocrinology* 1992;56:133–140. [PubMed: 1357569]
123. Jarry H, Leonhardt S, Schwarze T, Wuttke W. Preoptic rather than mediobasal hypothalamic amino acid neurotransmitter release regulates GnRH secretion during the estrogen-induced LH surge in the ovariectomized rat. *Neuroendocrinology* 1995;62:479–486. [PubMed: 8559279]
124. Jarry H, Leonhardt S, Wuttke W. Gamma-aminobutyric acid neurons in the preoptic/anterior hypothalamic area synchronize the phasic activity of the gonadotropin-releasing hormone pulse generator in ovariectomized rats. *Neuroendocrinology* 1991;53:261–267. [PubMed: 2041587]
125. Jarry H, Leonhardt S, Wuttke W. Laterality of the gonadotrophin releasing hormone pulse generator in the rat: interaction of neurons located in both preoptic areas as a basic feature. *Hum Reprod* 1993;8:7–12. [PubMed: 8276973]
126. Jarry H, Perschl A, Wuttke W. Further evidence that preoptic anterior hypothalamic GABAergic neurons are part of the GnRH pulse and surge generator. *Acta Endocrinol (Copenh)* 1988;118:573–579. [PubMed: 3041729]
127. Jennes L, Lin W, Lakhani S. Glutamatergic regulation of gonadotropin-releasing hormone neurons. *Prog Brain Res* 2002;141:183–192. [PubMed: 12508570]
128. Jung H, Shannon EM, Fritschy JM, Ojeda SR. Several GABAA receptor subunits are expressed in LHRH neurons of juvenile female rats. *Brain Res* 1998;780:218–229. [PubMed: 9507141]
129. Kang SH, Seong JY, Cho S, Cho H, Kim K. Acute increase of GABAergic neurotransmission exerts a stimulatory effect on GnRH gene expression in the preoptic/anterior hypothalamic area of ovariectomized, estrogen- and progesterone-treated adult female rats. *Neuroendocrinology* 1995;61:486–492. [PubMed: 7617125]
130. Kawakami S, Ichikawa M, Murahashi K, Hirunagi K, Tsukamura H, Maeda K. Excitatory amino acids act on the median eminence nerve terminals to induce gonadotropin-releasing hormone release in female rats. *Gen Comp Endocrinol* 1998a;112:372–382. [PubMed: 9843643]

131. Kawakami SI, Hirunagi K, Ichikawa M, Tsukamura H, Maeda KI. Evidence for terminal regulation of GnRH release by excitatory amino acids in the median eminence in female rats: A dual immunoelectron microscopic study. *Endocrinology* 1998b;139:1458–1461. [PubMed: 9492084]
132. Kimura F, Jinnai K. Bicuculline infusions advance the timing of luteinizing hormone surge in proestrous rats: comparisons with naloxone effects. *Horm Behav* 1994;28:424–430. [PubMed: 7729810]
133. Lamberts R, Vijayan E, Graf M, Mansky T, Wuttke W. Involvement of preoptic-anterior hypothalamic GABA neurons in the regulation of pituitary LH and prolactin release. *Exp Brain Res* 1983;52:356–362. [PubMed: 6653697]
134. Laube B, Kuhse J, Betz H. Evidence for a tetrameric structure of recombinant NMDA receptors. *J Neurosci* 1998;18:2954–2961. [PubMed: 9526012]
135. Laurie DJ, Wisden W, Seeburg PH. The distribution of thirteen GABAA receptor subunit mRNAs in the rat brain. III. Embryonic and postnatal development. *J Neurosci* 1992;12:4151–4172. [PubMed: 1331359]
136. Leonhardt S, Boning B, Luft H, Wuttke W, Jarry H. Activation of gene expression of the gamma-aminobutyric acid rather than the glutamatergic system in the preoptic area during the preovulatory gonadotropin surge of the rat. *Neuroendocrinology* 2000;71:8–15. [PubMed: 10644894]
137. Leonhardt S, Seong JY, Kim K, Thorun Y, Wuttke W, Jarry H. Activation of central GABAA-but not of GABAB-receptors rapidly reduces pituitary LH release and GnRH gene expression in the preoptic/anterior hypothalamic area of ovariectomized rats. *Neuroendocrinology* 1995;61:655–662. [PubMed: 7544879]
138. Leranath C, MacLusky NJ, Sakamoto H, Shanabrough M, Naftolin F. Glutamic acid decarboxylase-containing axons synapse on LHRH neurons in the rat medial preoptic area. *Neuroendocrinology* 1985;40:536–539. [PubMed: 3892354]
139. Leupen SM, Tobet SA, Crowley WF Jr. Kaila K. Heterogeneous expression of the potassium-chloride cotransporter KCC2 in gonadotropin-releasing hormone neurons of the adult mouse. *Endocrinology* 2003;144:3031–3036. [PubMed: 12810559]
140. Liposits Z, Merchenthaler I, Wetsel WC, Reid JJ, Mellon PL, Weiner RI, Negro-Vilar A. Morphological characterization of immortalized hypothalamic neurons synthesizing luteinizing hormone-releasing hormone. *Endocrinology* 1991;129:1575–1583. [PubMed: 1874189]
141. Lomeli H, Sprengel R, Laurie DJ, Kohr G, Herb A, Seeburg PH, Wisden W. The rat delta-1 and delta-2 subunits extend the excitatory amino acid receptor family. *FEBS Lett* 1993;315:318–322. [PubMed: 8422924]
142. Lopez FJ, Donoso AO, Negro-Vilar A. Endogenous excitatory amino acid neurotransmission regulates the estradiol-induced LH surge in ovariectomized rats. *Endocrinology* 1990;126:1771–1773. [PubMed: 2155110]
143. Loscher W, Wahnschaffe U, Rundfeldt C, Honack D, Hoppen HO. Regional alterations in brain amino acids during the estrous cycle of the rat. *Neurochem Res* 1992;17:973–977. [PubMed: 1508307]
144. Loucopoulos A, Ferin M, Vande Wiele RL, Dyrenfurth I, Linkie D, Yeh M, Jewelewicz R. Pulsatile administration of gonadotropin-releasing hormone for induction of ovulation. *Am J Obstet Gynecol* 1984;148:895–900. [PubMed: 6424475]
145. Luderer U, Strobl FJ, Levine JE, Schwartz NB. Differential gonadotropin responses to N-methyl-D,L-aspartate in metestrous, proestrous, and ovariectomized rats. *Biol Reprod* 1993;48:857–866. [PubMed: 8485252]
146. Luine VN, Grattan DR, Selmánoff M. Gonadal hormones alter hypothalamic GABA and glutamate levels. *Brain Res* 1997;747:165–168. [PubMed: 9042543]
147. MacDonald MC, Wilkinson M. Peripubertal treatment with N-methyl-D-aspartic acid or neonatally with monosodium glutamate accelerates sexual maturation in female rats, an effect reversed by MK-801. *Neuroendocrinology* 1990;52:143–149. [PubMed: 1980346]
148. MacDonald MC, Wilkinson M. Characterization and ontogenesis of N-Methyl-D-aspartate-evoked luteinizing hormone secretion in immature female rats. *Journal of Neuroendocrinology* 1992;4:223–229.

149. Maffucci, JA.; Gore, AC. Age-related changes in hormones and their receptors in animal models of female reproductive senescence. In: Conn, PM., editor. "Handbook of Models for the Study of Human Aging.". Academic Press, Inc.; 2005.
150. Maffucci JA, Walker DM, Ikegami A, Woller MJ, Gore AC. The NMDA receptor subunit NR2b affects LH release and GnRH gene expression in female rats, with modulation by estradiol. *Neuroendocrinology* 2008;87:129–141. [PubMed: 18025808]
151. Mahachoklertwattana P, Sanchez J, Kaplan SL, Grumbach MM. N-methyl-D-aspartate (NMDA) receptors mediate the release of gonadotropin-releasing hormone (GnRH) by NMDA in a hypothalamic GnRH neuronal cell line (GT1-1). *Endocrinology* 1994;134:1023–1030. [PubMed: 8119138]
152. Mahesh VB, Zamorano P, De Sevilla L, Lewis D, Brann DW. Characterization of ionotropic glutamate receptors in rat hypothalamus, pituitary and immortalized gonadotropin-releasing hormone (GnRH) neurons (GT1-7 cells). *Neuroendocrinology* 1999;69:397–407. [PubMed: 10364691]
153. Malyala A, Kelly MJ, Ronnekleiv OK. Estrogen modulation of hypothalamic neurons: activation of multiple signaling pathways and gene expression changes. *Steroids* 2005;70:397–406. [PubMed: 15862823]
154. Mano I, Teichberg VI. A tetrameric subunit stoichiometry for a glutamate receptor-channel complex. *Neuroreport* 1998;9:327–331. [PubMed: 9507977]
155. Manksy T, Mestres-Ventura P, Wuttke W. Involvement of GABA in the feedback action of estradiol on gonadotropin and prolactin release: hypothalamic GABA and catecholamine turnover rates. *Brain Res* 1982;231:353–364. [PubMed: 7198927]
156. Maragos WF, Penney JB, Young AB. Anatomic correlation of NMDA and 3H-TCP-labeled receptors in rat brain. *J Neurosci* 1988;8:493–501. [PubMed: 2828564]
157. Martinez de la Escalera G, Choi AL, Weiner RI. Biphasic gabaergic regulation of GnRH secretion in GT1 cell lines. *Neuroendocrinology* 1994;59:420–425. [PubMed: 8022517]
158. Marty A, Llano I. Excitatory effects of GABA in established brain networks. *Trends Neurosci* 2005;28:284–289. [PubMed: 15927683]
159. McCann SM, Vijayan E, Negro-Vilar A, Mizunuma H, Mangat H. Gamma aminobutyric acid (GABA), a modulator of anterior pituitary hormone secretion by hypothalamic and pituitary action. *Psychoneuroendocrinology* 1984;9:97–106. [PubMed: 6089244]
160. McCarthy MM. Estradiol and the developing brain. *Physiol Rev* 2008;88:91–124. [PubMed: 18195084]
161. McCarthy MM, Auger AP, Perrot-Sinal TS. Getting excited about GABA and sex differences in the brain. *Trends Neurosci* 2002;25:307–312. [PubMed: 12086749]
162. McCarthy MM, Davis AM, Mong JA. Excitatory neurotransmission and sexual differentiation of the brain. *Brain Res Bull* 1997;44:487–495. [PubMed: 9370215]
163. Meeker RB, Greenwood RS, Hayward JN. Glutamate receptors in the rat hypothalamus and pituitary. *Endocrinology* 1994;134:621–629. [PubMed: 7905409]
164. Meijs-Roelofs HM, Kramer P, van Leeuwen EC. The N-methyl-D-aspartate receptor antagonist MK-801 delays the onset of puberty and may acutely block the first spontaneous LH surge and ovulation in the rat. *J Endocrinol* 1991;131:435–441. [PubMed: 1838383]
165. Mellon PL, Windle JJ, Weiner RI. Immortalization of neuroendocrine cells by targeted oncogenesis. *Recent Prog Horm Res* 1991;47:69–93. discussion 93–96. [PubMed: 1745826]
166. Miller BH, Gore AC. N-Methyl-D-aspartate receptor subunit expression in GnRH neurons changes during reproductive senescence in the female rat. *Endocrinology* 2002;143:3568–3574. [PubMed: 12193572]
167. Miller MM, Joshi D, Billiar RB, Nelson JF. Loss of LH-RH neurons in the rostral forebrains of old female C57bl/6J mice. *Neurobiology of Aging* 1990;11:217–221. [PubMed: 2194136]
168. Mitsushima D, Shwe TT, Funabashi T, Shinohara K, Kimura F. GABA release in the medial preoptic area of cyclic female rats. *Neuroscience* 2002;113:109–114. [PubMed: 12123689]
169. Moenter SM, DeFazio RA. Endogenous gamma-aminobutyric acid can excite gonadotropin-releasing hormone neurons. *Endocrinology* 2005;146:5374–5379. [PubMed: 16123153]

170. Moguilevsky JA, Carbone S, Szwarcfarb B, Rondina D. Sexual maturation modifies the GABAergic control of gonadotrophin secretion in female rats. *Brain Res* 1991;563:12–16. [PubMed: 1664770]
171. Moguilevsky JA, Carbone S, Szwarcfarb B, Rondina D, Scacchi P. Hypothalamic excitatory amino acid system during sexual maturation in female rats. *J Steroid Biochem Mol Biol* 1995;53:337–341. [PubMed: 7626477]
172. Moguilevsky JA, Wuttke W. Changes in the control of gonadotrophin secretion by neurotransmitters during sexual development in rats. *Exp Clin Endocrinol Diabetes* 2001;109:188–195. [PubMed: 11453030]
173. Moragues N, Ciofi P, Lafon P, Tramu G, Garret M. GABAA receptor epsilon subunit expression in identified peptidergic neurons of the rat hypothalamus. *Brain Res* 2003;967:285–289. [PubMed: 12650990]
174. Moran J, Pasantes-Morales H, Redburn DA. Glutamate receptor agonists release [3H]GABA preferentially from horizontal cells. *Brain Res* 1986;398:276–287. [PubMed: 2879608]
175. Morello H, Caligaris L, Haymal B, Taleisnik S. Inhibition of proestrous LH surge and ovulation in rats evoked by stimulation of the medial raphe nucleus involves a GABA-mediated mechanism. *Neuroendocrinology* 1989;50:81–87. [PubMed: 2666873]
176. Morello H, Caligaris L, Haymal B, Taleisnik S. Daily variations in the sensitivity of proestrous LH surge in the inhibitory effect of intraventricular injection of 5-HT or GABA in rats. *Can J Physiol Pharmacol* 1992;70:447–451. [PubMed: 1498714]
177. Neal-Perry GS, Zeevalk GD, Santoro NF, Etgen AM. Attenuation of preoptic area glutamate release correlates with reduced luteinizing hormone secretion in middle-aged female rats. *Endocrinology* 2005;146:4331–4339. [PubMed: 15994345]
178. Neal-Perry GS, Zeevalk GD, Shu J, Etgen AM. Restoration of the luteinizing hormone surge in middle-aged female rats by altering the balance of GABA and glutamate transmission in the medial preoptic area. *Biol Reprod*. 2008 ePub ahead of print (DOI:10.1095/biolreprod.108.069831).
179. Ondo J, Mansky T, Wuttke W. In vivo GABA release from the medial preoptic area of diestrous and ovariectomized rats. *Exp Brain Res* 1982;46:69–72. [PubMed: 7200039]
180. Ottem EN, Godwin JG, Krishnan S, Petersen SL. Dual-phenotype GABA/glutamate neurons in adult preoptic area: sexual dimorphism and function. *J Neurosci* 2004;24:8097–8105. [PubMed: 15371511]
181. Ottem EN, Godwin JG, Petersen SL. Glutamatergic signaling through the N-methyl-D-aspartate receptor directly activates medial subpopulations of luteinizing hormone-releasing hormone (LHRH) neurons, but does not appear to mediate the effects of estradiol on LHRH gene expression. *Endocrinology* 2002;143:4837–4845. [PubMed: 12446611]
182. Owens DF, Kriegstein AR. Is there more to GABA than synaptic inhibition? *Nat Rev Neurosci* 2002;3:715–727. [PubMed: 12209120]
183. Parducz A, Perez J, Garcia-Segura LM. Estradiol induces plasticity of gabaergic synapses in the hypothalamus. *Neuroscience* 1993;53:395–401. [PubMed: 8492911]
184. Pin JP, Bockaert J. Two distinct mechanisms, differentially affected by excitatory amino acids, trigger GABA release from fetal mouse striatal neurons in primary culture. *J Neurosci* 1989;9:648–656. [PubMed: 2918382]
185. Ping L, Mahesh VB, Bhat GK, Brann DW. Regulation of gonadotropin-releasing hormone and luteinizing hormone secretion by AMPA receptors. Evidence for a physiological role of AMPA receptors in the steroid-induced luteinizing hormone surge. *Neuroendocrinology* 1997;66:246–253. [PubMed: 9349658]
186. Ping L, Mahesh VB, Brann DW. A physiological role for N-methyl-D-aspartic acid and non-N-methyl-D-aspartic acid receptors in pulsatile gonadotropin secretion in the adult female rat. *Endocrinology* 1994a;135:113–118. [PubMed: 7912182]
187. Ping L, Mahesh VB, Brann DW. Effect of NMDA and non-NMDA receptor antagonists on pulsatile luteinizing hormone secretion in the adult male rat. *Neuroendocrinology* 1995;61:226–234. [PubMed: 7898627]
188. Ping L, Mahesh VB, Wiedmeier VT, Brann DW. Release of glutamate and aspartate from the preoptic area during the progesterone-induced LH surge: in vivo microdialysis studies. *Neuroendocrinology* 1994b;59:318–324. [PubMed: 7911230]

189. Pinilla L, Gonzalez LC, Tena-Sempere M, Aguilar E. Interactions between GABAergic and aminoacidergic pathways in the control of gonadotropin and GH secretion in pre-pubertal female rats. *J Endocrinol Invest* 2002;25:96–100. [PubMed: 11929096]
190. Pinilla L, Tena-Sempere M, Aguilar E. Sexual differences in the role of kainate receptors in controlling gonadotrophin secretion in prepubertal rats. *J Reprod Fertil* 1998;113:269–273. [PubMed: 9861167]
191. Pohl CR, Lee LR, Smith MS. Qualitative changes in luteinizing hormone and prolactin responses to N-methyl-aspartic acid during lactation in the rat. *Endocrinology* 1989;124:1905–1911. [PubMed: 2647467]
192. Rabow LE, Russek SJ, Farb DH. From ion currents to genomic analysis: recent advances in GABAA receptor research. *Synapse* 1995;21:189–274. [PubMed: 8578436]
193. Reiter EO, Grumbach MM. Neuroendocrine control mechanisms and the onset of puberty. *Annu Rev Physiol* 1982;44:595–613. [PubMed: 6803663]
194. Richter TA, Terasawa E. Neural mechanisms underlying the pubertal increase in LHRH release in the rhesus monkey. *Trends Endocrinol Metab* 2001;12:353–359. [PubMed: 11551809]
195. Roberts CB, Hemond P, Suter KJ. Synaptic integration in hypothalamic gonadotropin releasing hormone (GnRH) neurons. *Neuroscience* 2008;154:1337–1351. [PubMed: 18556136]
196. Rohrbough J, Spitzer NC. Regulation of intracellular Cl⁻ levels by Na⁺-dependent Cl⁻ cotransport distinguishes depolarizing from hyperpolarizing GABAA receptor-mediated responses in spinal neurons. *J Neurosci* 1996;16:82–91. [PubMed: 8613812]
197. Romero MT, Silverman AJ, Wise PM, Witkin JW. Ultrastructural changes in gonadotropin-releasing hormone neurons as a function of age and ovariectomy in rats. *Neuroscience* 1994;58:217–225. [PubMed: 8159295]
198. Rosenmund C, Stern-Bach Y, Stevens CF. The tetrameric structure of a glutamate receptor channel. *Science* 1998;280:1596–1599. [PubMed: 9616121]
199. Roth C, Jung H, Kim K, Arias P, Moguilevsky J, Jarry H, Leonhardt S, Wuttke W. Involvement of gamma amino butyric acid (GABA) in the postnatal function of the GnRH pulse generator as determined on the basis of GnRH and GnRH-receptor gene expression in the hypothalamus and the pituitary. *Exp Clin Endocrinol Diabetes* 1997;105:353–358. [PubMed: 9439932]
200. Rubin BS, King JC. The number and distribution of detectable luteinizing hormone (LH)-releasing hormone cell bodies changes in association with the preovulatory LH surge in the brains of young but not middle-aged female rats. *Endocrinology* 1994;134:467–474. [PubMed: 8275960]
201. Rubin BS, King JC, Bridges RS. Immunoreactive forms of luteinizing hormone-releasing hormone in the brains of aging rats exhibiting persistent vaginal estrus. *Biol Reprod* 1984;31:343–351. [PubMed: 6383488]
202. Rubin BS, Lee CE, Ohtomo M, King JC. Luteinizing hormone-releasing hormone gene expression differs in young and middle-aged females on the day of a steroid-induced LH surge. *Brain Res* 1997;770:267–276. [PubMed: 9372229]
203. Scacchi P, Carbone S, Szwarcfarb B, Rondina D, Wuttke W, Moguilevsky JA. Interactions between GABAergic and serotonergic systems with excitatory amino acid neurotransmission in the hypothalamic control of gonadotropin secretion in prepubertal female rats. *Brain Res Dev Brain Res* 1998;105:51–58.
204. Scarbrough K, Wise PM. Age-related changes in pulsatile luteinizing hormone release precede the transition to estrous acyclicity and depend upon estrous cycle history. *Endocrinology* 1990;126:884–890. [PubMed: 2404750]
205. Seltzer AM, Donoso AO. Restraining action of GABA on estradiol-induced LH surge in the rat: GABA activity in brain nuclei and effects of GABA mimetics in the medial preoptic nucleus. *Neuroendocrinology* 1992;55:28–34. [PubMed: 1319004]
206. Seong JY, Lee YK, Lee CC, Kim K. NMDA receptor antagonist decreases the progesterone-induced increase in GnRH gene expression in the rat hypothalamus. *Neuroendocrinology* 1993;58:234–239. [PubMed: 7903433]
207. Shivers BD, Harlan RE, Morell JJ, Pfaff DW. Absence of oestradiol concentration in cell nuclei of LHRH-immunoreactive neurones. *Nature* 1983;304:345–347. [PubMed: 6348552]

208. Sim JA, Skynner MJ, Pape JR, Herbison AE. Late postnatal reorganization of GABA(A) receptor signalling in native GnRH neurons. *Eur J Neurosci* 2000;12:3497–3504. [PubMed: 11029619]
209. Sisk CL, Foster DL. The neural basis of puberty and adolescence. *Nat Neurosci* 2004;7:1040–1047. [PubMed: 15452575]
210. Smyth C, Wilkinson M. A critical period for glutamate receptor-mediated induction of precocious puberty in female rats. *J Neuroendocrinol* 1994;6:275–284. [PubMed: 7920593]
211. Spergel DJ, Krsmanovic LZ, Stojilkovic SS, Catt KJ. Glutamate modulates $[Ca^{2+}]_i$ and gonadotropin-releasing hormone secretion in immortalized hypothalamic GT1-7 neurons. *Neuroendocrinology* 1994;59:309–317. [PubMed: 7911229]
212. Spergel DJ, Krsmanovic LZ, Stojilkovic SS, Catt KJ. L-type Ca^{2+} channels mediate joint modulation by gamma-amino-butyric acid and glutamate of $[Ca^{2+}]_i$ and neuropeptide secretion in immortalized gonadotropin-releasing hormone neurons. *Neuroendocrinology* 1995;61:499–508. [PubMed: 7617127]
213. Spergel DJ, Kruth U, Hanley DF, Sprengel R, Seeburg PH. GABA- and glutamate-activated channels in green fluorescent protein-tagged gonadotropin-releasing hormone neurons in transgenic mice. *J Neurosci* 1999;19:2037–2050. [PubMed: 10066257]
214. Sullivan SD, Moenter SM. Neurosteroids alter gamma-aminobutyric acid postsynaptic currents in gonadotropin-releasing hormone neurons: a possible mechanism for direct steroidal control. *Endocrinology* 2003;144:4366–4375. [PubMed: 12960018]
215. Sullivan SD, Moenter SM. Gamma-aminobutyric acid neurons integrate and rapidly transmit permissive and inhibitory metabolic cues to gonadotropin-releasing hormone neurons. *Endocrinology* 2004;145:1194–1202. [PubMed: 14645118]
216. Sullivan SD, Moenter SM. GABAergic integration of progesterone and androgen feedback to gonadotropin-releasing hormone neurons. *Biol Reprod* 2005;72:33–41. [PubMed: 15342358]
217. Suter KJ, Wuarin JP, Smith BN, Dudek FE, Moenter SM. Whole-cell recordings from preoptic/hypothalamic slices reveal burst firing in gonadotropin-releasing hormone neurons identified with green fluorescent protein in transgenic mice. *Endocrinology* 2000;141:3731–3736. [PubMed: 11014229]
218. Suzuki M, Nishihara M, Takahashi M. Hypothalamic gonadotropin-releasing hormone gene expression during rat estrous cycle. *Endocr J* 1995;42:789–796. [PubMed: 8822321]
219. Szwarcfarb B, Carbone S, Stein ML, Medina J, Moguilevsky JA. Sexual differences in the effect of the GABAergic system on LH secretion and in the hypothalamic ontogenesis of GABA receptors in prepubertal rats. *Brain Res* 1994;646:351–355. [PubMed: 8069689]
220. Taleisnik S, Haymal B. Dual effect of electrochemical stimulation of the medial preoptic area on the release of LH: possible neurotransmitter involvement. *Neuroendocrinology* 1997;66:114–121. [PubMed: 9263208]
221. Temple JL, Wray S. Developmental changes in GABA receptor subunit composition within the gonadotrophin-releasing hormone-1 neuronal system. *J Neuroendocrinol* 2005;17:591–599. [PubMed: 16101898]
222. Terasawa E. Control of luteinizing hormone-releasing hormone pulse generation in nonhuman primates. *Cell Mol Neurobiol* 1995;15:141–164. [PubMed: 7648606]
223. Terasawa E. Role of GABA in the mechanism of the onset of puberty in non-human primates. *Int Rev Neurobiol* 2005;71:113–129. [PubMed: 16512348]
224. Terasawa E, Wiegand SJ, Bridson WE. A role for medial preoptic nucleus on afternoon of proestrus in female rats. *Am J Physiol* 1980;238:E533–E539. [PubMed: 7189967]
225. Tin Tin Win S, Mitsushima D, Shinohara K, Kimura F. Sexual dimorphism of GABA release in the medial preoptic area and luteinizing hormone release in gonadectomized estrogen-primed rats. *Neuroscience* 2004;127:243–250. [PubMed: 15219686]
226. Todman MG, Han SK, Herbison AE. Profiling neurotransmitter receptor expression in mouse gonadotropin-releasing hormone neurons using green fluorescent protein-promoter transgenics and microarrays. *Neuroscience* 2005;132:703–712. [PubMed: 15837132]
227. Unnerstall JR, Wamsley JK. Autoradiographic localization of high-affinity $[^3H]$ kainic acid binding sites in the rat forebrain. *Eur J Pharmacol* 1983;86:361–371. [PubMed: 6299756]

228. Urbanski HF, Fahy MM, Daschel M, Meshul C. N-methyl-D-aspartate receptor gene expression in the hamster hypothalamus and in immortalized luteinizing hormone-releasing hormone neurones. *J Reprod Fertil* 1994;100:5–9. [PubMed: 8182610]
229. Urbanski HF, Ojeda SR. Activation of luteinizing hormone-releasing hormone release advances the onset of female puberty. *Neuroendocrinology* 1987;46:273–276. [PubMed: 3309706]
230. Urbanski HF, Ojeda SR. A role for N-methyl-D-aspartate (NMDA) receptors in the control of LH secretion and initiation of female puberty. *Endocrinology* 1990;126:1774–1776. [PubMed: 1968384]
231. van den Pol AN. GABA immunoreactivity in hypothalamic neurons and growth cones in early development in vitro before synapse formation. *J Comp Neurol* 1997;383:178–188. [PubMed: 9182847]
232. van den Pol AN. Weighing the role of hypothalamic feeding neurotransmitters. *Neuron* 2003;40:1059–1061. [PubMed: 14687541]
233. van den Pol AN, Gao XB, Patrylo PR, Ghosh PK, Obrietan K. Glutamate inhibits GABA excitatory activity in developing neurons. *J Neurosci* 1998;18:10749–10761. [PubMed: 9852609]
234. van den Pol AN, Obrietan K, Cao V, Trombley PQ. Embryonic hypothalamic expression of functional glutamate receptors. *Neuroscience* 1995;67:419–439. [PubMed: 7545794]
235. Veneroni O, Cocilovo L, Muller EE, Cocchi D. Delay of puberty and impairment of growth in female rats given a non competitive antagonist of NMDA receptors. *Life Sci* 1990;47:1253–1260. [PubMed: 2147050]
236. Vicini S, Wang JF, Li JH, Zhu WJ, Wang YH, Luo JH, Wolfe BB, Grayson DR. Functional and pharmacological differences between recombinant N-methyl-D-aspartate receptors. *J Neurophysiol* 1998;79:555–566. [PubMed: 9463421]
237. Vijayan E, McCann SM. The effects of intraventricular injection of gamma-aminobutyric acid (GABA) on prolactin and gonadotropin release in conscious female rats. *Brain Res* 1978;155:35–43. [PubMed: 688015]
238. Virmani MA, Stojilkovic SS, Catt KJ. Stimulation of luteinizing hormone release by gamma-aminobutyric acid (GABA) agonists: mediation by GABAA-type receptors and activation of chloride and voltage-sensitive calcium channels. *Endocrinology* 1990;126:2499–2505. [PubMed: 2158428]
239. Wagner S, Castel M, Gainer H, Yarom Y. GABA in the mammalian suprachiasmatic nucleus and its role in diurnal rhythmicity. *Nature* 1997;387:598–603. [PubMed: 9177347]
240. Whiting PJ, McAllister G, Vassilatis D, Bonnert TP, Heavens RP, Smith DW, Hewson L, O'Donnell R, Rigby MR, Sirinathsinghji DJ, Marshall G, Thompson SA, Wafford KA, Vasilatis D. Neuronally restricted RNA splicing regulates the expression of a novel GABAA receptor subunit conferring atypical functional properties [corrected; erratum to be published]. *J Neurosci* 1997;17:5027–5037. [PubMed: 9185540]
241. Wiegand SJ, Terasawa E, Bridson WE, Goy RW. Effects of discrete lesions of preoptic and suprachiasmatic structures in the female rat. Alterations in the feedback regulation of gonadotropin secretion. *Neuroendocrinology* 1980;31:147–157. [PubMed: 6771669]
242. Wildt L, Hausler A, Marshall G, Hutchison JS, Plant TM, Belchetz PE, Knobil E. Frequency and amplitude of gonadotropin-releasing hormone stimulation and gonadotropin secretion in the rhesus monkey. *Endocrinology* 1981;109:376–385. [PubMed: 6788538]
243. Wintermantel TM, Campbell RE, Porteous R, Bock D, Grone HJ, Todman MG, Korach KS, Greiner E, Perez CA, Schutz G, Herbison AE. Definition of estrogen receptor pathway critical for estrogen positive feedback to gonadotropin-releasing hormone neurons and fertility. *Neuron* 2006;52:271–280. [PubMed: 17046690]
244. Wisden W, Seeburg PH. GABAA receptor channels: from subunits to functional entities. *Curr Opin Neurobiol* 1992;2:263–269. [PubMed: 1379501]
245. Wise PM. Estradiol-induced daily luteinizing hormone and prolactin surges in young and middle-aged rats: correlations with age-related changes in pituitary responsiveness and catecholamine turnover rates in microdissected brain areas. *Endocrinology* 1984;115:801–809. [PubMed: 6378599]

246. Witkin JW. Luteinizing hormone releasing hormone (LHRH) neurons in aging female rhesus macaques. *Neurobiol Aging* 1986;7:259–263. [PubMed: 3528888]
247. Woller MJ, Everson-Binotto G, Nichols E, Acheson A, Keen KL, Bowers CY, Terasawa E. Aging-related changes in release of growth hormone and luteinizing hormone in female rhesus monkeys. *J Clin Endocrinol Metab* 2002;87:5160–5167. [PubMed: 12414887]
248. Yamazaki M, Araki K, Shibata A, Mishina M. Molecular cloning of a cDNA encoding a novel member of the mouse glutamate receptor channel family. *Biochem Biophys Res Commun* 1992;183:886–892. [PubMed: 1372507]
249. Yin C, Ishii H, Tanaka N, Sakuma Y, Kato M. Activation of A-type gamma-amino butyric acid receptors excites gonadotrophin-releasing hormone neurones isolated from adult rats. *J Neuroendocrinol* 2008;20:566–575. [PubMed: 18363808]
250. Yin W, Gore AC. Neuroendocrine control of reproductive aging: roles of GnRH neurons. *Reproduction* 2006;131:403–414. [PubMed: 16514184]
251. Yin W, Mendenhall J, Bratton SB, Oung T, Janssen WGM, Morrison JH, Gore AC. Novel localization of NMDA receptors within neuroendocrine gonadotropin-releasing hormone terminals. *Exp Biol Med* 2007;232:662–673.
252. Zhang JH, Sato M, Araki T, Tohyama M. Postnatal ontogenesis of neurons containing GABAA alpha 1 subunit mRNA in the rat forebrain. *Brain Res Mol Brain Res* 1992;16:193–203. [PubMed: 1337929]
253. Zhang JH, Sato M, Tohyama M. Different postnatal development profiles of neurons containing distinct GABAA receptor beta subunit mRNAs (beta 1, beta 2, and beta 3) in the rat forebrain. *J Comp Neurol* 1991;308:586–613. [PubMed: 1650800]
254. Zuo Z, Mahesh VB, Zamorano PL, Brann DW. Decreased gonadotropin-releasing hormone neurosecretory response to glutamate agonists in middle-aged female rats on proestrus afternoon: a possible role in reproductive aging? *Endocrinology* 1996;137:2334–2338. [PubMed: 8641183]

Figure 1.

A) The anteroventral peroventricular nucleus is a region important to the integration of signals for proper female reproductive function. This region is located lateral to the third ventricle (iiiV) and rostral to the optic tract (OT), and is rich in cells immunopositive for the NMDAR subunits, including NR2b (shown here labeled by DAB-brown stain). A solid, black line highlights the border of the AVPV. The abundance of NR2b-immunoreactive cells can be seen in this representative middle-aged OVX rat treated with estradiol. **B)** NR2b-immunopositive cells from another representative OVX rat (in this case, young vehicle treated) is shown at higher magnification (DAB-brown stain labels the NR2b cells). The Nissl counterstain is seen in blue. **C.** Immunofluorescent labeling and confocal analysis of NR2b (green) and NR1 (red) in the AVPV of a representative rat (young, OVX plus estradiol plus progesterone) demonstrates the presence of colocalized (yellow) cells. The image size is 203 $\mu\text{m} \times 203 \mu\text{m}$.

Figure 2.

Simplified model for interactions among GnRH, GABA and glutamate neurons in the hypothalamus. For each of the prepubertal, adult, and aged life stages, a GnRH neuron is shown in yellow (terminals are shown containing large dense-core secretory vesicles of GnRH, in green); GABA neurons are colored tan, and glutamate neurons are in green. At the prepubertal stage, GnRH neurons are depicted in relatively greater proximity to GABA neurons, with relatively little glutamatergic inputs. In adulthood, the balance between glutamate and GABA onto GnRH cells has shifted to a predominantly glutamatergic stimulatory influence. At this age, GABA likely provides a mix of inhibitory and excitatory balance onto GnRH neurons, depending upon whether effects are direct or indirect. With senescence, this balance shifts in favor of inhibitory GABAergic inputs and a diminution of glutamatergic input to GnRH. Notably, at all ages, there are both direct effects of glutamate and GABA on GnRH cells as well as influences of glutamate and GABA cells on each other. Finally, while not shown, some cells may be both glutamatergic and GABAergic. Below each age is a brief summary of the relative effects of glutamate and GABA inputs on GnRH functions. Abbreviations: agon, agonist; antag, antagonist.

Table 1

Ionotropic glutamate receptor classes, and corresponding pharmacological agents

Ligand**Glutamate:** endogenous ligand, binds to all glutamate receptors

Receptor type	Receptor subunits	Agonist	Antagonist
NMDA receptor	NR1 NR2a, 2b, 2c, 2d NR3a, 3b	NMDA (N-methyl-D-aspartate), NMA (N-methyl-D,L-aspartate): specific agonists of NMDA receptors	MK-801: non-competitive NMDA receptor antagonist AP-5 or AP-7: competitive NMDA receptor antagonist Ifenprodil: Selective NR2b subunit antagonist
AMPA receptor	GluR1-4	AMPA (DL-α-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid): specific agonist of AMPA receptors	NBQX: competitive antagonist to AMPA receptors DNQX: competitive antagonist to AMPA and kainate receptors
Kainate receptor	GluR5-7 KA1, KA2	Kainate: specific agonist of Kainate receptors	DNQX: competitive antagonist to AMPA and kainate receptors

Other**Glutaminase:** Enzyme responsible for conversion of glutamine into glutamate**Vesicular glutamate transporter (VGLut):** Transporter of glutamate, the presence of which is thought to indicate a glutamatergic phenotype

Table 2

Effects of NMDA receptor agonists/antagonists on GnRH gene expression

Technique	Age, hormonal Status	Steroid hormone treatment	Drug Administration	Effect on GnRH gene expression
RNase protection assay	Intact female rat pups, postnatal days 0, 5, 10, 15	N/A	NMA (agonist) or MK-801 (antagonist)	Little effects of NMA or MK-801 on GnRH mRNA or primary transcript ^a
RNase protection assay	Intact pubertal rats, beginning day 25 through the day of VO	N/A	NMA twice daily	↑ GnRH mRNA ^b
Northern blot hybridization	Peripubertal (P28) OVX rats	Estradiol/progesterone	MK-801	↓ in GnRH mRNA in rostral hypothalamus ^c
RNase protection assay	a. Young intact (proestrus) b. Middle-aged intact	N/A	NMA	a. ↑ GnRH mRNA in POA b. ↓ GnRH mRNA in POA ^d
Quantitative RT-PCR	Proestrous female rats	N/A	MK-801	Attenuation of the increase in GnRH mRNA during the GnRH/LH surge ^e
<i>In situ</i> hybridization	OVX adult rats	Estradiol	a. NMDA b. MK-801	a. ↑ GnRH mRNA b. ↓ GnRH mRNA ^f
Real-time PCR	OVX young and middle-aged rats	Estradiol	Ifenprodil (NR2b antagonist)	Little effect at either age ^g

Abbreviations: NMA, N-methyl-D,L-aspartate; OVX, ovariectomized; PE, persistent estrus; VO, vaginal opening; ↓ decrease; ↑ increase.

^aFootnotes: Adams, M. M., Flagg, R. A., and Gore, A. C. (1999). Perinatal changes in hypothalamic N-methyl-D-aspartate receptors and their relationship to gonadotropin-releasing hormone neurons. *Endocrinology* **140**, 2288–2296;

^bGore, A. C., Wu, T. J., Rosenberg, J. J., and Roberts, J. L. (1996). Gonadotropin-releasing hormone and NMDA receptor gene expression and colocalization change during puberty in female rats. *J Neurosci* **16**, 5281–5289;

^cSeong, J. Y., Lee, Y. K., Lee, C. C., and Kim, K. (1993). NMDA receptor antagonist decreases the progesterone-induced increase in GnRH gene expression in the rat hypothalamus. *Neuroendocrinology* **58**, 234–239;

^dGore, A. C., Yeung, G., Morrison, J. H., and Oung, T. (2000b). Neuroendocrine aging in the female rat: the changing relationship of hypothalamic gonadotropin-releasing hormone neurons and N-methyl-D-aspartate receptors. *Endocrinology* **141**, 4757–4767;

^eSuzuki, M., Nishihara, M., and Takahashi, M. (1995). Hypothalamic gonadotropin-releasing hormone gene expression during rat estrous cycle. *Endocr J* **42**, 789–796;

^fOttm, E. N., Godwin, J. G., and Petersen, S. L. (2002). Glutamatergic signaling through the N-methyl-D-aspartate receptor directly activates medial subpopulations of luteinizing hormone-releasing hormone (LHRH) neurons, but does not appear to mediate the effects of estradiol on LHRH gene expression. *Endocrinology* **143**, 4837–4845;

^gMaffucci, J. A., Walker, D. M., Ikegami, A., Woller, M. J., and Gore, A. C. (2008). NMDA receptor subunit NR2b: effects on LH release and GnRH gene expression in young and middle-aged female rats, with modulation by estradiol. *Neuroendocrinology* **87**, 129–141.

Table 3

Ionotropic GABA receptor classes, and corresponding pharmacological agents

Ligand**GABA (gamma-aminobutyric acid):** endogenous ligand, binds to all GABA receptors

Receptor type	Receptor subunits	Agonist	Antagonist
GABA _A receptor	$\alpha 1-\alpha 6$ $\beta 1-\beta 3$ $\gamma 1-\gamma 3$ $\delta, \epsilon, \pi, \theta$	Muscimol; 4,5,6,7-Tetrahydroisoxazolo[5,4-c]pyridin-3-ol hydrochloride (THIP): GABA _A receptor agonists	Bicuculline: competitive antagonist to GABA _A receptors
GABA _B receptor	gb1a gb1b	Baclofen: GABA _B receptor agonist	Phaclofen: antagonist to GABA _B receptors

Other**Aminooxyacetic acid (AOAA):** GABA transaminase inhibitor, inhibits GABA breakdown**Glutamic acid decarboxylase (GAD):** Enzyme responsible for conversion of glutamate into GABA**GABA-transaminase (GVG):** Enzyme that inhibits uptake and recycling of GABA**GABA transporter (GAT):** Transport protein, localized to neuronal and glial cells; provides rapid removal of GABA from the synaptic cleft.**Vesicular GABA transporter (VGAT):** Transporter of GABA, the presence of which is thought to indicate a GABAergic Phenotype

GAD₆₇ mRNA expression shows regional specificity of expression in the hypothalamus throughout the GnRH/LH surge

Table 4

Animal Model	Technique	Preoptic Area	DBB/OVLT	MPN	ME/MBH
Adult, intact, proestrus	RNAse protection assay	Microdissections ^{**a}	PM ↓ ^a	No change ^a	No change ^a
Adult, intact, proestrus	<i>In situ</i> hybridization Histochemistry	Microdissections ^{**b}	No change ^b	↓PM ^b	Not examined ^b
Adult, OVX rats given estradiol	<i>In situ</i> hybridization histochemistry	Microdissections ^{**c}	AM ↑; PM ↓ ^c	AM ↑, PM ↓ ^c	Not examined ^c
Adult, intact, Proestrus	Competitive RT-PCR	PM ↑ ^d	Not examined ^{*d}	Not examined ^{*d}	Not examined ^d

Abbreviations: DBB/OVLT: Diagonal band of Broca, organum vasculosum of the lamina terminalis; ME/MBH: median eminence/medial basal hypothalamus; MPN: medial preoptic nucleus; OVX: ovariectomized; ↓ decrease; ↑ increase.

* Footnotes: Indicates that this study used a macrodissection of the preoptic area, in which this nucleus, as well as other hypothalamic nuclei, were included.

** Indicates that these studies used microdissections, examining single nuclei located in the preoptic area.

^a Grattan, D. R., Rocca, M. S., Strauss, K. I., Sagrillo, C. A., Selmanoff, M., and McCarthy, M. M. (1996). GABAergic neuronal activity and mRNA levels for both forms of glutamic acid decarboxylase (GAD65 and GAD67) are reduced in the diagonal band of Broca during the afternoon of proestrus. *Brain Res* **733**, 46–55;

^b Herbison, A. E., Augood, S. J., and McGowan, E. M. (1992). Expression of glutamic acid decarboxylase messenger RNA in rat medial preoptic area neurones during the oestrous cycle and after ovariectomy. *Brain Res Mol Brain Res* **14**, 310–316;

^c Curran-Rauhut, M. A., and Petersen, S. L. (2002). Regulation of glutamic acid decarboxylase 65 and 67 gene expression by ovarian steroids: Identification of two functionally distinct populations of GABA neurones in the preoptic area. *J Neuroendocrinol* **14**, 310–317;

^d Leonhardt, S., Bönig, B., Luft, H., Wuttke, W., and Jarry, H. (2000). Activation of gene expression of the gamma-aminobutyric acid rather than the glutamatergic system in the preoptic area during the preovulatory gonadotropin surge of the rat. *Neuroendocrinology* **71**, 8–15.

Table 5

Effects of GABA agonists/antagonists on GnRH gene expression

Technique	Age, ovarian status	Steroid hormone treatment	Drug administration	Effect on GnRH gene expression in POA
<i>In situ</i> Hybridization	Adult, OVX	N/A	a. Bicuculline (GABA _A antagonist) b. Tetrahydroisoxazolo[5,4- <i>c</i>]pyridin-3-ol hydrochloride c. Baclofen (GABA _B agonist) (all into POA)	a. ↑GnRH mRNA b. No effect c. ↓GnRH mRNA ^a
Competitive RT-PCR	Adult, OVX	N/A	a. Muscimol (GABA _A agonist) b. Baclofen (GABA _B agonist) (both into left lateral ventricle)	a. ↓GnRH mRNA b. No effect ^b
Northern blot	Adult, OVX	Estradiol + Progesterone	Baclofen (GABA _B agonist) (s.c. or into the third ventricle)	↑GnRH mRNA ^c
Northern blot	Adult, OVX	Estradiol + vehicle or progesterone	a. Muscimol (GABA _A agonist) b. Baclofen (GABA _B agonist) (both into third ventricle)	a. ↑GnRH mRNA (low dose only) b. ↑GnRH mRNA (dose-dependent) ^d

Abbreviations: OVX, ovariectomy; THIP, Tetrahydroisoxazolo[5,4-*c*]pyridin-3-olhydrochloride; POA, preoptic area; ↓decrease; ↑increase.

^a Bergen, H. T., Hejtmanick, J. F., and Pfaff, D. W. (1991). Effects of gamma-aminobutyric acid receptor agonists and antagonist on LHRH-synthesizing neurons as detected by immunocytochemistry and in situ hybridization. *Exp Brain Res* **87**, 46–56;

^b Leonhardt, S., Seong, J. Y., Kim, K., Thorun, Y., Wuttke, W., and Jarry, H. (1995). Activation of central GABAA-but not of GABAB-receptors rapidly reduces pituitary LH release and GnRH gene expression in the preoptic/anterior hypothalamic area of ovariectomized rats. *Neuroendocrinology* **61**, 655–662;

^c Cho, B. N., and Kim, K. (1997). Differential effect of baclofen on hypothalamic GnRH and pituitary LH beta gene expression in steroid-treated rats. *Mol Cells* **7**, 605–609;

^d Kang, S. H., Seong, J. Y., Cho, S., Cho, H., and Kim, K. (1995). Acute increase of GABAergic neurotransmission exerts a stimulatory effect on GnRH gene expression in the preoptic/anterior hypothalamic area of ovariectomized, estrogen- and progesterone-treated adult female rats. *Neuroendocrinology* **61**, 486–492.